

## Full Length Research Paper

# RNAi-based silencing of genes encoding the vacuolar-ATPase subunits a and c in pink bollworm (*Pectinophora gossypiella*)

Ahmed M. A. Mohammed

Agricultural Genetic Engineering Research Institute (AGERI), Agricultural Research Center, 9 Gamaa St., 12619 Giza, Egypt.

Received 9 August, 2016; Accepted 28 October, 2016

RNA interference is a post-transcriptional gene regulation mechanism that is predominantly found in eukaryotic organisms. RNAi demonstrated a successful gene silencing in insects that led to the development of novel approaches for insect pest management. In the current study, genes encoding vacuolar ATPase (V-ATPase) subunits a and c from the midgut of pink bollworm, *Pectinophora gossypiella*, were cloned and sequenced. The full length of V-ATPase subunits a and c cDNAs are 2526 and 1140 bp, respectively. The silencing effect of RNAi on these two genes was determined by microinjecting three dsRNA fragments into the thoracic region of pink bollworm larvae. Bioassay results revealed that 200 ng of dsRNAs silenced both genes causing mortality of 18.9 to 26.7%.

**Key words:** RNAi, dsRNA, Pink bollworm, vacuolar ATPase subunit a, vacuolar ATPase subunit c.

## INTRODUCTION

RNA interference (RNAi) is a mechanism of post-transcriptional regulation in higher eukaryotes inhibiting gene expression by RNA transcript degradation (Berezikov, 2011). It was first discovered in the nematode *Caenorhabditis elegans* by Fire et al. (1998) and, later, was exploited as an effective technique for a rapid analysis of gene function by phenotypical changes of the target silenced gene. In this technique, exogenous double stranded RNA (dsRNA) is used to significantly reduce the target gene transcript level (Fire et al., 1998). RNAi technology was harnessed in different applications including insect control, thus, was deployed as a powerful

tool in biological control for insect pest management (Gordon and Waterhouse, 2007; Price and Gatehouse, 2008; Zhu, 2013). A variety of methods are used to deliver dsRNA inside the cells such as soaking the animals in dsRNA-containing solution, microinjection and oral feeding (Baum et al., 2007; Chen et al., 2010; Liu et al., 2010; Rosa et al., 2012; Tian et al., 2009; Zhang et al., 2010; Zhao et al., 2011; Zhu et al., 2011).

An advanced progress of RNAi exploitation in the field of insect control was developed in 2007, by expressing insect specific-dsRNAs in transgenic plants. Transgenic corn expressing V-ATPase specific-dsRNAs showed a

E-mail: amohoammed00@yahoo.com.

Author(s) agree that this article remains permanently open access under the terms of the [Creative Commons Attribution License 4.0 International License](https://creativecommons.org/licenses/by/4.0/)

significant reduction in feeding damage caused by western corn rootworm, *Diabrotica virgifera virgifera* (Baum et al., 2007). Also, cotton was modified to harbor dsRNA targeting the cytochrome P450 gene (CYP6AE14) of the cotton bollworm, *Helicoverpa armigera* (Mao et al., 2007). Cotton bollworm larvae fed on transgenic cotton showed a reduction in growth rate and lower levels of CYP6AE14 transcript within the insect midgut. Thereafter, other RNAi-mediated transgenic plants were developed such as transgenic tobacco plants *Nicotiana tabacum*, that showed higher resistance level compared to non-transgenic against *H. armigera* (Xiong et al., 2013; Jin et al., 2015), *Spodoptera exigua* (Zhu et al., 2012) and *Bemisia tabaci* (Thakur et al., 2014).

The vacuolar proton pumps, V-ATPases, are ubiquitous holoenzyme among eukaryotes (Dow, 1999). It is a member of ATPases family (A, F and V) which are mainly responsible for ATP hydrolysis (Forgac, 2007). The main function of V-ATPases is acidifying a wide array of intracellular organelles and pump protons across the plasma membranes of numerous cell types (Nelson et al., 2000). V-ATPases utilize the energy derived from ATP hydrolysis to transport protons across intracellular and plasma membranes of eukaryotic cells. In the midgut of lepidopteran larvae, the V-ATPase in the apical cell membranes of the goblet cells plays a role in amino acid absorption, by energizing the plasma membrane through pumping H<sup>+</sup> ions/proton into the goblet lumen (Beyenbach and Wieczorek, 2006). V-ATPase is composed of two complexes; a peripheral, catalytic V<sub>1</sub> complex with subunits A<sub>3</sub>B<sub>3</sub>CDE<sub>γ</sub>FG<sub>γ</sub>H and a membrane-bound, proton-conducting V<sub>0</sub> complex with subunits a<sub>c</sub><sub>6</sub>de (Vitavska et al., 2003). The V-ATPase is down regulated during larval moult and starvation periods by the reversible dissociation of the enzyme into its two complexes (Sumner et al., 1995; Gräf et al., 1996) and, in turn, the V<sub>1</sub> complex level increases in the cytoplasm. Thus, biosynthesis of V ATPase subunits is down regulated and transcript levels of these subunits decrease gradually (Wieczorek et al., 2000).

Knock down of genes encoding different subunits belonging to insect V-ATPase have been reported in different insect species. In the current study, the effect of RNAi on V-ATPase genes encoding subunits a and c was determined in *P. gossypiella* by injecting the larval instar with V-ATPase subunits a and c-specific dsRNA fragments.

## MATERIALS AND METHODS

### Insect culture

*P. gossypiella* colony was reared on an artificial diet until pupation, at temperature 25±2°C and photoperiod of light:dark (16:8) hours (Bell and Joachim, 1976). A pair of neonate larvae was added to approximately 5 g diet in 35 ml glass vial covered by cotton plugs and pupae were collected in glass jars covered by filter paper as oviposition site.

### cDNA synthesis and target gene cloning

Total RNA was extracted from *P. gossypiella* midgut tissues using Triazol<sup>®</sup> (Invitrogen) according to manufacturer's instructions. First strand cDNA was prepared from total RNA using the Superscript II cDNA synthesis kit (Invitrogen) according to the manufacturer's instructions. Degenerate primers were designed based on the conserved regions of the V-ATPase subunits a and c sequences in NCBI database using the Vector NTI<sup>®</sup> software (Life technologies). The nucleotide sequences of degenerate, specific and RACE primers are presented in Table 1. One fragment of subunit a was amplified using degenerate primers; VATPavo2264FD / VATPavo2484RD and two more fragments were amplified using degenerate and specific primer sets VATPavo776FD/ VATPavo1559RS and VATPavo1439FD/VATPavo2416RS, respectively. On the other hand, two fragments of subunit c were amplified using degenerate primer set (VATPcvoFD/ VATPcvoRD) and degenerate primer VATPcvo146FD and specific primer VATPcvo901RS. The template cDNA was denatured at 95°C for 5 min followed by 25 cycles of 95°C for 30 s and annealing temperature at 55°C for another 30 s followed by 30 s at 72°C the PCR reaction was extended at 72°C for 7 min. The PCR product was cloned into a pGEM-T Easy vector (Promega, Madison, WI), sequenced using the Big Tri Dye sequencing kit (ABI Applied Biosystems) at Macrogen, Korea.

Both 5' and 3' ends were synthesized using First Choice<sup>®</sup> RLM-RACE kit (Ambion life technologies) following manufacturer's procedures. RACE adaptors and specific primers (Table 1) were used for two rounds of PCR to amplify both ends of subunits a and c with annealing temperatures of 57 and 65°C, respectively. For subunit a; the outer adaptors were mixed with specific primers VATPavo864RS and VATPavo2265 FS to amplify 5' and 3'-end, respectively. The second PCR round was performed using inner adaptors with specific primers VATPavo795 RS and VATPavo2431FS for 5' and 3'-end, respectively. On the other hand, outer and inner adaptors and specific primers VATPcvo244 RS and VATPcvo185 RS were used to amplify the 5'-end of subunit c. The 3'-end was amplified using specific primers VATPcvo819 FS and VATPcvo847 FS.

### Sequence, alignment, and phylogenetic analysis

The obtained sequences of V-ATPase subunits a and c were analyzed using the BLAST algorithm at NCBI for comparative analysis. Multiple alignments of both nucleotide and deduced amino acid sequences were performed by Vector NTI<sup>®</sup> advance 10 (Life Technologies). The deduced amino acid sequences were also scanned for motifs against the PROSITE database. A phylogenetic tree was constructed using the neighbor-joining method in MAGE version 6 (Tamura et al., 2013).

### dsRNA synthesis

The dsRNA fragments were generated using MEGA script<sup>®</sup> RNAi Kit (Ambion) according to manufacturer's instructions. Two dsRNA fragments; a-1 and a-2 (220 and 466 bp) were prepared to target the *P. gossypiella* V-ATPase subunit a transcript using the following primer sets "VATPavo2264F/VATPavo2484R" and "VATPavo1950F/VATPavo2416R". On the other hand, the gene encoding subunit c was knocked down by only one dsRNA fragment (740 bp) that was synthesized by primer set "VATPcvo161F/ VATPcvo901R" (Table 1).

### dsRNA injection

RNAi efficiency of gene silencing in *P. gossypiella* was determined

**Table 1.** Primers used in PCR, RACE and dsRNA synthesis.

Primer	Sequence
<b>Cloning primers</b>	
<b>Subunit a</b>	
VATPavo2264 FD	5'-GTGGGCDYTKRCKTGCBC-3'
VATPavo2484RD	5'-GAAYTTRCTCWKGAACCTCCACCC-3'
VATPavo1439FD	5'-TTCTTYGSYGGDCGTTACAT-3'
VATPavo2416 RS	5'-GACGAGGATGGCGAGCGTGAA-3'
VATPavo776 FD	5'-GTVTTYGTGGYBTTCTTCCA-3'
VATP avo1559 RS	5'-CGG AATGCGCCACGACGAGC-3'
<b>Subunit c</b>	
VATPcvoFD	5'-GCYCCYGGCGAYAAGACMTG-3'
VATPcvoRD	5'-GCCSAGRTCGBASAGSCCCAC-3'
VATPcvo146 FD	5'-TGGMACDGCCAAGTCVGGMACBGGT-3'
VATPcvo901 RS	5'-CATCCACTATAAGACAGAAAGCT-3'
<b>RACE primers</b>	
<b>Subunit a</b>	
5'-end	
VATPavo864 RS	5'-GTTGGAAGGCGGACACGGGTA-3'
VATPavo795 RS	5'-CTGCTCACCTTGAAGAACC-3'
3'-end	
VATPavo2265 FS	5'-CTTTCGCTGGCTCACGCAGAGTTATC-3'
VATPavo2431 FS	5'-TCTCGGCCTTCTCCACACCCT-3'
<b>Subunit c</b>	
5'-end	
VATPcvo244 RS	5'-ATAATACCCGCCATGACAACG-3'
VATPcvo185 RS	5'-CGACATAGCGGCGATACCCGT-3'
3'-end	
VATPcvo819 FS	5'-ACTGTGTACCACCATCTTTGC-3'
VATPcvo847 FS	5'-GGTAAAACCTGGCTTAAGTAC-3'
<b>dsRNA primers</b>	
<b>Subunit a</b>	
a-1	
VATPavo2264F	5'-TAATACGACTCACTATAGGGCTTTCGCTGGCTCACGCAGAGT-3'
VATPavo2484R	5'-TAATACGACTCACTATAGGGCAGGGTGTGGAGGAAGGCCGA-3'
a-2	
VATPavo1950F	5'-TAATACGACTCACTATAGGGGAAGATGGCTGCCACGAATA-3'
VATPavo2416R	5'-TAATACGACTCACTATAGGGGACGAGGATGGCGAGCGTG-3'
<b>Subunit c</b>	
VATPcvo161F	5'-TAATACGACTCACTATAGGGCGGCACGGGTATCGCCGCTAT-3'
VATPcvo901R	5'-TAATACGACTCACTATAGGGCATCCACTATAAGACAGAAAGC-3'

by microinjection of dsRNA into larval hemeolymph. The third larval instars were injected using Neuros Syringe model 1701RN controlled with dispenser (Hamelton). The dsRNA was diluted with injection buffer (0.1mM NaHPO<sub>4</sub> pH 6.8, 5 mM KCl) to final concentration of 1 µg/µl and 0.2 µl were injected into larvae between meso and meta-thoracic segments. The injection was repeated three times for each dsRNA fragment, each replica contained a group of 40 to 45 larvae. Control larvae were injected with injection buffer and treated as the same as experimental individuals. Larvae that died within the first for 24 h were removed and not counted. Larval mortality was recorded five days after

injection. Statistics of data was performed with Student's t-test in the Excel program.

## RESULTS

### Sequence analysis of the V-ATPase subunits a and c

The initial sequences of both transcripts were amplified using degenerate primers designed based on

homologous proteins. A partial sequence (220 bp) of V-ATPase subunit a was obtained from *P. gossypiella* by RT-PCR using degenerate primer set (VATPavo2264FD / VATPavo2484RD), and two longer fragments of 977 and 783 bp were isolated using two sets of combined degenerate and specific primers (VATPavo776FD/ VATPavo1559RS and VATPavo1439FD / VATPavo2416RS) (Supplementary Figure S-1). A 150 bp fragment of V-ATPase subunit c was also isolated based on degenerate primer set (VATPcvoFD/ VATPcvoRD) while a longer 755 bp sequence was amplified by degenerate and specific primers (VATPcvo146FD/VATPcvo901RS). However, the full lengths of the two V-ATPase transcripts were identified by cloning and sequencing the 5' and 3' ends (Supplementary Figure S-2). The RACE strategy demands two PCR reactions, the outer and the inner rounds using RACE adaptors with specific primers. Therefore, two specific primers were designed for each end (Table 1). Specific primers VATPavo864RS and VATPavo795 RS were used to amplify the 5'-end of subunit a, while the 3'-end was synthesized using VATPavo2265 FS and VATPavo2431FS primers. Likewise, the 5'-end of subunit c was identified using specific primers VATPcvo244 RS and VATPcvo185 RS and the 3'-end was amplified using specific primers VATPcvo819 FS and VATPcvo847 FS. The products of PCR and 5' and 3' RACE reactions were aligned to form a contig and finally the full lengths were assembled.

Full length sequence of subunit a (accession no. KU550964) consists of 2760 bp including open reading frame of 2526 bp encoding for 842 amino acids (aa) with 88% identical and 92% similarity to subunit a of *B. mori* and *Amyelois transitella*. The calculated molecular mass of the protein is 96.4 kDa with isoelectric point (pI) of 5.67. On the other hand, subunit c (accession no. KU550965) is 1140 bp containing an ORF of 480 bp and encoding a protein of 160aa with molecular weight of 16.2 kDa and pI of 8.96.

Searching the PROSITE database revealed no common motifs for both subunits sequences. The phylogenetic tree analysis shows that the V-ATPase subunit a of both *P. gossypiella* and *Tribolium castaneum* are located on the same branch as *Bactrocera dorsalis* with 94% bootstrap support (Figure 1a). On the other hand, the *P. gossypiella* V-ATPase subunit c is in close proximity with other lepidopteran species; *H. virescens*, *M. sexta* and *Plutella xylostella* by 50% bootstrap with distal scale length of 0.02 (Figure 1b). The similarity of PgVATPase subunit a and other cognate of other insect orders is presented in Figure 2 whereas subunit c belongs to highly conserved proteolipid protein family known as ductin (Pietrantonio and Gill, 1997). The amino acid sequence of PgVATPase subunit c is highly identical with other lepidopteran insects (Figure 3). The amino acid is 95% identical and 97% similar to that of *Heliothis virescens* and *Manduca sexta*.


## dsRNA injection

Third larval instars were injected with 200 ng dsRNA in between thoracic segments. Larval mortality within 24 h after injecting the dsRNA was neglected to make sure that mortality was not caused by injury from injections. Larval mortality was recorded between 24 and 96 h post injection. Although two dsRNAs fragments, a-1 and a-2, were designed to target the same site on the V-ATPase subunit a transcript, each caused different mortality to the injected larvae. The first fragment (a-1) is 220 bp in length (nucleotides 2264-2484), whereas the length of the second dsRNA (a-2) is 466 bp from 1950 to 2416 (Supplementary Figure S-3). Injection of a-1 fragment into the haemolymph causes larval mortality of 18.9%, with P value <0.05 compared to control larvae (5.7%). On the other hand, lethal effect of a-2 fragment results in 26.7% and shows significant difference with the control (P <0.05). Likewise, V-ATPase subunit c specific-dsRNA reveals similar effect on pink bollworm larvae with morbidity of 23.5% (P <0.05) (Figure 4a). Statistical analysis revealed no significance difference among the three dsRNAs treatments (P >0.05). Survived larvae from injection bioassays were kept on diet to observe larval development and pupation. Bodies' shrinkage and retardation of larval development are clearly shown of dsRNA-treated larvae due to starvation effect (Figure 4b). Some of these larvae failed to pupate due to deficiency in food absorption.

## DISCUSSION

V-ATPase plays a crucial role in the lepidopteran midgut keeping an alkaline environment in the midgut lumen and energizing secondary amino acid absorption. It is distributed with high density across the plasma membrane of the goblet cell with approximate ratio of 5000  $\mu\text{m}^{-2}$  (Vitavska et al., 2005). The dynamic function of V-ATPase was summarized by Osteresch et al. (2012). The headpiece A3B3 hexamer catalyses ATP binding and hydrolysis and the generated energy rotates the stalk that functions as a structural and functional connection between the  $V_1$  and  $V_0$ . The stalk is made of the central shaft "D and F subunits" that fills the central cavity of the headpiece (Ma et al., 2011) and the  $V_0$  subunits d with the proteolipid ring of c isoforms.  $\text{H}^+$  proton is translocated across the membrane through entering the proteolipid ring via the cytosolic half channel of subunit a. The proton binds to conserved glutamate of subunit c, upon rotation, it dissociates and leave into the lumen through the outer half channel of subunit a. The role played by V-ATPase in the insect midgut makes it a good candidate for gene silencing by RNAi.

In this study, we cloned genes encoding the membrane-bound  $V_0$  subunits a and c from the *P. gossypiella* midgut and dsRNA was *in vitro* synthesized and injected into the



**Figure 1.** Phylogenetic relationship of V-ATPase subunits. **(A)** V-ATPase subunit a. **(B)** V-ATPase subunit c. This un-rooted phylogenetic tree was constructed by the neighbor-joining method. Nodes indicate bootstrap calculated with 1000 replications support.

third larval instar. RNAi effectiveness depends on multiple factors including the amount of injected dsRNA (Chen et al., 2008). A concentration of 20 ng dsRNA/mg tissue was chosen for the current experiments comparable to previous assessments of dsRNAs on pink bollworm  $V_1$  subunits (Mohammed et al., 2015). Furthermore, high level of silencing is achieved at high doses of dsRNA (0.1 to 1  $\mu\text{g}/\text{mg}$ ) in some lepidopteran species (Terenius et al., 2011). Direct microinjection is one of the commonly used procedures for delivery of dsRNA into organisms (Bucher et al., 2002; Tomoyasu and Denell, 2004; Chen et al., 2008; Rong et al., 2013; Yao et al., 2013). Nevertheless, microinjection causes pressure on the insect body during injection and may result in wound that reduces insect survival. Therefore, deceased larvae within the first 24 h post injection were not counted. Although the  $V_0$  complex

plays a role in translocating the proton, only few reports on targeting  $V_0$  complex subunits were published. We therefore, cloned and targeted  $V_0$  subunits a and c genes and attempted to knockdown these genes by three dsRNA fragments.

Significant mortality was observed in dsRNA treatments compared to control, whereas there were no significant differences in the toxicity of the three dsRNAs. The dsRNA "a-2" was found to be most effective, followed by dsRNA "c" ( $P < 0.05$ ). The dsRNA "a-1" was the least toxic and caused mortality up to 19% only. The observed mortality of pink bollworm larvae by dsRNAs silencing  $V_0$  subunits is consistent with earlier reports with  $V_1$  subunits. Four  $V_1$  subunits (A, B, C and D) of the pink bollworm V-ATPase were knocked down by RNAi with mortality range of 23.5 to 40.5% (Mohammed et al.,

TcVATPase a KAPNPREDIDLEAHLEKTEGDIKELSESAVNLKSNYLELEIELKQVLEKIQAFFNEQDE  
 CqVATPase a RAPNPREDIDLEAHLEKTESEIMELSQNAVNLKSNYLELETELKHVLEKIQGFFFEQEG  
 BdVATPase a RAPNPREDIDLEAHLEKTENEILELAQNEVNLKSNYLELETELKRVLENTIQGFFSDQEV  
 PgVATPase a RAPNPREDIDLEAHLEKTENEILELSHNAVNLKQNYLELETELKHVLEKTEAFFVAQEE

TcVATPase a ANGLDSAHKALINDES-HNVSIRGRLGFVAGVINRERVPGFERMLWRISRGNVFLRQV  
 CqVATPase a S---GDTFRNNIIDDP-SNIQTRGRLGFVAGVIQREKVPGFERMLWRISRGNVFLRQA  
 BdVATPase a LN-LDSSNRGNADVD-VGPQNRGRLGFVAGVINRERVFAFERMLWRISRGNVFLKRS  
 PgVATPase a FG-MDPLTKSLISDETGGQAATRGRGLGFVAGVQREKVPGFERMLWRISRGNVFLRRA

TcVATPase a EIEKPLEDPATGNQLYKTVFVAFFQGEQLKTRIKKVCAGYHASLYACPSSIQERNEML  
 CqVATPase a ELEKPLEDPSTGNQIYKTVFAAFFQGEQLKTRIKKVCAGYHASLYPCPSAADEREEML  
 BdVATPase a DLDDPLKDPSTGHPIYKTVFVAFFQGEQLKNRIKKVCTGFHASMYPCPSSHTEREEMV  
 PgVATPase a ELDKPLEDPSTGNEIYKTVFVFFQGEQLKSRIKKVCTGFHASLYPCPSPNVERQDMV

TcVATPase a KGVCTRLEDLNLVLNQTDHRSRVLVSVAKELQNWVSVMSKMKAIYHTLNFNMDVTK  
 CqVATPase a KGVKTRLEDLSMVLNQTDHRSRVLSTVAKELPRWRIMVKKMKAIYHTLNFNMDVTK  
 BdVATPase a KGVTRLEDLKLVLSTQEDHRSRVLATVSKNLPSWSIMVKKMKAIYHTLNFNMDVTK  
 PgVATPase a KGVTRLEDLNMVLNQTRDHRQRVLVSVAKELPSWSIMVRKMKAIYHTLNFNMDVTK

TcVATPase a KCLIGECWVSSKDIPIVQKALS DGSSACGSSIPSFNLVINTINEDPPTFNRTNKFTRGE  
 CqVATPase a KCLIGECWVPVLDLPLIQKALS DGSAAVGSTIPSFNLVIEITSEAPPTFNRTNKFTRGE  
 BdVATPase a KCLIGECWVPTKDLVQKALS DGSAAVGSTIPSFNLVIDTINEQPPTFNRTNKFTRGE  
 PgVATPase a KCLIGECWVPVADLNVQKALADGSNACGSSIPSFNLCEITDEEPPTFNRTNKFTRGE

TcVATPase a QNLIDSYGVASYREANPALYTIITFPFLEAVMFGDVGHAMMALFGGYLVISEKKIMA  
 CqVATPase a QNLIDAYGVIASYREANPALYTIITFPFLEAVMFGDLGHGMMAAFGLWMTNERKLSA  
 BdVATPase a QNLIDAYGVASYREANPALYTCITFPFLEAVMFGDLGHGIIILALFGWMMYKSLQR  
 PgVATPase a QNLIDAYGVASYREANPALYTIITFPFLEAVMFGDFGHGATMMLFGAWMVCKEIVLSLSA

**Figure 2.** Deduced amino acid sequence between 117-406 aa of PgV-ATPase subunit a and aligned with *Tribolium castaneum* (Gene Bank™ accession number EEZ99071) (TcVATPase a) *Culex quinquefasciatus* EDS39016 (CqVATPase a) and *Bactrocera dorsalis* XP\_011211500 (BdVATPase a). The identical amino acids are shaded in black boxes.

PgVATPase c MSSPAADNPAYGPFVGMGAASAIIFSALGAAYGTAKSGTGIAAMSVMRPELIMKSII  
 HvVATPase c ----MAENPIYGPFFVGMGAASAIIFSALGAAYGTAKSGTGIAAMSVMRPELIMKSII  
 MsVATPase c ----MAENPIYGPFFVGMGAASAIIFSALGAAYGTAKSGTGIAAMSVMRPELIMKSII  
 PxVATPase c ---MSAENPIYGPFFVGMGAASAIIFSALGAAYGTAKSGTGIAAMSVMRPELIMKSII



PgVATPase c PVVMAGIIAIYGLVVAVLIAGAMDQPSAGYTLYKGFHILGAGLAVGFSGLAAGFAIGI  
 HvVATPase c PVVMAGIIAIYGLVVAVLIAGSLDAPSNNTLYKGFHILGAGLAVGFSGLAAGFAIGI  
 MsVATPase c PVVMAGIIAIYGLVVAVLIAGSLDSPSNNTLYRGFHILGAGLAVGFSGLAAGFAIGI  
 PxVATPase c PVVMAGIIAIYGLVVAVLIAGSLDSPAN-YTLYRGFHILGAGLAVGFSGLAAGFAIGI

PgVATPase c VGDAGVRGTAQQPRLFVGMILILIFAENVLGLYGLIVAIYLYTKQ  
 HvVATPase c VGDAGVRGTAQQPRLFVGMILILIFAENVLGLYGLIVAIYLYTKQ  
 MsVATPase c VGDAGVRGTAQQPRLFVGMILILIFAENVLGLYGLIVAIYLYTKQ  
 PxVATPase c VGDAGVRGTAQQPRLFVGMILILIFAENVLGLYGLIVAIYLYTKQ

**Figure 3.** Deduced amino acid sequence of PgV-ATPase subunit c and aligned with *Heliothis virescens* (HvVATPase c) and *Maduca sexta* CAB55500 (MsV-ATPase c) and *Plutella xylostella* XP\_011561281 (PxVATPase c) (PxV-ATPase c). The identical amino acids are shaded in black boxes.

2015). Current results suggest that silencing either  $V_1$  or  $V_0$  subunits have similar toxicity effects in pink bollworm.

However, other reports on other insect species focused on V-ATPase subunit A as a target for RNAi more than



(B)

**Figure 4.** The lethal effect of injecting dsRNA targeting V-ATPase subunits a and c transcripts into the haemolymph of pink bollworm larvae. The larval mortality was counted 120 h after injection. **(A)** Death bar demonstrate the mortality percentage of the injected larvae against either subunit a or c-specific dsRNAs as well as control larvae. The results are shown as the mean  $\pm$  SE. **(B)** Image of pink bollworm larvae were injected with V-ATPase specific-dsRNA (dsRNA); “i” dead larvae and “ii” retardation of larval development. Control larvae were injected with buffer showing normal development of control larvae (Cont.).

other subunits. As far as we know, transgenic plant knocking down V-ATPase was exclusively used to silence subunit A gene. Transgenic corn showed significant reduction in root damage by western corn rootworm (Baum et al., 2007). Thakur et al. (2014) developed transgenic tobacco expressing V-ATPase-A specific dsRNA, resulted in 62% reduction of the V-ATPase A transcripts level within whiteflies midguts. Transgenic plants showed high resistance to heavy infestation of

whiteflies compared to control plants. Recently, three genes encoding P450, chitin synthase B and V-ATPase A from the midgut of *H. armigera* were targeted by dsRNAs that are expressed in the chloroplast genome of transplastomic tobacco plants (Jin et al. 2015). The transcripts of these three genes were not detected in the midgut of larvae fed on tobacco leaves expressing dsRNA. As a result, the net weight and growth of fed larvae were retard and pupation rate was significantly

reduced.

Suppression of endogenous V-ATPase subunits genes showed variable results according to targeted subunit and insect species. The V<sub>1</sub> subunits A, D and E genes were targeted by oral feeding of specific dsRNA causing mortality against WCR (Baum et al., 2007). There was no apparent significant difference in the assessed dsRNAs. The V-ATPase A-dsRNA caused variable mortality; 97.5% in *B. tabaci* (Upadhyay et al., 2011), 27.3 to 54.5% in *Tetranychus urticae* (Kwon et al., 2013) and 35% in *B. dorsalis* (Li et al., 2011a). Likewise, the V-ATPase E transcript levels was reduced between 55 and 85% in *Leptinotarsa decemlineata* larvae after ingesting two dsRNA fragments (Ky et al., 2014), low silencing level detected in *M. sexta* (Whyard et al., 2009), and no observed response in *Nilaparvata lugens* (Li et al., 2011b). The V-ATPase B and D transcripts were targeted by both feeding and microinjection of dsRNAs in the corn plant hopper, *P. maidis* (Yao et al., 2013). Quantitative PCR analysis indicated a reduction of 27-fold of V-ATPase transcripts two days post injection, while ingestion of dsRNA resulted in only two fold reduction after six days of feeding. Higher mortality and lower fecundity as well as phenotypic deformation were observed in nymphs injected with 200 ng of either V-ATPase B or D dsRNA.

Different factors influence RNAi efficiency in insects such as; effective dose of dsRNA that efficiently knockdowns target transcripts (Kumar and Sarin, 2013), ability of insect cells to uptake dsRNA molecules and to involve them within the RNAi pathway (Terenius et al., 2011) and RNAi processing machinery within the cells and signal propagation across neighboring cells (Roignant et al., 2003; Miller et al., 2008). Also, length of the dsRNA, life-stage of the insect, and persistence of gene silencing, are important factors for successful RNAi application (Thakur et al., 2014).

Despite the low mortality achieved in the current study, deleterious effects on pink bollworms were noticed. V-ATPase is still a potential target for RNAi and could be deployed in the control of pink bollworm, if multiple genes could be targeted simultaneously by dsRNA, or by targeting different sites on the same target gene(s) causing an increase of larval mortality. RNAi is a promising strategy in insect pest management and further investigation is required to enhance its effect.

### Conflicts of Interests

The authors have not declared any conflict of interests.

### ACKNOWLEDGMENTS

I would like to thank the Science and Technology Development Fund (STDF), Ministry of Scientific Research, Egypt for partially funding this research through

project ID 723 and greatly acknowledge Mrs. Mervat R. Diab for her technical efforts.


### REFERENCES

- Baum JA, Bogaert T, Clinton W, Heck GR, Feldmann P, Ilagan O, Johnson S, Plaetinck G, Munyikwa T, Pleau M, Vaughn T, Roberts J (2007). Control of coleopteran insect pests through RNA interference. *Nat. Biotechnol.* 25:1322-1326.
- Bell RA, Joachim FG (1976). Techniques for Rearing Laboratory Colonies of Tobacco Hornworms and Pink Bollworms. *Ann. Entomol. Soc.* 69:365-373.
- Berezikov E (2011). Evolution of microRNA diversity and regulation in animals. *Nat. Rev. Genet.* 12:846-860.
- Beyenbach KW, Wieczorek H (2006). The V-type H<sup>+</sup>ATPase: molecular structure and function, physiological roles and regulation. *J. Exp. Biol.* 209:577-589.
- Bucher G, Scholten J, Klingler M (2002). Parental RNAi in *Tribolium* (Coleoptera). *Curr. Biol.* 12:R85-R86.
- Chen J, Zhang D, Yao Q, Zhang J, Dong X, Tian H, Zhang W (2010). Feeding-based RNA interference of a trehalose phosphate synthase gene in the brown planthopper, *Nilaparvata lugens*. *Insect Mol. Biol.* 19:777-786.
- Chen X, Tian H, Zaou L, Tang B, Hu J, Zhang W (2008). Disruption of *Spodoptera exigua* larval development by silencing chitin synthase gene with RNA interference. *Bull. Entomol. Res.* 98:613-619.
- Dow JAT (1999). The Multifunctional *Drosophila melanogaster* V-ATPase is encoded by a multigene family. *J. Bioenerg. Biomembr.* 31:75-83.
- Fire A, Xu SQ, Montgomery MK, Kostas SA, Driver SE, Mello CC (1998). Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*. *Nature* 39:806-811.
- Forgac M (2007). Vacuolar ATPases: rotary proton pumps in physiology and pathophysiology. *Nat. Rev. Mol. Cell Biol.* 8:917-929.
- Gordon KH, Waterhouse PM (2007). RNAi for insect-proof plants. *Nat. Biotechnol.* 25:1231-1232.
- Gräf R, Harvey WR, Wieczorek H (1996). Purification and properties of a cytosolic V1-ATPase. *J. Biol. Chem.* 271:20908-20913.
- Jin S, Singh ND, Li L, Zhang X, Daniell H (2015). Engineered chloroplast dsRNA silences *cytochrome p450 monooxygenase*, V-ATPase and *chitin synthase* genes in the insect gut and disrupts *Helicoverpa armigera* larval development and pupation. *Plant Biotechnol. J.* 13:435-446.
- Kumar A, Sarin NB (2013). RNAi: A promising approach to develop transgenic plants against geminiviruses and insects. *J. Plant Physiol. Pathol.* 1:1-6.
- Kwon DH, Park JH, Lee SH (2013). Screening of lethal genes for feeding RNAi by leaf disc-mediated systematic delivery of dsRNA in *Tetranychus urticae*. *Pestic. Biochem. Physiol.* 105:69-75.
- Ky F, Guo WC, Lü FG, Liu XP, Li GQ (2014). Response of the vacuolar ATPase subunit E to RNA interference and four chemical pesticides in *Leptinotarsa decemlineata* (Say). *Pestic. Biochem. Physiol.* 114:16-23.
- Li J, Chen Q, Lin Y, Jiang T, Wu G, Hua H (2011b). RNA interference in *Nilaparvatalugens* (Homoptera: Delphacidae) based on dsRNA ingestion. *Pest Manage. Sci.* 67:852-859.
- Li X, Zhang M, Zhang H (2011a). RNA interference of four genes in adult *Bactrocera dorsalis* by feeding their dsRNAs. *PLoS One* 6:e17788.
- Liu S, Ding Z, Zhang C, Yang B, Liu Z (2010). Gene knockdown by intro-thoracic injection of double-stranded RNA in the brown planthopper, *Nilaparvata lugens*. *Insect Biochem. Mol. Biol.* 40: 666-671.
- Ma B, Xiang Y, An L (2011). Structural bases of physiological functions and roles of the vacuolar H<sup>+</sup>-ATPase. *Cell Signal.* 23:1244-1256.
- Mao YB, Cai WJ, Wang JW, Hong GJ, Tao XY, Wang LJ, Huang YP, Chen XY (2007). Silencing a cotton bollworm P450 monooxygenase gene by plant-mediated RNAi impairs larval tolerance of gossypol. *Nat. Biotechnol.* 25:1307-1313.
- Miller SC, Brown SJ, Tomoyasu Y (2008). Larval RNAi in *Drosophila*?




- Dev. Genes Evol. 218:505-510.
- Mohammed A, Diab MR, Abd-Alla SMM, Hussien EHA (2015). RNA interference-mediated knockdown of vacuolar ATPase genes in pink bollworm, *Pectinophora gossypiella*. Int. J. Biol. Pharm. Allied Sci. 4:2641-2660.
- Nelson N, Perzov N, Cohen A, Hagai K, Padler V, Nelson H (2000). The cellular biology of proton-motive force generation by V-ATPases. J. Exp. Biol. 203:89-95.
- Osteresch C, Bender T, Grond S, Zezschwitz P, Kunze B, Jansen R, Huss M, Wiczorek H (2012). The binding site of the V-ATPase inhibitor apicularen is in the vicinity to those of bafilomycin and archazolid. J. Biol. Chem. 287:31866-31876.
- Pietrantonio PV, Gill SS (1997). Ductin, a component of the V-ATPase, is developmentally regulated in *Heliothis virescens* midgut, and anti-ductin antibodies label lateral membranes. Cell Tissue Res. 289:97-108.
- Price DR, Gatehouse JA (2008). RNAi-mediated crop protection against insects. Trends Biotechnol. 26:393-400.
- Roignant JY, Carré C, Mugat B, Szymczak D, Lepesant JA, Antoniewski C (2003). Absence of transitive and systemic pathways allows cell-specific and isoform-specific RNAi in *Drosophila*. RNA 9:299-308.
- Rong S, Li DQ, Zhang XY, Li S, Zhu KY, Guo YP, Ma EB, Zhang JZ (2013). RNA interference to reveal roles of  $\beta$ -N-acetyl glucosaminidase gene during molting process in *Locusta migratoria*. Insect Sci. 20:109-119.
- Rosa C, Kamita SG, Falk BW (2012). RNA interference is induced in the glassy winged sharpshooter *Homalodisca vitripennis* by actin dsRNA. Pest Manage. Sci. 68:995-1002.
- Sumner JP, Dow JAT, Earley FGP, Klein U, Jager D, Wiczorek H (1995). Regulation of plasma membrane V-ATPase activity by dissociation of peripheral subunit. J. Biol. Chem. 270:2649-2653.
- Tamura K, Stecher G, Peterson D, Filipowski A, Kumar S (2013). MEGA 6: Molecular evolutionary genetics analysis version 6.0. Mol. Biol. Evol. 30:2725-2729.
- Terenius O, Papanicolaou A, Garbutt JS, Eleftherianos I, Huvenne H, Kanginakudru S, Albrechtsen M, An C, Aymeric JL, Barthel A, Bebas P (2011). RNA interference in Lepidoptera: An overview of successful and unsuccessful studies and implications for experimental design. J. Insect Physiol. 57:231-245.
- Thakur N, Upadhyay SK, Verma PC, Chandrashekar K, Tuli R, Singh PK (2014). Enhanced whitefly resistance in transgenic tobacco plants expressing double stranded RNA of V-ATPase A gene. PLoS One 9:e87235.
- Tian H, Peng H, Yao Q, Chen H, Xie Q, Tang B, Zhang W (2009). Developmental control of a lepidopteran pest *Spodoptera exigua* by ingestion of bacteria expressing dsRNA of a non-midgut gene. Plos One 4:e6225.
- Tomoyasu Y, Denell RE (2004). Larval RNAi in *Tribolium* (Coleoptera) for analyzing adult development. Dev. Genes Evol. 214:575-578.
- Upadhyay SK, Chandrashekar K, Thakur N, Verma PC, Borgio JF, Singh PK, Tuli R (2011). RNA interference for the control of whiteflies (*Bemisia tabaci*) by oral route. J. Biosci. 36:153-161.
- Vitavska O, Wiczorek H, Merzendorfer H (2003). A novel role for subunit C in mediating binding of the H<sup>+</sup>V-ATPase to the actin cytoskeleton. J. Biol. Chem. 278:18499-18505.
- Vitavska O, Wiczorek H, Merzendorfer H (2005). The V-ATPase subunit C binds to polymeric F-actin as well as to monomeric G-actin and induces cross-linking of actin filaments. J. Biol. Chem. 280:1070-1076.
- Whyard S, Singh AD, Wong S (2009). Ingested double-stranded RNAs can act as species specific insecticides. Insect Biochem. Mol. Biol. 39:824-832.
- Wiczorek H, Grüber G, Harvey WR, Huss M, Merzendorfer H, Zeiske W (2000). Structure and regulation of insect plasma membrane H<sup>+</sup> V-ATPase. J. Exp. Biol. 203:127-135.
- Xiong Y, Zeng H, Zhang Y, Xu D, Qiu D (2013). Silencing the *HaHR3* gene by transgenic plant-mediated RNAi to disrupt *Helicoverpa armigera* development. Int. J. Biol. Sci. 9:370-381.
- Yao J, Rotenberg D, Afsharifar A, Barandoc-Alviar K, Whitfield AE (2013). Development of RNAi methods for *Peregrinus maidis*, the corn planthopper. PLoS One 8:e70243.
- Zhang X, Zhang J, Zhu KY (2010). Chitosan/double-stranded RNA nano-particle mediated RNA interference to silence chitin synthase genes through larval feeding in the African malaria mosquito (*Anopheles gambiae*). Insect Mol. Biol. 19:683-693.
- Zhao YY, Liu F, Yang G, You MS (2011). PsOr1, a potential target for RNA interference-based pest management. Insect Mol. Biol. 20:97-104.
- Zhu F, Xu JJ, Palli R, Ferguson J, Palli SR (2011). Ingested RNA interference for managing the populations of the Colorado potato beetle, *Leptinotarsa decemlineata*. Pest Manage. Sci. 67:175-182.
- Zhu JQ, Liu S, Ma Y, Zhang JQ, Qi HS, Wei ZJ, Yao Q, Zhang WQ, Li S (2012). Improvement of pest resistance in transgenic tobacco plants expressing dsRNA of an Insect-Associated gene EcR. PLoS One 7:e38572.
- Zhu KY (2013). RNA interference: A powerful tool in entomological research and a novel approach for insect pest management. Insect Sci. 20:1-3.


SUPPLEMENTARY FIGURES



**Figure S-1.** Cloning of VATPase subunits a and c. PCR products of subunit a; (1) 220 bp using degenerate primer set (VATPavo2264FD/VATPavo2484RD), (2) 977 bp using degenerate/specific primers (VATPavo1439FD/VATPavo2416RS), (3) 783 bp using (VATPavo776FD/ VATPavo1559RS) and subunit c (4) 150 bp using degenerate primer set (VATPcvoFD/ VATPcvoRD), (5) 755 bp using degenerate/specific primers (VATPcvo146FD/ VATPcvo901RS). (M) 1 Kb DNA marker.



**Figure S-2.** RACE reactions; (1) 5'-end of subunit a was amplified using outer 5 prime supplied with the kit with VATPavo864RS specific primer followed by PCR reaction using inner 5 prime with VATPavo795RS specific primer. (2) 5'-end of subunit c was amplified using outer and inner prime with specific primers VATPcvo244RS and VATPcvo185RS, respectively. (3) 3'-end of both subunits a and c were synthesized using outer and inner 3 prime supplied with the kit with specific primers VATPcvo819FS and VATPcvo847FS for subunit c, and with specific primers VATPavo2265 FS and VATPavo2431FS for subunit a.



**Figure S-3.** Synthesis of dsRNA; a-1 dsRNA fragment of 220 bp using VATPavo2264F/ VATPavo2484R primer set; a-2 dsRNA fragment of 466 bp using VATPavo1950F/VATPavo2416R and c-dsRNA fragment of 740 bp using VATPcvo161F/ VATPcvo901R. (M) 1 Kb DNA marker.