

Short Communication

Tetrapropylammonium bromochromate and tetrabutylammonium bromochromate $[\text{NR}_4]\text{CrO}_3\text{Br}$, (R= Pr, Bu): two new and efficient reagents for oxidation of alcohols

Shahriare Ghammamy^{1,2*}, Kheyrollah Mehrani¹, Hamid Afrand¹ and Maryam Hajighahramani¹

¹Department of Chemistry, Islamic Azad University, Ardabil Branch, Ardabil, Iran.

²Department of Chemistry, Faculty of Science, Imam Khomeini International University, Ghazvin, Iran.

Accepted 15 August, 2007

Tetrapropylammonium bromochromate (VI), $[\text{N}(\text{Pr})_4]\text{CrO}_3\text{Br}$, TPABC and tetrabutylammonium bromochromate (VI), $[\text{N}(\text{Bu})_4]\text{CrO}_3\text{Br}$, TBABC were prepared and used to quantitatively oxidizing of a number of organic substrates. These reagents are versatile reagents for the effective and selective oxidation of organic substrates, in particular for alcohols, under mild conditions.

Key words: Chromium (VI), reagent, tetrapropylammonium bromochromate, tetrabutylammonium bromochromate, oxidation, organic substrates, alcohols.

INTRODUCTION

Organic chemists must often choose from hundreds of oxidizing agents and reaction conditions to perform a desired oxidation without affecting other functional groups present or causing side reactions. Therefore, the search for new oxidizing agents is of interest to synthetic organic chemists. In recent years, significant improvements were achieved by the use of new oxidizing agents, (Maier et al., 1999; Bhandari et al, 2001; Meenahshisundaram and Soctaungam, 2001; Heravi et al., 1999; Chaubey et al. 2003; Bhattacharjee et al., 1982; Jeyanthi et al., 2002; Shirini et al., 2003; Srinivasan et al., 1997) such as tetramethylammonium fluorochromate (Mahjoub et al., 2003), 2,2'- bipyridinium chlorochromate, (Corey and Fleet, 1973), tributylammonium chlorochromate (Ghammamy and SeyedSadjadi, 2005), and 3,5-Dimethyl-pyrazolium fluorochromate (Bhattacharjee et al., 1987). Most of these reagents that have been developed so far suffer from at least one of the drawbacks such as high acidity, photosensitivity, instability, tedious work-up procedures, or requirement of large excess of reagents.

Therefore, the search for permanent new reagents persisted which have now led to the synthesis of tetra-propylammonium bromochromate (VI), $[\text{N}(\text{Pr})_4]\text{CrO}_3\text{Br}$, TPABC and tetrabutylammonium bromochromate (VI), $[\text{N}(\text{Bu})_4]\text{CrO}_3\text{Br}$, TBABC.

EXPERIMENTAL







All reagents and solvents are of reagent grades. ¹H and ¹³C NMR spectra were recorded using Bruker DRX-500 in CD₃CN solutions. IR spectra were recorded using a Shimadzu IR-420 spectrometer. The UV/Visible measurements were made on an Uvicon model 922 spectrometer. Melting points were obtained on an Electro-thermal 9100 apparatus. All separations and quantization of alcohols and aldehydes were performed using a Philips 4410 gas Chromatograph.

Preparation of Tetrapropylammonium bromochromate, $[\text{N}(\text{Pr})_4]\text{CrO}_3\text{Br}$

Chromium (VI) oxide (1 g, 10 mmol) was dissolved in dry acetonitrile (25) in a beaker and a stoichiometric amount of tetrapropylammonium Bromide (2.66 g, 10 mmol) was added under stirring at room temperature. Within 5 min, a clear orange solution formed which upon refrigerating gave solid TPABC, which was isolated by filtration. The solid was washed with hexane and dried under vacuum

*Corresponding author. E-mail: shghamami@ikiu.ac.ir or shghamamy@yahoo.com.

Table 1. Oxidations via TBABC and TPABC.

Substrate	Product	TBABC		TPABC	
		Time (min)	Yield (%)	Time (min)	Yield (%)
n-C ₃ H ₇ -OH	n-C ₂ H ₅ -CHO	130	90	120	88
2-C ₃ H ₇ -OH	2-C ₂ H ₅ -CHO	120	90	132	87
n-C ₄ H ₉ -OH	n-C ₃ H ₇ -CHO	100	89	95	85
2-C ₄ H ₉ -OH	2-C ₃ H ₇ -CHO	95	93	90	91
n-C ₅ H ₁₁ -OH	n-C ₄ H ₉ -CHO	97	87	104	86
n-C ₈ H ₁₇ -OH	n-C ₇ H ₁₅ -CHO	70	84	68	83
		145	94	155	89
		35	96	30	95
		5	92	5	90

for 1 h. C₁₂H₂₈BrCrO₃N: Calcd. %C, 39.93; %H, 7.65. Found: %C, 40.14; %H, 8.01. IR. (KBr): 844cm⁻¹ v₁(A₁) or v(CrO₃), 948cm⁻¹ v₄(E) or v(CrO₃). UV/Visible was all consistent with the TPABC structure. Electronic absorption at 452 nm, corresponding to 1a₂→9e (ε = 161 M⁻¹ cm⁻¹); 363 nm to 8e→9e (ε = 737 M⁻¹ cm⁻¹); and 261 nm to 12a₁→9e (ε = 1683 M⁻¹ cm⁻¹).

Preparation of tetrabutylammonium bromochromate, [N(Bu)₄]CrO₃Br

Tetrabutylammonium bromochromate(IV), [N(Bu)₄]CrO₃Br was prepared by dissolving Chromium (VI) oxide (1 g, 10 mmol) in MeCN and addition of this solution to a solution of tetrabutylammonium Bromide (3.22 g, 10 mmol) in MeCN under stirring at room temperature until an orange precipitate was formed. After 2 h stirring, the mixture was filtered. The solid was washed with hexane and dried under vacuum for 1 h. C₁₆H₃₆BrCrO₃N: Calcd. % C, 45.508; %H, 8.532. Found: %C, 46.40; %H, 8.91. IR. (KBr): 895cm⁻¹ v₁(A₁) or v(CrO₃), 949cm⁻¹ v₄(E) or u(CrO₃). UV/Visible ¹³C-NMR and ¹H-NMR were all consistent with the TBABC structure. Electronic absorption at 454 nm, corresponding to 1a₂→9e (ε = 130 M⁻¹ cm⁻¹); 363 nm to 8e→9e (ε = 670M⁻¹ cm⁻¹); and 260nm to 12a₁→9e (ε =1622 M⁻¹ cm⁻¹).

Oxidation of alcohols: general method

Tetraalkylammonium bromochromate (10 mmol) is added in one portion, to a stirred solution of each alcohol (25 ml) in CH₂Cl₂, at room temperature. The progresses of the reactions are monitored by TLC and UV/Vis spectrophotometry (at 363 nm). The mixture were stirred and refluxed for the time indicated in the Table 1 at room temperature, diluted with CH₂Cl₂ and filtered. Evaporation of solvent furnished the product. The molar ratio of substrate to oxidants was 1:1. The solution became homogeneous briefly before the black-brown reduced reagent precipitated. Products are charac-

terized by comparison with authentic samples (NMR, IR, TLC and mp/bp measurement).

RESULTS AND DISCUSSION

Tetraalkylammonium bromochromates could be easily prepared in good yield, quite stable when stored dry and in the absence of light, and are active as oxidizing agents for the conversion of alcohols to carbonyl compounds.

[N(Bu)₄]CrO₃Br and [N(Pr)₄]CrO₃Br were prepared by the reaction of relative salts with CrO₃ in 1:1 ratio in the acetonitrile solvent. (Scheme 1).



Scheme 1


They have been found that these reagents have certain advantages over similar oxidizing agents in terms of the amounts of oxidants and solvent required, and especially in the short reaction times required and in the higher yields of the product (Table 1).

The nature of the solvent does not appear to be particularly critical. Hydrocarbons, benzene, ethers and chlorinated hydrocarbons are equally effective, the practical choice being oriented by the solubility of the products and the desired reaction temperature. The chromium (VI) contents easily determined iodometrically. The IR spectra of TBABC and TPABC are similar to that of other bromochromates (Harmon et al., 1974). The IR spectrum of the CrO₃Br⁻ ion has been the subject of several publications

(Harmon et al., 1974; Nakamoto, 1978). The XY_3Z ions have six infrared active vibrations. These infrared active vibrations are $\nu_1(A_1)$ or $\nu(XY_3)$, $\nu_2(A_1)$ or $\nu(XZ)$, $\nu_3(A_1)$ or $\delta(XY_3)$, $\nu_4(E)$ or $\nu(XY_3)$, $\nu_5(E)$ or $\delta(XY_3)$ and $\nu_6(E)$ or $\rho_r(XY_3)$. In TBABC and TPABC, $\nu_1(A_1)$ or $\nu(XY_3)$, and $\nu_4(E)$ or $\nu(XY_3)$ are seen at 895 and 949 cm^{-1} , 844 and 948 cm^{-1} respectively.

In conclusion, the ready preparation of these reagents, their stability, nonhygroscopicity, the ease of the work up of the reaction mixtures, reasonable yields of products and reaction time make tetraalkylammonium bromo-chromates versatile and practical reagents for the oxidation of alcohols and useful additions to the presently available bench reagents in organic synthesis. TBABC and TPABC reagents are easy to handle, could be weighed and have no hazardous effects.

Tetraalkylammonium bromochromates in dichloro-methane also oxidize primary and secondary alcohols, respectively, the corresponding aldehydes or ketones in high yields (Table 1) (Scheme 2).



Scheme 2

During the reactions, the color of the oxidants change from orange to brown, providing visual means for ascertaining the progress of the oxidations. The results obtained with tetraalkylammonium bromochromates are very satisfactory and show the new reagents as valuable addition to the existing oxidizing agents.

Tetraalkylammonium bromochromates are soluble in water, dimethylformamide, acetonitrile and acetone; they are less soluble in dichloromethane and only sparingly soluble in benzene, carbon tetrachloride, chloroform and hexane.

The ^1H and ^{13}C NMR spectra data of the compounds in CD_3CN solvent confirm the proposed structures of the new reagents especially show the cationic parts. Important peaks in the ^1H and ^{13}C NMR spectra of each compound are discussed.

ACKNOWLEDGEMENT

The authors thank Dr. A. R. Mahjoub and Mr. K. Ghoddamy for valuable discussion.

REFERENCES

- Bhandari A, Sharma PK, Banerji KK (2001). Kinetics and mechanism of the oxidative regeneration of carbonyl compounds from oximes by pyridinium chlorochromate, Indian J. Chem. Sec. A, 40: 470.
- Bhattacharjee MN, Chaudhuri MK, Purkayastha S (1987). Some Aspects of Pyridinium Chlorochromate, $\text{C}_5\text{H}_5\text{CrO}_3\text{F}$ (PFC), Oxidations. Stoichiometry of Alcohols, Evidence for Oxygen Transfer, and Identity of the Reduced Chromium Species. Tetrahedron, 43: 5389.
- Bhattacharjee MN, Chaudhuri MK, Dasgupta HS (1982). Pyridinium Fluorochromate: A New and Efficient Oxidant for Organic Substrates. Synthesis, p. 588.
- Chaubey GS, Das S, Mahanti MK (2003). Quinolinium Dichromate Oxidation of Aliphatic Aldehydes: A Kinetic Study, Croatica Chemica Acta, 76: 287.
- Corey FJ, Fleet GWJ (2005). Chromium Trioxide-3,5-Dimethylpyrazole Complex as a Reagent for Oxidation of Alcohols to Carbonyl Compounds. Tetrahedron Lett, 45: 1973, 4499.
- Ghammamy S, SeyedSadjadi SA (2005). Tetrabutylammonium Fluorochromate(VI), (TBAFC): As a Mild and Efficient Reagent for Oxidation of Organic Substrates, J. Serb. Chem. Soc., 70: 1243.
- Harmon KM, Gennick I, Madeira SL (1974). Hydrogen bonding. IV. Correlation of infrared spectral properties with C-H...X hydrogen bonding and crystal habit in tetramethylammonium ion salts, J. Phys. Chem, 78: 2585.
- Heravi MM, Kiaakoojori R, Mirza-Aghayan M, Tabar-Hydar K Bolourtchian M (1999). An Aluminiumoxid adsorbiertes 3-Carboxypyridiniumchlorochromat als selektives Oxidationsmittel für primäre und sekundäre Alkohole, Monatsh. Chem. pp. 130 -481.
- Jeyanthi GF, Vijayakumar G, Elango KPJ (2002). The effect of solvent on the benzaldehydes by quinolinium chlorochromate in aqueous organic solvent media. Serb. Chem. Soc, 67: 803.
- Mahjoub AR, Ghammami S, Kassae MZ (2003). Tetramethylammonium fluorochromate(VI): a new and efficient oxidant for organic substrates. Tetrahedron Lett. 44: 4555.
- Maier G, Reisenauer HP, Marco MD, Angew (1999). Isonitroso hydrogen (hydroxy nitrene, Ion). Angew. Chem. Int. Ed, 38: 108.
- Meenahshisundaram S, Soctaugam R (2001). 3. Meenahshisundaram, S.; Soctaugam, R. Kinetics and mechanism of the quinolinium fluorochromate oxidation of some primary, secondary and unsaturated alcohols in acetonitrile. Collect. Czech. Chem. Commun., 66: 877.
- Shirini F, Mohammadpoor-Baltork I, Hejazi Z, Heravi P (2003). Caffeineinium Chlorochromate: A Mild and Efficient Reagent for Oxidation of Alcohols and Chemoselective Oxidative Cleavage of Oximes Bull. Korean Chem. Soc., 24: 517.
- Srinivasan R, Stanley P, Balasubramanian K (1997). Isoquinolinium Fluorochromate: A New and Efficient Oxidant for Organic Substrates, Synthetic Communications, 27: 2057.