

Full Length Research Paper

Bioactivity of *Hydnora africana* on selected bacterial pathogens: Preliminary phytochemical screening

Nethathe B. B.¹ and Ndip R. N.^{1,2*}

¹Microbial Pathogenicity and Molecular Epidemiology Research Group, Department of Biochemistry and Microbiology, Faculty of Science and Agriculture, University of Fort Hare, P/Bag X1314, Alice 5700, South Africa.

²Department of Biochemistry and Microbiology, Faculty of Science, University of Buea, Box 63, Buea, Cameroon.

Accepted 23 August, 2011

Medicinal plants contain a variety of chemical substances with important therapeutic properties that can be utilized in the treatment of human diseases. *Hydnora africana* is used in folklore remedies for the treatment of diarrhoea, dysentery, kidney and bladder complaints among other ailments; hence we assessed the *in vitro* antimicrobial activity of this plant against three bacterial species (*Helicobacter pylori* ATCC 43526, *Helicobacter pylori* PE 252C, *Staphylococcus aureus* NCTC 6571 and *Aeromonas hydrophila* ATCC 35654). The agar well diffusion method was used to determine the susceptibility of bacterial strains to crude extracts of the plant. The minimum inhibitory concentration (MIC₅₀) and minimum bactericidal concentration (MBC) of the active crude extracts were determined by the microdilution test. Ciprofloxacin (0.0125 mg/mL) was used as positive control. The presence of phytochemicals was also assessed using standard methods. Results were analyzed statistically by the one-way ANOVA test. *Hydnora africana* demonstrated antimicrobial activity against all the organisms with a mean zone diameter of inhibition ranging from 0 to 22 mm. The MIC₅₀ of the extracts ranged from 0.078 to 2.5 mg/mL and MBC ranged from 0.78 to 25 mg/mL. Phytochemical assay revealed the presence of alkaloids, tannins, flavonoids, saponins and steroids in the extracts. It is concluded that *H. africana* may contain compounds with therapeutic activity.

Key words: *Hydnora africana*, medicinal plant, minimum inhibitory concentration, minimum bactericidal concentration, phytochemicals.

INTRODUCTION

Plants have a great potential for producing new phytochemicals with profound antimicrobial activity against human pathogens. According to the world health organization (WHO), more than 80% of the world's population depends on traditional medicine for their primary health care needs (Duraipandiyar et al., 2006). There is an upsurge of resistant microbial strains to conventional antimicrobials necessitating the need for a search and development of new drugs to circumvent the problem (Panda et al., 2009).

Hydnora africana is a parasitic plant which is predominant in the dry and semi-arid parts of the Succulent

Karoo, Eastern Cape Karoo and the dry coastal thickets between the Eastern Cape and KwaZulu-Natal Provinces of South Africa (Asfaw et al., 1999). The plant belongs to the family *Hydnoraceae* (Musselman, 1991). It is renowned for its healing properties, and has been used by traditional medicine practitioners to treat ailments such as diarrhea, dysentery, kidney and bladder complaints (Van and Gericke, 2000). For example, infusions are used as face wash to treat acne by the Xhosa people of South Africa (Van and Gericke, 2000). More than 12 *Hydnora* species have been described with antimicrobial properties (Bolin and Musselman, 2009). However, there is paucity of data in South Africa on its antimicrobial activity as well as its chemical constituents, hence the present investigation against different Gram-positive and negative bacterial pathogens. *Helicobacter pylorus* is a Gram-negative, microaerophilic bacterium that causes

*Corresponding author. E-mail: ndip@ufh.ac.za; ndip3@yahoo.com. Tel: +27 782696191. Fax: +27 866224759.

duodenal ulceration, chronic gastritis and gastric cancer. (Ndip et al., 2008). An increasing number of infected individuals are found to harbour antibiotic-resistant strains (Ndip et al., 2008; Tanih et al., 2010). The emerging resistance to antibiotics, especially metronidazole and amoxicillin limits their use in the treatment of infections (Smith et al., 2001; Tanih et al., 2010). *Aeromonas hydrophila* is a heterotrophic, Gram-negative, rod shaped bacterium, mainly found in areas with a warm climate. This bacterium can also be found in fresh, salt, marine, estuarine, chlorinated, and un-chlorinated water (Villari et al., 2003). It causes gastroenteritis, cellulitis, myonecrosis and eczema. Its increasing resistance to agents such as chloramphenicol, florenicol, tetracycline, sulfonamide, nitrofurans derivatives, and ciprofloxacin used to eliminate and control impending infection requires attention (Wang et al., 2011). *Staphylococcus aureus* is a facultative anaerobic, Gram-positive coccus and the most common cause of staphylococcal infections (Cosgrove et al., 2009). It causes a range of illnesses from minor skin infections, such as pimples, impetigo, boils, cellulitis, folliculitis, carbuncles, scalded skin syndrome and abscesses, to life-threatening diseases such as pneumonia, meningitis, osteomyelitis, endocarditis, toxic shock syndrome, bacteremia and sepsis (Kluytmans et al., 1997). The treatment of choice for *S. aureus* infection is penicillin; but in most countries, penicillin-resistance as well as MRSA is extremely common (Neely and Maley, 2000; Nkwelang et al., 2009). This study therefore documents the antimicrobial activity of *H. africana* against the selected bacterial pathogens, and a preliminary assessment of the possible phytochemicals responsible for its action.

MATERIALS AND METHODS

Bacterial strains

The following standard strains of bacteria *Aeromonas hydrophila* ATCC 35654, *S. aureus* NCTC 6571, *Helicobacter pylori* ATCC 43526 and a local metronidazole-resistant strain of *H. pylori* PE 252C isolated in our laboratory (Tanih et al., 2010) were used. Cultures of the other organisms were resuscitated and maintained on nutrient agar slants at 4°C (Cheesbrough, 1982; Cowan and Steel, 2004), while *H. pylori* was suspended in 20% glycerol and stored at -80°C until used.

Preparation of plant extracts

Hydnora africana was selected based on ethnobotanical information and preliminary data obtained in our laboratory. It was identified in collaboration with botanists at the University of Venda, Limpopo Province, South Africa where voucher specimens (BP04) have been deposited.

The method described by Ndip et al. (2008) to prepare extracts was employed with modifications. The plant was harvested, air dried for 2 weeks and ground to fine powder using a blender (ATO MSE mix, 702732, England). Organic solvents including methanol, ethanol, acetone, ethyl acetate (100%) and water were used for extraction. Briefly, the dried plant material (2.8 kg) was macerated

in five fold excess of the solvent in extraction pots such that the level of the solvents was above that of the plant material. The slurry was put in a shaker incubator (Edison, N.J., USA) regulated at room temperature (RT) for 48 h then centrifuged at 300 rpm for 5 min (Model T-J-6 Beckman, USA) and filtered using filter papers of pore size 60^Å. The process was repeated twice for a total of three extractions (Okeleye et al., 2010). The combined extracts was concentrated in a rotavapor (BUCHI R461, Switzerland) and transferred to labelled vials and allowed to stand at RT to permit evaporation of residual solvents. A 3 gram sample of each plant extract was used for the preliminary bioassay, and 3 g kept in the extract bank for subsequent use. Stock solutions were prepared by dissolving the extracts in 10% Dimethyl Sulphoxide (DMSO).

Antibacterial susceptibility test

The agar well diffusion technique was employed as previously described by Dastouri et al. (2008). For *H. pylori*, Columbia base agar was prepared following the manufacturer's instructions, supplemented with 7% defibrinated horse blood and Skirrow's supplement (Oxoid, UK) while for *A. Hydrophila* and *S. aureus*, Muller-Hilton agar (Oxoid, UK) was prepared following the manufacturer's instructions. A 0.5 McFarland standard was prepared by the method of Koneman et al. (1992), and 5 mL put into a sterile test tube. An inoculum of each microorganism was prepared from subcultures of the bacterial suspension and estimated to contain 10⁸ colony forming units. Four to five colonies of the same morphological type were picked, emulsified in 0.9% physiological saline and used to evenly inoculate specific agar plates depending on the microorganisms. Wells were cut in each agar plate with a 6 mm cork borer. About 100 µL of the different concentrations (200, 100, 50 mg/mL) of the extract were put separately into each well, in each plate. Ciprofloxacin (0.0125 mg/mL) was used as a positive control. The plates were incubated at 37°C for 24 h for *A. hydrophila* and *S. aureus* and 3 to 5 days for *H. pylori* under microaerophilic conditions (Anaerocutt, Baringstoke, England); the diameter of the zone of inhibition was measured and recorded in millimeters. The experiment was repeated 2 times for each strain.

Determination of minimum inhibitory concentration (MIC₅₀)

MIC₅₀ determination was carried out by the microdilution test method in 96 well plates as earlier described (Banfi et al., 2003; Njume et al., 2010) with slight modifications. Briefly, Two-fold dilutions of the most potent extracts and antibiotic (ciprofloxacin) were prepared in the test wells in complete Brian Heart Infusion (BHI) broth (Oxoid, UK); the final extracts and antibiotic concentrations ranged from 0.0024 to 5 mg/mL. Twenty microlitres of each bacterial suspension was added to 180 µL of extract containing culture medium. Control wells were prepared with culture medium and bacterial suspension only. Also included was culture medium and extract only at different concentrations. An automatic ELISA micro plate reader (Model 680, Bio-Rad, Japan) adjusted to 620 nm was used to measure the absorbance of the plates before and after 24 h incubation. The absorbencies were compared to detect an increase or decrease in bacterial growth. The lowest concentration of the test extract resulting in inhibition of 50% of bacterial growth was recorded as the MIC.

Determination of minimum bactericidal concentration (MBC)

To determine the MBC, 0.2 mL of the contents of the MIC was serially diluted tenfold in 0.9% physiological saline (Ndip et al., 2007). A loop full was taken from each tube and inoculated onto BHI agar plates. The MBC was recorded as the lowest concen-

tration of the extract that gave complete inhibition of colony formation of the test bacteria at the latter cultivation.

Phytochemical screening of the extracts

A small portion of the dry extract was subjected to phytochemical test using previously established methods (Akinpelu et al., 2009) to test for alkaloids, tannins, flavonoids, steroids and saponins.

Test for alkaloids

Exactly 0.5 g of the plant extract was dissolved in 5 mL of 1% HCl on steam bath. A millilitre of the filtrate was treated with drops of Dragendorff's reagent. Turbidity or precipitation was taken as indicative of the presence of alkaloids.

Test for tannins

About 1 g of the extract was dissolved in 20 mL of distilled water and filtered. Two to three drops of 10% FeCl₃ were added to 2 mL of the filtrate. The production of a blackish-blue or blackish-green colouration was indicative of tannins. To another 2 mL of the filtrate was added 1 mL of bromine water. A precipitate was taken as positive for tannins.

Test for flavonoids

A 0.2 g of the extract was dissolved in 2 mL of methanol and heated. A chip of magnesium metal was added to the mixture followed by the addition of a few drops of concentrated HCl. The occurrence of a red or orange colouration was an indication of the flavonoids.

Test for saponins

Two grams of the extract was boiled in 20 mL of distilled water in a water bath and filtered (Acrodisc syringe filter pall, USA). Approximately 10 mL of the filtrate was mixed with 5 mL of distilled water and shaken vigorously for a stable persistent froth. The frosting was mixed with 3 drops of olive oil and shaken vigorously, then observed for the formation of an emulsion.

Test for steroids

About 0.5 g of the extract was dissolved in 3 mL of CHCl₃ and filtered. Concentrated H₂SO₄ was added to the filtrate to form a lower layer. A reddish brown colour was taken as positive for steroid ring.

Statistical analysis

Analysis was performed using the SPSS Version 17.0 (Illinois USA, 2009). The one way ANOVA test was used to determine if there was any statistically significant difference in the zone diameters of inhibition of the different solvent extracts; the MIC₅₀ of the extracts and the control antibiotic (ciproxacin). P-values < 0.05 were considered significant.

RESULTS

Extract yield

Different solvents including ethyl acetate, acetone,

ethanol, methanol and water were used for extraction because the type of solvent used may have an effect on the nature of the compounds extracted, the quantity extracted and the resulting bioactivity of the extract. Water extracted the highest quantity followed by methanol, and ethyl acetate the least (Figure 1).

Antimicrobial susceptibility testing

The extracts showed *in vitro* activity against the test bacterial pathogens with the exception of water. The mean zone diameter of inhibition ranged from 0 to 22 mm (Table 1). Acetone, methanol and ethyl acetate were the most active extracts for *S. aureus*, *A. hydrophila* with mean zone diameter of inhibition ranging from 13 to 22 mm, while for *H. pylori*, methanol and ethyl acetate extracts showed activity with mean zone diameter ranging from 14 to 21 mm. The most active crude extracts (methanol and ethyl acetate) against all test microorganisms were statistically significant (P < 0.05) compared to all other extracts, but not to ciprofloxacin (P > 0.05) the positive control, with mean zone diameter ranging from 14 to 17 mm. DMSO used as negative control, showed no activity. An inhibition zone of ≥ 12 mm was chosen as representative of bacterial susceptibility to the extracts. The breakpoint of ciprofloxacin (0.05 mg/mL) was taken at 21 mm (CLSI, 2008).

MIC and MBC determination

The active extracts were further assayed to determine their MIC₅₀ and MBC against the bacterial pathogens. Although only methanol and ethyl acetate extracts showed activity against *H. pylori*; methanol, acetone, ethanol and ethyl acetate extracts were active against *S. aureus* and *A. hydrophila*. Subsequently methanol and ethyl acetate extracts were used for the determination of MIC₅₀ and MBC for *H. pylori* (ATCC 43526 and PE 252C) and methanol, acetone, ethanol and ethyl acetate extracts were used to determine MIC₅₀ and MBC for *S. aureus* and *A. hydrophila*. The MIC₅₀ and MBC ranged from 0.078 to 2.5 mg/mL and 0.78 to 25 mg/mL, respectively for all studied microorganisms (Tables 2 and 3), while for ciprofloxacin it ranged from 0.0098 to 0.078 mg/mL and 0.78 to 0.098 mg/mL, respectively.

The Gram positive bacterium (*S. aureus*) was most susceptible to *H. africana* compared to the Gram negative bacteria (*A. hydrophila* and *H. pylori*). However, there was no statistically significant difference (P > 0.05) between the MIC₅₀ and MBC of the different solvents against Gram negative and Gram positive organisms.

Phytochemical analysis

Phytochemical analysis of the four extracts is




Figure 1. Quantity (grams) of *H. africana* flower extracted with different solvents. X-axis shows the different solvents used for extraction and Y-axis shows quantity extracted in grams.

Table 1. Antibacterial activity of extracts of *H. africana* against selected bacterial pathogens.

	Zone diameter at different concentration (mm)*																
	Methanol			water			Acetone mg/mL			Ethyl acetate			Ethanol			Cipro	
SBP	200	100	50	200	100	50	200	100	50	200	100	50	200	100	50	0.025	0.0125
<i>S.a</i>	17±2.1	21±2.1	22±2.1	0	0	0	20±0.7	19±1.4	22±3.5	16±0.7	18±0.7	19±0.7	14±1.4	16±1.4	17±1.4	17±0.7	17±0.7
<i>A.h</i>	17±0.7	16±0.7	15±1.4	0	0	0	17±2.1	17±1.4	18±1.4	14±1.4	13±1.4	15±0	13±0.7	16±1.4	14±0.7	17±1.4	17±1.4
<i>H.p1</i>	20±0.7	16±1.4	15±1.4	0	0	0	0	0	0	15±2.8	17±1.4	14±1.4	0	0	0	14±1.4	14±1.4
<i>H.p2</i>	17±2.1	21±2.1	18±0.7	0	0	0	0	0	0	16±0.7	19±1.4	17±0.7	0	0	0	15±0.7	15±0.7

SBP, selected bacterial pathogens; *S.a*, *S. aureus*; *A.h*, *A. hydrophila*; *H.P1*, *H. pylori* 43526; *H.p2*, *H. pylori* PE 252C; Cipro, ciprofloxacin; *, experiment was repeated twice and zone of inhibition recorded as mean zone diameter ±SD. Sensitivity zone ≥ 12 mm.

Table 2. MIC₅₀ of different solvents extract of *H. africana* and the antibiotic against selected bacterial pathogens.

SBP	Extracts and Antibiotic (mg/mL)				
	Methanol	Acetone	Ethanol	Ethyl acetate	Ciproxacillin
<i>S. aureus</i>	0.625	0.156	0.156	0.078	0.0098
<i>A. hydrophila</i>	0.3125	0.156	–	0.078	0.0098
<i>H. pylori43526</i>	2.5	ND	ND	1.25	0.078
<i>H. pylori252C</i>	–	ND	ND	2.5	0.078

SBP, selected bacterial pathogens; –, MIC₅₀ values not within susceptible range; ND, not determined.

Table 3. MBC (mg/mL) of different solvent extracts of *H. africana* and antibiotic against selected bacterial pathogens.

SBP	Extracts/ Antibiotic (mg/mL)				
	Methanol	Acetone	Ethanol	Ethyl acetate	Ciproxacillin
<i>S. aureus</i>	6.25	1.56	1.56	0.78	0.098
<i>A. hydrophila</i>	3.125	1.56	–	3.125	0.098
<i>H.pylori</i> 43526	25	–	–	12.5	0.78
<i>H.pylori</i> 252C	ND	–	–	ND	0.78

SBP, selected bacterial pathogens; –, MBC values not within susceptible range; ND, not determined.

Table 4. Phytochemical constituents of different solvent extracts of *H.africana*.

Phytochemicals	Solvent extracts			
	Methanol	Acetone	Ethanol	Ethyl acetate
Alkaloids	+++	++	+++	++
Saponins	+++	++	++	+++
Tannins	+++	+++	+++	+++
Flavonoids	+++	+++	+++	+++
Steroids	+++	+++	+++	+++

+++, Present in large quantity; ++, Present in moderate quantity.

in Table 4. The results revealed the presence of the following secondary metabolites: alkaloids, saponins, tannins, steroids and flavonoids, based on colour, haemolysis, turbidity, layers, emulsification and precipitation following the reactions.

DISCUSSION

The phytoconstituents of various plants have longed been known and their antimicrobial properties have been widely reported (Nostro et al., 2000; Roy et al., 2006). The antimicrobial activities of plant extracts have been linked to the presence of some bioactive compounds. These secondary metabolites also serve to protect the plants against bacterial, fungal and viral infections (De and Ifeoma, 2002; El-Mahmood and Amey, 2007). These bioactive compounds are known to work synergistically to produce various effects on human and animal subjects (Amagase, 2006). However, most reports on *H. africana* have focused mainly on the morphology of the plant, while information on its activity against hospital based pathogens is scanty (Bolin et al., 2009).

The extraction of active compounds from plant material and their activity depends on the type of solvent used in the extraction process (Parekh et al., 2005; Majhenic et al., 2007). In this study, it was observed that plant extractions with organic solvents provided stronger antibacterial activity than extraction with water. This study confirms the results of previous studies, which reported that water is not a suitable solvent for extraction of antibacterial compounds from medicinal plants compared

to organic solvents, such as methanol (Karaman et al, 2003; Moniharapon and Hashinaga, 2004; Parekh et al, 2005; Majhenic et al, 2007).

S. aureus and *A. hydrophila* were the most susceptible organisms to all solvent extracts except water, with mean zone diameter of inhibition ranging from 13 to 22 mm. Moreover *H. pylori* (43526 and PE 252C) were only susceptible to methanol and ethyl acetate with mean zone diameter that ranged from 14 to 21 mm. This variation in activity may be due to the solvent used for extraction, e.g., methanol and ethanol are used for alkaloid extraction; acetone for flavonoids and steroids and ethanol may also be used for sterols, polyphenols, and tannins (Büssing, 1996). However, lack of activity demonstrated by water extracts *in vitro* against the organisms, does not necessarily imply that they would demonstrate weak activities *in vivo*.

In line with the findings of this study, another study had demonstrated very potent antibacterial activity of *Hydnora abyssinica*. In their study, Saadabi and Ayoub (2009) screened crude extracts of the family *Hydnoraceae* and reported potent antibacterial activity against common pathogenic Gram-negative and positive bacteria including *Escherichia coli*, *Pseudomonas aeruginosa*, *Bacillus subtilis* and *Staphylococcus aureus*.

In the present study, MIC₅₀ and MBC recorded for *H. africana* against all studied microorganisms ranged from 0.078 to 2.5 mg/mL and 0.78 to 25 mg/mL, respectively. However, the methanol extract showed no inhibition at MIC₅₀ against *H. pylori* PE 252C. The MIC results confirm earlier findings by Nariman et al. (2004) who documented MIC ranges of 0.0037 to 2 mg/mL. Also, MIC values of

0.0625 to 0.5 mg/mL have been documented for East African medicinal plants against similar bacterial pathogens (Fabry et al., 1996). A Gram positive bacterium (*S. aureus*) was most susceptible to *H. africana* compared to the Gram negative bacteria (*A. hydrophila* and *H. pylori*). Most plants extracts are active against Gram positive bacteria; this has been attributed to the fact that the cell wall of Gram positive bacteria is easier to penetrate than the Gram negative bacteria which contains an outer membrane with a lipopoly-sacharide layer which is impermeable to certain antibiotics and antibacterial compounds (Fennell et al., 2004).

Phytochemical analysis of the extracts of *H. africana* revealed the presence of alkaloids, saponins, tannins, flavonoids and steroids (Table 4). These phytochemical compounds are known to be biologically active and thus aid the antimicrobial activities of plants. Alkaloids were one of the phytochemical compounds identified in this study. Most common biological properties of alkaloids are toxic against cells of foreign organisms, anti-inflammatory, anti-asthmatic, and anti-anaphylactic properties (Ganguly and Sainis, 2001; Staerk et al., 2002) and may be responsible for the observed activity.

The presence of flavonoids in crude extract of *H. africana* is important since they have been reported to exhibit antimicrobial, anti-inflammatory, anti-angionic, analgesic, anti-allergic, cytostatic, antioxidant, antitrypanosomal and antileishmanial properties (Ferguson, 2001; Hodek et al., 2002). Flavonoids in human diet may reduce the risk of various cancers, coronary heart diseases as well as preventing menopausal symptoms (Xu et al., 2000; Hodek et al., 2002; Tasdemir et al., 2006). Saponins and tannins were also reported in this study. Saponins are responsible for numerous pharmacological properties and are known to produce inhibitory effects on inflammation (Estrada et al., 2000). Tannins exert antimicrobial activities by iron deprivation, hydrogen bonding or specific interactions with vital proteins such as enzymes in microbial cells (Njume et al., 2009). Herbs that have tannins are astringent in nature and are used for treating intestinal disorders such as diarrhoea and dysentery (Dharmananda, 2003). Motar et al. (1985) revealed the importance of tannins for the treatment of inflamed or ulcerated tissues. Tannins were observed to have remarkable activity in cancer prevention (Li et al., 2003). Steroidal compounds were also present in the crude extracts of *H. africana*; they have drawn much interest in pharmacy due to their relationship with such compounds as sex hormones (Okwu, 2001). The findings of our study demonstrated the *in vitro* activities of the crude extracts of *H. africana* and provide preliminary evidence for the use of this plant in traditional medicine. This plant may provide new lead molecules, which could become starting materials for the synthesis of new drugs. Further study using bioassay-guided fractionation is necessary to isolate and characterize active compounds of the plant.

ACKNOWLEDGEMENTS

The authors are grateful to the National Research Foundation, South Africa (grant reference CSUR2008052900010), and the Govan Mbeki Research and Development Centre, University of Fort Hare, South Africa for funding this study. They are also grateful to Dr Samie A and Mr. Tshikhawe P for technical assistance.

REFERENCES

- Akinpelu DA, Olayinka AA, Okoh AI (2009). The bioactive potentials of two medicinal plants commonly used as folklore remedies among some tribes in West Africa. *Afr. J. Biotechnol.*, 8: 1660-1664.
- Amagase H (2006). Clarifying the real bioactive constituents of garlic. *J. Nutr.*, 136: 716-725.
- Asfaw D, Abebe D, Urga K (1999). Traditional medicine in Ethiopia: perspectives and developmental efforts. *J. Ethiop. Med. Practice*, 1(2): 114-17.
- Banfi E, Scialino G, Monti-Bragadin C (2003). Development of a microdilution method to evaluate Mycobacterium tuberculosis drug susceptibility. *J. Antimicrob. Chemother.*, 52: 796-800
- Bolin JFEM., Musselman LJ (2009). Pollination Biology of *Hydnora africana* Thunb. (Hydnoraceae) in Namibia: Brood_Site Mimicry with Insect Imprisonment. *Int. J. Plant Sci.*, 170(2): 157-163.
- Büssing A (1996). Induction of apoptosis by the mistletoe lectins: A review on the mechanisms of cytotoxicity mediated by *Viscum album* L. *Apoptosis*, 1: 25-32
- Cheesbrough M (1982). Medical laboratory Manual for tropical countries. Microbiology English language book service (ELBS), 11: 283-378.
- Clinical Laboratory Standards Institute (2008). Performance standards for antimicrobial susceptibility testing; disc diffusion supplemental tables. 28: M100-518, Wayne, PA.
- Cosgrove SE, Vigliani GA, Campion M (2009). Initial low-dose gentamicin for *Staphylococcus aureus* bacteremia and endocarditis is nephrotoxic. *Clin. Infect. Dis.*, 48(6): 713-731.
- Cowan ST, Steel KJ (2004). *In* Manual for identification of Medical Bacteria. 3rd Edition Cambridge pp. 146-157.
- Dastouri MR, Fakhirzadeh K, Shayeg J, Dolgari-Sharaf J, Valilou MR, Maheri-Sis N (2008). Evaluating antibacterial activity of the Iranian honey through MIC methanol on some dermal and intestinal pathogenic bacteria. *J. Ani. Vet. Adv.*, 7(4): 409-412.
- De N, Ifeoma E (2002). Antimicrobial effects of components of the bark extracts of neem (*Azadirachta indica* A. Juss). *Technol. Dev.*, 8: 23-26.
- Dharmananda S (2003). Gallnuts and the uses of Tannins in Chinese Medicine. In: Proceedings of institute for Traditional Medicine, Portland, Oregon pp. 89-105.
- Duraipandiyan V, Ayyanar M, Ignacimuthu S (2006). Antimicrobial activity of some ethnomedicinal plants used by Paliyar tribe from Tamil Nadu, India. *BMC Comple. Alter Med.*, 6: 35-41.
- EI-Mahmood AM, Amey JM (2007). *In vitro* antibacterial activity of *Parkia biglobosa* (Jacq) root bark extract against some microorganisms associated with urinary infections. *Afr. J. Biotechnol.*, 6(11): 1272-1275.
- Estrada A, Katselis GS, Laarveid B, Barl B (2000). Isolation and evaluation of immunological adjuvant activities of saponins from *Polygala senega* L. Comparative immunology. *Microbiol. Infect. Dis.*, 23: 27-43.
- Fabry W, Okemo P, Ansorg R (1996). Activity of East African medicinal plants against *Helicobacter pylori*. *Chemother*, 42: 315-317.
- Fennell CW, Lindsey KL, McGaw LJ, Sparg SG, Stafford, G.I., Elgorashi, E.E., Grace, O.M., van Staden, J. (2004). Assessing African medicinal plants for efficacy and safety: pharmacological screening and toxicology. *J. Ethnopharmacol.*, 94: 205-217.
- Ferguson LR (2001). Role of plant polyphenols in genomic stability. *Mut Res.*, 475: 89-111.

- Ganguly T, Sainis KB (2001). Inhibition of cellular immune response by *Tylophora indica* In experimental models. *Phytomedical*, 8(5): 348-355.
- Hodek P, Trefil P, Stiborova M (2002). Flavonoids-Potent and versatile biologically active compounds interacting with cytochrome P450. *Chemico-Biol. Int.*, 139(1): 1-21.
- Karaman I, Sahin F, Gulluce M, Ogutcu H, Sengul M, Adiguzel A (2003). Antimicrobial activity of aqueous and methanol extracts of *Juniperus oxycedrus* J. *Ethnopharmacol.*, 85: 231-235.
- Kluytmans J, van Belkum A, Verbrugh H (1997). Nasal carriage of *Staphylococcus aureus*: epidemiology, underlying mechanisms, and associated risk. *Clin. Microbiol. Rev.*, 10(3): 505-520.
- Koneman WE, Allen DS, Janda MW, Scherchenberger CP, Win WC Jr. (1992). Antimicrobial susceptibility testing. Textbook of diagnostic microbiology 4th edition. JB Lippincott company pp. 624, 629- 637.
- Li H, Wang Z, Liu Y (2003). Review in the studies on tannins activity of cancer prevention and anticancer. *Zhong-Yao-Cai*, 26(6): 444-448.
- Majhenic L, Kerget MS, Knez Z (2007). Antioxidant and antimicrobial activity of guarana seed extracts. *Food Chem.*, 104: 1258-1268.
- Moniharapon E, Hashinag AF (2004). Antimicrobial activity of Atung (*Parinari glaberrimum* Hassk) fruit extract. *Pak. J. Biol. Sci.*, 7(6): 1057-1061.
- Motar MLR, Thomas G, BarbosaFillo JM (1985). Effects of *Anacardium occidentale* stem bark extract on *in vivo* inflammatory models. *J. Ethnopharmacol.*, 95(2-3): 139-142.
- Musselman LJ (1991). The genus *Hydnora* (*Hydnoraceae*). In Ransom J K, Musselman LJ, Worsham AD, Parker C, editors Proceedings of the fifth international symposium on parasitic weeds, CIMMYT, Nairobi, Kenya. pp. 247-250.
- Nariman F, Eftekhari F, Habibi Z, Falsafi F (2004). Anti-*Helicobacter pylori* activities of six Iranian plants, *Helicobacter*, 9: 146–151.
- Ndip RN, Ajongfac AN, Mbulah SM, Tanih NF, Akoachere J-F TK, Ndip LM, Luma HN, Wirmum C, Ngwa F, Efange SMN (2007). *In vitro* anti-*Helicobacter pylori* activity of *Lycopodium cernuum* (Linn) Pic. *Serm. Afri. J. Biotechnol.*, 7(22): 3989-3994.
- Ndip RN, Malange TAE, Mbulah SM, Lumab HN, Malongue A, Ndip LM, Nyongbela K, Wirmum C, Efange SMN (2008). *In vitro* anti-*Helicobacter pylori* activity of extracts of selected medicinal plants from North West Cameroon. *J. Ethnopharmacol.*, 114: 452–457
- Neely AN, Maley MP (2000). Survival of enterococci and staphylococci on hospital fabrics and plastics. *J. Clin. Microbiol.*, 38(2): 724-726.
- Njume C, Afolayan AJ, Clarke AM, Ndip RN (2010). Crude ethanolic extracts of *Garcinia Kola* seeds Heckel (Guttiferae) prolong the lag phase of *Helicobacter pylori*: inhibitory and bacterial potential. *J. Med. Foods*, 14(7-8): 822-827.
- Njume CJ, Afolayan AJ, Ndip RN (2009). An overview of antimicrobial resistance and the future of medicinal plants in the treatment of *Helicobacter pylori* infections. *Afri. J. Pharm. Pharmacol.*, (3)13: 685-699.
- Nkwelang G, Akoachere J-F TK, Kamga LH, Nfoncham ED, Ndip RN (2009). *Staphylococcus aureus* isolates from clinical and environmental samples in a semi-rural area of Cameroon. Phenotypic characterization of isolates. *Afr. J. Microbiol. Res.*, 3(11): 731-736.
- Nostro A, Germano MP, D'Angelo V, Marino A, Cannatelli MA (2000). Extraction methods and bioautography for evaluation of medicinal plant antimicrobial activity. *Lett. Appl. Microbiol.*, 30(5): 379.
- Okeleye BI, Samie A, Bessong PO, Mkwetshana NF, Green E, Clarke AM, Ndip RN (2010). Crude Ethyl Acetate Extract of the Stem Bark of *Peltophorum africanum* (Sond, Fabaceae) possesses *in vitro* inhibitory and bactericidal activity against clinical isolates of *Helicobacter pylori*. *J. Med. Plants Res.*, 4(14): 1432 - 1440
- Okwu DE (2001). Evaluation of the chemical composition of indigenous species flavouring agent. *Global J. Pure Appl. Sci.*, 39: 69-72.
- Panda SK, Thatoi HN, Dutta SK (2009). Antibacterial activity and phytochemical screening of leaf and bark extracts of *Vitex negundo* L. from simlipal biosphere reserve, Orissa. *J. Med. Plants Res.*, 3(4): 299-300.
- Parekh J, Jadeja D, Chanda S (2005). Efficacy of aqueous and methanol extracts of some medicinal plants for potential antibacterial activity. *Turk. J. Biol.*, 29: 203-210.
- Roy J, Shakaya DM, Callery PS, Thomas JG (2006). Chemical constituents and antimicrobial activity of a traditional herbal medicine containing garlic and black cumin. *Afri. J. Trad. Compl. Alter. Med.*, 3 (20): 1-7.
- Saadabi AMA, Ayoub SMH (2009). Comparative bioactivity of hydnora abyssinica A.Braun against different groups of fungi and bacteria. *J. Med. Plants Res.*, 3(4): 262-265.
- Smith SI, Oyediji KS, Arigbabu AO, Atimomo C, Coker AO (2001). High amoxicillin resistance in *Helicobacter pylori* from gastritis and peptic ulcer patients in Western Nigeria. *J. Gastroenterol.*, 36: 67-68.
- Staerk D, Lykkeberg AK, Christensen J, Budnik BA, Abe F, Jaroszewski JW (2002). *In vitro* Cytotoxic activity of phenanthroindolizidine alkaloids from *Cynanchum vincetoxicum* and *Tylophora tanaka* against drug-sensitive and multidrug-resistant cancer cells. *J. Nat. Prod.*, 65(9): 1299-1302.
- Tanih NF, Okeleye BI, Naidoo N, Clarke AM, Mkwetshana N, Green E, Ndip RN (2010). Marked susceptibility of South African *Helicobacter pylori* strains to ciprofloxacin and amoxicillin: Clinical implications. *S. Afri. Med. J.*, 100: 49-52.
- Tasdemir D, Kaiser M, Brun R, Yardley V, Schmidt TJ, Tosun F, Reudi P (2006). Antitrypanosomal and antileishmanial activities of flavonoids and their analogues: *In vitro*, *in vivo*, structure-activity relationship, and quantitative structure-activity relationship studies. *Antimicrob. Agents Chemother.*, 50(4): 1352-1364.
- Van Wyk B-E, Gericke N (2000). People's plants. A guide to useful plants of southern Africa. Briza Publications, Pretoria pp 34-37.
- Villari P, Crispino M, Montuori P, Boccia S (2003). Molecular typing of *Aeromonas* isolates in natural mineral waters. *Appl. Environ. Microbiol.*, 69: 697-701.
- Wang EW, Warren DK, Ferris VM, Casabar E, Nussenbaum B (2011). Leech – transmitted ciprofloxacin-resistant *Aeromonas hydrophila*. *Arch. Otolaryngol, Head Neck Surg.*, 137(2): 190-193.
- Xu HX, Wan M, Dong H, But PPH, Foo LY (2000). Inhibitory activity of Flavonoids and tannins against HIV-1 protease. *Biol. Pharmacol. Bull.*, 23: 1072-1076.