

# International Journal of Plant Physiology and Biochemistry

Volume 6 Number 2, February 2014

ISSN 2141-2162



*Academic  
Journals*

## ABOUT IJPPB

The **International Journal of Plant Physiology and Biochemistry (IJPPB)** (ISSN 2141-2162) is published Monthly (one volume per year) by AcademicJournals.

**International Journal of Plant Physiology and Biochemistry (IJPPB)** provides rapid publication (monthly) of articles in all areas of the subject such as plant hormones, seed biology, plant DNA repair, Concepts of target cells in plant differentiation etc.

The Journal welcomes the submission of manuscripts that meet the general criteria of significance and scientific excellence. Papers will be published shortly after acceptance. All articles published in IJPPB are peer-reviewed.

## Submission of Manuscript

Please read the **Instructions for Authors** before submitting your manuscript. The manuscript files should be given the last name of the first author

[Click here to Submit manuscripts online](#)

If you have any difficulty using the online submission system, kindly submit via this email [ijppb@academicjournals.org](mailto:ijppb@academicjournals.org).

With questions or concerns, please contact the Editorial Office at [ijppb@academicjournals.org](mailto:ijppb@academicjournals.org).

## Editors

**Prof Dr. Ishrak Khafagi**

*Faculty of Science, Suez Canal University,  
Ismailia, Egypt*

**Prof. Mohamed Mahgoub Azooz**

*Biology Department-Faculty of Science- King Faisal  
University, Saudi Arabia.*

**Dr. Bansal Parveen**

*National Institute of Ayurvedic Pharmaceutical  
Research  
Moti Bagh Road, Patiala-(Punjab) India.*

**Prof. Bechan Sharma**

*University of Allahabad  
Department of Biochemistry, University of  
Allahabad, Faculty of Science, Allahabad-211002,  
India.*

## Editor Board

**Prof. Weimin Zhang**

*Guangdong Institute of Microbiology  
100 Central Xian lie Road, Guangzhou, Guangdong  
510070, China.*

**Dr. Xu Suxia**

*Fujian Institute of Subtropical Botany,  
780-800, Jiahe Road, Xiamen, China361006, China.*

**Dr. Adaku Vivien Iwueke**

*Federal University of Technology, Owerri  
Department of Biochemistry, Federal University of  
Technology, Owerri  
Nigeria.*

**Ass. Prof. Turgay CELIK**

*Gulhane Military Medical Academy,  
School of Medicine, Department of Cardiology,  
Turkey.*

**Dr. Topik Hidayat**

*Department of Biology Education  
Indonesia University of Education (UPI)  
Jalan Dr. Setiabudhi 229 Bandung 40154 Indonesia  
Indonesia.*

**Dr. Tariq Mahmood**

*Quaid-i-Azam University,  
Department of Plant Sciences, Quaid-i-Azam  
University, Islamabad, Pakistan.*

**Dr. Neveen B. Talaat**

*Department of Plant Physiology, Faculty of  
Agriculture, Cairo University, Egypt.*

**Dr. Sudhamoy Mandal**

*Fulbright Visiting Fellow  
Department of Plant Pathology  
University of Nebraska Lincoln  
USA*

**Asso. Prof. Chankova Stephka Georgieva**

*Central Laboratory of General Ecology, Bulg Acad  
Sci 1113 Sofia, 2 Gagarin str, Bulgaria.*

**Shijie Han**

*Center of Forestry, Institute of Applied Ecology,  
Chinese Academy of Sciences,  
72 Wenhua Road, Shenyang City, Liaoning  
Province 110016, PR China.*

**Szu-Chuan Shen**

*Department of Medical Nutrition,  
I-Shou University  
Yanchao Township,  
Kaohsiung County 824, Taiwan.*

**SEDDIK KHENNOUF**

*Dept of BIOLOGY, Faculty of Science  
University Ferhat Abbas, SETIF, 19000, ALGERIA.*

**Saranyu Khammuang**

*Maharakham University  
Department of Chemistry, Faculty of Science  
Thailand.*

**Samir Ranjan Sikdar**

*Bose Institute  
P-1/12, C.I.T. Scheme VII M, Kolkata 700 054,  
India.*

**Dr. M. Abdul Salam**

*Department of Agronomy, College of Agriculture  
Kerala Agricultural University, Vellayani 695 522,  
Trivandrum, Kerala.*

**Dr.Saeed Aminzadeh**

*National Institute of Genetic Engineering and  
Biotechnology (NIGEB)  
Shahrak-e-Pajooresh Km 15, Tehran-Karaj  
Highway, Tehran, I.R.Iran.*

**Dr.Ruzica Stricevic**

*Faculty of Agriculture-University of Belgrade  
Nemanjina 6, Zemun, 11080, Serbia.*

**Rumbos Christos**

*University of Thessaly  
Fytokoy Str, 384 46  
Volos, Greece.*

**Dr. Özge Zencir**

*Kemah Vocational Training School,  
Erzincan University, Kemah,  
Erzincan, Turkey.*

**RIYAZALI ZAFARALI SAYYED**

*SI P Arts, GBP Science & STSKVS Comm. College  
SHAHADA Dist Nandurbar, Maharashtra INDIA.*

**Raul Rodriguez-Herrera**

*Universidad Autonoma de Coahuila  
School of Chemistry  
Blvd. V. Carranza y González Lobo s/n Col República  
Saltillo Coahuila México.*

**Dr. A. H. M. Mahbubur Rahman**

*Rajshahi University, Bangladesh.  
Department of Botany, University of Rajshahi,  
Bangladesh.*

**Paul S. Marley**

*Department of Crop Protection, IAR/FOA  
Ahmadu Bello University, P.M.B. 1044, Samaru,  
Zaria, Nigeria.*

**Patrick Addo-Fordjour**

*Kwame Nkrumah University Of Science And  
Technology (Knust), Kumasi,  
Department Of Theoretical And Applied Biology,  
Ghana.*

**Battu.Prasanna Reddy**

*Nosch Labs Pvt Ltd  
Hyderabad, India.*

**Noureddine Benkeblia**

*UWI - Department of Life Sciences  
Mona Campus, Kingston 7, Jamaica.*

**Keutgen, Norbert**

*Uniwersytet Technologiczno-Przyrodniczy  
im. Jana i Jędrzeja Śniadeckich w Bydgoszczy  
Kadra Katedry Fizjologii Roslin (Institute of Plant  
Physiology)  
ul. Bernardynska 6/8, 85-029 Bydgoszcz, Poland.*

**Nicholas E. Korres**

*University College Cork, Environmental Research  
Institute.  
Lee Road, Cork, Ireland.*

**Dr Naveen Kumar**

*University of Florida  
2685 SR 29 N SWFREC/IFAS/UFL, Immokalee,  
FL34142, USA.*

**Dr Modala Venkateswarlu**

*Seribiotech research Laboratory, Kodathi  
Carmelaram post, Bangalore.*

**Mirza Hasanuzzaman**

*Department of Agronomy, Faculty of Agriculture,  
Sher-e-Bangla Agricultural University, Dhaka-  
1207, Bangladesh.*

**Maybelle Gaballah**

*National Research Centre, El Behoos street, Dokki,  
Cairo.*

**Mauro Guida Santos**

*Universidade Federal de Pernambuco  
Street Moraes Rego – CDU – CCB – Botany  
Department, s/n. 50670-901. Pernambuco State,  
Brazil.*

**Marcelo Rodrigo Pace**

*University of Sao Paulo  
Rua do Matão, 277, Cidade Universitária São  
Paulo, Brazil.*

**Marcelo Francisco Pompelli**

*Federal University of Pernambuco  
Department of Botany, Profª Moraes Rego Av.,  
Recife – PE – Brazil, 50670-901.*

**Luca Catalina Mariana**

*University of Bucharest, Faculty of Biology, Dept of  
Biochemistry and Molecular Biology  
Spl. Independentei, no.91-95, Bucharest 5,  
Romania.*

**Lin Wang**

*Institute of Biostatistics, Fudan University  
220 Handan Road, Shanghai 200433, China  
genetics, microbiology  
China.*

**Li Qiang**

*Institute of karst geology, MLR  
50 Qixing Road,  
China.*

**Dr. Ayanakumar Kumar**

*C.Abdul Hakeem College of Engg. & Tech.,  
Melvisharam-632 509, Vellore Dist, Tamil Nadu,  
INDIA.*

**P.Krishnamoorthy**

*P.G. AND RESEARCH DEPARTMENT OF ZOOLOGY  
RAJAH SERFOJI GOVT. COLLEGE.  
India.*

**Hare Krishna**

*Central Institute of Temperate Horticulture-  
Regional Station,  
Mukteshwar-263 138, District- Nainital,  
Uttarakhand, India.*

**K.G. Mandal**

*Directorate of Water Management (formerly Water  
Technology Centre for Eastern Region)  
Indian Council of Agricultural Research  
C.S. Pur, Bhubaneswar-751023, ORISSA, INDIA.*

**Dr. Jukta Adhikari**

*Presidency College  
86/1, College Street, Kolkata – 700 073, India.*

**Jorge Teixeira**

*Botany Department, Faculty of Sciences,, University  
of Porto,  
Edifício FC4, Rua do Campo Alegre, S/N, 4169-007  
Porto, Portugal.*

**Johnson Toyin Fasinmirin**

*Federal University of Technology, Akure, Nigeria  
Department of Agricultural Engineering, FUT,  
P.M.B. 704, Akure, Ondo State, Nigeria..*

**Joel K. Ransom**

*North Dakota State University  
166 Loftsgard Hall, Department of Plant Sciences,  
NDSU Dept. 7670, PO Box 6050, Fargo, ND 58108-  
6050.*

**João Claudio Damasceno de Sá**

*UENF  
Av. José Carlos Pereira Pinto, 39. Pq. Vicente Dias.  
Campos RJ. Brazil.*

**Jalal Jalali Sendi**

*University of Guilan  
Department of Plant Protection, university of  
Guilan, Rasht, Iran.*



**Lúri Drumond Louro**

*Universidade Federal do Espírito Santo  
Rua Horácio Andrade de Carvalho, 210, Victoria,  
ES, 29052-620, Brazil.*

**Hong Bo Guo**

*Northwest A and F University  
22 Xinong, Yangling 712100, Shaanxi, PR China.*

**Harsukh P. Gajera**

*Junagadh Agricultural University  
Department of Biochemistry, College of  
Agriculture, JAU, Junagadh- 362 001, Gujarat,  
India.*

**Hanan Abdel Fattah El-Sadawy**

*National Research Center  
El-Buhoth St., Dokki, Giza, Egypt.*

**Assist. Prof .Azime KÜÇÜKGÜL GÜLEÇ**

*Tunceli University Fisheries Faculty 62000,  
Tunceli/TURKEY.*

**Greg T. Hannig**

*DuPont  
1090 Elkton Road Newark, DE 19711.*

**Gilberto Santos Andrade**

*Instituto de Biotecnologia Aplicada a  
Agropecuária (BIOAGRO), Departamento de  
Biologia Animal, Universidade Federal de Viçosa,  
Viçosa, MG 36571-000, Brazil.*

**Dr. T. Muthukumar**

*Department of Botany, Bharathiar University  
Coimbatore -641 046, Tamilnadu, India.*

**Kunjupillai Vijayan**

*Institute of Plant and Microbial Biology  
Academia Sinica, Taipei, Taiwan-115, ROC.  
Taiwan.*

**Badre Alam**

*National Research Centre For Agroforestry  
Gwalior Road, Jhansi-284003, U.P., India.*

**Abeer Essam El-Din Mahmoud**

*Biochemistry Department  
Genetic Engineering & Biotechnology Division  
National Research Center El Tahrir St., El Dokki  
12622, Cairo, Egypt.*

**Qazi Fariduddin**

*Aligarh Muslim University  
Department of Botany, Aligarh 202 002, India.*

**Darmawan Darma**

*Faculty of Agriculture, Andalas University  
Kampus Limau Manis Padang-25163, Indonesia.*

**Barbara Chaves**

*Institute for Agricultural and Fisheries Research.*

**Sudhamoy Mandal**

*Central Horticultural Experiment Station (ICAR)  
Aiginia, Bhubaneswar, PIN-751019.*

**Cavit Bircan**

*Adnan Menderes University  
Faculty of Agriculture  
Department of Food Engineering  
09100/Aydin/Turkey.*

**Carlos Alberto Ortega-Ojeda**

*Central University of Ecuador. Faculty of Agriculture  
Sciences. Quito, Ecuador  
Calle 12 # 29 B - 78, Apto. 102 F, Unidad Residencial  
Colseguros, Cali, Colombia.*

**Brian Wade Jamandre**

*National Taiwan University  
Rm. 622, life science bldg., NTU, no.1, sec.4,  
Roosevelt rd. Taipei 10617, Taiwan (ROC).*

**Bitá Naseri**

*Agricultural Research Institute  
Department of Plant Protection, Agricultural  
Research Institute, PO Box 45195474, Zanjan, Iran..*

**Behzad Kaviani**

*Adeyemi Oluyomi Stephen  
Bells University of Technology  
Chemical Sciences Department, Km 8 Ididroko  
Road, Ota, Ogun State,  
Nigeria.*

**Ajayi Adedayo Olajide**

*Adekunle Ajasin University  
Dept. of Microbiology, P.M.B 01, Akungba-Akoko,  
Ondo State, Nigeria.*

**Alexandre Igor Azevedo Pereira (Pereira, A.I.A.)**  
*Universidade Federal de Viçosa, Departamento de Biologia Animal, Programa de Pós-Graduação em Entomologia. 36570-000, Viçosa, Minas Gerais State, Brazil.*

**Gilberto Santos Andrade**  
*Instituto de Biotecnologia Aplicada a Agropecuária (BIOAGRO), Departamento de Biologia Animal, Universidade Federal de Viçosa, Viçosa, MG 36571-000, Brazil.*

**Pradeep. A.R., Ph.D**  
*Seribiotech Research Laboratory  
Carmelaram.P.O; Bangalore, INDIA.*

**Azamal Husen**  
*University of Gondar  
Department of Biology, Faculty of Natural Sciences, University of Gondar  
P.O. Box #196, Gondar, Ethiopia.*

**Muhammad Aslam**  
*University College of Agriculture, Bahauddin Zakariya University  
Multan 60800, Pakistan.*

**Autumn J. Smith**  
*Sam Houston State University, Texas.*

**La Sara**  
*Haluoleo University  
Kampus Baru Tridharma, Kendari,  
Southeast Sulawesi, Indonesia.*

**Aliyu Mohammed**  
*Department of Human Physiology, ABU, Zaria.  
Nigeria.*

**Prof. EL-Said Ahmed AL-Sayed Ragab**  
*National Research Institute of Astronomy and Geophysics, 11421, Helwan, Egypt.*

**Shnoudy Anwar Bakhoum**  
*National Institute of Oceanography & Fisheries (NIOF), Egypt.*

**Antonio Americo Barbosa Viana**  
*Embrapa Recursos Genéticos e Biotecnologia PBI-LPP1  
PqEB Final W/5 Norte, Brasília, DF – Brazil*

**Dr. Shirish Rajmalwar**  
*National Research Laboratory for Conservation, Shirish Rajmalwar, LIG Plot No. 43, Mhada colony, Wardha – 442001, (MS) India.*

**Dr. Amaresh Chandra**  
*Universidade Federal de Viçosa, Departamento de Biologia Animal, Programa de Pós-Graduação em Entomologia. 36570-000, Viçosa, Minas Gerais State, Brazil.*

**Dr. Atul Kumar**  
*GB PANT University of Agriculture & Technology  
Department of Basic Science, College of Forestry & Hill Agriculture, HILL CAMPUS, PO Ranichauri, Tehri Garhwal, Uttarakhand State, India.*

**Prof. Levenko Boris**  
*Natl. Botanical Gardens, NAS of Ukraine  
01014 Kiev, 1 Timiryasevska st. Ukraine.*

**Dr. Dionisio G. Alvindia**  
*Bureau of Postharvest Research and Extension  
CLSU Compound, Science City of Munoz, Nueva Ecija 3120, Philippines.*

**Dr. Bhoopander Giri**  
*University of Delhi  
Department of Botany, SSNC (University of Delhi)  
Alipur, Delhi 110036, India.*

**Dr. Anjali Sood**  
*University of Delhi  
Department of Botany, University of Delhi, Delhi-110 007, INDIA.*

**Dr. A. K. Verma**  
*G.B. Pant University of Agriculture & Technology, Pantnagar, Department of Biochemistry, College of Basic Sciences,  
India.*

**Dr. Anjana Jajoo**  
*School of Life Science, Devi Ahilya University,  
Indore, DAVV  
Khandwa Road campus, Indore 452 017, M.P.,  
India.*

**Dr. Deepak Ganjewala**  
*Vellore Institute of Technology University  
55 Thennaraam Street, Vellore-632 014 (T.N.),  
India.*

**Dr. Geetha Govind**

*Max-Planck-Institute for Chemical Ecology  
Hans-Knöll Straße 8, 07745 Jena, Germany.*

**Dr. Hossam El-Din Saad El-Beltagi**

*Biochemistry Department, Faculty of Agriculture,  
Cairo University, Giza, Egypt, P.O.Box 12613  
Egypt.*

**Prof. Dr. Md. Shahidul Haque**

*Dept. of Biochemistry and Molecular Biology  
University of Rajshahi, Rajshahi-6205, Bangladesh  
Bangladesh.*

**DR. P.K.NAGAR**

*Retired Senior Scientist, IHBT, Palampur, (H.P.),  
B.21/115-10A Batuk Dham Colony, Kamachha,  
Varanasi 221 010, INDIA.*

**Dr. Satyawati Sharma**

*Indian Institute of Technology  
Centre for Rural Development & Technology, IIT  
Delhi-110016  
Biomass Production on waste land, India.*

**Dr. Uğur Çakılcıoğlu**

*Firat University Elazığ/TURKEY  
Cumhuriyet M. Malatya C. No:50/A.*

**Prof. Abdelrhani Elachqar**

*Faculty of Sciences Dhar El Mahraz, Fez, Morocco  
BP 1796, Fès-Atlas, Fès, Maroc, Morocco.*

**Ass. Prof. Jianfeng Xu**

*Arkansas State University  
PO Box 639, State University, AR 72467 USA.*

**Ass.Prof. Jin Xu**

*Center for Agricultural Resources Research,  
Institute of Genetics and Developmental Biology,  
Chinese Academy of Sciences  
Huaizhong RD 286, Shijiazhuang, HeBei, China.*

**José Carlos Rebuglio Velloso Ph.D**

*PARANÁ STATE UNIVERSITY OF PONTA GROSSA  
(Universidade Estadual de Ponta Grossa – UEPG)  
General Carlos Cavalcanti Avenue, 4748,  
Uvaranas, Ponta Grossa/PR – PO box 84030-900*

**Dr. Krouma Abdelmajid**

*Centre of Biotechnology, Borj Cedria Ecopark  
BP 901, Hammam-Lif 2050, Tunisia  
College of Science and Arts, Qassim University, BP 53,  
Al-Rass 3330353, Qassim, Saudi Arabia  
Saudi Arabia*

**Dr. Majid Rostami**

*Malayer University  
Department of Agriculture and Natural Resources,  
Postal code: 65719- 95863, University of Malayer  
Malayer, Iran.*

**Dr. Mohammad Nasir Khan**

*Aligarh Muslim University, Aligarh, INDIA  
Plant Physiology Section, Department of Botany,  
Aligarh Muslim University, Aligarh-202 002, U.P.,  
India.*

**Prof. N.K.Matta**

*Kurukshetra University  
Department of Botany, Kurukshetra University,  
Kurukshetra 136119, INDIA.*

**Dr. Naceur Djebali**

*Centre of Biotechnology Borj-Cedria (CBBC)  
BP 901, Hammam-Lif 2050 Tunisia.*

**Dr. Nader Chaparzadeh**

*Azərbaycan University of Tarbiat Moallem, Tabriz,  
Iran.*

**Nautiyal Prakash Chandra**

*Directorate Of Groundnut Research (Icar)  
Post box, No. 5, Ivnagar Road, Junagadh-362001,  
Gujarat, India.*

**Prof. Hussein Fawzy Hussein Abouzienna**

*National Research Center  
Botany Department, National Research Center,  
Elbhos Street, Dokki, Cairo, Egypt.*

**Dr. D.E. Chandrashekar Rao**

*National Research Council Canada / Plant  
Biotechnology Institute (NRC-PBI)  
110 Gymnasium Place / Saskatoon, Saskatchewan  
S7N 0W9  
Canada.*



**Dr. S.R Madhan Shankar**

*PSG College of Arts & Science  
Civil Aerodrome Post, Coimbatore-641 014, India.*

**Prof. Dr. Safdar Hussain Shah**

*Institute of Biotechnology and Genetic Engineering  
NWFP, Agricultural University Peshawar, Pakistan.*

**Prof. Dr. Md. Shahidul Haque**

*Dept. of Biochemistry and Molecular Biology  
University of Rajshahi, Rajshahi-6205, Bangladesh.*

**Dr. Sivakumar Swaminathan**

*Iowa State University (ISU)  
G-319, Agronomy Department, ISU, Ames, Iowa -  
50011, USA*

**Dr. Subrahmanyam Desiraju**

*Directorate of Rice Research (ICAR)  
Plant Physiology Division, Rajendranagar,  
Hyderabad-500030, A.P. India.*

**Dr. Tariq Aziz Dr. Deepak Ganjewala**

*University of Agriculture, Faisalabad, Sub-Campus  
Depalpur, Dist. Okara, Pakistan.*

**Dr. Thangavel Palaniswamy**

*SUN YAT-SEN UNIVERSITY  
GUANGZHOU, PR CHINA.*

**Yi-Ping Chen Ph.D**

*Institute of Earth Environment,  
Chinese Academy of Science Fenghui S.R, 10,  
Xi'an Hi-Tech Zone, Xi'an, China.*

**Saha Prasenjit**

*The Samuel Roberts Noble Foundation  
2510 Sam Noble Parkway, Ardmore, Ok USA.*

**Abdul Khaliq Ph.D**

*Department of Agronomy  
University of Agriculture  
Faisalabad 38040, Pakistan.*

**Dr. Arafat Abdel Hamed Abdel Latef**

*Assistant Professor of Plant physiology  
Botany Department  
Faculty of Science at Qena  
South Valley University  
Egypt.*

**Dr. Ahmad Bybordi**

*Research Center of Agriculture and Natural  
Resources of East Azarbaijan  
Member of Scientific Board of Research Center of  
Agriculture and Natural Resources of East  
Azarbaijan,  
Tabriz. Iran.*

**Dr. Arijit Sinhababu**

*Bankura Christian College (under –The University of  
Burdwan)  
Department of Botany, Bankura Christian College,  
P.O. + Dist. Bankura, Pin.-722101, West, Bengal,  
India.*

**Dr. Maria Alejandra Equiza**

*University of Alberta,  
4-51 Earth Sciences Building, Dept. Renewable  
Resources, University of Alberta, Edmonton, AB T6G  
2E3,  
Canada.*

**Dr. Suphla Bajpai Gupta**

*Indian Institute of integrative Medicine –CSIR,  
Scientist, Plant biotechnology division, Canal Road,  
Jammu, Jammu & Kashmir, India-180001. India.*

**Dr. Linga R Gutha**

*Washington State University,  
2410 N Bunn Road, Prosser, WA 99350,  
USA.*

**Dr. Medhat Mekhail Tawfik**

*National Research Center,  
El Bohooth Str. Dokki, Giza. Egypt, PO Box 12311,  
Egypt.*

**Dr. Rafiq Islam**

*The Ohio State University South Centers,  
1864 Shyville Road, Piketon, OH 45661.*

**Dr. Rakesh Kumar**

*V.S.P. Govt. P.G. College, Kairana, Muzaffarnagar  
(Uttar Pradesh),  
Department of Botany, V.S.P. Govt. P.G. College,  
Kairana, Muzaffarnagar (Uttar Pradesh), India-  
247774.*

**Dr. Ivan Sestari**

*University of São Paulo,  
Av. Pádua Dias, 11: CP 9. CEP 13418-900.*

**D.Sc. Rachel Fatima Gagliardi**

*State University of Rio de Janeiro,  
Rua São Francisco Xavier, 524 – PHLC sala 602.*

**Dr. Ullas Pedmale**

*Salk Institute for Biological Studies,  
10010 N Torrey Pines RD, La Jolla, CA 92037.*

**Dr. Allah Bakhsh Dr. Deepak Ganjewala**

*Department of Field Crops, Faculty of Agriculture,  
University of Ankara,  
Apartment No. 12/10, Sanatorym Caddesi, Kalaba,  
Kecioren, Ankara,  
Turkey.*

**Dr. Atilgan Atilgan**

*Suleyman Demirel University, Agriculture Faculty,  
Department of Agricultural Structures and  
Irrigation, Isparta,  
Turkey.*

**Mr. Andrej Pilipovic**

*University of Novi Sad – Institute of Lowland  
Forestry and Environment,  
Antona Cehova 13, 21000 Novi Sad,  
Serbi.*

**Dr. Zulfiqar Ahmad Saqib**

*Institute of Soil and Environmental Sciences,  
University of Agriculture, Faisalabad,  
Civil Line Road, Faisalabad,  
Pakistan.*

**MS. C. Mehrnoush Eskandari Torbaghan**

*North Khorasan Agricultural & Natural Resource  
Research Center (NKANRRC)  
P.O. Box: 94155-1416, No. 52, Hassan Kallate  
Alley, Tarbiyat St., Mother Sq. Bojnourd,  
Iran.*

**Dr. Vinod Kumar**

*Department of Zoology & Environmental Science,  
Gurukula Kangri University, Haridwar-249404  
(UK), India.*

**Dr . Panda Tribhubana**

*Kalahandi Institute for Tribiology and  
Ethnobiology(KITE),  
At-Jilingdar, PO-Deydar, Dist-Kalahandi, Odisha,  
India,766014, India*

**Dr. Sabarinath Sundaram**

*Institute of Developmental and Molecular Biology,  
Texas A&M University,  
Biological Sciences Building West Suite 403.*

**Dr. Diogo Pineda Rivelli**

*University of São Paulo,  
Av. Prof. Lineu Prestes 580, São Paulo, SP, 05508-000.*

**Dr. Qiang Wang**

*Virginia Tech,  
427 Latham Hall.*

**Dr. Foteini Hassiotou**

*University of Western Australia,  
35 Stirling Highway, Crawley, WA 6009,  
Australia.*

**Dr. Nivedita Sahu**

*Indian Institute of Chemical Technology,  
Chemical Biology Laboratory  
(NaturalProductChemistry), Uppal Road,  
Hyderabad-500607.*

**Dr. Mohammad Anwar Hossain**

*Bangladesh Agricultural University,  
Assistant Professor, Dept. of Genetics and Plant  
Breeding, Bangladesh Agricultural University,  
Mymensingh-2202, Bangladesh.*

**Dr. Ahmad Ali**

*National Institute of Pharmaceutical Education &  
Research,  
Dept of Biotechnology, NIPER, Jandaha Road,  
Hajipur, Bihar, India, Pin – 844102, India.*

**Mr. Karthikkumar V**

*Annamalai University,  
Department of Biochemistry & Biotechnology.*

**Dr.K.Rajendiran**

*Dept of plant science, Tagore Govt. college,  
9, 4th cross, Tagore Nagar, Pondicherry – 605 008,  
India.*

**Dr. V.Balakrishnan**

*K.S.Rangasamy College of Technology,  
Department of Biotechnology, KSR Kalvi  
nagar, Tiruchengode-637215, Tamilnadu,  
India.*

**Dr. NourAli Sajedi**

*Department of Agronomy and plant Breeding,  
Islamic Azad University, Arak Branch, Arak,  
Iran.*

**(Dr) Ms. Rachel Predeepa**

*Not Applicable ,  
2/387 Gokul Nagar, Kannanenthal Madurai.*

**Dr. Rajendra Gyawali**

*Department of Pharmacy and Biology, Kathmandu  
University,  
Dhulikhel,  
Nepal.*

**Ms. Rocheli de Souza**

*UFRGS,  
Porto Alegre,  
Brazil.*

**Dr. Om Prakash Verma**

*Sam Higginbottom Institute of Agriculture,  
Technology & Sciences (Formerly Allahabad  
Agricultural Institute), Allahabad, U.P.,  
Department of Molecular & Cellular Engineering,  
Jacob School of Biotechnology & Bioengineering,  
India.*

**Dr. Ashwani Kumar**

*JMIT, Radaur,  
Department of Biotechnology, JMIT, Radaur-  
135133, Haryana, India.*

**Dr. Sarfaraz F. A. Al-Bamarny**

*University of Duhok, College of Agriculture, Dept.  
of Horticulture,  
Duhok, Iraqi Kurdistan Region,  
Iraq.*

**Prof. Wafaa Mohamed Shukry Abdel Meamem**

*Dammam University - Saudi Arabia,  
Faculty of Science for Girl.  
Biology Department,  
P.O.Box: 838 Dammam 31113,  
Saudi Arabia.*

**Dr. Stephka G. Chankova**

*Institute of Biodiversity and ecosystem Research,  
BAS,  
2 Gagarin str, 1113 Sofia, Bulgaria.*

**Dr. Nana Ewusi-Mensah**

*Kwame Nkrumah University of Science and  
Technology,  
Dept. of Crop and Soil Sciences, Faculty of  
Agriculture, KNUST, Kumasi.*

**Dr. Mukesh Lokanath Chavan**

*K.r.c. College of horticulture, arabhavi 591 310,  
karnataka,  
University of horticultlura sciences, bagakot,  
India.*

**Dr. Maiti Parthapratim**

*Dept. of Botany Midnapore College, Midnapore-  
721101, Paschim Medinipur, West Bengal, India.*

**Mr. Mohammad Anwar Hossain**

*Kagawa University (Present), Bangladesh  
Agricultural University (Permanent)  
Lab. of Plant Stress Responses, Faculty of  
Agriculture, Kagawa University, Miki-cho, Kita-gun,  
Kagawa 761 0795, Japan.*

**Dr. Antonia Tathiana Batista Dutra**

*Universidade Federal do Ceará,  
Av. Humberto Monte s/n – Pici Bloco 907,  
laboratório 1080.*

**Dr. Kuntal Das**

*St. John's Pharmacy College,  
#6, II Main, 9th Cross, Vijayanagar, Bangalore-104.  
India.*

**Dr. Amitava Rakshit**

*Banaras Hindu University,  
Department of Soil Science & Agril Chemistry.*

**Dr. Kranthi Kiran Mandadi**

*Texas A&M University,  
2132 TAMU, Peterson-Rm408, College Station,  
Texas-77840,  
USA.*

**Dr. Monica Butnariu**

*Banat's University of Agricultural Sciences and  
Veterinary Medicine from Timisoara,  
Chemistry and Vegetal Biochemistry Department,  
Calea Aradului no.119, 300645 Timisoara,  
Romania.*

**Dr. Ahmad Bybordi**

*East Azarbaijan Research Center for Agriculture and Natural Resources, Tabriz, Iran.*

**Dr. Haiwei Gu**

*903 Fifth St., West Lafayette, IN 47906.*

**Dr. Hu Yanbo**

*Northeast Forestry University, 26# Hexing Road, Xiangfang District, Harbin city, 150040, P.R., China.*

**Dr. Arash Kianianmomeni**

*Institute of Biology / Humboldt-University Berlin, Invalidenstr. 42.*

**Dr. Zvonko Pacanoski**

*Faculty for Agriculture Sciences and Food, Boul. Aleksandar Makedonski bb, 1000 Skopje, R.of Macedonia..*

**Dr. Lingjuan Zheng**

*Department of Organismic Biology, University of Salzburg, Hellbrunnenstraße 34, 5020, Salzburg, Austria.*

**Dr. Md. Mokter Hossain**

*Department of Horticulture, Bangladesh Agricultural University, Mymensingh-2202, Bangladesh.*

**Dr. Forouzandeh Soltani**

*Department of Horticultural Sciences, College of Agriculture and Natural Resources, University, of Tehran, Daneshkadeh Street, Karaj 31587-11167, Iran.*

**Dr. M.C.Harish**

*Bharathiar University, Department of Biotechnology, Coimbatore, India.*

**Dr. Zong-shen Zhang**

*School of Biological Engineering, Dalian Polytechnic University, Qinggongyuan, Ganjingzi District, Dalian, China, postcode 116034.*

**Prof. T. V. Ramana Rao**

*B R Doshi School of Biosciences, Sardar Patel University, Vallabh Vidyanagar, Gujarat,India.*

**Dr. Sanjeev Chandel**

*Baba Isher Singh Institute of Sciences & Technology, Gagra (Moga), Punjab, India.*

**Dr. Kuladip Jana**

*Bose Institute Centenary Campus, P 1/12, C.I.T. Scheme VIIM, Kolkata-700 054, India.*

**Prof. Ljubinko Jovanovic**

*University Educons, Faculty for ecological agriculture, Sremska Kamenica, Vojvode Putnika 87, Serbia.*

**Dr. Luis F. Goulao**

*Instituto de Investigacao Cientifica Tropical [Tropical Research Institute] Eco-Bio / IICT, Av. da Republica - Quinta do Marques, 2784-505 Oeiras, Portugal.*

**Dr. Lucky K. Attri**

*College of Punjabi University Patiala, E-41, Sector-14, Panjab University, Chandigarh.*

**Prof. Bassam Taha Yasseen**

*Looking for appropriate employer, Flat 307 Point Red, 146 Midland Road, Luton, LU2 0BL, UK.*

**Dr. Massimo Piccotto**

*Tecna S.r.l., Area Science Park, Loc. Padriciano, 99, I-34149 Trieste, Italy.*

# Instructions for Author

**Electronic submission** of manuscripts is strongly encouraged, provided that the text, tables, and figures are included in a single Microsoft Word file (preferably in Arial font).

The **cover letter** should include the corresponding author's full address and telephone/fax numbers and should be in an e-mail message sent to the Editor, with the file, whose name should begin with the first author's surname, as an attachment.

## Article Types

Three types of manuscripts may be submitted:

**Regular articles:** These should describe new and carefully confirmed findings, and experimental procedures should be given in sufficient detail for others to verify the work. The length of a full paper should be the minimum required to describe and interpret the work clearly.

**Short Communications:** A Short Communication is suitable for recording the results of complete small investigations or giving details of new models or hypotheses, innovative methods, techniques or apparatus. The style of main sections need not conform to that of full-length papers. Short communications are 2 to 4 printed pages (about 6 to 12 manuscript pages) in length.

**Reviews:** Submissions of reviews and perspectives covering topics of current interest are welcome and encouraged. Reviews should be concise and no longer than 4-6 printed pages (about 12 to 18 manuscript pages). Reviews are also peer-reviewed.

## Review Process

All manuscripts are reviewed by an editor and members of the Editorial Board or qualified outside reviewers. Authors cannot nominate reviewers. Only reviewers randomly selected from our database with specialization in the subject area will be contacted to evaluate the manuscripts. The process will be blind review.

Decisions will be made as rapidly as possible, and the journal strives to return reviewers' comments to authors as fast as possible. The editorial board will re-review manuscripts that are accepted pending revision. It is the goal of the AJFS to publish manuscripts within weeks after submission.

## Regular articles

All portions of the manuscript must be typed double-spaced and all pages numbered starting from the title page.

**The Title** should be a brief phrase describing the contents of the paper. The Title Page should include the authors' full names and affiliations, the name of the corresponding author along with phone, fax and E-mail information. Present addresses of authors should appear as a footnote.

**The Abstract** should be informative and completely self-explanatory, briefly present the topic, state the scope of the experiments, indicate significant data, and point out major findings and conclusions. The Abstract should be 100 to 200 words in length. Complete sentences, active verbs, and the third person should be used, and the abstract should be written in the past tense. Standard nomenclature should be used and abbreviations should be avoided. No literature should be cited.

Following the abstract, about 3 to 10 key words that will provide indexing references should be listed.

A list of non-standard **Abbreviations** should be added. In general, non-standard abbreviations should be used only when the full term is very long and used often. Each abbreviation should be spelled out and introduced in parentheses the first time it is used in the text. Only recommended SI units should be used. Authors should use the solidus presentation (mg/ml). Standard abbreviations (such as ATP and DNA) need not be defined.

**The Introduction** should provide a clear statement of the problem, the relevant literature on the subject, and the proposed approach or solution. It should be understandable to colleagues from a broad range of scientific disciplines.

**Materials and methods** should be complete enough to allow experiments to be reproduced. However, only truly new procedures should be described in detail; previously published procedures should be cited, and important modifications of published procedures should be mentioned briefly. Capitalize trade names and include the manufacturer's name and address. Subheadings should be used. Methods in general use need not be described in detail.



**Results** should be presented with clarity and precision. The results should be written in the past tense when describing findings in the authors' experiments. Previously published findings should be written in the present tense. Results should be explained, but largely without referring to the literature. Discussion, speculation and detailed interpretation of data should not be included in the Results but should be put into the Discussion section.

**The Discussion** should interpret the findings in view of the results obtained in this and in past studies on this topic. State the conclusions in a few sentences at the end of the paper. The Results and Discussion sections can include subheadings, and when appropriate, both sections can be combined.

**The Acknowledgments** of people, grants, funds, etc should be brief.

**Tables** should be kept to a minimum and be designed to be as simple as possible. Tables are to be typed double-spaced throughout, including headings and footnotes. Each table should be on a separate page, numbered consecutively in Arabic numerals and supplied with a heading and a legend. Tables should be self-explanatory without reference to the text. The details of the methods used in the experiments should preferably be described in the legend instead of in the text. The same data should not be presented in both table and graph form or repeated in the text.

**Figure legends** should be typed in numerical order on a separate sheet. Graphics should be prepared using applications capable of generating high resolution GIF, TIFF, JPEG or Powerpoint before pasting in the Microsoft Word manuscript file. Tables should be prepared in Microsoft Word. Use Arabic numerals to designate figures and upper case letters for their parts (Figure 1). Begin each legend with a title and include sufficient description so that the figure is understandable without reading the text of the manuscript. Information given in legends should not be repeated in the text.

**References:** In the text, a reference identified by means of an author's name should be followed by the date of the reference in parentheses. When there are more than two authors, only the first author's name should be mentioned, followed by 'et al'. In the event that an author cited has had two or more works published during the same year, the reference, both in the text and in the reference list, should be identified by a lower case letter like 'a' and 'b' after the date to distinguish the works.

Examples:

Abayomi (2000), Agindotan et al. (2003), (Kelebeni, 1983), (Usman and Smith, 1992), (Chege, 1998;

1987a,b; Tijani, 1993,1995), (Kumasi et al., 2001) References should be listed at the end of the paper in alphabetical order. Articles in preparation or articles submitted for publication, unpublished observations, personal communications, etc. should not be included in the reference list but should only be mentioned in the article text (e.g., A. Kingori, University of Nairobi, Kenya, personal communication). Journal names are abbreviated according to Chemical Abstracts. Authors are fully responsible for the accuracy of the references.

Examples:

Chikere CB, Omoni VT and Chikere BO (2008). Distribution of potential nosocomial pathogens in a hospital environment. *Afr. J. Biotechnol.* 7: 3535-3539.

Moran GJ, Amii RN, Abrahamian FM, Talan DA (2005). Methicillinresistant *Staphylococcus aureus* in community-acquired skin infections. *Emerg. Infect. Dis.* 11: 928-930.

Pitout JDD, Church DL, Gregson DB, Chow BL, McCracken M, Mulvey M, Laupland KB (2007). Molecular epidemiology of CTXM-producing *Escherichia coli* in the Calgary Health Region: emergence of CTX-M-15-producing isolates. *Antimicrob. Agents Chemother.* 51: 1281-1286.

Pelczar JR, Harley JP, Klein DA (1993). *Microbiology: Concepts and Applications.* McGraw-Hill Inc., New York, pp. 591-603.

### Short Communications

Short Communications are limited to a maximum of two figures and one table. They should present a complete study that is more limited in scope than is found in full-length papers. The items of manuscript preparation listed above apply to Short Communications with the following differences: (1) Abstracts are limited to 100 words; (2) instead of a separate Materials and Methods section, experimental procedures may be incorporated into Figure Legends and Table footnotes; (3) Results and Discussion should be combined into a single section.

Proofs and Reprints: Electronic proofs will be sent (e-mail attachment) to the corresponding author as a PDF file. Page proofs are considered to be the final version of the manuscript. With the exception of typographical or minor clerical errors, no changes will be made in the manuscript at the proof stage.

**Fees and Charges:** Authors are required to pay a \$550 handling fee. Publication of an article in the International Journal of Plant Physiology and Biochemistry is not contingent upon the author's ability to pay the charges. Neither is acceptance to pay the handling fee a guarantee that the paper will be accepted for publication. Authors may still request (in advance) that the editorial office waive some of the handling fee under special circumstances

**Copyright: © 2014, Academic Journals.**

All rights Reserved. In accessing this journal, you agree that you will access the contents for your own personal use but not for any commercial use. Any use and or copies of this Journal in whole or in part must include the customary bibliographic citation, including author attribution, date and article title.

Submission of a manuscript implies: that the work described has not been published before (except in the form of an abstract or as part of a published lecture, or thesis) that it is not under consideration for publication elsewhere; that if and when the manuscript is accepted for publication, the authors agree to automatic transfer of the copyright to the publisher.

**Disclaimer of Warranties**

In no event shall Academic Journals be liable for any special, incidental, indirect, or consequential damages of any kind arising out of or in connection with the use of the articles or other material derived from the IJPPB, whether or not advised of the possibility of damage, and on any theory of liability.

This publication is provided "as is" without warranty of any kind, either expressed or implied, including, but not limited to, the implied warranties of merchantability, fitness for a particular purpose, or non-infringement. Descriptions of, or references to, products or publications does not imply endorsement of that product or publication. While every effort is made by Academic Journals to see that no inaccurate or misleading data, opinion or statements appear in this publication, they wish to make it clear that the data and opinions appearing in the articles and advertisements herein are the responsibility of the contributor or advertiser concerned. Academic Journals makes no warranty of any kind, either express or implied, regarding the quality, accuracy, availability, or validity of the data or information in this publication or of any other publication to which it may be linked.

**ARTICLES**

**Changes in antioxidants and pungency in response to foliar applications of Mn and Fe in onion**

Manas Denre, A. Bhattacharya, S. Pal, A. Chakravarty, A. Chattopadhyay and D. Mazumdar

Full Length Research Paper

## Changes in antioxidants and pungency in response to foliar applications of Mn and Fe in onion

Manas Denre<sup>1\*</sup>, A. Bhattacharya<sup>1</sup>, S. Pal<sup>1</sup>, A. Chakravarty<sup>1</sup>, A. Chattopadhyay<sup>2</sup> and D. Mazumdar<sup>3</sup>

<sup>1</sup>Department of Agricultural Biochemistry, Bidhan Chandra Krishi Viswavidyalaya, Mohanpur, 741252, Nadia, West Bengal, India.

<sup>2</sup>AICRP on Vegetable Crops, Directorate of Research, Bidhan Chandra Krishi, Viswavidyalaya, Kalyani, Nadia, West Bengal, India.

<sup>3</sup>Department of Agricultural Statistics, Bidhan Chandra Krishi Viswavidyalaya, Mohanpur, 741252, Nadia, West Bengal, India.

Accepted 11 February 2014

The aim of the present work was to study the changes in antioxidants and pungency in response to foliar applications of Mn [manganese: Mn<sub>1</sub>(0%); Mn<sub>2</sub>(0.5%); Mn<sub>3</sub>(1.0%)] and Fe [iron: Fe<sub>1</sub>(0%); Fe<sub>2</sub>(0.5%); Fe<sub>3</sub>(1.0%)] in onion genotype Sukhsagar (*Allium cepa* L.); a field experiment in factorial randomized block design with three replications was conducted in research farm of Bidhan Chandra Krishi Viswavidyalaya, Kalyani, Nadia, West Bengal, India. In this experiment, we observed that the total and free phenol contents were significantly higher than control in both the single (0.5%) and double (1.0%) doses of Fe alone. Again, the activity of enzymatic antioxidant super oxide dismutase (SOD) increased with increasing doses of Fe alone, while the maximum peroxidase (POD) activity was obtained in the treatment combination of double doses (1.0%) of both Fe and Mn. The highest dose of Mn (1%) in combination with all doses of Fe (0, 0.5 and 1.0%) exhibited the highest set of values of catalase (CAT) activity. The activity of 2,2-diphenyl-β-picrylhydrazyl radical scavenging activity (DPPHRAC) assay increased with increasing doses of Fe. The lowest degree of lipid peroxidation was observed in the case of 1% Mn in combination with 1% Fe, which also exhibited the highest pungency. The average values of all the parameters along with the results of PCA, the treatment Mn<sub>3</sub>Fe<sub>3</sub> (the double dose or 1.0% both of Fe and Mn) was found to be the most efficient to ensure the antioxidant properties of the genotype Sukhsagar.

**Key words:** *Allium cepa* L., manganese, iron, antioxidants, pungency.

### INTRODUCTION

From time immemorial, onion (*Allium cepa* L.) has been used as common food and also for the treatment of many diseases. The first citation of this plant is found in the Codex Ebers (1550 BC), an Egyptian medical papyrus reporting several therapeutic formulas based on onions as useful remedy for a variety of diseases such as heart problems, headache, bites, worms and tumors (Block, 1985). Egyptians thought onions aided endurance and

assumed large quantities of them. Raw plants were routinely given to asthmatics and to those people suffering from bronchial-pulmonary complaints. Later on, these food plants were known by Greeks and Romans. Who used them as important healing agents and are still being used by most of the people of the Mediterranean area (Griffiths et al., 2002). Processed onions are applied as a poultice to indolent boils, bruises, wounds, to

\*Corresponding author. E-mail: manas.denre0803@gmail.com.

relieve hot sensations and applied to the navel for dysentery and fever (Brewster and Rabinowitch, 1990). The different clinical studies have shown the benefit of onions in the reduction of the risk of cardiovascular diseases by inducing lowering of serum cholesterol and blood pressure (Steiner and Lin, 1994). They have liver protective (Dion and Milner, 1996), immune enhancement and anti-infection (Lau, 1989), anti-stress and anti-fatigue (Kawashima, 1986), anti-cancer and cancer preventive effects (Balasenthil et al., 2001; Dion and Milner, 1997; Pinto et al., 1997). There are also several other reports for anti-carcinogenic, antibiotic, antibacterial, antifungal and antioxidant properties of onion (Benkeblia, 2005).

Aerobic organisms are constantly exposed to one or more systems that generate reactive oxygen species (ROS), including the superoxide radical anion ( $O_2^{\cdot-}$ ), hydroxyl radical ( $\cdot OH$ ), hydrogen peroxide ( $H_2O_2$ ), various peroxy radicals ( $ROO^{\cdot}$ ), and singlet oxygen ( $^1O_2$ ). These ROS are highly reactive and can damage living cells if formed in significant amounts. To avoid cellular damage by ROS, most biological systems have developed defense mechanisms, antioxidants that convert ROS to unreactive derivatives. Antioxidants are broadly defined as molecules that, when present at low concentrations, as compared to an oxidizable substrate, significantly delay or prevent oxidation of that substrate (Gutteridge, 1994). These include enzymes, frequently known as preventive antioxidants, that is, superoxide dismutase (SOD), catalase (CAT) and peroxidase (POD), which prevent or inhibit formation of ROS, and chain-breaking antioxidants, that is, vitamin C and phenolics, which scavenge oxygen radicals and break radical chain sequences.

When cells are exposed to an unusually high load of oxidants and free radicals, natural defense mechanisms may not be sufficient for eradication of free radical-induced deleterious effects. These include oxidation of nucleic acids, proteins, lipids and carbohydrates; cell death; tissue injury; and development of disease processes such as atherosclerosis, carcinogenesis, cirrhosis, fibrosis and inflammation, aging and aging-related disorders in humans (Halliwell and Gutteridge, 1989). Supplementation of antioxidants may ameliorate harmful effects of the oxidative processes in organisms (Logani and Davies, 1980).

Onions (*Allium cepa* L.) have an important status among vegetables because of their particular flavour and ability to enhance the flavour of other foods. Many sulfur compounds contribute to flavour synthesis and the development of flavour precursors. Onion flavour arises from the enzymatic decomposition of S-alk(en)yl cysteine sulfoxides (ACSOs) when onion tissue is disrupted. The products of the decomposition process, such as lachrymatory factor and thiosulfinates, present onion's unique flavour (Randle, 1997).

Generally, undamaged onions are odourless but, when onion tissues are cut or disrupted, the enzyme alliinase is

released to hydrolyze the S-alk(en)yl cysteine sulfoxides to produce pyruvate, ammonia and volatile sulfur compounds. The volatile sulfur compounds play the most important role in the flavour chemistry (Lancaster and Boland, 1990). In 1992, Block pointed out that alliin (S-alk(en)yl cysteine sulfoxides) are the major flavour precursors to produce the odour and flavour of the *Allium* species when cells are damaged. Schnug (1993) opined that alliin is the most important sulfur-containing compounds of the secondary metabolism of the genus *Allium*.

Although, fragmentary reports are there on the content of the antioxidants but information on the activity of antioxidative enzymes is lacking. Therefore it is important to have detailed information about the content of non-enzymatic antioxidants and the activity of enzymatic antioxidants in onion. The incidences of micronutrients have received a great deal of importance in crop production during recent years because of the widespread occurrences of their deficiencies from different parts of the country. Researchers from almost all the states in the country have also reported significant responses of many crops to micronutrient fertilization.

Iron is an essential element for plant growth and development, since it is fundamental for the proper functioning of numerous metabolic and enzymatic processes. Some studies estimates indicate that a large number of diverse materials can serve as sources of plant nutrients. The majority of nutrient input to agriculture comes from commercial mineral fertilizers. Mineral fertilizers are considered to play a significant but lesser role in nutrient contribution, leaving aside their beneficial effects on soil physicochemical and biological properties. Foliar feeding is a relatively new and controversial technique of feeding plants by applying liquid fertilizer directly to their leaves.

Iron is the component of iron-sulfur (Fe-S) proteins and also non-heme iron proteins which are required during photosynthesis, respiration and  $N_2$  fixation (Taiz and Zeiger, 1991). It is also reported that, it regulates the chlorophyll biosynthesis (Marschner, 1995). About 80% leaf iron is localized in chloroplast and 60% of iron is involved in regular functioning of electron transport chain (Terry and Abadía, 1986). Thus under iron deficiency, the level of chlorophyll fluorescence (Morales et al., 1991) and photosynthetic activity of the plants are decreased markedly (Iturbe-Ormaeche et al., 1995). Iron is a component of several metalloenzymes, including catalase and peroxidase (Valenzuela et al., 1992).

Enzymatic activities of certain antioxidant enzymes like superoxide dismutase (SOD) and peroxidase (POD) are altered under such conditions. Superoxide dismutase is the major scavenger of superoxide ( $O_2^{\cdot-}$ ) in plant system and its enzymatic reaction leads to formation of  $H_2O_2$  and  $O_2$  (Fridovich, 1995). The hydrogen peroxide ( $H_2O_2$ ) is decomposed in the presence of peroxidase (Minotti and Aust, 1987). Hence, under iron deficient condition normal



physiological and biochemical processes in the plant system are severely hampered.

Divalent manganese ions ( $Mn^{2+}$ ) is converted to  $Mn^{3+}$  or  $Mn^{4+}$  easily, therefore in the plant manganese plays an important role on oxidation and reduction processes, as electron transport in photosynthesis. Moreover manganese acts as an activator of several enzymes (more than 35 different enzymes), which involve oxidation reactions, carboxylation, carbohydrates metabolism, phosphorus reactions and citric acid cycle. Of the most important enzymes, protein-manganese in Photosystem II and superoxide dismutase can be pointed. There is more than 90% of superoxide dismutase in chloroplasts with about 4 to 5% of it in mitochondria (Jackson et al., 1978; Millaleo et al., 2010; Mukhopadhyay and Sharma, 1991; Uehara et al., 1974). Manganese also activates several enzymes of the shikimic acids pathway and subsequent pathways, leading to the biosynthesis of aromatic amino acids, such as tyrosine, various secondary products, such as lignin, flavonoids, as well as indol acetic acid (IAA) (Burnell, 1988; Hughes and Williams, 1988). For example,  $Mn^{2+}$  affects phenylalanine ammonia-lyase (PAL) and stimulates peroxidases required for lignin biosynthesis.

The literature on Mn and Fe fertilization of onion is scarce. Bar-Akiva (1971) reported that the peroxidase and catalase activities have been distinguished by Fe and Mn deficiency in Citrus, as peroxidase activity is lowered by Fe deficiency and increased by Mn deficiency in cucumber and tomato (Valenzuela and Romero, 1968). Therios et al. (2005) also reported that peroxidase levels are positively influenced by the increment of Fe and the decline of Mn, in young and old leaves of citrus. The aim of this present study was to assess the changes in antioxidants (non-enzymatic antioxidants: ascorbic acid, phenolics; enzymatic antioxidants: antioxidant superoxide dismutase (SOD), peroxidase (POD), catalase (CAT); antioxidant activities: 2,2-diphenyl- $\beta$ -picrylhydrazyl radical scavenging activity (DPPHRAC), molybdate reducing antioxidant potential (MRAP), lipid peroxidation (LP) and pungency (PAD) in response to foliar applications of Mn and Fe in onion.

## MATERIALS AND METHODS

### Field experiment

The seedling was grown in nursery beds prepared in a sandy loam soil. The beds were 1.5 m long and 1.0 m wide. Weathered cow dung manure, 4 kg·m<sup>-2</sup>, was mixed into the beds. Beds were drenched with formaldehyde (4%) and covered with polythene sheets for ten days to avoid damping off disease. Seedling was treated with Thiram (3 g·kg<sup>-1</sup> of seed) (Gujarat Pesticides Private Limited, Gandhinagar, Gujarat, India) prior to sowing. Onion seeds of the genotype Sukhsagar the most popular cultivar in West Bengal was obtained from AICRP on Vegetable Crops and were sown at 0.5 cm depth and 5 cm apart and covered with finely sieved leaf mold. After sowing, beds were covered with straw until germination and hand watered regularly. Seedlings were raised as

before and hardened by withholding water 4 days before transplanting.

The main field was of sandy loam soil. Land preparation was done with 3-4 ploughings and beds were prepared. Fertilizer at the rate of 60N-60P-60K kg·ha<sup>-1</sup> was applied before transplanting and incorporated well into the soil. The source of nitrogen (N) was urea, that of phosphate (P) was single superphosphate and potash (K) was muriate of potash. Additional nitrogen fertilizer at the rate of 60 N kg·ha<sup>-1</sup> was applied 21 days after transplanting. The 40 days old seedlings were transplanted on muddy flat surface soil in 5 cm deep furrows with 15 cm spacing between rows and 10 cm within plants in plots. The plot size was 2.25 × 2.0 m. The experiment consisting of nine treatments including control (only water sprayed) were arranged in a factorial randomized block design with three replications of each treatment at the In-check Research Farm, Bidhan Chandra Krishi Viswavidyalaya, Kalyani, Nadia, West Bengal, India. The detail treatments are summarized in Table 1.

Since onion is a shallow rooted crop, supplementary irrigations (12 weekly irrigations providing approximately 5 cm of water were applied with the first being before transplanting) were given. There were four light hoeings within 9 weeks of transplanting. The application of fungicide Chlorothalonil at 2 g·L<sup>-1</sup> with sticker Sandovit at 1 ml·L<sup>-1</sup> was applied four to five times beginning from 15 days after transplanting and at 15 days intervals to control purple blotch disease [*Alternaria porri* (Ellis) Cif.].

Onion bulbs from each plot were harvested at 127-128 days after transplanting (when more than 80% plant leaves were dried). Sample (15-20 pieces of bulbs) from each replication was collected after harvesting. Fresh materials were washed, dried with soft tissue, and chopped with a sharp knife into small pieces before analysis of the content of non-enzymatic antioxidants, activity of enzymatic antioxidants and antioxidant activity under different systems of assay [MRAP, DPPH radical scavenging activity and lipid peroxidation] as well as pyruvic acid development (PAD) which is a measure of pungency.

### Chemical analysis

#### Analysis of ascorbic acid content (AAC)

One gram of finely chopped onion tissue was extracted with 20 ml of 4% oxalic acid following maceration in a pestle and mortar and the material centrifuged for 30 min at 10,000 g. Ascorbic acid content was determined using the dichlorophenol indophenol titration procedure (AOAC).

#### Analysis of total phenol content (TPC) and free phenol content (TPC)

Total phenol was extracted in 50% methanolic 1.2 N HCl and free phenol was extracted in 50% aqueous methanol by boiling one gram of finely chopped tissue for 1.5 h at 80-90°C following the method of Vinson et al. (1998) and subsequent analysis was with the Folin-Ciocalteu reagent using gallic acid as standard.

#### Analysis of super oxide dismutase (SOD) activity

One gram finely chopped bulb tissue extracts to inhibit photochemical reduction of nitroblue tetrazolium (NBT) in riboflavin light NBT system (Beauchamp and Fridovich, 1971) was used to determine super oxide dismutase (SOD) activity. Absorbance at 560 nm was recorded and percentage inhibitory activity was calculated as  $[(A_0 - A_e) / A_0] \times 100$  where  $A_0$  = absorbance without extract and  $A_e$  = absorbance with extract.

**Table 1.** Detail doses of treatments used as foliar spray for onion.

Treatment	Manganese (Mn) (%)	Iron (Fe) (%)
T <sub>1</sub> (Mn <sub>1</sub> Fe <sub>1</sub> )	0.0	0.0
T <sub>2</sub> (Mn <sub>1</sub> Fe <sub>2</sub> )	0.0	0.5
T <sub>3</sub> (Mn <sub>1</sub> Fe <sub>3</sub> )	0.0	1.0
T <sub>4</sub> (Mn <sub>2</sub> Fe <sub>1</sub> )	0.5	0.0
T <sub>5</sub> (Mn <sub>2</sub> Fe <sub>2</sub> )	0.5	0.5
T <sub>6</sub> (Mn <sub>2</sub> Fe <sub>3</sub> )	0.5	1.0
T <sub>7</sub> (Mn <sub>3</sub> Fe <sub>1</sub> )	1.0	0.0
T <sub>8</sub> (Mn <sub>3</sub> Fe <sub>2</sub> )	1.0	0.5
T <sub>9</sub> (Mn <sub>3</sub> Fe <sub>3</sub> )	1.0	1.0

The source of Mn was MnSO<sub>4</sub>.H<sub>2</sub>O and that of Fe was FeSO<sub>4</sub>.7H<sub>2</sub>O. Each spraying was done four times with sticker starting from 30 days after transplanting and subsequent ones at an interval of 10 days during vegetative stage.

#### **Analysis of peroxidase (POD) activity**

One gram of fresh onion tissue from each treatment was macerated in a pestle and mortar and extracted with 10 ml tris-HCL buffer (pH 7.6) to determine POD activity. Triturated samples were centrifuged at 10,000 g for 30 min at 4°C and supernatants were assessed for enzyme activity. The POD was estimated by mixing 0.1 ml chilled enzyme extract with 2.8 ml reaction mixture (0.5% *o*-dianisidine dissolved in methanol, 0.28 ml Na-acetate buffer, 2.4 ml water). The reaction was initiated by adding 0.1 ml H<sub>2</sub>O<sub>2</sub> (30%). Changes in absorbance at 430 nm were monitored at 1 min intervals for up to 3 min. POD activity was measured using *o*-dianisidine as the substrate (Bhattacharya et al., 2010) and expressed as mM *o*-dianisidine oxidized·min<sup>-1</sup>·g<sup>-1</sup> of fresh tissue using the extinction coefficient 1.13 × 10<sup>4</sup>·M<sup>-1</sup>·cm<sup>-1</sup>.

#### **Analysis of catalase (CAT) activity**

Two grams of fresh tissue was macerated in a pre-chilled pestle and mortar and extracted with 10 ml phosphate buffer (pH 7.0) to determine CAT activity. Triturated samples were centrifuged at 10,000 g for 30 min at 4°C and supernatants were assessed for enzyme activity. An assay mixture of 3 ml of phosphate buffer, 2 ml of H<sub>2</sub>O<sub>2</sub>, and 1 ml of enzyme source were pipette into a porcelain crucible and incubated for 1 min at 20°C to estimate CAT activity. The reaction was stopped with addition of 10 ml of 0.7N H<sub>2</sub>SO<sub>4</sub> and the reaction mixture titrated against 0.01 N KMnO<sub>4</sub> to determine residual H<sub>2</sub>O<sub>2</sub> until a faint pink color that persisted for at least 15 s was seen. One unit of CAT activity was defined as amount of enzyme that destroyed 1 μmol of H<sub>2</sub>O<sub>2</sub>·min<sup>-1</sup>·g<sup>-1</sup> of fresh tissue. Changes in activity were measured using the method of Kar and Mishra (1976).

#### **Analysis of total antioxidant activity**

A 0.5 gram freshly chopped onion tissue was extracted by macerating with 10 ml distilled water for estimation of total antioxidant activity. Tubes containing extract and reagent solution (0.6 M sulfuric acid, 28 mM sodium phosphate, and 4 mM ammonium molybdate) were incubated at 95°C for 90 min. After incubation the mixture cooled to room temperature, absorbance of each solution was measured at 695 nm against a blank (Prieto et al., 1999). Antioxidant capacity was expressed as gallic acid

equivalent.

#### **Analysis of 2, 2-diphenyl-β-picrylhydrazyl radical scavenging activity (DPPHRAC)**

The scavenging effect of onion for stable 2,2-diphenyl-β-picrylhydrazyl (DPPH) radical was monitored according to the method of Braca et al. (2001). One gram of fresh, finely chopped onion tissue was extracted with 10 ml distilled water. A 0.2 ml aqueous extract was added to 6 ml of a 0.004% methanolic solution of DPPH. Absorbance at 517 nm was recorded after 30 min and percentage inhibitory activity was calculated as [(A<sub>0</sub> - A<sub>e</sub>)/A<sub>0</sub>] × 100 where A<sub>0</sub> = absorbance without extract and A<sub>e</sub> = absorbance with extract.

#### **Analysis of lipid peroxidation (LP)**

Lipid peroxidation was measured as the amount of malondialdehyde (MDA) produced by the thiobarbituric acid (TBA) reaction as described by Dhindsa et al. (1981). A one gram sample of chopped fresh onion tissue was homogenized in 2 ml of 20% trichloroacetic acid (TCA) followed by centrifugation at 5000 g. A 20% TCA solution containing 0.5% TBA and 4% butylated hydroxytoluene was added to a 1 ml aliquot before heating at 95°C for 30 min and then cooled in an ice bath. The contents were centrifuged at 10,000 g and absorbance read at 532 nm. The concentration of MDA was calculated using an extinction coefficient of 155·mM<sup>-1</sup>·cm<sup>-1</sup>.

#### **Analysis of pyruvic acid development (PAD)**

Pungency of onion bulbs was determined as pyruvic acid (Anthon and Barrett, 2003; Schwimmer and Weston, 1961). Pyruvic acid concentration was determined using the method of Schwimmer and Weston (1961). One gram sample of each treatment was crushed in a pestle and mortar and incubated with 2, 4-dinitrophenylhydrazine and the absorbance read at 420 nm on a spectrophotometer for total pyruvic acid concentration, determined against a sodium pyruvate standard curve.

#### **Statistical analysis**

Data were subjected to ANOVA of a factorial randomized block design and by Duncan's multiple range tests, to determine differences among means. Principal component analysis (PCA), as the method of identifying the factor dimension of the data, was used to summarize the treatment informing in a reduced number of factors for selection of the best performing treatment. Statistical analyses were done using SPSS Professional Statistics ver. 7.5 (SPSS Inc., Irvine, California).

## **RESULTS AND DISCUSSION**

### **Non-enzymatic antioxidants**

#### **Ascorbic acid and phenol**

Ascorbic acid is one of the most important hydrophilic antioxidants that scavenge harmful free radicals and other ROS and also regenerate other antioxidants like tocopherol to its functional state. In the present

**Table 2.** Effect of Mn and Fe on antioxidants content and antioxidant activity together with pyruvic acid development in onion.

Treatment	AAC (mg/100 g)	TPC (mg/100 g GAE)	FPC (mg/100 g GAE)	SOD (IC <sub>50</sub> :mg/ml)	POD (mM o-dianisidine oxidized/g/min)	CAT (units)	MRAP (mg/g GAE)	DPPHRAC (IC <sub>50</sub> : mg/ml)	LP (nmol/ 100 g)	PAD (μmol/g)
<b>Mn</b>										
1	11.11	115.28	73.30a <sup>a</sup>	5.02	0.26b	0.98b	64.22a	15.19a	64.11a	8.13b
2	11.32	121.02	50.31c	5.01	0.24c	0.94b	42.46b	13.13b	66.33a	8.44b
3	12.63	115.87	54.12b	5.49	0.41a	1.04a	41.98b	13.10b	60.78b	10.75a
<b>Fe</b>										
1	11.76	107.55b	52.25c	5.93a	0.26c	0.97	53.60a	16.93a	63.67	8.35b
2	11.76	121.52a	56.87b	4.94b	0.33a	1.00	38.53b	12.92b	63.89	9.25a
3	11.54	123.10a	68.61a	4.65b	0.31b	0.99	56.53a	11.57c	63.67	9.72a
<b>Mn<sub>1</sub></b>										
Fe <sub>1</sub>	10.45	117.81	62.26	5.23	0.11	0.91	56.49	14.67	58.33	7.39
Fe <sub>2</sub>	11.11	112.97	73.06	5.17	0.35	1.03	65.27	16.93	68.67	9.29
Fe <sub>3</sub>	11.76	115.06	84.59	4.64	0.33	0.99	70.88	13.98	65.33	7.70
<b>Mn<sub>2</sub></b>										
Fe <sub>1</sub>	11.11	99.22	35.82	5.95	0.27	0.96	37.67	18.52	66.67	8.10
Fe <sub>2</sub>	12.41	135.05	59.41	4.39	0.28	0.95	26.90	8.52	63.67	9.41
Fe <sub>3</sub>	10.45	128.81	55.71	4.70	0.17	0.92	62.82	12.35	68.67	7.81
<b>Mn<sub>3</sub></b>										
Fe <sub>1</sub>	13.72	105.64	58.67	6.61	0.41	1.05	66.63	17.60	66.00	9.56
Fe <sub>2</sub>	11.76	116.53	38.14	5.24	0.37	1.04	23.41	13.32	59.33	9.04
Fe <sub>3</sub>	12.41	125.42	65.55	4.62	0.44	1.05	35.89	8.38	57.00	13.66
<b>Analysis of variance (F values)</b>										
Mn	NS	NS	102.661**	NS	652.155**	18.220**	17.907**	27.500**	11.792**	41.944**
Fe	NS	10.240**	48.117**	14.065**	109.367**	NS	10.357**	147.843**	NS	9.905**
Mn x Fe	NS	5.854**	31.031**	NS	238.373**	3.659*	11.233**	82.056**	14.054**	22.838**

\*Significant at 5%, \*\*significant at 1%, NS = non significant; <sup>a</sup>Values followed by the same letter in a column are not significant at  $P < 0.05$ , Duncan's multiple range test (DMRT).

experiment, application of Mn and Fe through different treatments could produce no significant

differences in the content of ascorbic acid (Table 2). Therefore, no significant effects could be

observed by application of Mn and Fe on the content of ascorbic acid. Phenols are ubiquitous

phytochemicals that contribute largely to antioxidant potential of any plant. In the present experiment, total (TPC) and free (FPC) phenols were estimated to assess the contents of conjugated and unconjugated (free) phenols in the genotype Sukhsagar of onion. Manganese application alone failed to produce any significant effect in TPC.

On the contrary, the Mn treatment alone produced significant differences among the values of FPC, which might be due to activation of enzymes of the Shikimic acid pathway and subsequent pathways and activates the biosynthesis of flavonoids that are an important class of phenolics (Marschner, 1995). However, application of Fe produced values of both TPC and FPC with significant differences while the FPC in Mn treatment decreased with increasing doses and also with respect to control (Mn<sub>1</sub>), both TPC and FPC increased significantly following Fe application. The highest values of TPC and FPC were exhibited by the treatments Mn<sub>2</sub>Fe<sub>2</sub> and Mn<sub>1</sub>Fe<sub>3</sub> (Table 2) respectively while the treatment Mn<sub>2</sub>Fe<sub>1</sub> exhibited the lowest values of both TPC and FPC.

## Enzymatic antioxidants

### SOD, POD and CAT

SOD are present in all aerobic organisms and convert O<sub>2</sub><sup>•-</sup> to H<sub>2</sub>O<sub>2</sub>, which is ultimately neutralized to water, a safe product, by CAT. Despite SODs have several cofactors like Cu and Zn, Fe and Mn; Mn-SOD is not widely distributed in various families of higher plants (Sandmann and Boger, 1983). In the present experiment, Mn alone showed no significant effect on SOD whereas Fe application resulted in a significantly decreasing IC<sub>50</sub> values indicative of increasing SOD activity with increasing dose of Fe, which seems to be quite likely because SOD has a Fe cofactor.

Regarding POD activity, significant effects were observed in application of both the micronutrients. The highest dose of Mn (Mn<sub>3</sub>) resulted in a pretty higher value of POD activity as compared to control (Mn<sub>1</sub>). However, application of Fe increased the activity at single dose (Fe<sub>2</sub>), and decreased with response to the highest dose (Fe<sub>3</sub>).

The fact that Mn activates POD leading to biosynthesis of lignin is already established (Marschner, 1995). The initial increase of POD activity with application of Fe is also expected because POD has a heme prosthetic group (Marschner, 1995). The maximum POD activity was observed in the case of the treatment Mn<sub>3</sub>Fe<sub>3</sub>.

CAT is another antioxidative enzyme, which has heme prosthetic group. CAT detoxifies the H<sub>2</sub>O<sub>2</sub> produced from the action of SOD on O<sub>2</sub><sup>•-</sup> and also from different other sources. In the present experiment, Mn alone only produced a significant increase of CAT activity at the highest dose. The lowest dose, however, could produce no signifi-

cant differences with the value obtained in the control. Presence of no significant differences between the CAT activity value as effectuated by Fe application, was contrary to the observation expected. The interaction effect also shows that the highest set of values was presented by the treatments Mn<sub>3</sub>Fe<sub>1</sub>, Mn<sub>3</sub>Fe<sub>2</sub> and Mn<sub>3</sub>Fe<sub>3</sub>.

## Antioxidant activities

### MRAP, DPPHRAC and LP

Antioxidant activity was analyzed under different systems of assay. In our experiment, MRAP, DPPHRAC and LP are the three different systems. The MRAP and DPPHRAC yielded significant differences between the treatments, whereas in the case of LP there were significant differences only in Mn treatments alone. Mn doses alone resulted in significantly decreasing values of MRAP with respect to Mn<sub>1</sub>, whereas the Fe<sub>1</sub> and Fe<sub>3</sub> had produced no significant differences in MRAP values, but the single dose of Fe (Fe<sub>2</sub>) resulted in a value lesser than the control (Fe<sub>1</sub>). The highest value of MRAP assay was offered by the combination of Mn<sub>3</sub>Fe<sub>1</sub> and the lowest by Mn<sub>3</sub>Fe<sub>2</sub>.

The DPPHRAC assay produced IC<sub>50</sub> values with no significant difference between the Mn treatments alone but there was significant difference with the control or Mn<sub>1</sub>. In the case of Fe treatments alone the values were gradually decreased significantly indicating increase in radical scavenging activity with increasing doses of Fe. The most efficient treatment combination with respect to DPPHRAC was Mn<sub>3</sub>Fe<sub>3</sub> followed by Mn<sub>2</sub>Fe<sub>2</sub>.

Regarding LP, the treatment Mn<sub>3</sub> only resulted in significantly higher value of MDA concentration with respect to Mn<sub>1</sub>. The Fe treatments could produce no significant effects on LP. With regards to LP the most efficient combination was Mn<sub>3</sub>Fe<sub>3</sub> and the combination with the highest MDA concentration was Mn<sub>2</sub>Fe<sub>3</sub>.


## Pungency

### PAD

The pungency of onion is one of its culinary properties. It is measured by the development of pyruvic acid produced upon rupturing the cells. The single dose of Mn (Mn<sub>2</sub>) and the absence of Mn (Mn<sub>1</sub>) resulted in no significant differences whereas Mn<sub>3</sub> exhibited a significantly high value of PAD. The Fe treatments alone (Fe<sub>2</sub> and Fe<sub>3</sub>) produced no significant differences between them but the values these treatments produced were significantly higher than that of Fe<sub>1</sub>. The highest value of PAD was exhibited by Mn<sub>3</sub>Fe<sub>3</sub> followed by Mn<sub>3</sub> alone.

**Table 3.** Results of principal component analysis (PCA) for effects of Mn and Fe on antioxidants content and antioxidant activity together with pyruvic acid development in onion.

Principal component	Eigen value	Variance (%)	Cumulative variance (%)
Eigen values and variance accounted for (%) by PCA based on correlation matrix			
1	3.50	35.01	35.01
2	3.19	31.91	66.92
3	1.79	17.89	84.81
Variables	PC1	PC2	PC3
Factor loadings due to PCs with Eigen values greater than 1			
Ascorbic acid	0.60	0.59	0.13
Total phenol (GAE)	0.47	-0.80	0.17
Free phenol (GAE)	0.09	-0.15	0.94
SOD	-0.36	0.83	-0.28
POD	0.69	0.66	0.17
CAT	0.60	0.72	0.17
MRAP (GAE)	-0.54	0.18	0.77
DPPHRAC	-0.73	0.66	-0.05
LP	-0.59	0.22	0.38
PAD	0.88	0.18	0.04



**Figure 1.** Scatter diagram of the regression factor scores for the first and second components produced by PCA.

### Principal component analysis

Principal component analysis was used to obtain a simplified view of the relationship between the variables and the PCA component loadings for the first three factors are presented in Table 3. These factors were


chosen because the Eigen values were greater than 1.0, which explained 84.81% of total variance.

The first component (PC1) alone explained 35% of total variance in which AAC, TPC, FPC, POD, CAT and PAD restrict SOD, MRAP, DPPHRAC and LP. The negative values of SOD, DPPHRAC and LP are desirable. So, on the basis of PC1 the treatment  $Mn_3Fe_3$ , followed by  $Mn_2Fe_2$ ,  $Mn_3Fe_2$ , and  $Mn_3Fe_1$  can be selected as having all the desirable characters apart from MRAP.

The second component (PC2) explained an additional 32% of total variance in which all the variables except phenols assumed positive values. Therefore, AAC, POD, CAT, MRAP, SOD, DPPHRAC and LP and PAD are increased along the Y axis in the scatterplot (Figure 1) while TPC, FPC are decreased. Here also, the increase in  $IC_{50}$  values of SOD and DPPHRAC and MDA concentration of LP are not desirable. Therefore, on the basis of PC2 the treatments  $Mn_1Fe_2$ ,  $Mn_3Fe_2$ ,  $Mn_3Fe_3$ ,  $Mn_1Fe_3$  and  $Mn_2Fe_2$  can be selected as having the better combinations of all the antioxidants. Considering PC1 and PC2, the treatment  $Mn_3Fe_3$  followed by  $Mn_3Fe_2$  and  $Mn_2Fe_2$  can be selected as performers.

The third component (PC3) explained another 18% of total variance, where all the variables except SOD and DPPHRAC are positively loaded indicating the higher values of all other variables are associated with lower values of SOD and DPPHRAC. This situation is desirable except for LP in which lower values are expected to increase the shelf-life of the crop. Considering PC1 and PC3 (Figure 2), the treatment combination  $Mn_3Fe_3$  followed by  $Mn_3Fe_1$  and  $Mn_2Fe_2$  can be selected to





**Figure 2.** Scatter diagram of the regression factor scores for the first and third components produced by PCA.

ensure all the desirable qualities with respect to anti-oxidant parameters are present in the genotype Sukhsagar following application of manganese and iron.

## Conclusions

From the average values of all the parameters and the results of three principal component analyses, it is suggested that the most efficient treatment was the combination of the double doses of both Mn (1.0%) and Fe (1.0%) to ensure all the desirable antioxidant qualities together with pungency development of onion.

## ACKNOWLEDGEMENT

We would like to thank the Bidhan Chandra Krishi Viswavidyalaya for supporting this work.

## REFERENCES

Anthon GE, Barrett DM (2003). Modified method for the determination of pyruvic acid with dinitrophenylhydrazine in the assessment of onion pungency. *J. Sci. Food Agric.* 83(12):1210-1213.  
 Association of Official Analytical Chemists (1984). Official methods of

analysis. 14<sup>th</sup> ed. Association of Official Analytical Chemists, Washington, D.C.  
 Balasenthil S, Ramachandran CR, Nagini S (2001). Prevention of 4-nitroquinoline 1-oxide-induced rat tongue carcinogenesis by garlic. *Fitoterapia* 72(5):524-531.  
 Bar-Akiva A (1971). Functional aspects of mineral nutrients in use for evaluation of plant nutrient requirement. In: Samish, RM (eds.), *Recent Advances in Plant Nutrition*. Gordon and Breach, New York, 1:115-142.  
 Beauchamp C, Fridovich I (1971). Superoxide dismutase: Improved assays and an assay applicable to acrylamide gels. *Anal. Biochem.* 44(1):276-287.  
 Benkeblia N (2005). Free radical scavenging capacity and antioxidant properties of some selected onion (*Allium cepa* L.) and garlic (*Allium sativum* L.) extracts. *Braz. Arch. Biol. Technol.* 48(5): 753-759.  
 Bhattacharya A, Chattopadhyay A, Muzumdar D, Chakravarty A, Pal S (2010). Antioxidant constituents and enzyme activities in chilli pepper. *Int. J. Veg. Sci.* 16(3):201-211.  
 Block E (1985). The chemistry of garlic and onions. *Sci. Am.* 252(3):114-119.  
 Block E (1992). The organosulfur Chemistry of the genus *Allium*. Implications for the organic chemistry of sulfur. *Angewandte Chemie, International Education*. English. 31:1135-1178.  
 Braca A, Tommasi ND, Bari LD, Pizza C, Politi M, Morelli I (2001). Antioxidant principles from *Bauhinia terapotensis*. *J. Nat. Prod.* 64(7):892-895.  
 Burnell JN (1988). The biochemistry of manganese in plants. In: *Manganese in Soils and Plants*. eds. Graham RD, Hannam RJ, Uren NC, Dordrecht (Netherland), Kluwer Academic, 33:125-137.  
 Dhindsa RS, Dhindsa PP, Thorpe TA (1981). Leaf senescence: correlated with increased level of membrane permeability and lipid

- peroxidation and decreased level of superoxide dismutase and catalase. *J. Exp. Bot.* 32(1):93-101.
- Dion ME, Milner JA (1996). Formation and bioactivation of nitrosomorpholine is inhibited by S-allyl cysteine. *FASEB J.* 10:A498.
- Dion ME, Milner JA (1997). Garlic inhibits cytochrome P450 2E1 mediated chlorzoxazone metabolism. *FASEB J.* 11:A370.
- Fridovich I (1995). Superoxide radical and superoxide dismutases. *Ann. Rev. Biochem.* 64:97-112.
- Griffiths G, Trueman L, Crowther T, Thomas B, Smith B (2002). Onions - a global benefit to health. *Phytother. Res.* 16(7):603-615.
- Gutteridge JMC (1994). Free radicals and Aging. *Rev. Clin. Gerontol.* 4(4):279-288.
- Halliwell B, Gutteridge JMC (1989). Protection against oxidants in biological system: the superoxide theory of oxygen toxicity. In: Halliwell, B and Gutteridge, JM (eds.). *Free Radic. Biol. Med.*, Oxford University Press, Oxford. pp. 86-187.
- Hughes NP, Williams RJP (1988). An introduction to manganese biological chemistry. In: *Manganese in Soils and Plants*. eds. Graham RD, Hannam RJ, Uren NC, Dordrecht (Netherland), Kluwer Academic. 33:7-19.
- Iturbe-Ormaeche I, Moran JF, Arrese-Igor C, Cogorcena Y, Klucas RV, Becana M (1995). Activated oxygen and antioxidant defenses in iron deficient pea plants. *Plant Cell Environ.* 18(4):421-429.
- Jackson C, Dench J, Moore AL, Halliwell B, Foyer CH, Hall DO (1978). Subcellular localisation and identification of superoxide dismutase in the leaves of higher plants. *Eur. J. Biochem.* 91(2):339-44.
- Kar M, Mishra D (1976). Catalase, peroxidase, and polyphenoloxidase activities during rice leaf senescence. *Plant Physiol.* 57(2):315-319.
- Kawashima H (1986). Anti-fatigue effect of Aged Garlic Extract in Athletic Club Students. *Clin. Report*, 20(16):111-127.
- Lancaster JE, Boland MJ (1990). Flavour Biochemistry. In: Rabinowitch, H.D. and Brewster, J. L. (eds.): *Onions and allied crops*. CRC Press, Boca Raton, Ill, 3:33-71
- Lau BHS (1989). Detoxifying Radioprotective and Phagocyte-enhancing Effects of Garlic. *Int. Clin. Nutr. Rev.* 9(1):27-31.
- Logani MK, Davies RE (1980). Lipid oxidation: biologic effects and antioxidants, a review. *Lipids*, 15(6):485-495.
- Marschner H (1995). Functions of mineral nutrients: micronutrients. Iron. In: *Mineral nutrition of higher plants*. Academic Press, Cambridge, UK, pp. 313-324.
- Millaleo R, Reyes-Díaz M, Ivanov AG, Mora ML, Alberdi M (2010). Manganese as essential and toxic element for plants: transport, accumulation and resistance mechanisms. *J. Soil Sci. Plant Nutr.* 10(4):470-481.
- Minotti G, Aust SD (1987). The requirement for iron (III) in the initiation of lipid peroxidation by iron (II) and hydrogen peroxide. *J. Biol. Chem.*
- Morales F, Abadía A, Abadía J (1991). Chlorophyll fluorescence and photon yield of oxygen evolution in iron-deficient sugar beet (*Beta vulgaris* L.) leaves. *Plant Physiol.* 97(3):886-889.262(3):1098-1104.
- Mukhopadhyay MJ, Sharma A (1991). Manganese in cell metabolism of higher plants. *Bot. rev.* 57(2):117-149.
- Panchal SC, Bhatnagar R, Momin RA, Chauhan NP (2001). Capsaicin and ascorbic acid content of chilli as influenced by cultural practices. *Capsicum and Eggplant Newslett.* 20:19-22.
- Pinto JT, Qiao CH, Xing J, Suffoletto BP, Rivlin RS, Heston WDW (1997). Garlic constituents modify expression of biomarkers for human prostatic carcinoma cells. *FASEB J.* 11:A439.
- Prieto P, Pineda M, Aguilar M (1999). Spectrophotometric quantitation of antioxidant capacity through the formation of a phosphomolybdenum complex: Specific application to the determination of vitamin E. *Anal. Biochem.* 269(2):337-341.
- Rabinowitch HD, Brewster JL (1990). *Onions and allied crops*. Publisher: Boca Raton, Fla.: CRC Press, pp. 74-105.
- Randle WM (1997). Onion flavour chemistry and factors influencing flavor intensity. In: RISCH, S. J. and C.-T. HO (eds.): *Spices - Flavour chemistry and antioxidant properties*. ACS-Symposium series, 660:41-52.
- Sandmann G, Boger P (1983). The enzymological function of heavy metals and their role in electron transfer process of plants. In: *encyclopedia of plant physiology new series*. Lauchli A and Bielecki RL (eds.), 15:563-596.
- Schwimmer S, Weston WJ (1961). Enzymatic development of pyruvic acid in onion as a measure of pungency. *J. Agric. Food Chem.* 9(4):301-304.
- Steiner M, Lin RI (1994). Cardiovascular and lipid changes in response to aged garlic extract ingestion. *J. Am. Coll. Nutr.* 13(5):524.
- Taiz L, Zeiger E (1991). *Plant physiology*. Redwood City: The Benjamin/Cummings Publ Co Inc. p. 565.
- Terry N, Abadía J (1986). Function of iron in chloroplasts. *J. Plant Nutri.* 9(3-7):609-646.
- Therios I, Chouliaras V, Bizas D, Boukouvalas S, Palioura E, Hatzidimitriou E, Basioukas D, Gioldasis V, Stavrou N (2005). Changes in Leaf Biochemical and Physiological Indices due to Iron Deficiency in Citrus. *Agro Thesis*, 3(1):18-25.
- Uehara K, Fujimoto S, Taniguchi T, Nakai K (1974). Studies on violet-colored acid phosphatase of sweet potato. II. Enzymatic properties and amino acid composition. *J. Biochem.* 75(3):639-649.
- Valenzuela JL, Romero L (1968). Biochemical indicators and iron index for appraisal of mineral status in leaves of cucumber and tomato. *J. Plant Nutr.* 11:1177-1184.
- Valenzuela JL, Sanchez A, del-Rio A, Lopez-Cantarero I, Romero L (1992). Influence of plant age on mature leaf iron parameters. *J. Plant Nutr.* 15(10):2035-2043.
- Vinson JA, Hap Y, Su X, Zubik L (1998). Phenolic antioxidant quantity and quality in foods: vegetables. *J. Agric. Food Chem.* 46(9):3630-3634.

# International Journal of Plant Physiology and Biochemistry

Related Journals Published by Academic Journals

- *African Journal of Plant Science*
- *Journal of Microbiology and Antimicrobials*
- *International Journal for Biotechnology and Molecular Biology Research*
- *Journal of Botany and Plant Pathology*
- *International Journal of Biodiversity and Conservation*

**academic**Journals