

The background of the cover is a photograph of a tree trunk with a large, dark, irregular hole. The tree bark is brown and textured. The hole is surrounded by lush green foliage, including leaves of various shapes and sizes, some with serrated edges. The overall scene is outdoors, likely in a forest or garden.

International Journal of Plant Physiology and Biochemistry

Volume 6 Number 4, May 2014

ISSN 2006-9871



*Academic
Journals*

ABOUT IJPPB

The **International Journal of Plant Physiology and Biochemistry (IJPPB)** (ISSN 2006-9871) is published Monthly (one volume per year) by AcademicJournals.

International Journal of Plant Physiology and Biochemistry (IJPPB) provides rapid publication (monthly) of articles in all areas of the subject such as plant hormones, seed biology, plant DNA repair, Concepts of target cells in plant differentiation etc.

The Journal welcomes the submission of manuscripts that meet the general criteria of significance and scientific excellence. Papers will be published shortly after acceptance. All articles published in IJPPB are peer-reviewed.

Submission of Manuscript

Please read the **Instructions for Authors** before submitting your manuscript. The manuscript files should be given the last name of the first author

[Click here to Submit manuscripts online](#)

If you have any difficulty using the online submission system, kindly submit via this email ijppb@academicjournals.org.

With questions or concerns, please contact the Editorial Office at ijppb@academicjournals.org.

Editors

Prof Dr. Ishrak Khafagi

*Faculty of Science, Suez Canal University,
Ismailia, Egypt*

Prof. Mohamed Mahgoub Azooz

*Biology Department-Faculty of Science- King Faisal
University, Saudi Arabia.*

Dr. Bansal Parveen

*National Institute of Ayurvedic Pharmaceutical
Research
Moti Bagh Road, Patiala-(Punjab) India.*

Prof. Bechan Sharma

*University of Allahabad
Department of Biochemistry, University of
Allahabad, Faculty of Science, Allahabad-211002,
India.*

Editor Board

Prof. Weimin Zhang

*Guangdong Institute of Microbiology
100 Central Xian lie Road, Guangzhou, Guangdong
510070, China.*

Dr. Xu Suxia

*Fujian Institute of Subtropical Botany,
780-800, Jiahe Road, Xiamen, China361006, China.*

Dr. Adaku Vivien Iwueke

*Federal University of Technology, Owerri
Department of Biochemistry, Federal University of
Technology, Owerri
Nigeria.*

Ass. Prof. Turgay CELIK

*Gulhane Military Medical Academy,
School of Medicine, Department of Cardiology,
Turkey.*

Dr. Topik Hidayat

*Department of Biology Education
Indonesia University of Education (UPI)
Jalan Dr. Setiabudhi 229 Bandung 40154 Indonesia
Indonesia.*

Dr. Tariq Mahmood

*Quaid-i-Azam University,
Department of Plant Sciences, Quaid-i-Azam
University, Islamabad, Pakistan.*

Dr. Neveen B. Talaat

*Department of Plant Physiology, Faculty of
Agriculture, Cairo University, Egypt.*

Dr. Sudhamoy Mandal

*Fulbright Visiting Fellow
Department of Plant Pathology
University of Nebraska Lincoln
USA*

Asso. Prof. Chankova Stephka Georgieva

*Central Laboratory of General Ecology, Bulg Acad
Sci 1113 Sofia, 2 Gagarin str, Bulgaria.*

Shijie Han

*Center of Forestry, Institute of Applied Ecology,
Chinese Academy of Sciences,
72 Wenhua Road, Shenyang City, Liaoning
Province 110016, PR China.*

Szu-Chuan Shen

*Department of Medical Nutrition,
I-Shou University
Yanchao Township,
Kaohsiung County 824, Taiwan.*

SEDDIK KHENNOUF

*Dept of BIOLOGY, Faculty of Science
University Ferhat Abbas, SETIF, 19000, ALGERIA.*

Saranyu Khammuang

*Maharakham University
Department of Chemistry, Faculty of Science
Thailand.*

Samir Ranjan Sikdar

*Bose Institute
P-1/12, C.I.T. Scheme VII M, Kolkata 700 054,
India.*

Dr. M. Abdul Salam

*Department of Agronomy, College of Agriculture
Kerala Agricultural University, Vellayani 695 522,
Trivandrum, Kerala.*

Dr.Saeed Aminzadeh

*National Institute of Genetic Engineering and
Biotechnology (NIGEB)
Shahrak-e-Pajooresh Km 15, Tehran-Karaj
Highway, Tehran, I.R.Iran.*

Dr.Ruzica Stricevic

*Faculty of Agriculture-University of Belgrade
Nemanjina 6, Zemun, 11080, Serbia.*

Rumbos Christos

*University of Thessaly
Fytokoy Str, 384 46
Volos, Greece.*

Dr. Özge Zencir

*Kemah Vocational Training School,
Erzincan University, Kemah,
Erzincan, Turkey.*

RIYAZALI ZAFARALI SAYYED

*SI P Arts, GBP Science & STSKVS Comm. College
SHAHADA Dist Nandurbar, Maharashtra INDIA.*

Raul Rodriguez-Herrera

*Universidad Autonoma de Coahuila
School of Chemistry
Blvd. V. Carranza y González Lobo s/n Col República
Saltillo Coahuila México.*

Dr. A. H. M. Mahbubur Rahman

*Rajshahi University, Bangladesh.
Department of Botany, University of Rajshahi,
Bangladesh.*

Paul S. Marley

*Department of Crop Protection, IAR/FOA
Ahmadu Bello University, P.M.B. 1044, Samaru,
Zaria, Nigeria.*

Patrick Addo-Fordjour

*Kwame Nkrumah University Of Science And
Technology (Knust), Kumasi,
Department Of Theoretical And Applied Biology,
Ghana.*

Battu.Prasanna Reddy

*Nosch Labs Pvt Ltd
Hyderabad, India.*

Noureddine Benkeblia

*UWI - Department of Life Sciences
Mona Campus, Kingston 7, Jamaica.*

Keutgen, Norbert

*Uniwersytet Technologiczno-Przyrodniczy
im. Jana i Jędrzeja Śniadeckich w Bydgoszczy
Kadra Katedry Fizjologii Roslin (Institute of Plant
Physiology)
ul. Bernardynska 6/8, 85-029 Bydgoszcz, Poland.*

Nicholas E. Korres

*University College Cork, Environmental Research
Institute.
Lee Road, Cork, Ireland.*

Dr Naveen Kumar

*University of Florida
2685 SR 29 N SWFREC/IFAS/UFL, Immokalee,
FL34142, USA.*

Dr Modala Venkateswarlu

*Seribiotech research Laboratory, Kodathi
Carmelaram post, Bangalore.*

Mirza Hasanuzzaman

*Department of Agronomy, Faculty of Agriculture,
Sher-e-Bangla Agricultural University, Dhaka-
1207, Bangladesh.*

Maybelle Gaballah

*National Research Centre, El Behoos street, Dokki,
Cairo.*

Mauro Guida Santos

*Universidade Federal de Pernambuco
Street Moraes Rego – CDU – CCB – Botany
Department, s/n. 50670-901. Pernambuco State,
Brazil.*

Marcelo Rodrigo Pace

*University of Sao Paulo
Rua do Matão, 277, Cidade Universitária São
Paulo, Brazil.*

Marcelo Francisco Pompelli

*Federal University of Pernambuco
Department of Botany, Profª Moraes Rego Av.,
Recife – PE – Brazil, 50670-901.*

Luca Catalina Mariana

*University of Bucharest, Faculty of Biology, Dept of
Biochemistry and Molecular Biology
Spl. Independentei, no.91-95, Bucharest 5,
Romania.*

Lin Wang

*Institute of Biostatistics, Fudan University
220 Handan Road, Shanghai 200433, China
genetics, microbiology
China.*

Li Qiang

*Institute of karst geology,MLR
50 Qixing Road,
China.*

Dr. Ayanakumar Kumar

*C.Abdul Hakeem College of Engg. & Tech.,
Melvisharam-632 509, Vellore Dist, Tamil Nadu,
INDIA.*

P.Krishnamoorthy

*P.G. AND RESEARCH DEPARTMENT OF ZOOLOGY
RAJAH SERFOJI GOVT. COLLEGE.
India.*

Hare Krishna

*Central Institute of Temperate Horticulture-
Regional Station,
Mukteshwar-263 138, District- Nainital,
Uttarakhand, India.*

K.G. Mandal

*Directorate of Water Management (formerly Water
Technology Centre for Eastern Region)
Indian Council of Agricultural Research
C.S. Pur, Bhubaneswar-751023, ORISSA, INDIA.*

Dr. Jukta Adhikari

*Presidency College
86/1, College Street, Kolkata – 700 073, India.*

Jorge Teixeira

*Botany Department, Faculty of Sciences,, University
of Porto,
Edifício FC4, Rua do Campo Alegre, S/N, 4169-007
Porto, Portugal.*

Johnson Toyin Fasinmirin

*Federal University of Technology, Akure, Nigeria
Department of Agricultural Engineering, FUT,
P.M.B. 704, Akure, Ondo State, Nigeria..*

Joel K. Ransom

*North Dakota State University
166 Loftsgard Hall, Department of Plant Sciences,
NDSU Dept. 7670, PO Box 6050, Fargo, ND 58108-
6050.*

João Claudio Damasceno de Sá

*UENF
Av. José Carlos Pereira Pinto, 39. Pq. Vicente Dias.
Campos RJ. Brazil.*

Jalal Jalali Sendi

*University of Guilan
Department of Plant Protection, university of
Guilan, Rasht, Iran.*

Lúri Drumond Louro

*Universidade Federal do Espírito Santo
Rua Horácio Andrade de Carvalho, 210, Victoria,
ES, 29052-620, Brazil.*

Hong Bo Guo

*Northwest A and F University
22 Xinong, Yangling 712100, Shaanxi, PR China.*

Harsukh P. Gajera

*Junagadh Agricultural University
Department of Biochemistry, College of
Agriculture, JAU, Junagadh- 362 001, Gujarat,
India.*

Hanan Abdel Fattah El-Sadawy

*National Research Center
El-Buhoth St., Dokki, Giza, Egypt.*

Assist. Prof .Azime KÜÇÜKGÜL GÜLEÇ

*Tunceli University Fisheries Faculty 62000,
Tunceli/TURKEY.*

Greg T. Hannig

*DuPont
1090 Elkton Road Newark, DE 19711.*

Gilberto Santos Andrade

*Instituto de Biotecnologia Aplicada a
Agropecuária (BIOAGRO), Departamento de
Biologia Animal, Universidade Federal de Viçosa,
Viçosa, MG 36571-000, Brazil.*

Dr. T. Muthukumar

*Department of Botany, Bharathiar University
Coimbatore -641 046, Tamilnadu, India.*

Kunjupillai Vijayan

*Institute of Plant and Microbial Biology
Academia Sinica, Taipei, Taiwan-115, ROC.
Taiwan.*

Badre Alam

*National Research Centre For Agroforestry
Gwalior Road, Jhansi-284003, U.P., India.*

Abeer Essam El-Din Mahmoud

*Biochemistry Department
Genetic Engineering & Biotechnology Division
National Research Center El Tahrir St., El Dokki
12622, Cairo, Egypt.*

Qazi Fariduddin

*Aligarh Muslim University
Department of Botany, Aligarh 202 002, India.*

Darmawan Darma

*Faculty of Agriculture, Andalas University
Kampus Limau Manis Padang-25163, Indonesia.*

Barbara Chaves

Institute for Agricultural and Fisheries Research.

Sudhamoy Mandal

*Central Horticultural Experiment Station (ICAR)
Aiginia, Bhubaneswar, PIN-751019.*

Cavit Bircan

*Adnan Menderes University
Faculty of Agriculture
Department of Food Engineering
09100/Aydin/Turkey.*

Carlos Alberto Ortega-Ojeda

*Central University of Ecuador. Faculty of Agriculture
Sciences. Quito, Ecuador
Calle 12 # 29 B - 78, Apto. 102 F, Unidad Residencial
Colseguros, Cali, Colombia.*

Brian Wade Jamandre

*National Taiwan University
Rm. 622, life science bldg., NTU, no.1, sec.4,
Roosevelt rd. Taipei 10617, Taiwan (ROC).*

Bitá Naseri

*Agricultural Research Institute
Department of Plant Protection, Agricultural
Research Institute, PO Box 45195474, Zanjan, Iran..*

Behzad Kaviani

*Adeyemi Oluyomi Stephen
Bells University of Technology
Chemical Sciences Department, Km 8 Ididroko
Road, Ota, Ogun State,
Nigeria.*

Ajayi Adedayo Olajide

*Adekunle Ajasin University
Dept. of Microbiology, P.M.B 01, Akungba-Akoko,
Ondo State, Nigeria.*

Alexandre Igor Azevedo Pereira (Pereira, A.I.A.)
Universidade Federal de Viçosa, Departamento de Biologia Animal, Programa de Pós-Graduação em Entomologia. 36570-000, Viçosa, Minas Gerais State, Brazil.

Gilberto Santos Andrade
Instituto de Biotecnologia Aplicada a Agropecuária (BIOAGRO), Departamento de Biologia Animal, Universidade Federal de Viçosa, Viçosa, MG 36571-000, Brazil.

Pradeep. A.R., Ph.D
*Seribiotech Research Laboratory
Carmelaram.P.O; Bangalore, INDIA.*

Azamal Husen
*University of Gondar
Department of Biology, Faculty of Natural Sciences, University of Gondar
P.O. Box #196, Gondar, Ethiopia.*

Muhammad Aslam
*University College of Agriculture, Bahauddin Zakariya University
Multan 60800, Pakistan.*

Autumn J. Smith
Sam Houston State University, Texas.

La Sara
*Haluoleo University
Kampus Baru Tridharma, Kendari,
Southeast Sulawesi, Indonesia.*

Aliyu Mohammed
*Department of Human Physiology, ABU, Zaria.
Nigeria.*

Prof. EL-Said Ahmed AL-Sayed Ragab
National Research Institute of Astronomy and Geophysics, 11421, Helwan, Egypt.

Shnoudy Anwar Bakhoum
National Institute of Oceanography & Fisheries (NIOF), Egypt.

Antonio Americo Barbosa Viana
*Embrapa Recursos Genéticos e Biotecnologia PBI-LPP1
PqEB Final W/5 Norte, Brasília, DF – Brazil*

Dr. Shirish Rajmalwar
National Research Laboratory for Conservation, Shirish Rajmalwar, LIG Plot No. 43, Mhada colony, Wardha – 442001, (MS) India.

Dr. Amaresh Chandra
Universidade Federal de Viçosa, Departamento de Biologia Animal, Programa de Pós-Graduação em Entomologia. 36570-000, Viçosa, Minas Gerais State, Brazil.

Dr. Atul Kumar
*GB PANT University of Agriculture & Technology
Department of Basic Science, College of Forestry & Hill Agriculture, HILL CAMPUS, PO Ranichauri, Tehri Garhwal, Uttarakhand State, India.*

Prof. Levenko Boris
*Natl. Botanical Gardens, NAS of Ukraine
01014 Kiev, 1 Timiryasevka st. Ukraine.*

Dr. Dionisio G. Alwindia
*Bureau of Postharvest Research and Extension
CLSU Compound, Science City of Munoz, Nueva Ecija 3120, Philippines.*

Dr. Bhoopander Giri
*University of Delhi
Department of Botany, SSNC (University of Delhi)
Alipur, Delhi 110036, India.*

Dr. Anjali Sood
*University of Delhi
Department of Botany, University of Delhi, Delhi-110 007, INDIA.*

Dr. A. K. Verma
*G.B. Pant University of Agriculture & Technology, Pantnagar, Department of Biochemistry, College of Basic Sciences,
India.*

Dr. Anjana Jajoo
*School of Life Science, Devi Ahilya University,
Indore, DAVV
Khandwa Road campus, Indore 452 017, M.P.,
India.*

Dr. Deepak Ganjewala
*Vellore Institute of Technology University
55 Thennaraam Street, Vellore-632 014 (T.N.),
India.*

Dr. Geetha Govind

*Max-Planck-Institute for Chemical Ecology
Hans-Knöll Straße 8, 07745 Jena, Germany.*

Dr. Hossam El-Din Saad El-Beltagi

*Biochemistry Department, Faculty of Agriculture,
Cairo University, Giza, Egypt, P.O.Box 12613
Egypt.*

Prof. Dr. Md. Shahidul Haque

*Dept. of Biochemistry and Molecular Biology
University of Rajshahi, Rajshahi-6205, Bangladesh
Bangladesh.*

DR. P.K.NAGAR

*Retired Senior Scientist, IHBT, Palampur, (H.P.),
B.21/115-10A Batuk Dham Colony, Kamachha,
Varanasi 221 010, INDIA.*

Dr. Satyawati Sharma

*Indian Institute of Technology
Centre for Rural Development & Technology, IIT
Delhi-110016
Biomass Production on waste land, India.*

Dr. Uğur Çakılcıoğlu

*Firat University Elazığ/TURKEY
Cumhuriyet M. Malatya C. No:50/A.*

Prof. Abdelrhani Elachqar

*Faculty of Sciences Dhar El Mahraz, Fez, Morocco
BP 1796, Fès-Atlas, Fès, Maroc, Morocco.*

Ass. Prof. Jianfeng Xu

*Arkansas State University
PO Box 639, State University, AR 72467 USA.*

Ass.Prof. Jin Xu

*Center for Agricultural Resources Research,
Institute of Genetics and Developmental Biology,
Chinese Academy of Sciences
Huaizhong RD 286, Shijiazhuang, HeBei, China.*

José Carlos Rebuglio Velloso Ph.D

*PARANÁ STATE UNIVERSITY OF PONTA GROSSA
(Universidade Estadual de Ponta Grossa – UEPG)
General Carlos Cavalcanti Avenue, 4748,
Uvaranas, Ponta Grossa/PR – PO box 84030-900*

Dr. Krouma Abdelmajid

*Centre of Biotechnology, Borj Cedria Ecopark
BP 901, Hammam-Lif 2050, Tunisia
College of Science and Arts, Qassim University, BP 53,
Al-Rass 3330353, Qassim, Saudi Arabia
Saudi Arabia*

Dr. Majid Rostami

*Malayer University
Department of Agriculture and Natural Resources,
Postal code: 65719- 95863, University of Malayer
Malayer, Iran.*

Dr. Mohammad Nasir Khan

*Aligarh Muslim University, Aligarh, INDIA
Plant Physiology Section, Department of Botany,
Aligarh Muslim University, Aligarh-202 002, U.P.,
India.*

Prof. N.K.Matta

*Kurukshetra University
Department of Botany, Kurukshetra University,
Kurukshetra 136119, INDIA.*

Dr. Naceur Djebali

*Centre of Biotechnology Borj-Cedria (CBBC)
BP 901, Hammam-Lif 2050 Tunisia.*

Dr. Nader Chaparzadeh

*Azarbaijan University of Tarbiat Moallem, Tabriz,
Iran.*

Nautiyal Prakash Chandra

*Directorate Of Groundnut Research (Icar)
Post box, No. 5, Ivnagar Road, Junagadh-362001,
Gujarat, India.*

Prof. Hussein Fawzy Hussein Abouzienna

*National Research Center
Botany Department, National Research Center,
Elbhos Street, Dokki, Cairo, Egypt.*

Dr. D.E. Chandrashekar Rao

*National Research Council Canada / Plant
Biotechnology Institute (NRC-PBI)
110 Gymnasium Place / Saskatoon, Saskatchewan
S7N 0W9
Canada.*

Dr. S.R Madhan Shankar

*PSG College of Arts & Science
Civil Aerodrome Post, Coimbatore-641 014, India.*

Prof. Dr. Safdar Hussain Shah

*Institute of Biotechnology and Genetic Engineering
NWFP, Agricultural University Peshawar, Pakistan.*

Prof. Dr. Md. Shahidul Haque

*Dept. of Biochemistry and Molecular Biology
University of Rajshahi, Rajshahi-6205, Bangladesh.*

Dr. Sivakumar Swaminathan

*Iowa State University (ISU)
G-319, Agronomy Department, ISU, Ames, Iowa -
50011, USA*

Dr. Subrahmanyam Desiraju

*Directorate of Rice Research (ICAR)
Plant Physiology Division, Rajendranagar,
Hyderabad-500030, A.P. India.*

Dr. Tariq Aziz Dr. Deepak Ganjewala

*University of Agriculture, Faisalabad, Sub-Campus
Depalpur, Dist. Okara, Pakistan.*

Dr. Thangavel Palaniswamy

*SUN YAT-SEN UNIVERSITY
GUANGZHOU, PR CHINA.*

Yi-Ping Chen Ph.D

*Institute of Earth Environment,
Chinese Academy of Science Fenghui S.R, 10,
Xi'an Hi-Tech Zone, Xi'an, China.*

Saha Prasenjit

*The Samuel Roberts Noble Foundation
2510 Sam Noble Parkway, Ardmore, Ok USA.*

Abdul Khaliq Ph.D

*Department of Agronomy
University of Agriculture
Faisalabad 38040, Pakistan.*

Dr. Arafat Abdel Hamed Abdel Latef

*Assistant Professor of Plant physiology
Botany Department
Faculty of Science at Qena
South Valley University
Egypt.*

Dr. Ahmad Bybordi

*Research Center of Agriculture and Natural
Resources of East Azarbaijan
Member of Scientific Board of Research Center of
Agriculture and Natural Resources of East
Azarbaijan,
Tabriz. Iran.*

Dr. Arijit Sinhababu

*Bankura Christian College (under –The University of
Burdwan)
Department of Botany, Bankura Christian College,
P.O. + Dist. Bankura, Pin.-722101, West, Bengal,
India.*

Dr. Maria Alejandra Equiza

*University of Alberta,
4-51 Earth Sciences Building, Dept. Renewable
Resources, University of Alberta, Edmonton, AB T6G
2E3,
Canada.*

Dr. Suphla Bajpai Gupta

*Indian Institute of integrative Medicine –CSIR,
Scientist, Plant biotechnology division, Canal Road,
Jammu, Jammu & Kashmir, India-180001. India.*

Dr. Linga R Gutha

*Washington State University,
2410 N Bunn Road, Prosser, WA 99350,
USA.*

Dr. Medhat Mekhail Tawfik

*National Research Center,
El Bohooth Str. Dokki, Giza. Egypt, PO Box 12311,
Egypt.*

Dr. Rafiq Islam

*The Ohio State University South Centers,
1864 Shyville Road, Piketon, OH 45661.*

Dr. Rakesh Kumar

*V.S.P. Govt. P.G. College, Kairana, Muzaffarnagar
(Uttar Pradesh),
Department of Botany, V.S.P. Govt. P.G. College,
Kairana, Muzaffarnagar (Uttar Pradesh), India-
247774.*

Dr. Ivan Sestari

*University of São Paulo,
Av. Pádua Dias, 11: CP 9. CEP 13418-900.*

D.Sc. Rachel Fatima Gagliardi

State University of Rio de Janeiro,
Rua São Francisco Xavier, 524 – PHLC sala 602.

Dr. Ullas Pedmale

Salk Institute for Biological Studies,
10010 N Torrey Pines RD, La Jolla, CA 92037.

Dr. Allah Bakhsh Dr. Deepak Ganjewala

Department of Field Crops, Faculty of Agriculture,
University of Ankara,
Apartment No. 12/10, Sanatorym Caddesi, Kalaba,
Kecioren, Ankara,
Turkey.

Dr. Atilgan Atilgan

Suleyman Demirel University, Agriculture Faculty,
Department of Agricultural Structures and
Irrigation, Isparta,
Turkey.

Mr. Andrej Pilipovic

University of Novi Sad – Institute of Lowland
Forestry and Environment,
Antona Cehova 13, 21000 Novi Sad,
Serbi.

Dr. Zulfiqar Ahmad Saqib

Institute of Soil and Environmental Sciences,
University of Agriculture, Faisalabad,
Civil Line Road, Faisalabad,
Pakistan.

MS. C. Mehrnoush Eskandari Torbaghan

North Khorasan Agricultural & Natural Resource
Research Center (NKANRRC)
P.O. Box: 94155-1416, No. 52, Hassan Kallate
Alley, Tarbiyat St., Mother Sq. Bojnourd,
Iran.

Dr. Vinod Kumar

Department of Zoology & Environmental Science,
Gurukula Kangri University, Haridwar-249404
(UK), India.

Dr . Panda Tribhubana

Kalahandi Institute for Tribiology and
Ethnobiology(KITE),
At-Jilingdar, PO-Deydar, Dist-Kalahandi, Odisha,
India, 766014, India

Dr. Sabarinath Sundaram

Institute of Developmental and Molecular Biology,
Texas A&M University,
Biological Sciences Building West Suite 403.

Dr. Diogo Pineda Rivelli

University of São Paulo,
Av. Prof. Lineu Prestes 580, São Paulo, SP, 05508-000.

Dr. Qiang Wang

Virginia Tech,
427 Latham Hall.

Dr. Foteini Hassiotou

University of Western Australia,
35 Stirling Highway, Crawley, WA 6009,
Australia.

Dr. Nivedita Sahu

Indian Institute of Chemical Technology,
Chemical Biology Laboratory
(NaturalProductChemistry), Uppal Road,
Hyderabad-500607.

Dr. Mohammad Anwar Hossain

Bangladesh Agricultural University,
Assistant Professor, Dept. of Genetics and Plant
Breeding, Bangladesh Agricultural University,
Mymensingh-2202, Bangladesh.

Dr. Ahmad Ali

National Institute of Pharmaceutical Education &
Research,
Dept of Biotechnology, NIPER, Jandaha Road,
Hajipur, Bihar, India, Pin – 844102, India.

Mr. Karthikkumar V

Annamalai University,
Department of Biochemistry & Biotechnology.

Dr.K.Rajendiran

Dept of plant science, Tagore Govt. college,
9, 4th cross, Tagore Nagar, Pondicherry – 605 008,
India.

Dr. V.Balakrishnan

K.S.Rangasamy College of Technology,
Department of Biotechnology, KSR Kalvi
nagar, Tiruchengode-637215, Tamilnadu,
India.

Dr. NourAli Sajedi

*Department of Agronomy and plant Breeding,
Islamic Azad University, Arak Branch, Arak,
Iran.*

(Dr) Ms. Rachel Predeepa

*Not Applicable ,
2/387 Gokul Nagar, Kannanenthal Madurai.*

Dr. Rajendra Gyawali

*Department of Pharmacy and Biology, Kathmandu
University,
Dhulikhel,
Nepal.*

Ms. Rocheli de Souza

*UFRGS,
Porto Alegre,
Brazil.*

Dr. Om Prakash Verma

*Sam Higginbottom Institute of Agriculture,
Technology & Sciences (Formerly Allahabad
Agricultural Institute), Allahabad, U.P.,
Department of Molecular & Cellular Engineering,
Jacob School of Biotechnology & Bioengineering,
India.*

Dr. Ashwani Kumar

*JMIT, Radaur,
Department of Biotechnology, JMIT, Radaur-
135133, Haryana, India.*

Dr. Sarfaraz F. A. Al-Bamarny

*University of Duhok, College of Agriculture, Dept.
of Horticulture,
Duhok, Iraqi Kurdistan Region,
Iraq.*

Prof. Wafaa Mohamed Shukry Abdel Meamem

*Dammam University - Saudi Arabia,
Faculty of Science for Girl.
Biology Department,
P.O.Box: 838 Dammam 31113,
Saudi Arabia.*

Dr. Stephka G. Chankova

*Institute of Biodiversity and ecosystem Research,
BAS,
2 Gagarin str, 1113 Sofia, Bulgaria.*

Dr. Nana Ewusi-Mensah

*Kwame Nkrumah University of Science and
Technology,
Dept. of Crop and Soil Sciences, Faculty of
Agriculture, KNUST, Kumasi.*

Dr. Mukesh Lokanath Chavan

*K.r.c. College of horticulture, arabhavi 591 310,
karnataka,
University of horticultlura sciences, bagakot,
India.*

Dr. Maiti Parthapratim

*Dept. of Botany Midnapore College, Midnapore-
721101, Paschim Medinipur, West Bengal, India.*

Mr. Mohammad Anwar Hossain

*Kagawa University (Present), Bangladesh
Agricultural University (Permanent)
Lab. of Plant Stress Responses, Faculty of
Agriculture, Kagawa University, Miki-cho, Kita-gun,
Kagawa 761 0795, Japan.*

Dr. Antonia Tathiana Batista Dutra

*Universidade Federal do Ceará,
Av. Humberto Monte s/n – Pici Bloco 907,
laboratório 1080.*

Dr. Kuntal Das

*St. John's Pharmacy College,
#6, II Main, 9th Cross, Vijayanagar, Bangalore-104.
India.*

Dr. Amitava Rakshit

*Banaras Hindu University,
Department of Soil Science & Agril Chemistry.*

Dr. Kranthi Kiran Mandadi

*Texas A&M University,
2132 TAMU, Peterson-Rm408, College Station,
Texas-77840,
USA.*

Dr. Monica Butnariu

*Banat's University of Agricultural Sciences and
Veterinary Medicine from Timisoara,
Chemistry and Vegetal Biochemistry Department,
Calea Aradului no.119, 300645 Timisoara,
Romania.*

Dr. Ahmad Bybordi

East Azarbaijan Research Center for Agriculture and Natural Resources, Tabriz, Iran.

Dr. Haiwei Gu

903 Fifth St., West Lafayette, IN 47906.

Dr. Hu Yanbo

Northeast Forestry University, 26# Hexing Road, Xiangfang District, Harbin city, 150040, P.R., China.

Dr. Arash Kianianmomeni

Institute of Biology / Humboldt-University Berlin, Invalidenstr. 42.

Dr. Zvonko Pacanoski

Faculty for Agriculture Sciences and Food, Boul. Aleksandar Makedonski bb, 1000 Skopje, R.of Macedonia..

Dr. Lingjuan Zheng

Department of Organismic Biology, University of Salzburg, Hellbrunnenstraße 34, 5020, Salzburg, Austria.

Dr. Md. Mokter Hossain

Department of Horticulture, Bangladesh Agricultural University, Mymensingh-2202, Bangladesh.

Dr. Forouzandeh Soltani

Department of Horticultural Sciences, College of Agriculture and Natural Resources, University, of Tehran, Daneshkadeh Street, Karaj 31587-11167, Iran.

Dr. M.C.Harish

Bharathiar University, Department of Biotechnology, Coimbatore, India.

Dr. Zong-shen Zhang

School of Biological Engineering, Dalian Polytechnic University, Qinggongyuan, Ganjingzi District, Dalian, China, postcode 116034.

Prof. T. V. Ramana Rao

B R Doshi School of Biosciences, Sardar Patel University, Vallabh Vidyanagar, Gujarat,India.

Dr. Sanjeev Chandel

Baba Isher Singh Institute of Sciences & Technology, Gagra (Moga), Punjab, India.

Dr. Kuladip Jana

Bose Institute Centenary Campus, P 1/12, C.I.T. Scheme VIIM, Kolkata-700 054, India.

Prof. Ljubinko Jovanovic

University Educons, Faculty for ecological agriculture, Sremska Kamenica, Vojvode Putnika 87, Serbia.

Dr. Luis F. Goulao

Instituto de Investigacao Cientifica Tropical [Tropical Research Institute] Eco-Bio / IICT, Av. da Republica - Quinta do Marques, 2784-505 Oeiras, Portugal.

Dr. Lucky K. Attri

College of Punjabi University Patiala, E-41, Sector-14, Panjab University, Chandigarh.

Prof. Bassam Taha Yasseen

Looking for appropriate employer, Flat 307 Point Red, 146 Midland Road, Luton, LU2 0BL, UK.

Dr. Massimo Piccotto

Tecna S.r.l., Area Science Park, Loc. Padriciano, 99, I-34149 Trieste, Italy.

Instructions for Author

Electronic submission of manuscripts is strongly encouraged, provided that the text, tables, and figures are included in a single Microsoft Word file (preferably in Arial font).

The **cover letter** should include the corresponding author's full address and telephone/fax numbers and should be in an e-mail message sent to the Editor, with the file, whose name should begin with the first author's surname, as an attachment.

Article Types

Three types of manuscripts may be submitted:

Regular articles: These should describe new and carefully confirmed findings, and experimental procedures should be given in sufficient detail for others to verify the work. The length of a full paper should be the minimum required to describe and interpret the work clearly.

Short Communications: A Short Communication is suitable for recording the results of complete small investigations or giving details of new models or hypotheses, innovative methods, techniques or apparatus. The style of main sections need not conform to that of full-length papers. Short communications are 2 to 4 printed pages (about 6 to 12 manuscript pages) in length.

Reviews: Submissions of reviews and perspectives covering topics of current interest are welcome and encouraged. Reviews should be concise and no longer than 4-6 printed pages (about 12 to 18 manuscript pages). Reviews are also peer-reviewed.

Review Process

All manuscripts are reviewed by an editor and members of the Editorial Board or qualified outside reviewers. Authors cannot nominate reviewers. Only reviewers randomly selected from our database with specialization in the subject area will be contacted to evaluate the manuscripts. The process will be blind review.

Decisions will be made as rapidly as possible, and the journal strives to return reviewers' comments to authors as fast as possible. The editorial board will re-review manuscripts that are accepted pending revision. It is the goal of the AJFS to publish manuscripts within weeks after submission.

Regular articles

All portions of the manuscript must be typed double-spaced and all pages numbered starting from the title page.

The Title should be a brief phrase describing the contents of the paper. The Title Page should include the authors' full names and affiliations, the name of the corresponding author along with phone, fax and E-mail information. Present addresses of authors should appear as a footnote.

The Abstract should be informative and completely self-explanatory, briefly present the topic, state the scope of the experiments, indicate significant data, and point out major findings and conclusions. The Abstract should be 100 to 200 words in length. Complete sentences, active verbs, and the third person should be used, and the abstract should be written in the past tense. Standard nomenclature should be used and abbreviations should be avoided. No literature should be cited.

Following the abstract, about 3 to 10 key words that will provide indexing references should be listed.

A list of non-standard **Abbreviations** should be added. In general, non-standard abbreviations should be used only when the full term is very long and used often. Each abbreviation should be spelled out and introduced in parentheses the first time it is used in the text. Only recommended SI units should be used. Authors should use the solidus presentation (mg/ml). Standard abbreviations (such as ATP and DNA) need not be defined.

The Introduction should provide a clear statement of the problem, the relevant literature on the subject, and the proposed approach or solution. It should be understandable to colleagues from a broad range of scientific disciplines.

Materials and methods should be complete enough to allow experiments to be reproduced. However, only truly new procedures should be described in detail; previously published procedures should be cited, and important modifications of published procedures should be mentioned briefly. Capitalize trade names and include the manufacturer's name and address. Subheadings should be used. Methods in general use need not be described in detail.

Results should be presented with clarity and precision. The results should be written in the past tense when describing findings in the authors' experiments. Previously published findings should be written in the present tense. Results should be explained, but largely without referring to the literature. Discussion, speculation and detailed interpretation of data should not be included in the Results but should be put into the Discussion section.

The Discussion should interpret the findings in view of the results obtained in this and in past studies on this topic. State the conclusions in a few sentences at the end of the paper. The Results and Discussion sections can include subheadings, and when appropriate, both sections can be combined.

The Acknowledgments of people, grants, funds, etc should be brief.

Tables should be kept to a minimum and be designed to be as simple as possible. Tables are to be typed double-spaced throughout, including headings and footnotes. Each table should be on a separate page, numbered consecutively in Arabic numerals and supplied with a heading and a legend. Tables should be self-explanatory without reference to the text. The details of the methods used in the experiments should preferably be described in the legend instead of in the text. The same data should not be presented in both table and graph form or repeated in the text.

Figure legends should be typed in numerical order on a separate sheet. Graphics should be prepared using applications capable of generating high resolution GIF, TIFF, JPEG or Powerpoint before pasting in the Microsoft Word manuscript file. Tables should be prepared in Microsoft Word. Use Arabic numerals to designate figures and upper case letters for their parts (Figure 1). Begin each legend with a title and include sufficient description so that the figure is understandable without reading the text of the manuscript. Information given in legends should not be repeated in the text.

References: In the text, a reference identified by means of an author's name should be followed by the date of the reference in parentheses. When there are more than two authors, only the first author's name should be mentioned, followed by 'et al'. In the event that an author cited has had two or more works published during the same year, the reference, both in the text and in the reference list, should be identified by a lower case letter like 'a' and 'b' after the date to distinguish the works.

Examples:

Abayomi (2000), Agindotan et al. (2003), (Kelebeni, 1983), (Usman and Smith, 1992), (Chege, 1998;

1987a,b; Tijani, 1993,1995), (Kumasi et al., 2001) References should be listed at the end of the paper in alphabetical order. Articles in preparation or articles submitted for publication, unpublished observations, personal communications, etc. should not be included in the reference list but should only be mentioned in the article text (e.g., A. Kingori, University of Nairobi, Kenya, personal communication). Journal names are abbreviated according to Chemical Abstracts. Authors are fully responsible for the accuracy of the references.

Examples:

Chikere CB, Omoni VT and Chikere BO (2008). Distribution of potential nosocomial pathogens in a hospital environment. *Afr. J. Biotechnol.* 7: 3535-3539.

Moran GJ, Amii RN, Abrahamian FM, Talan DA (2005). Methicillinresistant *Staphylococcus aureus* in community-acquired skin infections. *Emerg. Infect. Dis.* 11: 928-930.

Pitout JDD, Church DL, Gregson DB, Chow BL, McCracken M, Mulvey M, Laupland KB (2007). Molecular epidemiology of CTXM-producing *Escherichia coli* in the Calgary Health Region: emergence of CTX-M-15-producing isolates. *Antimicrob. Agents Chemother.* 51: 1281-1286.

Pelczar JR, Harley JP, Klein DA (1993). *Microbiology: Concepts and Applications.* McGraw-Hill Inc., New York, pp. 591-603.

Short Communications

Short Communications are limited to a maximum of two figures and one table. They should present a complete study that is more limited in scope than is found in full-length papers. The items of manuscript preparation listed above apply to Short Communications with the following differences: (1) Abstracts are limited to 100 words; (2) instead of a separate Materials and Methods section, experimental procedures may be incorporated into Figure Legends and Table footnotes; (3) Results and Discussion should be combined into a single section.

Proofs and Reprints: Electronic proofs will be sent (e-mail attachment) to the corresponding author as a PDF file. Page proofs are considered to be the final version of the manuscript. With the exception of typographical or minor clerical errors, no changes will be made in the manuscript at the proof stage.

Fees and Charges: Authors are required to pay a \$550 handling fee. Publication of an article in the International Journal of Plant Physiology and Biochemistry is not contingent upon the author's ability to pay the charges. Neither is acceptance to pay the handling fee a guarantee that the paper will be accepted for publication. Authors may still request (in advance) that the editorial office waive some of the handling fee under special circumstances

Copyright: © 2014, Academic Journals.

All rights Reserved. In accessing this journal, you agree that you will access the contents for your own personal use but not for any commercial use. Any use and or copies of this Journal in whole or in part must include the customary bibliographic citation, including author attribution, date and article title.

Submission of a manuscript implies: that the work described has not been published before (except in the form of an abstract or as part of a published lecture, or thesis) that it is not under consideration for publication elsewhere; that if and when the manuscript is accepted for publication, the authors agree to automatic transfer of the copyright to the publisher.

Disclaimer of Warranties

In no event shall Academic Journals be liable for any special, incidental, indirect, or consequential damages of any kind arising out of or in connection with the use of the articles or other material derived from the IJPPB, whether or not advised of the possibility of damage, and on any theory of liability.

This publication is provided "as is" without warranty of any kind, either expressed or implied, including, but not limited to, the implied warranties of merchantability, fitness for a particular purpose, or non-infringement. Descriptions of, or references to, products or publications does not imply endorsement of that product or publication. While every effort is made by Academic Journals to see that no inaccurate or misleading data, opinion or statements appear in this publication, they wish to make it clear that the data and opinions appearing in the articles and advertisements herein are the responsibility of the contributor or advertiser concerned. Academic Journals makes no warranty of any kind, either express or implied, regarding the quality, accuracy, availability, or validity of the data or information in this publication or of any other publication to which it may be linked.

ARTICLES

Antioxidant defence mechanisms in chickpea (*Cicer arietinum* L.) under copper and arsenic toxicity

Shashi Kumar, Surender Kumar, Pravin Prakash and Mahendra Singh

Full Length Research Paper

Antioxidant defence mechanisms in chickpea (*Cicer arietinum* L.) under copper and arsenic toxicity

Shashi Kumar¹, Surender Kumar^{1*}, Pravin Prakash¹ and Mahendra Singh²¹Department of Plant Physiology, Institute of Agricultural Sciences, BHU, Varanasi-221 005, India.²Department of Agronomy, Banaras Hindu University, Varanasi- 221 005, India.

Received 6 May, 2014; Accepted 19 May 2014

Chickpea (*Cicer arietinum* L.) is one of the oldest pulse crops cultivated from ancient time both in Asia and Europe. India is the largest producer of chickpea in the world, sharing 65 and 70% of production and total global area, respectively. The inhibition of plant growth and crop productivity by excess heavy metals is a global agricultural problem. Therefore, a pot experiment was conducted at wire house of the Department of Plant Physiology, Institute of Agricultural Sciences, Banaras Hindu University, Varanasi during *Rabi* season of 2009-10, using a complete randomized design with four replications and five treatments. Our experimental findings reveal that excessive amount of copper (Cu) (200 mg/kg) and arsenic (As) (200 mg/kg) in soil caused a mark reduction in nitrate reductase enzyme activity at both growth stages. However, a little increment was found in proline content, peroxidase and superoxide dismutase activity under higher doses of Cu and As at both growth stages. The study aimed to get more information on physiological changes in activity of antioxidant enzymes under heavy metal toxicity.

Key words: Antioxidant enzymes, chickpea, heavy metals, nitrate reductase, proline content.

INTRODUCTION

Chickpea (*Cicer arietinum* L.) is one of the oldest pulse crops cultivated from ancient time both in Asia and Europe. India is the largest producer of chickpea in the world, sharing 65 and 70% of production and total global area, respectively. It is grown in tropical, sub-tropical and temperate regions as winter season crop. Chickpea contains 19-21% protein and 60% carbohydrate, so it provides an excellent quality of dietary protein for overcoming future food shortages. Heavy metals are metals with density higher than 5 g cm⁻³. Fifty three out of

the ninety naturally occurring elements have density higher than 5 g cm⁻³ but not all of them are of biological importance. Based on their solubility under physiological conditions, seventeen heavy metals may be available for living cells and are of importance to organism and ecosystem (Weast, 1984). Among these metals, Fe, Mo, Zn, and Mn are important as micro nutrient: Ni, Co, Cu, V, W and Cr are toxic elements with high or low importance as trace elements. However, As, Hg, Ag, Cd and Pb have no known function as nutrients and seem to

*Corresponding author. E-mail: surender.bhu@gmail.com.

be more or less toxic to plants and microbial population (Godbold and Hatlerman, 1985). Heavy metals enter the food chain from industrial sources, natural resources and geochemical cycle. Heavy metals are extremely phytotoxic in nature and the danger is aggravated by their almost indefinite persistence in the environment.

The inhibition of plant growth and crop productivity by excessive heavy metals is a global agricultural problem. The most widespread visual evidence of metal phytotoxicity is reduction in plant growth as metal toxicity increases. However, different metals have different sites of action within the plant. Copper as heavy metal plays a vital role in deteriorating soil biotic and abiotic factor and disrupting plants metabolism. There are several factors that increase copper concentration up to toxic level in the soils such as soil chemical characteristics, agricultural manufacturing, metalliferous mining, smelting sites, waste disposal practices, use of metal containing pesticides, sewage sludge, etc.

Copper as a micronutrient plays a role in normal metabolism in higher plants; it participates in the reduction of electron transfer in photosynthesis in the form of plastocyanin. However, copper at high level becomes strongly phototoxic to cell and causes inhibition of plant or even death (Mocquot et al., 1996). Studies from plant species demonstrate that excess copper in plant growth medium induces formation of ROS in treated tissues; copper induced generation of hydrogen peroxide is directly correlated with damage to protein and lipids (Murphy and Taiz, 2000). Photosynthesis is also sensitive to excess copper; the pigment and protein components of photosynthesis membranes are the targets. Excess copper causes an elevated susceptibility to photo inhibition of Ps-II *in vitro* and *in vivo*. It has long been known that high concentration of copper when added to the incubation medium of isolated thylakoids inhibits PS-II electron transfer activity on the acceptor sites. In addition, copper toxicity is related to the disturbances in the uptake of essential elements (Assche and Clijsters, 1990). Arsenic can be potential activator of the biosynthesis of phytochelatins (pcs) in several plants (Maitani et al., 1996; Schmogger et al., 2000). So far, little has been known about its effect on the synthesis of homophytochelatins (hpcs) in plants. Work done by previous workers suggests that there is increase in oxidative stress under heavy metal toxicity. Therefore, there is need to study the effect of heavy metals such as copper and arsenic on important metabolic enzymes and antioxidant enzymes activity. Hence, an investigation was undertaken to study the antioxidant defence mechanisms in chickpea (*C. arietinum* L.) under copper and arsenic toxicity.

MATERIALS AND METHODS

The present investigation was carried out with a variety of chickpea

(Radhey) obtained from the Department of Genetics and Plant Breeding, Institute of Agricultural Sciences, Banaras Hindu University Varanasi during Rabi season of 2009-10, using a complete randomized design with 4 replications and 5 treatments. The details of the treatment are: control, copper sulphate (100 and 200 mg/kg), sodium arsenate (100 and 200 mg/kg) soil. The seeds were sown in plastic pots containing 3 kg pulverized farm soil mixture with farm yard manure in the ratio of 3:1. The sowing was done on 26th November, 2009. Fertilizer was applied at the ratio of 20, 35 and 25 ppm of N, P and K, respectively. There were four pots for each replication and four healthy plants were kept in each pot. Antioxidant enzyme activities were done on two separate plants from each pot at two different growth stages through destructive techniques.

For estimation of nitrate reductase activity, leaves from uniformly grown seedlings in a homogeneous population were selected for enzyme induction. The enzyme activity was assayed *in vitro* by the method of Srivastava (1975) in the first fully expanded leaf. Peroxidase activity was assayed in leaf tissue at early stage after imposing stress in control and treated plants. The enzyme assay was performed according to the protocol of Kar and Mishra (1976). Superoxide dismutase activity was assayed in leaf tissue at early stage after imposing stress in control and treated plants using the method of Dhindsa et al. (1981). Proline content was determined in leaf sample after imposing stress in control and treated plants using the method of Bates et al. (1973).

Statistical analysis

Simple CRD design was followed in all observations. Analysis of variance was performed on the data at each stage as described by Panse and Sukhatme (1967). Least significant difference (LSD) was calculated at 5% probability level.

RESULTS AND DISCUSSION

Nitrate reductase activity

Significant and greater reduction in nitrate reductase (NR) activity was observed at higher doses of arsenic and copper (Table 1). Arsenic showed greater depression in NR activity as compared to copper but there was non-significant difference at the same doses of arsenic and copper. Sodium arsenate of 200 mg/kg recorded the least NR activity and significantly differed from control and lower doses of copper and arsenic. Non-significant differences were observed between copper sulphate of 100 mg/kg and sodium arsenate of 100 mg/kg at 30 and 45 days after sowing (DAS). There was 65.0% reduction in NR activity as observed in sodium arsenate (200 mg/kg) as compared to the control.

Proline content

Proline content increased from 30 DAS to 45 DAS under stressful environment (Table 1). Highest significant increase in proline content was observed in sodium arsenate of 200 mg/kg (65.8%) as compared to copper sulphate of 100 and 200 mg/kg. But copper sulphate of 100 and 200 mg/kg was at par at 30 DAS. Greater increase in proline content was observed at 30 DAS as

Table 1. Effect of copper and arsenic on nitrate reductase, peroxidase, proline and superoxide dismutase activity in chickpea at 30 and 45 days after sowing.

Treatment	Dose (ppm)	Nitrate reductase ($\mu\text{moles NO}_2^- \text{ g}^{-1}$ fresh weight hour^{-1})			Peroxidase (U g^{-1} fresh weight min^{-1})			Proline ($\mu\text{g g}^{-1}$ fresh weight)			Superoxide dismutase (U g^{-1} fresh weight min^{-1})		
		I	II	Mean	I	II	Mean	I	II	Mean	I	II	Mean
Control	-	0.34	0.58	0.46	14.23	14.45	14.34	23.48	25.38	24.43	153.65	168.50	161.0
Copper Sulphate	100	0.21 (-38.2)	0.51 (-12.0)	0.36	15.38 (+8.0)	15.92 (+10.1)	15.65	32.36 (+37.8)	34.34 (+35.0)	33.35	186.80 (+21.7)	198.36 (+17.7)	192.5
Copper Sulphate	200	0.13 (-61.7)	0.15 (-74.0)	0.14	16.43 (+15.4)	17.13 (+18.5)	16.78	34.25 (+45.9)	36.28 (+42.9)	35.27	260.05 (+69.2)	291.96 (+73.2)	276.0
Sodium Arsenate	100	0.18 (-47.0)	0.38 (-34.0)	0.28	15.13 (+6.3)	15.56 (+7.6)	15.30	36.83 (+56.9)	38.94 (+53.4)	37.89	352.82 (+129.6)	380.89 (+126.0)	366.8
Sodium Arsenate	200	0.08 (-65.0)	0.09 (-84.0)	0.08	17.33 (+21.7)	17.87 (+23.6)	17.60	38.92 (+65.8)	41.29 (+62.6)	40.11	391.61 (+154.8)	410.05 (+143.3)	400.8
Mean		1.12	1.30	-	4.70	5.18	-	33.17	35.25	-	287.06	288.62	-
SEM \pm		0.03	0.05	-	0.33	0.27	-	1.52	1.04	-	8.32	7.3	-
C.D. at 5%		0.19	0.14	-	0.98	0.81	-	4.47	3.12	-	24.94	21.7	-

Figure in parenthesis represents percentage increase (+) and decrease (-) over control, I and II means data recorded at 30 and 45 days after sowing.

compared to 45 DAS except for sodium arsenate of 200 mg/kg. Arsenic and copper at higher doses resulted in higher production of proline in plants. 100 and 200 mg/kg copper sulphate and 100 mg/kg sodium arsenate did not show significant differences between them at both growth stages, but there was significant differences observed in the control.

Peroxidase activity

There was significant increase in peroxidase activity observed with increasing doses of arsenic and copper (Table 1). Mean peroxidase activity recorded at both growing stages was least in control (14.23 U g^{-1} fresh weight min^{-1}) and highest in sodium arsenate of 200 mg/kg (17.87 U g^{-1} fresh weight min^{-1}) in all the treatments. Higher doses of arsenate significantly differed from all

other treatments with respect to peroxidase activity. Similarly, all the treatment varied significantly for peroxidase activity except control and copper sulphate of 100 mg/kg at 30 DAS. The highest increase in peroxidase activity was recorded in sodium arsenate of 200 mg/kg (23.6%) at 45 DAS as compared to the control.

Superoxide dismutase activity

Changes in superoxide dismutase (SOD) activity are presented in Table 1. SOD activity increased significantly with increasing doses of arsenic and copper at both growth stages. Arsenic had greater increase in SOD activity as compared to copper. There was significant difference for SOD activity in all the treatments at both growth stages. Highest increase in SOD activity was recorded in

sodium arsenate of 200 mg/kg (54.8%) as compared to control. Mean SOD activity was recorded as least in control (153.65 U g^{-1} fresh weight min^{-1}) and highest in sodium arsenate (200 mg/kg) (410.05 U g^{-1} fresh weight min^{-1}).

DISCUSSION

The inhibition of plant growth and crop production by excess heavy metals such as copper, cadmium, arsenic, or lead in contaminated-soil is a global agricultural problem. In this study, an effort was made to study the effect of two highly toxic metal pollutants of soil (arsenic and copper) on different antioxidant enzymes activities in chickpea at 30 and 45 days after imposing the heavy metal stress treatment. It was observed that arsenic had more pronounced toxic effects

in terms of reduction in NR activity and increased proline content, peroxidase activity and SOD activity in comparison with copper at the same concentrations. Gupta et al. (2004) also reported more inhibitory effect of arsenic and cadmium on plants as compared to Cu, Zn and Ni. They have concluded that growth inhibition is specific for metal species rather than their total uptake by the tissues.

The presence of arsenic and copper significantly inhibited the activity of the enzyme nitrate reductase (NR) and the degree was dependent on the concentration of the metal used. This may be due to inhibition or metabolic dysfunction of the enzyme protein. Moreover, metals also have an impact on the activity of plasma membrane bound proton pump and the fluidity of the membrane restricting the uptake of nitrate, the inducer and the substrate of NR.

Arsenic and copper stress also resulted in an increase in the level of proline in the leaves of chickpea plant. There was more increase in proline content at higher doses of the heavy metal. The accumulation of free proline in response to heavy metal is widespread among plant. The accumulation of proline is a gene regulated process which is the consequence of the over expression of the gene involved in its biosynthesis and depression of those involved in its degradation, in the plants under stress. The functional significance of proline accumulation under heavy metal stress might include water balance maintenance, scavenging of hydroxyl radical or metal chelation.

The activity of two antioxidant enzymes peroxidase and superoxide dismutase (SOD) was found to increase on imposition of arsenic and copper stress. There was increased activity of these enzymes at higher doses of heavy metal stress. Heavy metals are known to cause the production of ROS, that is, the formation of superoxide (O_2^-), hydroxyl (OH^\cdot) radicals, hydrogen peroxide (H_2O_2) and singlet oxygen (O_2^1) in plants (Foyer et al., 1997). Mohan et al. (1997) suggested that the increase of peroxidase activity may be an effect of accelerated senescence, connected with enhanced formation of hydrogen peroxide (H_2O_2) or secondary metabolites such as phenolic compound. It has been demonstrated in *Zea mays* that catalase and SOD were stimulated after exposure to arsenic (Mylona et al., 1998). Arsenic accumulated in the plant tissue stimulates peroxidase synthesis during the early phases of plant development, long before the appearance of visible changes (Miteva and Peycheva, 1999; Stoena et al., 2003). There are also reports from several plant species establishing that Cu causes oxidative stress and increases the activities of anti-oxidative enzymes (Luna et al., 1994; Gupta, 1999).

Conflict of Interests

The author(s) have not declared any conflict of interests.

REFERENCES

- Assche VF, Clisjters H (1990). Effect of metals on enzyme activity in plants. *Plant Cell Environ.* 13:195-206.
- Bates LS, Waldren RP, Tare ID (1973). Rapid determination of free proline for water-stress studies. *Plant Soil.* 39:205-207.
- Dhindsa RS, Plumb-Dhindsa P, Thorpe TA (1981). Leaf senescence, correlated with increased levels of membrane permeability and lipid peroxidation, and decreased levels of superoxide dismutase and catalase. *J. Exp. Bot.* 32:93-101.
- Foyer CH (1997). Oxygen metabolism and electron transport in photosynthesis. In *molecular biology of free radical scavenging enzyme*. Cold Spring Harbor Laboratory, New York. 54:234-237.
- Godbold DL, Huttermann A (1985). Effect of zinc, cadmium and mercury on root elongation of *Picea abies* (Karst.) seedlings, and the significance of these metals to forest die-back. *Environ. Pollut. (series A)*. 38:375-381.
- Gupta DK (2004). Changes in the levels of phytochelatin and related metal-binding peptides in chickpea seedling exposed to arsenic and different heavy metal ions. *J. Plant Res.* 117:253-256.
- Gupta MA, Cuypers J, Vangronsveld, Clijsters H (1999). Copper affects the enzymes of the ascorbate-glutathione cycle and its related metabolites in the roots of *Phaseolus vulgaris*. *Physiol. Plant.* 106:262-267.
- Kar M, Mishra D (1976). Catalase, peroxidase and polyphenoloxidase activities during rice leaf senescence. *Plant Physiol.* 57:315-319.
- Luna CM, González CA, Trippi VS (1994). Oxidative damage caused by an excess of copper in oat leaves. *Plant Cell Physiol.* 35:11-15.
- Maitani T, Kubota H, Sato K, Yamada T (1996). The composition of metals bound to class III metallothionein (phytochelatin and its desglycyl peptid) induced by various metals in root cultures of *Rubia tictorum*. *Plant Physiol.* 110:1145-1150.
- Miteva E, Peycheva S (1999). Arsenic accumulation and effect on peroxidase activity in green bean and tomatoes. *Bulgarian J. Agric. Sci.* 5:737-740.
- Mocquot B, Vangernsveld J, Clijsters H, Mench M (1996). Copper toxicity in young maize (*Zea mays* L.) plants effects on growth, mineral and chlorophyll contents and enzymes activities. *Plants soil.* 182:287-300.
- Mohan MS, Nair A, Bhagwat TG, Krishna, Yano M (1997). Genome mapping, molecular markers and marker-assisted selection in crop plants. *Mol. Breed.* 3:87-103.
- Mylona PV, Polidoros AN, Scandalios, JG (1998). Modulation of antioxidant responses by arsenic in maize. *Free Radical Biol. Med.* 25:576-585.
- Panse VG, Sukhatme PV (1967). *Statistical methods for agricultural workers*. ICAR publications, New Delhi. 167-174.
- Schmoger MEV, Oven M, Grill E (2000). Detoxification of arsenic by phytochelatin in plants. *Plant Physiol.* 122:793-801.
- Srivastava HS (1974). *In vitro* activity of nitrate reductase in maize seedlings. *Indian J. Biochem. Biophys.* 11: 230-232.
- Stoena N, Berova M, Zlatev ZL (2003). Physiological response of maize (*Zea mays* L.) to different level of as contamination. *Biologia Plant.* (In print).
- Weast C (1984). *Handbook of geochemistry and physics*. 50th, Edn. Chemical Rubber Co., Cleveland, Ohio.



International Journal of Plant Physiology and Biochemistry

Related Journals Published by Academic Journals

- *African Journal of Plant Science*
- *Journal of Microbiology and Antimicrobials*
- *International Journal for Biotechnology and Molecular Biology Research*
- *Journal of Botany and Plant Pathology*
- *International Journal of Biodiversity and Conservation*

academicJournals