

# International Journal of Plant Physiology and Biochemistry

Volume 6 Number 6, July 2014

ISSN 2006-9871



*Academic  
Journals*

## ABOUT IJPPB

The **International Journal of Plant Physiology and Biochemistry (IJPPB)** (ISSN 2006-9871) is published Monthly (one volume per year) by AcademicJournals.

**International Journal of Plant Physiology and Biochemistry (IJPPB)** provides rapid publication (monthly) of articles in all areas of the subject such as plant hormones, seed biology, plant DNA repair, Concepts of target cells in plant differentiation etc.

The Journal welcomes the submission of manuscripts that meet the general criteria of significance and scientific excellence. Papers will be published shortly after acceptance. All articles published in IJPPB are peer-reviewed.

## Submission of Manuscript

Please read the **Instructions for Authors** before submitting your manuscript. The manuscript files should be given the last name of the first author

[Click here to Submit manuscripts online](#)

If you have any difficulty using the online submission system, kindly submit via this email [ijppb@academicjournals.org](mailto:ijppb@academicjournals.org).

With questions or concerns, please contact the Editorial Office at [ijppb@academicjournals.org](mailto:ijppb@academicjournals.org).

## Editors

**Prof Dr. Ishrak Khafagi**

*Faculty of Science, Suez Canal University,  
Ismailia, Egypt*

**Prof. Mohamed Mahgoub Azooz**

*Biology Department-Faculty of Science- King Faisal  
University, Saudi Arabia.*

**Dr. Bansal Parveen**

*National Institute of Ayurvedic Pharmaceutical  
Research  
Moti Bagh Road, Patiala-(Punjab) India.*

**Prof. Bechan Sharma**

*University of Allahabad  
Department of Biochemistry, University of  
Allahabad, Faculty of Science, Allahabad-211002,  
India.*

## Editor Board

**Prof. Weimin Zhang**

*Guangdong Institute of Microbiology  
100 Central Xian lie Road, Guangzhou, Guangdong  
510070, China.*

**Dr. Xu Suxia**

*Fujian Institute of Subtropical Botany,  
780-800, Jiahe Road, Xiamen, China361006, China.*

**Dr. Adaku Vivien Iwueke**

*Federal University of Technology, Owerri  
Department of Biochemistry, Federal University of  
Technology, Owerri  
Nigeria.*

**Ass. Prof. Turgay CELIK**

*Gulhane Military Medical Academy,  
School of Medicine, Department of Cardiology,  
Turkey.*

**Dr. Topik Hidayat**

*Department of Biology Education  
Indonesia University of Education (UPI)  
Jalan Dr. Setiabudhi 229 Bandung 40154 Indonesia  
Indonesia.*

**Dr. Tariq Mahmood**

*Quaid-i-Azam University,  
Department of Plant Sciences, Quaid-i-Azam  
University, Islamabad, Pakistan.*

**Dr. Neveen B. Talaat**

*Department of Plant Physiology, Faculty of  
Agriculture, Cairo University, Egypt.*

**Dr. Sudhamoy Mandal**

*Fulbright Visiting Fellow  
Department of Plant Pathology  
University of Nebraska Lincoln  
USA*

**Asso. Prof. Chankova Stephka Georgieva**

*Central Laboratory of General Ecology, Bulg Acad  
Sci 1113 Sofia, 2 Gagarin str, Bulgaria.*

**Shijie Han**

*Center of Forestry, Institute of Applied Ecology,  
Chinese Academy of Sciences,  
72 Wenhua Road, Shenyang City, Liaoning  
Province 110016, PR China.*

**Szu-Chuan Shen**

*Department of Medical Nutrition,  
I-Shou University  
Yanchao Township,  
Kaohsiung County 824, Taiwan.*

**SEDDIK KHENNOUF**

*Dept of BIOLOGY, Faculty of Science  
University Ferhat Abbas, SETIF, 19000, ALGERIA.*

**Saranyu Khammuang**

*Maharakham University  
Department of Chemistry, Faculty of Science  
Thailand.*

**Samir Ranjan Sikdar**

*Bose Institute  
P-1/12, C.I.T. Scheme VII M, Kolkata 700 054,  
India.*

**Dr. M. Abdul Salam**

*Department of Agronomy, College of Agriculture  
Kerala Agricultural University, Vellayani 695 522,  
Trivandrum, Kerala.*

**Dr.Saeed Aminzadeh**

*National Institute of Genetic Engineering and  
Biotechnology (NIGEB)  
Shahrak-e-Pajooresh Km 15, Tehran-Karaj  
Highway, Tehran, I.R.Iran.*

**Dr.Ruzica Stricevic**

*Faculty of Agriculture-University of Belgrade  
Nemanjina 6, Zemun, 11080, Serbia.*

**Rumbos Christos**

*University of Thessaly  
Fytokoy Str, 384 46  
Volos, Greece.*

**Dr. Özge Zencir**

*Kemah Vocational Training School,  
Erzincan University, Kemah,  
Erzincan, Turkey.*

**RIYAZALI ZAFARALI SAYYED**

*SI P Arts, GBP Science & STSKVS Comm. College  
SHAHADA Dist Nandurbar, Maharashtra INDIA.*

**Raul Rodriguez-Herrera**

*Universidad Autonoma de Coahuila  
School of Chemistry  
Blvd. V. Carranza y González Lobo s/n Col República  
Saltillo Coahuila México.*

**Dr. A. H. M. Mahbubur Rahman**

*Rajshahi University, Bangladesh.  
Department of Botany, University of Rajshahi,  
Bangladesh.*

**Paul S. Marley**

*Department of Crop Protection, IAR/FOA  
Ahmadu Bello University, P.M.B. 1044, Samaru,  
Zaria, Nigeria.*

**Patrick Addo-Fordjour**

*Kwame Nkrumah University Of Science And  
Technology (Knust), Kumasi,  
Department Of Theoretical And Applied Biology,  
Ghana.*

**Battu.Prasanna Reddy**

*Nosch Labs Pvt Ltd  
Hyderabad, India.*

**Noureddine Benkeblia**

*UWI - Department of Life Sciences  
Mona Campus, Kingston 7, Jamaica.*

**Keutgen, Norbert**

*Uniwersytet Technologiczno-Przyrodniczy  
im. Jana i Jędrzeja Śniadeckich w Bydgoszczy  
Kadra Katedry Fizjologii Roslin (Institute of Plant  
Physiology)  
ul. Bernardynska 6/8, 85-029 Bydgoszcz, Poland.*

**Nicholas E. Korres**

*University College Cork, Environmental Research  
Institute.  
Lee Road, Cork, Ireland.*

**Dr Naveen Kumar**

*University of Florida  
2685 SR 29 N SWFREC/IFAS/UFL, Immokalee,  
FL34142, USA.*

**Dr Modala Venkateswarlu**

*Seribiotech research Laboratory, Kodathi  
Carmelaram post, Bangalore.*

**Mirza Hasanuzzaman**

*Department of Agronomy, Faculty of Agriculture,  
Sher-e-Bangla Agricultural University, Dhaka-  
1207, Bangladesh.*

**Maybelle Gaballah**

*National Research Centre, El Behoos street, Dokki,  
Cairo.*

**Mauro Guida Santos**

*Universidade Federal de Pernambuco  
Street Moraes Rego – CDU – CCB – Botany  
Department, s/n. 50670-901. Pernambuco State,  
Brazil.*

**Marcelo Rodrigo Pace**

*University of Sao Paulo  
Rua do Matão, 277, Cidade Universitária São  
Paulo, Brazil.*

**Marcelo Francisco Pompelli**

*Federal University of Pernambuco  
Department of Botany, Profª Moraes Rego Av.,  
Recife – PE – Brazil, 50670-901.*

**Luca Catalina Mariana**

*University of Bucharest, Faculty of Biology, Dept of  
Biochemistry and Molecular Biology  
Spl. Independentei, no.91-95, Bucharest 5,  
Romania.*

**Lin Wang**

*Institute of Biostatistics, Fudan University  
220 Handan Road, Shanghai 200433, China  
genetics, microbiology  
China.*

**Li Qiang**

*Institute of karst geology,MLR  
50 Qixing Road,  
China.*

**Dr. Ayanakumar Kumar**

*C.Abdul Hakeem College of Engg. & Tech.,  
Melvisharam-632 509, Vellore Dist, Tamil Nadu,  
INDIA.*

**P.Krishnamoorthy**

*P.G. AND RESEARCH DEPARTMENT OF ZOOLOGY  
RAJAH SERFOJI GOVT. COLLEGE.  
India.*

**Hare Krishna**

*Central Institute of Temperate Horticulture-  
Regional Station,  
Mukteshwar-263 138, District- Nainital,  
Uttarakhand, India.*

**K.G. Mandal**

*Directorate of Water Management (formerly Water  
Technology Centre for Eastern Region)  
Indian Council of Agricultural Research  
C.S. Pur, Bhubaneswar-751023, ORISSA, INDIA.*

**Dr. Jukta Adhikari**

*Presidency College  
86/1, College Street, Kolkata – 700 073, India.*

**Jorge Teixeira**

*Botany Department, Faculty of Sciences,, University  
of Porto,  
Edifício FC4, Rua do Campo Alegre, S/N, 4169-007  
Porto, Portugal.*

**Johnson Toyin Fasinmirin**

*Federal University of Technology, Akure, Nigeria  
Department of Agricultural Engineering, FUT,  
P.M.B. 704, Akure, Ondo State, Nigeria..*

**Joel K. Ransom**

*North Dakota State University  
166 Loftsgard Hall, Department of Plant Sciences,  
NDSU Dept. 7670, PO Box 6050, Fargo, ND 58108-  
6050.*

**João Claudio Damasceno de Sá**

*UENF  
Av. José Carlos Pereira Pinto, 39. Pq. Vicente Dias.  
Campos RJ. Brazil.*

**Jalal Jalali Sendi**

*University of Guilan  
Department of Plant Protection, university of  
Guilan, Rasht, Iran.*

**Lúri Drumond Louro**

*Universidade Federal do Espírito Santo  
Rua Horácio Andrade de Carvalho, 210, Victoria,  
ES, 29052-620, Brazil.*

**Hong Bo Guo**

*Northwest A and F University  
22 Xinong, Yangling 712100, Shaanxi, PR China.*

**Harsukh P. Gajera**

*Junagadh Agricultural University  
Department of Biochemistry, College of  
Agriculture, JAU, Junagadh- 362 001, Gujarat,  
India.*

**Hanan Abdel Fattah El-Sadawy**

*National Research Center  
El-Buhoth St., Dokki, Giza, Egypt.*

**Assist. Prof .Azime KÜÇÜKGÜL GÜLEÇ**

*Tunceli University Fisheries Faculty 62000,  
Tunceli/TURKEY.*

**Greg T. Hannig**

*DuPont  
1090 Elkton Road Newark, DE 19711.*

**Gilberto Santos Andrade**

*Instituto de Biotecnologia Aplicada a  
Agropecuária (BIOAGRO), Departamento de  
Biologia Animal, Universidade Federal de Viçosa,  
Viçosa, MG 36571-000, Brazil.*

**Dr. T. Muthukumar**

*Department of Botany, Bharathiar University  
Coimbatore -641 046, Tamilnadu, India.*

**Kunjupillai Vijayan**

*Institute of Plant and Microbial Biology  
Academia Sinica, Taipei, Taiwan-115, ROC.  
Taiwan.*

**Badre Alam**

*National Research Centre For Agroforestry  
Gwalior Road, Jhansi-284003, U.P., India.*

**Abeer Essam El-Din Mahmoud**

*Biochemistry Department  
Genetic Engineering & Biotechnology Division  
National Research Center El Tahrir St., El Dokki  
12622, Cairo, Egypt.*

**Qazi Fariduddin**

*Aligarh Muslim University  
Department of Botany, Aligarh 202 002, India.*

**Darmawan Darma**

*Faculty of Agriculture, Andalas University  
Kampus Limau Manis Padang-25163, Indonesia.*

**Barbara Chaves**

*Institute for Agricultural and Fisheries Research.*

**Sudhamoy Mandal**

*Central Horticultural Experiment Station (ICAR)  
Aiginia, Bhubaneswar, PIN-751019.*

**Cavit Bircan**

*Adnan Menderes University  
Faculty of Agriculture  
Department of Food Engineering  
09100/Aydin/Turkey.*

**Carlos Alberto Ortega-Ojeda**

*Central University of Ecuador. Faculty of Agriculture  
Sciences. Quito, Ecuador  
Calle 12 # 29 B - 78, Apto. 102 F, Unidad Residencial  
Colseguros, Cali, Colombia.*

**Brian Wade Jamandre**

*National Taiwan University  
Rm. 622, life science bldg., NTU, no.1, sec.4,  
Roosevelt rd. Taipei 10617, Taiwan (ROC).*

**Bitá Naseri**

*Agricultural Research Institute  
Department of Plant Protection, Agricultural  
Research Institute, PO Box 45195474, Zanjan, Iran..*

**Behzad Kaviani**

*Adeyemi Oluyomi Stephen  
Bells University of Technology  
Chemical Sciences Department, Km 8 Ididroko  
Road, Ota, Ogun State,  
Nigeria.*

**Ajayi Adedayo Olajide**

*Adekunle Ajasin University  
Dept. of Microbiology, P.M.B 01, Akungba-Akoko,  
Ondo State, Nigeria.*

**Alexandre Igor Azevedo Pereira (Pereira, A.I.A.)**  
*Universidade Federal de Viçosa, Departamento de Biologia Animal, Programa de Pós-Graduação em Entomologia. 36570-000, Viçosa, Minas Gerais State, Brazil.*

**Gilberto Santos Andrade**  
*Instituto de Biotecnologia Aplicada a Agropecuária (BIOAGRO), Departamento de Biologia Animal, Universidade Federal de Viçosa, Viçosa, MG 36571-000, Brazil.*

**Pradeep. A.R., Ph.D**  
*Seribiotech Research Laboratory  
Carmelaram.P.O; Bangalore, INDIA.*

**Azamal Husen**  
*University of Gondar  
Department of Biology, Faculty of Natural Sciences, University of Gondar  
P.O. Box #196, Gondar, Ethiopia.*

**Muhammad Aslam**  
*University College of Agriculture, Bahauddin Zakariya University  
Multan 60800, Pakistan.*

**Autumn J. Smith**  
*Sam Houston State University, Texas.*

**La Sara**  
*Haluoleo University  
Kampus Baru Tridharma, Kendari,  
Southeast Sulawesi, Indonesia.*

**Aliyu Mohammed**  
*Department of Human Physiology, ABU, Zaria.  
Nigeria.*

**Prof. EL-Said Ahmed AL-Sayed Ragab**  
*National Research Institute of Astronomy and Geophysics, 11421, Helwan, Egypt.*

**Shnoudy Anwar Bakhoum**  
*National Institute of Oceanography & Fisheries (NIOF), Egypt.*

**Antonio Americo Barbosa Viana**  
*Embrapa Recursos Geneticos e Biotecnologia PBI-LPP1  
PqEB Final W/5 Norte, Brasília, DF – Brazil*

**Dr. Shirish Rajmalwar**  
*National Research Laboratory for Conservation, Shirish Rajmalwar, LIG Plot No. 43, Mhada colony, Wardha – 442001, (MS) India.*

**Dr. Amaresh Chandra**  
*Universidade Federal de Viçosa, Departamento de Biologia Animal, Programa de Pós-Graduação em Entomologia. 36570-000, Viçosa, Minas Gerais State, Brazil.*

**Dr. Atul Kumar**  
*GB PANT University of Agriculture & Technology  
Department of Basic Science, College of Forestry & Hill Agriculture, HILL CAMPUS, PO Ranichauri, Tehri Garhwal, Uttarakhand State, India.*

**Prof. Levenko Boris**  
*Natl. Botanical Gardens, NAS of Ukraine  
01014 Kiev, 1 Timiryasevka st. Ukraine.*

**Dr. Dionisio G. Alwindia**  
*Bureau of Postharvest Research and Extension  
CLSU Compound, Science City of Munoz, Nueva Ecija 3120, Philippines.*

**Dr. Bhoopander Giri**  
*University of Delhi  
Department of Botany, SSNC (University of Delhi)  
Alipur, Delhi 110036, India.*

**Dr. Anjali Sood**  
*University of Delhi  
Department of Botany, University of Delhi, Delhi-110 007, INDIA.*

**Dr. A. K. Verma**  
*G.B. Pant University of Agriculture & Technology, Pantnagar, Department of Biochemistry, College of Basic Sciences,  
India.*

**Dr. Anjana Jajoo**  
*School of Life Science, Devi Ahilya University,  
Indore, DAVV  
Khandwa Road campus, Indore 452 017, M.P.,  
India.*

**Dr. Deepak Ganjewala**  
*Vellore Institute of Technology University  
55 Thennaraam Street, Vellore-632 014 (T.N.),  
India.*

**Dr. Geetha Govind**

*Max-Planck-Institute for Chemical Ecology  
Hans-Knöll Straße 8, 07745 Jena, Germany.*

**Dr. Hossam El-Din Saad El-Beltagi**

*Biochemistry Department, Faculty of Agriculture,  
Cairo University, Giza, Egypt, P.O.Box 12613  
Egypt.*

**Prof. Dr. Md. Shahidul Haque**

*Dept. of Biochemistry and Molecular Biology  
University of Rajshahi, Rajshahi-6205, Bangladesh  
Bangladesh.*

**DR. P.K.NAGAR**

*Retired Senior Scientist, IHBT, Palampur, (H.P.),  
B.21/115-10A Batuk Dham Colony, Kamachha,  
Varanasi 221 010, INDIA.*

**Dr. Satyawati Sharma**

*Indian Institute of Technology  
Centre for Rural Development & Technology, IIT  
Delhi-110016  
Biomass Production on waste land, India.*

**Dr. Uğur Çakılcıoğlu**

*Firat University Elazığ/TURKEY  
Cumhuriyet M. Malatya C. No:50/A.*

**Prof. Abdelrhani Elachqar**

*Faculty of Sciences Dhar El Mahraz, Fez, Morocco  
BP 1796, Fès-Atlas, Fès, Maroc, Morocco.*

**Ass. Prof. Jianfeng Xu**

*Arkansas State University  
PO Box 639, State University, AR 72467 USA.*

**Ass.Prof. Jin Xu**

*Center for Agricultural Resources Research,  
Institute of Genetics and Developmental Biology,  
Chinese Academy of Sciences  
Huaizhong RD 286, Shijiazhuang, HeBei, China.*

**José Carlos Rebuglio Velloso Ph.D**

*PARANÁ STATE UNIVERSITY OF PONTA GROSSA  
(Universidade Estadual de Ponta Grossa – UEPG)  
General Carlos Cavalcanti Avenue, 4748,  
Uvaranas, Ponta Grossa/PR – PO box 84030-900*

**Dr. Krouma Abdelmajid**

*Centre of Biotechnology, Borj Cedria Ecopark  
BP 901, Hammam-Lif 2050, Tunisia  
College of Science and Arts, Qassim University, BP 53,  
Al-Rass 3330353, Qassim, Saudi Arabia  
Saudi Arabia*

**Dr. Majid Rostami**

*Malayer University  
Department of Agriculture and Natural Resources,  
Postal code: 65719- 95863, University of Malayer  
Malayer, Iran.*

**Dr. Mohammad Nasir Khan**

*Aligarh Muslim University, Aligarh, INDIA  
Plant Physiology Section, Department of Botany,  
Aligarh Muslim University, Aligarh-202 002, U.P.,  
India.*

**Prof. N.K.Matta**

*Kurukshetra University  
Department of Botany, Kurukshetra University,  
Kurukshetra 136119, INDIA.*

**Dr. Naceur Djebali**

*Centre of Biotechnology Borj-Cedria (CBBC)  
BP 901, Hammam-Lif 2050 Tunisia.*

**Dr. Nader Chaparzadeh**

*Azərbaycan University of Tarbiat Moallem, Tabriz,  
Iran.*

**Nautiyal Prakash Chandra**

*Directorate Of Groundnut Research (Icar)  
Post box, No. 5, Ivnagar Road, Junagadh-362001,  
Gujarat, India.*

**Prof. Hussein Fawzy Hussein Abouzienna**

*National Research Center  
Botany Department, National Research Center,  
Elbhos Street, Dokki, Cairo, Egypt.*

**Dr. D.E. Chandrashekar Rao**

*National Research Council Canada / Plant  
Biotechnology Institute (NRC-PBI)  
110 Gymnasium Place / Saskatoon, Saskatchewan  
S7N 0W9  
Canada.*



**Dr. S.R Madhan Shankar**

*PSG College of Arts & Science  
Civil Aerodrome Post, Coimbatore-641 014, India.*

**Prof. Dr. Safdar Hussain Shah**

*Institute of Biotechnology and Genetic Engineering  
NWFP, Agricultural University Peshawar, Pakistan.*

**Prof. Dr. Md. Shahidul Haque**

*Dept. of Biochemistry and Molecular Biology  
University of Rajshahi, Rajshahi-6205, Bangladesh.*

**Dr. Sivakumar Swaminathan**

*Iowa State University (ISU)  
G-319, Agronomy Department, ISU, Ames, Iowa -  
50011, USA*

**Dr. Subrahmanyam Desiraju**

*Directorate of Rice Research (ICAR)  
Plant Physiology Division, Rajendranagar,  
Hyderabad-500030, A.P. India.*

**Dr. Tariq Aziz Dr. Deepak Ganjewala**

*University of Agriculture, Faisalabad, Sub-Campus  
Depalpur, Dist. Okara, Pakistan.*

**Dr. Thangavel Palaniswamy**

*SUN YAT-SEN UNIVERSITY  
GUANGZHOU, PR CHINA.*

**Yi-Ping Chen Ph.D**

*Institute of Earth Environment,  
Chinese Academy of Science Fenghui S.R, 10,  
Xi'an Hi-Tech Zone, Xi'an, China.*

**Saha Prasenjit**

*The Samuel Roberts Noble Foundation  
2510 Sam Noble Parkway, Ardmore, Ok USA.*

**Abdul Khaliq Ph.D**

*Department of Agronomy  
University of Agriculture  
Faisalabad 38040, Pakistan.*

**Dr. Arafat Abdel Hamed Abdel Latef**

*Assistant Professor of Plant physiology  
Botany Department  
Faculty of Science at Qena  
South Valley University  
Egypt.*

**Dr. Ahmad Bybordi**

*Research Center of Agriculture and Natural  
Resources of East Azarbaijan  
Member of Scientific Board of Research Center of  
Agriculture and Natural Resources of East  
Azarbaijan,  
Tabriz. Iran.*

**Dr. Arijit Sinhababu**

*Bankura Christian College (under –The University of  
Burdwan)  
Department of Botany, Bankura Christian College,  
P.O. + Dist. Bankura, Pin.-722101, West, Bengal,  
India.*

**Dr. Maria Alejandra Equiza**

*University of Alberta,  
4-51 Earth Sciences Building, Dept. Renewable  
Resources, University of Alberta, Edmonton, AB T6G  
2E3,  
Canada.*

**Dr. Suphla Bajpai Gupta**

*Indian Institute of integrative Medicine –CSIR,  
Scientist, Plant biotechnology division, Canal Road,  
Jammu, Jammu & Kashmir, India-180001. India.*

**Dr. Linga R Gutha**

*Washington State University,  
2410 N Bunn Road, Prosser, WA 99350,  
USA.*

**Dr. Medhat Mekhail Tawfik**

*National Research Center,  
El Bohooth Str. Dokki, Giza. Egypt, PO Box 12311,  
Egypt.*

**Dr. Rafiq Islam**

*The Ohio State University South Centers,  
1864 Shyville Road, Piketon, OH 45661.*

**Dr. Rakesh Kumar**

*V.S.P. Govt. P.G. College, Kairana, Muzaffarnagar  
(Uttar Pradesh),  
Department of Botany, V.S.P. Govt. P.G. College,  
Kairana, Muzaffarnagar (Uttar Pradesh), India-  
247774.*

**Dr. Ivan Sestari**

*University of São Paulo,  
Av. Pádua Dias, 11: CP 9. CEP 13418-900.*

**D.Sc. Rachel Fatima Gagliardi**

*State University of Rio de Janeiro,  
Rua São Francisco Xavier, 524 – PHLC sala 602.*

**Dr. Ullas Pedmale**

*Salk Institute for Biological Studies,  
10010 N Torrey Pines RD, La Jolla, CA 92037.*

**Dr. Allah Bakhsh Dr. Deepak Ganjewala**

*Department of Field Crops, Faculty of Agriculture,  
University of Ankara,  
Apartment No. 12/10, Sanatorym Caddesi, Kalaba,  
Kecioren, Ankara,  
Turkey.*

**Dr. Atilgan Atilgan**

*Suleyman Demirel University, Agriculture Faculty,  
Department of Agricultural Structures and  
Irrigation, Isparta,  
Turkey.*

**Mr. Andrej Pilipovic**

*University of Novi Sad – Institute of Lowland  
Forestry and Environment,  
Antona Cehova 13, 21000 Novi Sad,  
Serbi.*

**Dr. Zulfiqar Ahmad Saqib**

*Institute of Soil and Environmental Sciences,  
University of Agriculture, Faisalabad,  
Civil Line Road, Faisalabad,  
Pakistan.*

**MS. C. Mehrnoush Eskandari Torbaghan**

*North Khorasan Agricultural & Natural Resource  
Research Center (NKANRRC)  
P.O. Box: 94155-1416, No. 52, Hassan Kallate  
Alley, Tarbiyat St., Mother Sq. Bojnourd,  
Iran.*

**Dr. Vinod Kumar**

*Department of Zoology & Environmental Science,  
Gurukula Kangri University, Haridwar-249404  
(UK), India.*

**Dr . Panda Tribhubana**

*Kalahandi Institute for Tribiology and  
Ethnobiology(KITE),  
At-Jilingdar, PO-Deydar, Dist-Kalahandi, Odisha,  
India,766014, India*

**Dr. Sabarinath Sundaram**

*Institute of Developmental and Molecular Biology,  
Texas A&M University,  
Biological Sciences Building West Suite 403.*

**Dr. Diogo Pineda Rivelli**

*University of São Paulo,  
Av. Prof. Lineu Prestes 580, São Paulo, SP, 05508-000.*

**Dr. Qiang Wang**

*Virginia Tech,  
427 Latham Hall.*

**Dr. Foteini Hassiotou**

*University of Western Australia,  
35 Stirling Highway, Crawley, WA 6009,  
Australia.*

**Dr. Nivedita Sahu**

*Indian Institute of Chemical Technology,  
Chemical Biology Laboratory  
(NaturalProductChemistry), Uppal Road,  
Hyderabad-500607.*

**Dr. Mohammad Anwar Hossain**

*Bangladesh Agricultural University,  
Assistant Professor, Dept. of Genetics and Plant  
Breeding, Bangladesh Agricultural University,  
Mymensingh-2202, Bangladesh.*

**Dr. Ahmad Ali**

*National Institute of Pharmaceutical Education &  
Research,  
Dept of Biotechnology, NIPER, Jandaha Road,  
Hajipur, Bihar, India, Pin – 844102, India.*

**Mr. Karthikkumar V**

*Annamalai University,  
Department of Biochemistry & Biotechnology.*

**Dr.K.Rajendiran**

*Dept of plant science, Tagore Govt. college,  
9, 4th cross, Tagore Nagar, Pondicherry – 605 008,  
India.*

**Dr. V.Balakrishnan**

*K.S.Rangasamy College of Technology,  
Department of Biotechnology, KSR Kalvi  
nagar, Tiruchengode-637215, Tamilnadu,  
India.*

**Dr. NourAli Sajedi**

*Department of Agronomy and plant Breeding,  
Islamic Azad University, Arak Branch, Arak,  
Iran.*

**(Dr) Ms. Rachel Predeepa**

*Not Applicable ,  
2/387 Gokul Nagar, Kannanenthal Madurai.*

**Dr. Rajendra Gyawali**

*Department of Pharmacy and Biology, Kathmandu  
University,  
Dhulikhel,  
Nepal.*

**Ms. Rocheli de Souza**

*UFRGS,  
Porto Alegre,  
Brazil.*

**Dr. Om Prakash Verma**

*Sam Higginbottom Institute of Agriculture,  
Technology & Sciences (Formerly Allahabad  
Agricultural Institute), Allahabad, U.P.,  
Department of Molecular & Cellular Engineering,  
Jacob School of Biotechnology & Bioengineering,  
India.*

**Dr. Ashwani Kumar**

*JMIT, Radaur,  
Department of Biotechnology, JMIT, Radaur-  
135133, Haryana, India.*

**Dr. Sarfaraz F. A. Al-Bamarny**

*University of Duhok, College of Agriculture, Dept.  
of Horticulture,  
Duhok, Iraqi Kurdistan Region,  
Iraq.*

**Prof. Wafaa Mohamed Shukry Abdel Meamem**

*Dammam University - Saudi Arabia,  
Faculty of Science for Girl.  
Biology Department,  
P.O.Box: 838 Dammam 31113,  
Saudi Arabia.*

**Dr. Stephka G. Chankova**

*Institute of Biodiversity and ecosystem Research,  
BAS,  
2 Gagarin str, 1113 Sofia, Bulgaria.*

**Dr. Nana Ewusi-Mensah**

*Kwame Nkrumah University of Science and  
Technology,  
Dept. of Crop and Soil Sciences, Faculty of  
Agriculture, KNUST, Kumasi.*

**Dr. Mukesh Lokanath Chavan**

*K.r.c. College of horticulture, arabhavi 591 310,  
karnataka,  
University of horticultlura sciences, bagakot,  
India.*

**Dr. Maiti Parthapratim**

*Dept. of Botany Midnapore College, Midnapore-  
721101, Paschim Medinipur, West Bengal, India.*

**Mr. Mohammad Anwar Hossain**

*Kagawa University (Present), Bangladesh  
Agricultural University (Permanent)  
Lab. of Plant Stress Responses, Faculty of  
Agriculture, Kagawa University, Miki-cho, Kita-gun,  
Kagawa 761 0795, Japan.*

**Dr. Antonia Tathiana Batista Dutra**

*Universidade Federal do Ceará,  
Av. Humberto Monte s/n – Pici Bloco 907,  
laboratório 1080.*

**Dr. Kuntal Das**

*St. John's Pharmacy College,  
#6, II Main, 9th Cross, Vijayanagar, Bangalore-104.  
India.*

**Dr. Amitava Rakshit**

*Banaras Hindu University,  
Department of Soil Science & Agril Chemistry.*

**Dr. Kranthi Kiran Mandadi**

*Texas A&M University,  
2132 TAMU, Peterson-Rm408, College Station,  
Texas-77840,  
USA.*

**Dr. Monica Butnariu**

*Banat's University of Agricultural Sciences and  
Veterinary Medicine from Timisoara,  
Chemistry and Vegetal Biochemistry Department,  
Calea Aradului no.119, 300645 Timisoara,  
Romania.*

**Dr. Ahmad Bybordi**

*East Azarbaijan Research Center for Agriculture and Natural Resources, Tabriz, Iran.*

**Dr. Haiwei Gu**

*903 Fifth St., West Lafayette, IN 47906.*

**Dr. Hu Yanbo**

*Northeast Forestry University, 26# Hexing Road, Xiangfang District, Harbin city, 150040, P.R., China.*

**Dr. Arash Kianianmomeni**

*Institute of Biology / Humboldt-University Berlin, Invalidenstr. 42.*

**Dr. Zvonko Pacanoski**

*Faculty for Agriculture Sciences and Food, Boul. Aleksandar Makedonski bb, 1000 Skopje, R.of Macedonia..*

**Dr. Lingjuan Zheng**

*Department of Organismic Biology, University of Salzburg, Hellbrunnenstraße 34, 5020, Salzburg, Austria.*

**Dr. Md. Mokter Hossain**

*Department of Horticulture, Bangladesh Agricultural University, Mymensingh-2202, Bangladesh.*

**Dr. Forouzandeh Soltani**

*Department of Horticultural Sciences, College of Agriculture and Natural Resources, University, of Tehran, Daneshkadeh Street, Karaj 31587-11167, Iran.*

**Dr. M.C.Harish**

*Bharathiar University, Department of Biotechnology, Coimbatore, India.*

**Dr. Zong-shen Zhang**

*School of Biological Engineering, Dalian Polytechnic University, Qinggongyuan, Ganjingzi District, Dalian, China, postcode 116034.*

**Prof. T. V. Ramana Rao**

*B R Doshi School of Biosciences, Sardar Patel University, Vallabh Vidyanagar, Gujarat,India.*

**Dr. Sanjeev Chandel**

*Baba Isher Singh Institute of Sciences & Technology, Gagra (Moga), Punjab, India.*

**Dr. Kuladip Jana**

*Bose Institute Centenary Campus, P 1/12, C.I.T. Scheme VIIM, Kolkata-700 054, India.*

**Prof. Ljubinko Jovanovic**

*University Educons, Faculty for ecological agriculture, Sremska Kamenica, Vojvode Putnika 87, Serbia.*

**Dr. Luis F. Goulao**

*Instituto de Investigacao Cientifica Tropical [Tropical Research Institute] Eco-Bio / IICT, Av. da Republica - Quinta do Marques, 2784-505 Oeiras, Portugal.*

**Dr. Lucky K. Attri**

*College of Punjabi University Patiala, E-41, Sector-14, Panjab University, Chandigarh.*

**Prof. Bassam Taha Yasseen**

*Looking for appropriate employer, Flat 307 Point Red, 146 Midland Road, Luton, LU2 0BL, UK.*

**Dr. Massimo Piccotto**

*Tecna S.r.l., Area Science Park, Loc. Padriciano, 99, I-34149 Trieste, Italy.*

# Instructions for Author

**Electronic submission** of manuscripts is strongly encouraged, provided that the text, tables, and figures are included in a single Microsoft Word file (preferably in Arial font).

The **cover letter** should include the corresponding author's full address and telephone/fax numbers and should be in an e-mail message sent to the Editor, with the file, whose name should begin with the first author's surname, as an attachment.

## Article Types

Three types of manuscripts may be submitted:

**Regular articles:** These should describe new and carefully confirmed findings, and experimental procedures should be given in sufficient detail for others to verify the work. The length of a full paper should be the minimum required to describe and interpret the work clearly.

**Short Communications:** A Short Communication is suitable for recording the results of complete small investigations or giving details of new models or hypotheses, innovative methods, techniques or apparatus. The style of main sections need not conform to that of full-length papers. Short communications are 2 to 4 printed pages (about 6 to 12 manuscript pages) in length.

**Reviews:** Submissions of reviews and perspectives covering topics of current interest are welcome and encouraged. Reviews should be concise and no longer than 4-6 printed pages (about 12 to 18 manuscript pages). Reviews are also peer-reviewed.

## Review Process

All manuscripts are reviewed by an editor and members of the Editorial Board or qualified outside reviewers. Authors cannot nominate reviewers. Only reviewers randomly selected from our database with specialization in the subject area will be contacted to evaluate the manuscripts. The process will be blind review.

Decisions will be made as rapidly as possible, and the journal strives to return reviewers' comments to authors as fast as possible. The editorial board will re-review manuscripts that are accepted pending revision. It is the goal of the AJFS to publish manuscripts within weeks after submission.

## Regular articles

All portions of the manuscript must be typed double-spaced and all pages numbered starting from the title page.

**The Title** should be a brief phrase describing the contents of the paper. The Title Page should include the authors' full names and affiliations, the name of the corresponding author along with phone, fax and E-mail information. Present addresses of authors should appear as a footnote.

**The Abstract** should be informative and completely self-explanatory, briefly present the topic, state the scope of the experiments, indicate significant data, and point out major findings and conclusions. The Abstract should be 100 to 200 words in length. Complete sentences, active verbs, and the third person should be used, and the abstract should be written in the past tense. Standard nomenclature should be used and abbreviations should be avoided. No literature should be cited.

Following the abstract, about 3 to 10 key words that will provide indexing references should be listed.

A list of non-standard **Abbreviations** should be added. In general, non-standard abbreviations should be used only when the full term is very long and used often. Each abbreviation should be spelled out and introduced in parentheses the first time it is used in the text. Only recommended SI units should be used. Authors should use the solidus presentation (mg/ml). Standard abbreviations (such as ATP and DNA) need not be defined.

**The Introduction** should provide a clear statement of the problem, the relevant literature on the subject, and the proposed approach or solution. It should be understandable to colleagues from a broad range of scientific disciplines.

**Materials and methods** should be complete enough to allow experiments to be reproduced. However, only truly new procedures should be described in detail; previously published procedures should be cited, and important modifications of published procedures should be mentioned briefly. Capitalize trade names and include the manufacturer's name and address. Subheadings should be used. Methods in general use need not be described in detail.

**Results** should be presented with clarity and precision. The results should be written in the past tense when describing findings in the authors' experiments. Previously published findings should be written in the present tense. Results should be explained, but largely without referring to the literature. Discussion, speculation and detailed interpretation of data should not be included in the Results but should be put into the Discussion section.

**The Discussion** should interpret the findings in view of the results obtained in this and in past studies on this topic. State the conclusions in a few sentences at the end of the paper. The Results and Discussion sections can include subheadings, and when appropriate, both sections can be combined.

**The Acknowledgments** of people, grants, funds, etc should be brief.

**Tables** should be kept to a minimum and be designed to be as simple as possible. Tables are to be typed double-spaced throughout, including headings and footnotes. Each table should be on a separate page, numbered consecutively in Arabic numerals and supplied with a heading and a legend. Tables should be self-explanatory without reference to the text. The details of the methods used in the experiments should preferably be described in the legend instead of in the text. The same data should not be presented in both table and graph form or repeated in the text.

**Figure legends** should be typed in numerical order on a separate sheet. Graphics should be prepared using applications capable of generating high resolution GIF, TIFF, JPEG or Powerpoint before pasting in the Microsoft Word manuscript file. Tables should be prepared in Microsoft Word. Use Arabic numerals to designate figures and upper case letters for their parts (Figure 1). Begin each legend with a title and include sufficient description so that the figure is understandable without reading the text of the manuscript. Information given in legends should not be repeated in the text.

**References:** In the text, a reference identified by means of an author's name should be followed by the date of the reference in parentheses. When there are more than two authors, only the first author's name should be mentioned, followed by 'et al'. In the event that an author cited has had two or more works published during the same year, the reference, both in the text and in the reference list, should be identified by a lower case letter like 'a' and 'b' after the date to distinguish the works.

Examples:

Abayomi (2000), Agindotan et al. (2003), (Kelebeni, 1983), (Usman and Smith, 1992), (Chege, 1998;

1987a,b; Tijani, 1993,1995), (Kumasi et al., 2001) References should be listed at the end of the paper in alphabetical order. Articles in preparation or articles submitted for publication, unpublished observations, personal communications, etc. should not be included in the reference list but should only be mentioned in the article text (e.g., A. Kingori, University of Nairobi, Kenya, personal communication). Journal names are abbreviated according to Chemical Abstracts. Authors are fully responsible for the accuracy of the references.

Examples:

Chikere CB, Omoni VT and Chikere BO (2008). Distribution of potential nosocomial pathogens in a hospital environment. *Afr. J. Biotechnol.* 7: 3535-3539.

Moran GJ, Amii RN, Abrahamian FM, Talan DA (2005). Methicillinresistant *Staphylococcus aureus* in community-acquired skin infections. *Emerg. Infect. Dis.* 11: 928-930.

Pitout JDD, Church DL, Gregson DB, Chow BL, McCracken M, Mulvey M, Laupland KB (2007). Molecular epidemiology of CTXM-producing *Escherichia coli* in the Calgary Health Region: emergence of CTX-M-15-producing isolates. *Antimicrob. Agents Chemother.* 51: 1281-1286.

Pelczar JR, Harley JP, Klein DA (1993). *Microbiology: Concepts and Applications.* McGraw-Hill Inc., New York, pp. 591-603.

### Short Communications

Short Communications are limited to a maximum of two figures and one table. They should present a complete study that is more limited in scope than is found in full-length papers. The items of manuscript preparation listed above apply to Short Communications with the following differences: (1) Abstracts are limited to 100 words; (2) instead of a separate Materials and Methods section, experimental procedures may be incorporated into Figure Legends and Table footnotes; (3) Results and Discussion should be combined into a single section.

Proofs and Reprints: Electronic proofs will be sent (e-mail attachment) to the corresponding author as a PDF file. Page proofs are considered to be the final version of the manuscript. With the exception of typographical or minor clerical errors, no changes will be made in the manuscript at the proof stage.

**Fees and Charges:** Authors are required to pay a \$550 handling fee. Publication of an article in the International Journal of Plant Physiology and Biochemistry is not contingent upon the author's ability to pay the charges. Neither is acceptance to pay the handling fee a guarantee that the paper will be accepted for publication. Authors may still request (in advance) that the editorial office waive some of the handling fee under special circumstances

**Copyright: © 2014, Academic Journals.**

All rights Reserved. In accessing this journal, you agree that you will access the contents for your own personal use but not for any commercial use. Any use and or copies of this Journal in whole or in part must include the customary bibliographic citation, including author attribution, date and article title.

Submission of a manuscript implies: that the work described has not been published before (except in the form of an abstract or as part of a published lecture, or thesis) that it is not under consideration for publication elsewhere; that if and when the manuscript is accepted for publication, the authors agree to automatic transfer of the copyright to the publisher.

**Disclaimer of Warranties**

In no event shall Academic Journals be liable for any special, incidental, indirect, or consequential damages of any kind arising out of or in connection with the use of the articles or other material derived from the IJPPB, whether or not advised of the possibility of damage, and on any theory of liability.

This publication is provided "as is" without warranty of any kind, either expressed or implied, including, but not limited to, the implied warranties of merchantability, fitness for a particular purpose, or non-infringement. Descriptions of, or references to, products or publications does not imply endorsement of that product or publication. While every effort is made by Academic Journals to see that no inaccurate or misleading data, opinion or statements appear in this publication, they wish to make it clear that the data and opinions appearing in the articles and advertisements herein are the responsibility of the contributor or advertiser concerned. Academic Journals makes no warranty of any kind, either express or implied, regarding the quality, accuracy, availability, or validity of the data or information in this publication or of any other publication to which it may be linked.

# International Journal of Plant Physiology and Biochemistry

Table of Content: Volume 6 Number 6, July 2014

## ARTICLES

**The determination of vitamin C, total phenol and antioxidant activity of some commonly cooking spices crops used in West Bengal**

Manas Denre



Full Length Research Paper

## The determination of vitamin C, total phenol and antioxidant activity of some commonly cooking spices crops used in West Bengal

Manas Denre

Department of Agricultural Biochemistry, Bidhan Chandra Krishi Viswavidyalaya, Mohanpur, Nadia-741252, West Bengal, India.

Received 6 June, 2014; Accepted 23 July, 2014

In the present study, vitamin C, total phenols and antioxidant activity of some common cooking spices crops used in West Bengal were analyzed. Ten (10) spices (onion, chilli, garlic, ginger, turmeric, mustard seed, cumin seed, clove, cardamom and cassia leaf) were selected in order to determine the concentration of ascorbic acid (AA), total phenol (TP) and antioxidant activity as DPPH radical scavenging activity (DPPHRAC). The results obtained show that the values of different variables varied from 5.55 to 0.08 mg g<sup>-1</sup> (AA), 21.55 to 7.67 GAE mg g<sup>-1</sup> (TP) and 0.18 to 5.99 IC<sub>50</sub> value: mg ml<sup>-1</sup> (DPPHRAC), respectively. There were negative correlations between TPC and IC<sub>50</sub> value of DPPHRAC.

**Key words:** Vitamin C, total phenols, antioxidant, spices crops.

### INTRODUCTION

Food provides not only essential nutrients needed for life but also other bioactive compounds for health promotion and disease prevention. Thus, consumers demand for natural foods with antioxidant property, which enhances health and food preservation. There is a great demand for antioxidants that are derived from natural resources. Also, there is increased demand for natural dietary products which produce antioxidants in the body (Barlow, 1990; Rice-Evans et al., 1997).

Antioxidants prevent oxidative damage caused by free radicals; they interfere with the oxidation process by reacting with free radicals, chelating, and catalytic metals and also by acting as oxygen scavengers (Shahid and

Wanasundara, 1992; Buyukokuroglu et al., 2001). One of the paradoxes of life on this planet is that the molecule that sustains aerobic life, oxygen, is not only fundamentally essential for energy metabolism and respiration, but also has implication in many diseases and degenerative conditions (Marx, 1985). A common element in such diverse human disorders such as ageing, arthritis, cancer, Lou Gehrig's disease and many others is the involvement of partially reduced form of oxygen. Oxygen can accept single electron to form unstable derivatives referred to as Reactive Oxygen Species (ROS) or Active Oxygen Species (AOS) produced during various metabolic cellular processes. ROS include free

E-mail: sanama06020803@sify.com.

Author(s) agree that this article remain permanently open access under the terms of the [Creative Commons Attribution License 4.0 International License](http://creativecommons.org/licenses/by/4.0/)

radicals such as superoxide radical anion ( $O_2^{\cdot-}$ ), hydroxyl radical ( $\cdot OH$ ), various peroxy radicals ( $ROO^{\cdot}$ ) and non-free radicals such as hydrogen peroxide ( $H_2O_2$ ), singlet oxygen ( $^1O_2$ ) and hypochlorous acid ( $HOCl$ ) (Halliwell, 1995; Odukoya et al., 2005). Normal range for the production of ROS helps in the regulation of cell proliferation, intercellular signalization, phagocytosis and also synthesis of biologically active compounds. But hyper production of ROS develops oxidative stress which causes oxidative damage to various biomolecules including proteins, lipids, lipoproteins and DNA (Farber, 1994). This oxidative damage is a critical etiological factor implicated in several chronic human diseases such as cardiovascular diseases, diabetes, tumors, rheumatoid arthritis, epilepsy, mutagenesis, carcinogenesis, arterio-sclerosis, Alzheimer's disease, tissue injury (Halliwell and Gutteridge, 1989; Alho and Leinonen, 1999).

Spices are various parts (bud, fruit, seed, bark, rhizome, leaf and bulb) of plants used as a flavoring or seasoning, although many can also be used as herbal medicine. The term 'spice' originated from the Latin word 'species', meaning of specific kind. A closely related term, 'herb' is used to distinguish plant parts having the same functions. Spices have many functions in food. Primarily they are used for flavoring food products; in addition they are also used for preservation of food and for providing nutritional and health benefits (Nazeeem, 1995).

The two terms may be used for the same plants, that is, 'spices' and 'herb'. They have been investigated in recent scientific developments throughout the world, one due to their flavor, color, nutritional, health, or preservative effects and another due to their potent antioxidant activities. As per literature survey, so many researchers have reported that chilli, onion, garlic, ginger, turmeric, mustard seed, cumin seed, clove, cardamom and cassia leaf are used as common spices cum herb crops from ancient times.

Apart from providing basic nutrition, these crops are well known for their health benefits; antioxidant constituents and antioxidant activities have been reported in chilli (Bhattacharya et al., 2010; Biswas et al., 2011; Denre et al., 2013a, 2014), onion (Benkeblia, 2005; Denre, et al., 2011), ginger (Hinneburg et al., 2006; Kim et al., 2007; Kota et al., 2008; Suhaj and Horvathova, 2007), garlic (Agarwal, 1996; Banerjee et al., 2003; Denre et al., 2013b) turmeric (Kaur and Kapoor, 2002; Tangkanakul et al., 2009), mustard seed (Dande and Manchala, 2011; Dubie et al., 2013), cumin seed (Dua et al., 2012; Nadeem and Riaz, 2012), clove (Lee and Shibamoto, 2001; Suhaj and Horvathova, 2007; Devi et al., 2012), cardamom (Devi et al., 2012) and cassia leaf (Yong et al., 2012). Foods rich in antioxidants play a role in preventing cardiovascular diseases, cancers (Garber et al., 2002), neurodegenerative diseases (Di-Matteo and Esposito, 2003), inflammation and problems caused by cell and cutaneous aging (Ames et al., 1993). The aim of the present work was to study the determination of

vitamin C, total phenols and antioxidant activity of some commonly cooking used spices crops, which may be added in its right (proper value) while cooking.

## MATERIALS AND METHODS

The 10 spices crops [Chilli (*Capsicum annum* L.), onion (*Allium capa* L.), garlic (*Allium sativum* L.), Ginger (*Zingiber officinale* L.), turmeric (*Curcuma Longa* L.), mustard seed (*Brassica Juncea*), cumin seed (*cuminum cyminum*), clove (*Syzygium aromaticum* L.), cardamom (*Elattaria cardamomum*), cassia leaf (*Cinnamomum cassia* L.)] were collected from the market (Shyampur market, Uluberia, Howrah, West Bengal, India) for the experiment. All the collected samples were washed, air dried in shade and then samples were oven dried at 40°C for 96 h until constant weight was gained. These dried materials were prepared for chemical analyses by grinding into a fine powder using an electric grinder. The powder of each sample was stored at three replications in an air-tight cellophane bag as stock sample in a refrigerator (4°C) until further analysis. All the analytical work was performed at Agricultural Biochemistry laboratory, Department of Agricultural Biochemistry, Bidhan Chandra Krishi Viswavidyalaya, Mohanpur, Nadia-741252, West Bengal, India.

### Determination of ascorbic acid content (AAC)

Ascorbic acid was determined by the 2, 6-dichlorophenol indophenol (DCPIP) titration procedures based on the method of Casanas et al. (2002). The 0.1 g powdered sample was extracted with 20 ml of 4% oxalic acid. Then the material was centrifuged at 10,000 rpm for 30 min. 10 ml of sample's aliquot and 10 ml of 4% oxalic acid were taken in a conical flask and titrated against 2,6-dichlorophenol indophenol (DCPIP) dye ( $V_2$ ) until the appearance of a faint pink colour that persisted for a few minutes. Another 5 ml of 100 ppm solution of ascorbic acid and 10 ml of 4% oxalic acid were taken and also titrated against DCPIP dye ( $V_1$ ). The ascorbic acid content ( $mg\ g^{-1}$ ) was determined using the following formula:

$$\text{Amount of ascorbic acid (mg g}^{-1}\text{)} = \frac{0.5 \times V_2}{V_1 \times 10 \times 0.1}$$

$V_1$  = dye consumed by 0.5 mg ascorbic acid;  $V_2$  = dye consumed by 10 ml of test solution.

### Determination of total phenols content (TPC)

The total phenols content in spices crops were determined by the Folin-Ciocalteu reagent (Vinson et al., 1998) using gallic acid as standards. The 0.1 g powdered samples was extracted with 15 ml of 1.2 M HCl in 50% aqueous methanol and heated at 85°C for 2 h to measure the conjugated plus unconjugated ('total') phenols. After cooling the extracted material was centrifuged at 10,000 rpm for 30 min. The supernatant was decanted off in a beaker and evaporated to dryness. The crude extract was diluted to 25 ml with distilled water. For estimation of the total phenols content 0.5 ml of sample's aliquot, 2.5 ml of distilled water and 0.5 ml Folin-Ciocalteu reagent were pipetted out in a test tube. After 3 min, 2 ml of 10% sodium carbonate was added. Then the test tubes were kept on water bath and maintained at 60-70°C for 5 min. The solution was cooled and the absorbance was read at 650 nm. The total phenols content was measured ( $mg\ g^{-1}$ ) using a calibration curve against gallic acid equivalent.

**Table 1.** Concentration of vitamin C, total phenols and antioxidant activity of some commonly cooking spices crops used in West Bengal.

Spices crop	Scientific name	Part used	AAC <sup>n</sup> (mg g <sup>-1</sup> )	TPC (mg g <sup>-1</sup> )	DPPH (IC <sub>50</sub> value: mg ml <sup>-1</sup> )
Onion	<i>Allium capa</i> L.	Bulb	0.62±0.09	7.67±0.55	2.28±0.08
Chilli	<i>Capsicum annuum</i> L.	Pod	5.55±0.55	8.30±0.67	1.26±0.15
Garlic	<i>Allium sativum</i> L.	Bulb	0.37±0.08	14.30±0.69	1.12±0.03
Ginger	<i>Zingiber officinale</i> L.	Rhizome	0.48±0.03	15.52±0.96	1.11±0.02
Turmeric	<i>Curcuma Longa</i> L.	Rhizome	0.75±0.06	13.00±0.51	5.99±0.68
Mustard	<i>Brassica Juncea</i>	Seed	0.08±0.03	13.79±0.79	0.66±0.05
Cumin	<i>cuminum cyminum</i>	Seed	0.09±0.02	20.32±1.32	2.40±0.07
Clove	<i>Syzygium aromaticum</i> L.	Bud	0.14±0.05	21.55±0.67	0.26±0.03
Cardamom	<i>Elattaria cardamomum</i>	Pod	0.27±0.06	16.00±1.00	0.18±0.05
Cassia	<i>Cinnamomum cassia</i> L.	Leaf	0.76±0.12	10.51±0.51	1.09±0.09

Data are mean ± standard deviation values.

#### Determination of 1,1-diphenyl-2-picrylhydrazyl radical scavenging activity (DPPHRAC)

The 1,1-diphenyl-2-picrylhydrazyl (DPPH) radical scavenging activity was performed based on the method of Braca et al. (2001). In order to determine the DPPH radical scavenging activity present in spices crops, 0.5 g powdered samples was extracted with 10 ml of distilled water. Extracted sample was centrifuged at 10,000 rpm for 15 min. Each 6.4 ml reaction mixture containing 0.1-0.4 ml of sample's aliquot, 0.1-0.4 ml of distilled water, and 6 ml of 0.004% DPPH were taken in tubes and shaken by hand. They were kept for 30 min at room temperature in dark place. Absorbance of the reaction mixture was read at 517 nm against the blank. The reaction mixture lacking sample developed the most intense colour. The colour decreased with increasing volume of extract added. The ability to scavenge the DPPH radical was calculated using the following equation: DPPH radical scavenging activity (% inhibition) =  $[(A_0 - A_e)/A_0] \times 100$ , where  $A_e$  was the absorbance of sample and  $A_0$  was the absorbance of the control. The % inhibition value was plotted against concentration of the sample extract. From the graph, value of the sample extract producing 50% inhibition (IC<sub>50</sub> value) was calculated.

#### Statistical analysis

Data were analyzed statistically by using Daniel's XL Toolbox 6.52 software for analysis of variance and correlation.

## RESULTS AND DISCUSSION

### Ascorbic acid

Ascorbic acid is one of the most powerful antioxidants (Smirnoff, 1996; Noctor and Foyer, 1998; Arrigoni and de Tullio, 2000; Horemans et al., 2000b; Smirnoff, 2000) that scavenge harmful free radicals and other ROS; it also regenerates other antioxidants like tocopherol to its functional state. In the present study (Table 1), the wide remarkable variation in ascorbic acid concentration was shown among the spices crops that varied from 5.55 to 0.08 mg g<sup>-1</sup>; and ranking from high to low concentration was chilli>cassia leaf>turmeric>onion>ginger>garlic>cardamom>clove>cumin seed>mustard seed, respectively.

The recommended daily allowance of ascorbic acid is 60 mg in USA. However, Australia and United Kingdom recommend 30 to 40 mg; Russia recommends 100 mg. Higher intake of ascorbic acid has been linked to lower risk of cardiovascular diseases (Simon et al., 1988) and several types of cancers (Howe et al., 1990) by forming N-nitroso compounds in stomach and by stimulating the immune system (Byers and Perry, 1992). Vitamin C deficiency exacerbates atherogenesis in animal models. Risk of oesophageal, pancreatic and lung cancer also appears to be lower in people that take ample vitamin C (Wargovich, 2000).

### Total phenols

Phenolics are diverse secondary metabolites that are abundant in plant tissues (Grace and Logan, 2000). Antioxidant properties of phenols arise from their high reactivity as hydrogen or electron donors, from the ability of the phenol derived radical to stabilize and delocalize the unpaired electron (Chain breaking function) and from their ability to chelate transition metal ions (termination of the Fenton reaction) (Rice-Evans et al., 1997). Another mechanism underlying the antioxidative properties of phenolics is the ability of flavonoids to alter peroxidation kinetics by modifying the lipid picking order and decreasing the fluidity of the membranes (Arora et al., 2000). These changes could sterically hinder diffusion of free radicals and restrict peroxidation reaction. In the present experiment (Table 1), the total phenols contents ranged from 21.55 to 7.67 GAE mg g<sup>-1</sup> with decreasing order: clove>cumin

seed>cardamom>ginger>garlic>mustard seed>turmeric>cassia leaf>chilli>onion, respectively. The health benefit of phenolics is linked primarily to their antioxidant potential. Phenolics are effective antioxidants because the radicals products of these molecules are resonance stabilized and thus relatively stable. To

**Table 2.** Correlation matrix among variables.

Variable	AAC	TPC
AAC		
TPC	-0.541	
DPPHRAC (IC <sub>50</sub> value)	0.005	-0.194

overcome the potential hazard from oxidative damage in the body, consumption of a diet rich in antioxidant phenolics including flavonoids and phenolic acids is considered the first line of defense against highly reactive toxicants. Scalbert and Williamson (2000) estimated that the total human intake of phenolics is about 1 g day<sup>-1</sup> which consists of two-thirds flavonoids and one-third phenolic acids.

### DPPHRAC

DPPH analysis is one of the best-known, accurate, and frequently employed methods for evaluating antioxidant activity (Zhou and Yu, 2004). It is a stable free radical because of its spare electron delocalization over the whole molecule. In the present experiment (Table 1), it is revealed that the DPPH radicals scavenging activity ranges from 0.18 to 5.99 mg ml<sup>-1</sup> following the downward activity order: Cardamom>clove>mustard seed>cassia leaf>ginger>garlic>chilli>onion>turmeric, respectively. The smaller the IC<sub>50</sub> value the greater the radical scavenging activity and reducing ability of the extracts.

### Correlation among variables

Table 2 did not show any significant correlation among the variables, but apparently there were negative correlations between AAC with TPC and TPC with IC<sub>50</sub> values of DPPHRAC. Interestingly, the negative correlation of TPC with IC<sub>50</sub> value of DPPHRAC indicated that the higher total phenol content of extracts resulted in lower IC<sub>50</sub> value of DPPHRAC. That means the higher total phenol content of extracts resulted in higher antioxidant (DPPH radical scavenging activity) activity, because the lower IC<sub>50</sub> value indicated the higher DPPH radical scavenging activity in extract of spices crops. Maizura et al. (2011) reported that the higher total phenolic content of extracts resulted in higher antioxidant (DPPH radical scavenging activity) activity in extracts of kesum, ginger and turmeric.

### Conclusion

Based on the mean and ANOVA results, there is a wide variation among variables in the ten selected cooking species crops. The maximum values of ascorbic acid,

total phenol and DPPH radical activity were exhibited as 5.55 mg g<sup>-1</sup> (chilli), 21.55 mg g<sup>-1</sup> (clove) and 0.18 IC<sub>50</sub> value: mg ml<sup>-1</sup> (Cardamom), respectively. Thus, it is crucial to determine not only their antioxidant capacity but also their characterization potential in promoting health.

### Conflict of Interests

The author(s) have not declared any conflict of interests.

### ACKNOWLEDGEMENTS

The author is grateful to the Department of Agricultural Biochemistry, Bidhan Chandra Krishi Viswavidyalaya, Mohanpur, Nadia-741252, West Bengal, and India for its support in carrying out the research work.

### REFERENCES

- Agarwal KC (1996). Therapeutic actions of garlic constituents. *Med. Res. Rev.* 16:111-124.
- Alho H, Leinonen J (1999). "Total antioxidant activity measured by chemiluminescence methods," *Meth. Enzymol.* 299:3-15.
- Ames BM, Shigenaga MK, Hagen TM (1993). Oxidants, antioxidants and the degenerative diseases of aging. *Proc. Nat. Acad. Sci. United States of America*, 90(17):7915-7922.
- Arora A, Byrem TM, Nair MG, Strasburg GM (2000). Modulation of liposomal membrane fluidity by flavonoids and isoflavonoids. *Arch. Biochem. Biophys.* 373:102-109.
- Arrigoni O, De-Tullio MC (2000). The role of ascorbic acid in cell metabolism: between gene directed functions and unpredictable chemical reactions. *J. Pl. Physiol.* 157:481-488.
- Banerjee SK, Mukherjee PK, Maulik SK (2003). Garlic as an antioxidant: The good, the bad and the ugly. *Phytother. Res.* 17:97-106.
- Barlow SN (1990). "Toxicological aspects of antioxidants used as food additives", Elsevier. p. 253.
- Benkeblia N (2005). Free radical scavenging capacity and antioxidant properties of some selected onion (*Allium cepa* L.) and garlic (*Allium sativum* L.) extracts. *Braz. Arch. Biol. Technol.* 48(5):753-759.
- Bhattacharya A, Chattopadhyay A, Muzumdar D, Chakravarty A, Pal S (2010). Antioxidant constituents and enzyme activities in chilli pepper. *Int. J. Veg. Sci.* 16(3):201-211.
- Biswas A, Bhattacharya A, Chattopadhyay A, Chakravarty A, Pal S (2011). Antioxidants and Antioxidant Activity in Green Pungent Peppers. *Int. J. Veg. Sci.* 17:224-232.
- Braca A, De-Tommasi N, Di-Bary L, Pizzi C, Polifi M, Morelli I (2001). Antioxidant principles from *Bauhinia terapotensis*. *J. Nat. Prod.* 64:892-895.
- Buyukokuroglu ME, Gulcin I, Oktay M, Kufrevioglu OI (2001). *In vitro* antioxidant properties of dantrolene sodium. *Pharmacol. Res.* 44:491-494.
- Byers T, Perry G (1992). Dietary Carotenes, Vitamin C, and Vitamin E as protective antioxidants in human cancers. *Ann. Rev. Nutr.* 12:139-159.
- Casanas R, Gonzalez M, Rodriguez E, Marrero A, Diaz C (2002). Chemometric studies of chemical compounds in five cultivars of potatoes from Tenerife. *J. Agric. Food Chem.* 50(7): 2076-2082.
- Dande S, Manchala R (2011). Antioxidant and Phenolic Content of Nuts, Oil Seeds, Milk and Milk Products Commonly Consumed in India. *Food Nutr. Sci.* 2:422-427.
- Denre M, Bandopadhyay PK, Chakravarty A, Pal S, Bhattacharya A (2013a). Changes in some biochemical characteristics in response to foliar applications of chelator and micronutrients in green pungent pepper. *Int. J. Plant Physiol. Biochem.* 5(2):25-35.

- Denre M, Bhattacharya A, Pal S, Chakravarty A, Chattopadhyay A, Mazumdar D (2014). Changes in antioxidants and pungency in response to foliar applications of Mn and Fe in onion. *Int. J. Plant Physiol. Biochem.* 6(2):19-27.
- Denre M, Pal S, Chattopadhyay A, Mazumdar D, Chakravarty A, Bhattacharya A, (2011). Antioxidants and pungency of onion. *Int. J. Veg. Sci.* 17:233-245.
- Denre M, Pal S, Mazumdar D, Chakravarty A, Bhattacharya A (2013b). Variation in antioxidants and antioxidant activity in garlic cultivars. *Int. J. Veg. Sci.* 19:374-383.
- Devi SA, Umasankar ME, Babu S (2012). A comparative study of antioxidant properties in common Indian spices. *Int. Res. J. Pharmacy*, 3(5):465-468.
- Di-Matteo V, Esposito E (2003). Biochemical and therapeutic effects of antioxidants in the treatment of Alzheimer's disease, Parkinson's disease, and amyotrophic lateral sclerosis. *Curr. Drug Targets- CNS Neurolog. Disord.* 2(2):95-107.
- Dua A, Gupta SK, Mittal A, Mahajan R. (2012). A Study of Antioxidant Properties and Antioxidant Compounds of Cumin (*Cuminum cyminum*). *Int. J. Pharmacol. Biolog. Arch.* 3(5):1110-1116.
- Dubie J, Slancik A, Morra M, Nindo C (2013). Antioxidant extraction from mustard (*Brassica Juncea*) seed meal using high intensity ultrasound. *J. Food Sci.* 78(4):E542-E548.
- Farber JL (1994). Mechanisms of cell injury by activated oxygen. *Environ. Health Perspect.* 102:17-24.
- Garber SL, Rintala DH, Holmes SA, Rodriguez GP, Friedman J (2002). A structured educational model to improve pressure ulcer prevention knowledge in veterans with spinal cord dysfunction. *J. Rehabil. Res. Dev.* 39(5):575-588.
- Grace, S. and Logan, B.A. (2000). Energy dissipation and radical scavenging by the plant phenylpropanoid pathway. *Philosophical Transactions of the Royal Society of London B*, 355:1499-1510.
- Halliwell B (1995), "How to characterize an antioxidant: an update". *Biochem. Soc. Symp.* 61:73-101.
- Halliwell B, Gutteridge JMC (1989). Protection against oxidants in biological system: the superoxide theory of oxygen toxicity. *In: Halliwell, B. and Gutteridge, J. M. (eds.). Free Radic. Biol. Med.* pp: 86-123. Clarendon Press, Oxford.
- Hinneburg I, Damien Dorman HJ, Hiltunen R (2006). Antioxidant activities of extracts from selected culinary herbs and spices. *Food Chem.* 97: 122- 129.
- Horemans N, Foyer CH, Potters G, Asard H (2000b). Ascorbate function and associated transport systems in plants. *Pl. Physiol. Biochem.* 38: 531-540.
- Howe GR, Jain M, Miller AB (1990). Dietary factors and risk of pancreatic cancer: results of a Canadian population based case control study. *Int. J. Cancer*, 45: 604-608.
- Kaur C, Kapoor HC (2002). Anti-oxidant activity and total phenolic content of some Asian vegetables. *Int. J. Food Sci. Technol.* 37: 153-161
- Kim JK, Kim Y, Na KM, Surh YJ, Kim TY (2007). Gingerol prevents UVB-induced ROS production and COX-2 expression *in vitro* and *in vivo*. *Free Radic. Res.* 41: 603-614.
- Kota N, Krishna P, Polasa K (2008). Alterations in antioxidant status of rats following intake of ginger through diet. *Food Chem.* 106: 991-996.
- Lee K, Shibamoto T (2001). Antioxidant property of aroma extract isolated from clove buds [*Syzygium aromaticum* (L.) Merr. et Perry]. *Food Chem.* 74:443-448.
- Maizura M, Aminah A, Wan-Aida WM (2011). Total phenolic content and antioxidant activity of kesum (*Polygonum minus*), ginger (*Zingiber officinale*) and turmeric (*Curcuma longa*) extract. *Int. Food Res. J.* 18:529-534.
- Marx JL (1985). Oxygen free radicals linked to many diseases. *Sci.* 235:529-531.
- Nadeem M, Riaz A (2012). Cumin (*Cuminum cyminum*) as a potential source of antioxidants. *Pak. J. FOOD Sci.* 22(2): 101-107.
- Nazeem PA (1995). The spices of India. *The Herb, Spice Medicinal Plant Digest* 13(1):1-5.
- Noctor G, Foyer CH (1998). Ascorbate and glutathione: keeping active oxygen under control. *Ann. Rev. Pl. Physiol. Plant Mol. Biol.* 49: 249-279.
- Odukoya AO, Ilori OO, Sofidiya MO, Aniunoh OA, Lawal BM, Tade IO (2005). Antioxidant activity of Nigerian dietary spices. *EJEAFChe.* ISSN:1579-4377.
- Rice-Evans CA, Miller NJ, Paganga G (1997). Antioxidant properties of phenolic compounds. *Trends Plant Sci.* 2:152-159.
- Scalbert A, Willinson G (2000). Dietary intake and bio-availability of polyphenols. *J. Nutr.* 130: 20735-20855.
- Shahidi F, Wanasundara PD (1992). Phenolic antioxidants. *Cri. Rev. Food Sci. Nutr.* 32: 67-103.
- Simon JP, Hudes ES, Brown WS (1988). Serum ascorbic acid and cardiovascular disease prevalence in US adults. *Epidemiology* 9:316-321.
- Smirnoff N (1996). The function and metabolism of ascorbic acid in plants. *Ann. Bot.* 78:661-669.
- Smirnoff N (2000). Ascorbic acid: metabolism and functions of a multifaceted molecule. *Curr. Opin. Pl. Biol.* 3:229-235.
- Suhaj M, Horvathova J (2007). Changes in antioxidant activity induced by irradiation of clove (*Syzygium aromaticum*) and ginger (*Zingiber officinale*). *J. Food Nutr. Res.* 46(3):112-122.
- Tangkanakul P, Auttaviboonkul P, Niyomwit B, Lowviton N, Charoenthamawat P, Trakoontivakorn G (2009). Antioxidant capacity, total phenolic content and nutritional composition of Asian foods after thermal processing. *Int. Food Res. J.* 16:571-580.
- Vinson JA, Hao Y, Su X, Zubik L (1995). Phenol antioxidant quantity and quality in foods: vegetables. *J. Agric. Food Chem.* 46(9):3630-3634.
- Wargovich MJ (2000). Anticancer properties of fruits and vegetables. *Hort. Sci.* 35:573-575.
- Yang CH, Li RX, Chuang LY (2012). Antioxidant Activity of Various parts of *Cinnamomum cassia* extracted with different extraction methods. *Molecules* 17:7294-7304.
- Zhou K, Yu L (2004). Effects of extraction solvent on wheat bran antioxidant activity estimation. *LWT-Food Sci. Technol.* 37(7):717-721.

# International Journal of Plant Physiology and Biochemistry

## *Related Journals Published by Academic Journals*

- *African Journal of Plant Science*
- *Journal of Microbiology and Antimicrobials*
- *International Journal for Biotechnology and Molecular Biology Research*
- *Journal of Botany and Plant Pathology*
- *International Journal of Biodiversity and Conservation*

**academicJournals**