

Biotechnology and Molecular Biology Reviews

Volume 10 Number 1, January 2015

ISSN 1538-2273



*Academic
Journals*

ABOUT BMBR

The **Biotechnology and Molecular Biology Reviews (BMBR)** (ISSN 1538-2273) is published Monthly (one volume per year) by Academic Journals.

Biotechnology and Molecular Biology Reviews (BMBR), a new broad-based review journal, is an open access journal that was founded on two key tenets: (1) to publish the most exciting, cutting-edge reviews in all areas of applied biochemistry, industrial microbiology, genomics and proteomics, and metabolic engineering, and (2) to provide the most rapid turn-around time possible for reviewing and publishing. It is our hope these articles will serve teaching and reference tools.

Submission of Manuscript

Please read the **Instructions for Authors** before submitting your manuscript. The manuscript files should be given the last name of the first author

[Click here to Submit manuscripts online](#)

If you have any difficulty using the online submission system, kindly submit via this email ajb@academicjournals.org.

With questions or concerns, please contact the Editorial Office at bmb@academicjournals.org.

Editor-In-Chief

P. Ravindra, Ph.D.

*School of Engineering &
IT University Malaysia Sabah
88999, Kota Kinabalu, Sabah,
Malaysia*

Editor

David Maina Menge, Ph.D.

*University of California, Irvine
College of Health Sciences/Program in Public Health
Hewitt Hall Room 3501; Irvine, CA 92697-4050, USA.*

Evans Kaimoyo, Ph.D.

*Plant Microbe-interactions Laboratory,
Boyce Thompson Institute of Plant Research, Ithaca,
NY, 14853, USA.*

Solomon Olawale Odemuyiwa, Ph.D.

*Pulmonary Research Group Department of Medicine
550 Heritage Medical Research Centre
University of Alberta
Edmonton
Canada T6G 2S2*

Vikash Kumar Dubey

*Department of Biotechnology Room No: 1N 102
Indian Institute of Technology, Guwahati
Guwahati- 781039, Assam, India.*

Beng Ti Tey

*Department of Chemical and Environmental
Engineering,
Faculty of Engineering, Universiti Putra, Malaysia*

Behera B.C.

*Agharkar Research Institute
Pune-411004, India*

Editorial Board

Dr. Daiana P.

*Stolf University of Toronto, Toronto,
Canada*

Dr. Stephane Chevaliez

*Department of Virology and INSERM,
Henri Mondor Hospital,
avenue du Maréchal de l'atter de Tassigby
Cretéil*

Dr. Mohammad Asgharzadeh

*Tuberculosis and Lung Disease Research Center and
Biotechnology Research Center,
Tabriz University of Medical Sciences,
Iran*

Dr. Mukul Das

*Food Toxicology Division Industrial Toxicology Research
Centre
Mahatma Gandhi Marg Lucknow, India*

Dr. Jian-Zhong Liu

*Biotechnology Research Center,
Zhongshan (Sun Yat-Sen) University Guangzhou,
China*

Prof. Peter J. Reilly

*Dept of Chemical and Biological Eng
Anson Iowa State University*

Prof. Mohammad Miransari

*Department of Soil Science,
College of Agricultural Sciences,
Shahed University,
Tehran, Iran*

Dr. Chhandak Basu

*School of Biological Sciences
Ross Hall University of Northern Colorado Greeley,
Colorado
USA*

Prof. Anil Kumar

*School of Biotechnology Devi Ahilya University,
Khandwa Road, Indore,
India*

Dr. Ahmed M Malki

*Alexandria University,
Faculty of Science,
Biochemistry department,
Alexandria,
Egypt*

Dr. Christopher Brigham

*Sinskey Laboratory
Massachusetts Institute of Technology
Cambridge, USA*

Prof. Mahmoud Saker

*National Research Center,
El Behouse St., Dokki, Cairo,
Egypt*

Prof. Karl Bayer

*Institute of Applied Microbiology,
University of Natural Resources and Applied Life
Sciences, Muthgasse Austria.*

Dr. Hector Budman

*University of Waterloo Waterloo,
Ontario*

Prof. Mohammad Miransari

*Shahed University,
Tehran, Iran*

Prof. R.P Singh

*University of Roorkee, Roorkee
India*

Prof. Jane B. Lian

*University of Massachusetts Medical School 55
Lake Avenue
North Worcester*

Dr. Vicki Ann Luna

University of South Florida USA

Dr. Helene F Rosenberg

*National Institute of Allergy and Infectious
Diseases
Rockville Pike, Bethesda,
USA*

Dr. Silvia Bautista-Baños

*National Polytechnic Institute Yautepec,
Morelos Cuernavaca,
México*

Dr. Stephen Bakiamoh

*Michigan Biotechnology Institute International
3900 Collins Road
Lansing, MI 48909, USA*

Dr. RA Siddique

*Department of Veterinary Biochemistry.
College of Veterinary Science and Animal Husbandry.
Navsari Agricultural University,
Navsari Gujarat, India*

Dr. Eijiro Miyak

*Health Technology Research Center,
National Institute of Advanced Industrial Science and
Technology (AIST) Takamatsu,
Japan*

Dr. Carla Marchetti

*Istituto di Biofisica.
Consiglio Nazionale delle Ricerche. via De Marini,
Genova,
Italy*

Dr. T Anjana Devi

*Centre for Chemical Biology,
Indian Institute of chemical technology
Habsiguda Hyderabad
Andhra Pradesh, India*

Dr. Moytri Roy-Chowdhury

*Washington State University- Pullman,
Washington
USA*

Dr. Poluri Krishna Mohan

*University of Texas
Department of Biochemistry and Molecular Biology
and The Sealy Center for Structural Biology and
Molecular Biophysics
Medical Branch Galveston, Texas*

Dr. Gao Guo

*School of Dentistry,
University of California, Los Angeles (UCLA)
10833 Le Conte Avenue Los Angeles, CA 90095, USA*

Dr. Alireza Valdiani

*Department of Biochemistry,
Faculty of Biotechnology and Biomolecular
Sciences,
Universiti Putra Malaysia,
43400 UPM Serdang,
Selangor DE, Malaysia*

Dr. S. A. Anitha Christy

*Genomic Medicine
Methodist Hospital Research Institute
8.330, Bertner Ave
Houston, TX 77030*

Dr. Sridhar Boppana

*Department of Pediatrics,
Center for Advanced Biotechnology and Medicine
(CABM),
University of Medicine and Dentistry of New Jersey
(UMDNJ),
679 Hoes Lane West, Piscataway,
New Jersey- 08854. USA*

Dr. Jeffy George

*Department of Microbiology and Immunology F.
Edward Hébert School of Medicine Uniformed
Services University of the Health Sciences
4301 Jones Bridge Road,
Bethesda, MD 20814*

Dr. Ajamaluddin Malik

*Protein Research Chair Department of Biochemistry
College of Science
Building no 5, room no 2A46
P.O. Box 2455
King Saud University
Riyadh 11451
Kingdom of Saudi Arabia*

Dr. Deepshikha Pande Katare

*Amity Institute of Biotechnology,
AUUP, Noida, India*

Dr. Haopeng Wang

*Arthur Weiss Lab UCSF-HHMI
513 Parnassus Ave, S-1024 Bay3,
BOX 0795, San Francisco, CA, 94143-0795*

Dr. Mohammad Shoeb

*University of Texas Medical Branch,
Galveston-TX,
USA*

Dr. Bechan Sharma

*Department of Biochemistry,
University of Allahabad, Allahabad, 211 002,
India*

Dr. Sarita Saraswati

*Department of Pharmacology, College of Medicine,
King Saud University, Riyadh
Kingdom of Saudi Arabia*

Dr. Santosh R. Mohanty

*Soil Microbiology Department of Soil Biology
Indian Institute of Soil Science,
Indian Council of Agricultural Research (ICAR)
Bearsia Road, Nabibagh, Bhopal, MP,
India*

Dr. Yule Yue Wang

*Biotechnology and Medicinal Biochemistry
Division of Life Science
The Hong Kong University of Science & Technology
Clear Water Bay, Hong Kong*

Dr. Yogender Pal Khosa

*Department of Microbiology,
University of Delhi South Campus,
New Delhi – 110021*

Dr. Wei Li Cai

*Eunice Kennedy Shriver National Institute of Child
Health and Human Development
National Institutes of Health,
18 Library Drive, MSC-5430
Bethesda, MD 20892*

Dr. Alireza Valdiani

*Department of Biochemistry,
Faculty of Biotechnology and Biomolecular
Sciences,
Universiti Putra Malaysia (UPM),
43400 UPM Serdang
Malaysia*

Instructions for Author

Electronic submission of manuscripts is strongly encouraged, provided that the text, tables, and figures are included in a single Microsoft Word file (preferably in Arial font).

The **cover letter** should include the corresponding author's full address and telephone/fax numbers and should be in an e-mail message sent to the Editor, with the file, whose name should begin with the first author's surname, as an attachment.

Article Types

Three types of manuscripts may be submitted:

Regular articles: These should describe new and carefully confirmed findings, and experimental procedures should be given in sufficient detail for others to verify the work. The length of a full paper should be the minimum required to describe and interpret the work clearly.

Short Communications: A Short Communication is suitable for recording the results of complete small investigations or giving details of new models or hypotheses, innovative methods, techniques or apparatus. The style of main sections need not conform to that of full-length papers. Short communications are 2 to 4 printed pages (about 6 to 12 manuscript pages) in length.

Reviews: Submissions of reviews and perspectives covering topics of current interest are welcome and encouraged. Reviews should be concise and no longer than 4-6 printed pages (about 12 to 18 manuscript pages). Reviews are also peer-reviewed.

Review Process

All manuscripts are reviewed by an editor and members of the Editorial Board or qualified outside reviewers. Authors cannot nominate reviewers. Only reviewers randomly selected from our database with specialization in the subject area will be contacted to evaluate the manuscripts. The process will be blind review.

Decisions will be made as rapidly as possible, and the journal strives to return reviewers' comments to authors as fast as possible. The editorial board will re-review manuscripts that are accepted pending revision. It is the goal of the AJFS to publish manuscripts within weeks after submission.

Regular articles

All portions of the manuscript must be typed double-spaced and all pages numbered starting from the title page.

The Title should be a brief phrase describing the contents of the paper. The Title Page should include the authors' full names and affiliations, the name of the corresponding author along with phone, fax and E-mail information. Present addresses of authors should appear as a footnote.

The Abstract should be informative and completely self-explanatory, briefly present the topic, state the scope of the experiments, indicate significant data, and point out major findings and conclusions. The Abstract should be 100 to 200 words in length. Complete sentences, active verbs, and the third person should be used, and the abstract should be written in the past tense. Standard nomenclature should be used and abbreviations should be avoided. No literature should be cited.

Following the abstract, about 3 to 10 key words that will provide indexing references should be listed.

A list of non-standard **Abbreviations** should be added. In general, non-standard abbreviations should be used only when the full term is very long and used often. Each abbreviation should be spelled out and introduced in parentheses the first time it is used in the text. Only recommended SI units should be used. Authors should use the solidus presentation (mg/ml). Standard abbreviations (such as ATP and DNA) need not be defined.

The Introduction should provide a clear statement of the problem, the relevant literature on the subject, and the proposed approach or solution. It should be understandable to colleagues from a broad range of scientific disciplines.

Materials and methods should be complete enough to allow experiments to be reproduced. However, only truly new procedures should be described in detail; previously published procedures should be cited, and important modifications of published procedures should be mentioned briefly. Capitalize trade names and include the manufacturer's name and address. Subheadings should be used. Methods in general use need not be described in detail.

Results should be presented with clarity and precision. The results should be written in the past tense when describing findings in the authors' experiments. Previously published findings should be written in the present tense. Results should be explained, but largely without referring to the literature. Discussion, speculation and detailed interpretation of data should not be included in the Results but should be put into the Discussion section.

The Discussion should interpret the findings in view of the results obtained in this and in past studies on this topic. State the conclusions in a few sentences at the end of the paper. The Results and Discussion sections can include subheadings, and when appropriate, both sections can be combined.

The Acknowledgments of people, grants, funds, etc should be brief.

Tables should be kept to a minimum and be designed to be as simple as possible. Tables are to be typed double-spaced throughout, including headings and footnotes. Each table should be on a separate page, numbered consecutively in Arabic numerals and supplied with a heading and a legend. Tables should be self-explanatory without reference to the text. The details of the methods used in the experiments should preferably be described in the legend instead of in the text. The same data should not be presented in both table and graph form or repeated in the text.

Figure legends should be typed in numerical order on a separate sheet. Graphics should be prepared using applications capable of generating high resolution GIF, TIFF, JPEG or Powerpoint before pasting in the Microsoft Word manuscript file. Tables should be prepared in Microsoft Word. Use Arabic numerals to designate figures and upper case letters for their parts (Figure 1). Begin each legend with a title and include sufficient description so that the figure is understandable without reading the text of the manuscript. Information given in legends should not be repeated in the text.

References: In the text, a reference identified by means of an author's name should be followed by the date of the reference in parentheses. When there are more than two authors, only the first author's name should be mentioned, followed by 'et al'. In the event that an author cited has had two or more works published during the same year, the reference, both in the text and in the reference list, should be identified by a lower case letter like 'a' and 'b' after the date to distinguish the works.

Examples:

Abayomi (2000), Agindotan et al. (2003), (Kelebeni, 1983), (Usman and Smith, 1992), (Chege, 1998;

1987a,b; Tijani, 1993,1995), (Kumasi et al., 2001) References should be listed at the end of the paper in alphabetical order. Articles in preparation or articles submitted for publication, unpublished observations, personal communications, etc. should not be included in the reference list but should only be mentioned in the article text (e.g., A. Kingori, University of Nairobi, Kenya, personal communication). Journal names are abbreviated according to Chemical Abstracts. Authors are fully responsible for the accuracy of the references.

Examples:

Chikere CB, Omoni VT and Chikere BO (2008). Distribution of potential nosocomial pathogens in a hospital environment. *Afr. J. Biotechnol.* 7: 3535-3539.

Moran GJ, Amii RN, Abrahamian FM, Talan DA (2005). Methicillinresistant *Staphylococcus aureus* in community-acquired skin infections. *Emerg. Infect. Dis.* 11: 928-930.

Pitout JDD, Church DL, Gregson DB, Chow BL, McCracken M, Mulvey M, Laupland KB (2007). Molecular epidemiology of CTXM-producing *Escherichia coli* in the Calgary Health Region: emergence of CTX-M-15-producing isolates. *Antimicrob. Agents Chemother.* 51: 1281-1286.

Pelczar JR, Harley JP, Klein DA (1993). *Microbiology: Concepts and Applications.* McGraw-Hill Inc., New York, pp. 591-603.

Short Communications

Short Communications are limited to a maximum of two figures and one table. They should present a complete study that is more limited in scope than is found in full-length papers. The items of manuscript preparation listed above apply to Short Communications with the following differences: (1) Abstracts are limited to 100 words; (2) instead of a separate Materials and Methods section, experimental procedures may be incorporated into Figure Legends and Table footnotes; (3) Results and Discussion should be combined into a single section.

Proofs and Reprints: Electronic proofs will be sent (e-mail attachment) to the corresponding author as a PDF file. Page proofs are considered to be the final version of the manuscript. With the exception of typographical or minor clerical errors, no changes will be made in the manuscript at the proof stage.

Fees and Charges: Authors are required to pay a \$550 handling fee. Publication of an article in the Biotechnology and Molecular Biology Reviews is not contingent upon the author's ability to pay the charges. Neither is acceptance to pay the handling fee a guarantee that the paper will be accepted for publication. Authors may still request (in advance) that the editorial office waive some of the handling fee under special circumstances

Copyright: © 2015, Academic Journals.

All rights Reserved. In accessing this journal, you agree that you will access the contents for your own personal use but not for any commercial use. Any use and or copies of this Journal in whole or in part must include the customary bibliographic citation, including author attribution, date and article title.

Submission of a manuscript implies: that the work described has not been published before (except in the form of an abstract or as part of a published lecture, or thesis) that it is not under consideration for publication elsewhere; that if and when the manuscript is accepted for publication, the authors agree to automatic transfer of the copyright to the publisher.

Disclaimer of Warranties

In no event shall Academic Journals be liable for any special, incidental, indirect, or consequential damages of any kind arising out of or in connection with the use of the articles or other material derived from the BMBR, whether or not advised of the possibility of damage, and on any theory of liability.

This publication is provided "as is" without warranty of any kind, either expressed or implied, including, but not limited to, the implied warranties of merchantability, fitness for a particular purpose, or non-infringement. Descriptions of, or references to, products or publications does not imply endorsement of that product or publication. While every effort is made by Academic Journals to see that no inaccurate or misleading data, opinion or statements appear in this publication, they wish to make it clear that the data and opinions appearing in the articles and advertisements herein are the responsibility of the contributor or advertiser concerned. Academic Journals makes no warranty of any kind, either express or implied, regarding the quality, accuracy, availability, or validity of the data or information in this publication or of any other publication to which it may be linked.

Biotechnology and Molecular Biology Reviews

Table of Contents: Volume 10 Number 1 January, 2015

ARTICLES

Immunomodulatory effects of carbohydrates and advanced glycation end products

Kankia, H. I.

Review

Immunomodulatory effects of carbohydrates and advanced glycation end products

Kankia, H. I.

Department of Biochemistry, Umaru Musa Yar'adua University, Katsina, Katsina State, Nigeria.

Received 12 November, 2014; Accepted 17 December, 2014

Carbohydrates which comprise of monosaccharides, disaccharides and polysaccharides are heterogeneous complex structure in living systems and are found to bind to other organic molecules such as proteins and lipids. Advanced glycation end products (AGEs) are also heterogeneous group of molecules that accumulate in plasma and other body fluids and in cell and tissues. This review focuses on the impact of polysaccharides and AGEs on immune function. A number of *in vitro* and *in vivo* studies have demonstrated that polysaccharides and AGEs modulate both innate and adaptive immune responses. Polysaccharides such as lectins are reported to activate innate immune cells and T-helper cells leading to the up-regulation of T-lymphocytes, B-lymphocytes and macrophages and release of different cytokine profiles such as tumour necrosis factor (TNF- α), interleukin (IL)-1, IL-6, IL-8, IL-10, IL-12, interferon gamma and beta (IFN- γ and IFN- β 2). Both lectins which are non-enzymatic proteins present in plants and animals that preferentially bind to specific carbohydrate structures and AGE which triggers inflammatory response by binding to RAGE do so there by mediating inflammation known as a key underlying cause in the development of vascular complications leading to an increased expression of cytokines, growth factors and adhesion molecules with mediate and immune response.

Key words: Immunomodulation, carbohydrates, advanced glycation end products (AGEs), receptors, lectins, *in vitro* and *in vivo* studies.

INTRODUCTION

Immunomodulation through natural substances could be considered an alternative for the prevention and cure of various diseases. There is growing evidence suggesting that polysaccharides from natural plants can significantly enhance immune system responses. Plant polysaccharides are known to possess various biological and pharmacological activities hence are regarded as promising immunomodulatory agents which are relatively nontoxic and so have no significant side effects (Kim et al., 2011;

Jin-Kyung et al., 2011; Tang et al., 2012; Xia et al., 2012). It has been demonstrated that most of the polysaccharides are activators of immune cells through their capacity to enhance production of immune mediators, such as reactive oxygen species (ROS), nitric oxide (NO), TNF- α and IL-10 and also their immunomodulatory characteristics are mainly dependent on their unique molecular structures which activate varied surface receptors (Jeurink et al., 2008; Weng et al.,

*Corresponding author. E-mail: irohamza2000@yahoo.com or 1217323@chester.ac.uk.

2011).

Macrophages and lymphocytes are important immune cells which play a pivotal role in immune responses by releasing cytokines that plays a critical role in mediating signal transduction and stimulating the immune defence system (Chen et al., 2008; Schepetkin and Quinn, 2006; Zha et al., 2007). Cytokines which are soluble glycoproteins ought to critically involve in the immune response. The functions of these proteins are diverse and include roles in normal humoral and T-lymphocyte mediated immune response (Schepetkin and Quinn, 2006; Zha et al., 2007).

The killer cell immunoglobulin-like receptors (KIR) and C-type lectin-like receptors are the two major families of receptors expressed by human NK cells which recognize class I MHC molecules. The receptors within each family have similar features with highly homologous extracellular domains attached to various transmembrane regions and cytoplasmic tails. However, the inhibitory receptors belonging to both KIR and C-type lectin-like receptor families carry a pair of ITAMs in their cytoplasmic tails (Baba et al., 2000). Moreover, due to their early production of cytokines and chemokines and their ability to lyse target cells without prior sensitisation NK cells are crucial components of the innate immune system, providing a first line of defence against infectious agents. Hence, they can be activated non-specifically by several cytokines, including IL-2, IL-15 and IL-18 (Middleton et al., 2002).

Numbers of pathologic conditions are accompanied by changes in cytokine levels and by disturbances in the cytokine-mediated interplay between innate and acquired immune responses (Stanilova et al., 2005). IL-2 is found to be an essential cytokine for T lymphocyte growth. IFN- γ promotes the differentiation of Th1 cells and functions principally on macrophages to enhance antimicrobial properties. IL-4 is crucial for the differentiation of naive Th cells into the Th2 cells that promote humoral immunity and also has a central role in the pathogenesis of allergic inflammation (Chen et al., 2008). A study by Zha et al. (2007) using *Dendrobium huoshanense* in an *in vitro* study showed that the polysaccharide of HPS-1B23 was able to stimulate IFN- γ and TNF- α production in the culture medium of splenocytes and macrophages, respectively with no observed toxicity effect of the samples (Zha et al., 2007).

In the human body, the immune system is divided into the innate and adaptive immunities. Innate immunity refers to a non-specific immune response in encountering a particular antigen, while the adaptive immunity shows a high specificity to antigens invaded and is responsible for a memory response after the first exposure of the same antigen. These two systems do not work separately, they act closely together to help fight against the external threats by triggering a series of immune responses (Shao et al., 2004; Wong et al., 2011).

Carbohydrates are storage components in living system

and comprise monosaccharides, disaccharides and polysaccharides (Khodse et al., 2007). Their role in biological processes makes them to be considered as vital biomolecules (Ghazarian et al., 2011; Hu and Xu, 2011). Carbohydrates decorate the surfaces of all cells (glycocalyx) in the form of polysaccharides, glycoproteins, glycolipids and other glycoconjugates. The saccharide binding proteins mediate cell-cell interactions in numerous biological processes, such as blood coagulation, immune response, viral infection, pathological processes, inflammation, embryogenesis and cellular signal transfer (Han et al., 2011; Hu and Xu, 2011), thus they are recognized as ligands for carbohydrate recognition proteins (for example, lectins) (Hu and Xu, 2011; Nakamura et al., 2013; Wu et al., 2006).

AGEs are heterogeneous group of molecules that accumulate in plasma and other body fluids and in cell and tissues (Kasper and Funk, 2001). They are main pathogenic agent of advanced age, diabetes, atherosclerosis, renal failure and other chronic diseases. Researches has shown that AGEs are risk factors of disease as are known to be generated by the non-enzymatic reaction of amino groups with different biomolecules such as proteins and lipids (Hirasawa et al., 2011; Kasper and Funk, 2001; Wu et al., 2011). AGEs accumulate in the living system due to an increased level of sugars and reactive dicarbonyl compounds such as glucose, fructose, deoxyglucose, glyoxal, methylglyoxal and triosephosphates and their formation is irreversible, causing protease-resistant cross-linking of peptides and proteins, which can lead to protein deposition and amyloidosis (Krautwald and Münch, 2010).

CARBOHYDRATE STRUCTURE AND FUNCTION

Organic molecules in living systems are classified into four major groups: proteins, lipids, nucleic acids and carbohydrates. However carbohydrates are by far the most abundant organic molecules found in nature, and nearly all organisms synthesize and metabolize carbohydrates. It has the empirical formula $C_nH_{2n}O_n$, where n is ≥ 3 and glucose is the common monosaccharide that may be linked to form a variety of other macromolecules. Carbohydrates appear to form a heterogeneous complex structure in living systems by binding to other organic molecules such as proteins and lipids by the action of glycosyltransferases and glycosidases. The combined activity levels of these enzymes in the endoplasmic reticulum (ER) and the Golgi apparatus, and perhaps on the cell surface, determine the glycosylation patterns (carbohydrate domains) of glycolipids and glycoproteins (Ghazarian et al., 2011).

The structural features of a polysaccharide are defined by molecular weight, monosaccharide composition, sequence of monosaccharide, configuration and position of glycosidic linkages, type and polymerization degree of




Figure 1. Showing carbohydrate heterogeneity found on cell surface glycoproteins and glycolipids. Sialic acids (Sia) are usually found at the terminal residue of O-linked and N-linked glycans of glycoproteins and glycolipids. Glc, glucose; Gal, galactose; Man, mannose; Fuc, fucose; GalNAc, N-acetylgalactosamine; GlcNAc, N-acetylglucosamine. Illustration prepared by HaikeGhazarian using Corel Draw X4 (Ghazarian et al., 2011).

branch, spatial configuration, etc (Wu et al., 2011). The extent to which polysaccharides exerts their bioactivities is closely related to their chemical composition, molecular weight, branching, chain conformation and water solubility. The basic understanding of both the primary and secondary structures for the polysaccharides is essential for the successful interpretation of their bioactivities, and therefore, the monosaccharide composition analysis of polysaccharides is the most important step for the further discovery of its physicochemical properties, structure and structure-bioactivity relationship (Huang et al., 2007; Xie et al., 2013a).

Possible structures are O- or N-linked glycans and composition varies, giving a wide variety of structures able to bind to many different ligands (Figure 1). However, it is the secondary structure that determines the biological activity.

Most of the polysaccharides, especially those found in the cell wall of the plant are mainly composed of monosaccharides such as xylose (xyl), arabinose (ara), mannose (man), sorbinose (sor), ribose (rib), Fuc, Glc, Gal, GlcNAc and GalNAc, along with small proportions of rhamnose (rha), fructose (fur), and 4-O-methyl galactosamine. The glycosyl linkages of these monosaccharide residues are those determined to give an insight into the structure of polysaccharides (Hsieh et al., 2008; Wang et al., 2012).

CELLULAR RECEPTORS FOR CARBOHYDRATES

Certain carbohydrates found on circulating cells, proteins and pathogens are recognized by specific lectin receptors involved in clearance or immunity (Nangia-Makker et al., 2002). Cell-surface carbohydrates, composed of several different kinds of monosaccharides, interact with many types of lectins as part of a communication signalling network between cells. Since the biogenesis of the cell-surface carbohydrates is tightly regulated, alterations in glycosylation patterns may play key roles in governing the invasive and metastasis potentials of many types of cancers. Viral virulence as well as bacterial and parasite infections are also mediated by host-cell surface carbohydrates, especially their terminal saccharides, which are recognized by invading pathogens (Hyun et al., 2007).

Complex carbohydrates (polysaccharides) are attached to the majority of secreted proteins and cover the surface of cells as a dense layer. However their importance, availability and structural variations reflect a key role in many biological functions. Lectins which are carbohydrate-binding proteins that are able to detect subtle differences between complex carbohydrate structures, decipher glycodes recognizing certain sugars to carry out various functions like cell attachment, migration or invasion and that is why among all of the variety of




Figure 2. Myeloid C-type lectin receptors (CLRs) in innate immunity. A, In innate immunity, CLRs such as DC-SIGN, mannose receptor (MR), or DEC-205 serve as pattern-recognition receptors and bind to glycan structures present on pathogens such as bacteria, viruses, or parasites. B, Myeloid CLRs share one or more homologous carbohydrate-recognition domains (CRDs) that bind to carbohydrates often in a Ca^{2+} -dependent manner.

adhesion molecules, cell-surface carbohydrate structures have been a focus of many investigations (Nangia-Makker et al., 2002).

Lectins are found throughout nature in plants, bacteria and animals. The major types of animal lectins include galectins, C-type, I-type, P-type and R-type lectins. Most lectins contain a carbohydrate recognition domain (CRD) that binds with high specificity to the outermost sugars of *N*-linked and *O*-linked glycans as well as glycosphingolipids (Sørensen et al., 2012).

Among other members, C-type lectin receptors (CLRs) are considered to be suitable target in innate immunity and cell specific drug delivery due to their endocytotic or cell activation properties. They represent a large receptor family including collectins, selectins, lymphocyte lectins and proteoglycan and share a structurally homologous carbohydrate-recognition domain (CRD) as well as often binding to carbohydrates in a Ca^{2+} -dependent pattern. Myeloid CLRs such as DC-SIGN, mannose receptor (MR), DEC-205 and others are crucial to initiate immune responses against various pathogens including bacteria, viruses, parasites and fungi (Figure 2a). However, some CLRs serve solely as phagocytic receptors whereas others activate signalling pathway by inducing the activation of tyrosine kinases such as spleen tyrosine kinase (SYK) through classical immunoreceptor tyrosine-based activation motifs (ITAMs) or hemITAMs (Figure 2b) (Lepenies et al., in press, accessed 2013).

The signaling pathway elicited upon CLR engagement

depends on their cytoplasmic signaling motifs. Some CLRs contain an intracellular hemITAM or are associated with ITAM-bearing adaptor proteins leading to the activation of the tyrosine kinase Syk. In contrast, other CLRs contain an intracellular ITIM that induces the recruitment of phosphatases such as SHP-1 and SHP-2. Another group of myeloid CLRs including DC-SIGN or MR does not possess intracellular signaling motifs, thus signaling is independent of Syk or phosphatases (Lepenies et al., in press, accessed 2013).

CARBOHYDRATES IN CELLULAR FLUIDS

Cellular fluids, especially extracellular have been isolated from various bodily fluids and used as biological markers for various diseases among which are cancer and diabetes mellitus (Tzimagiorgis et al., 2011). Carbohydrates appeared to be basic to virtually every aspect of extracellular circulation, fulfilling roles from purely structural to mediating the highly specific recognition events that underlie cell-cell communication. Sugars are naturally chosen by nature to suitably mediate such an extraordinary variety of interactions. They are linked and similar to nucleotides and amino acids, also are able to form branched structures with stereospecific linkages, features that confer exponential capacity for structural diversity. Extracellular sugars are often elaborated with covalent modifications, such as sulfation, acetylation and

and phosphorylation that impart further structural variety (Bowman and Bertozzi, 1999).

Several conjugates including glycolipids and glycol-proteins can adhere to each other by *trans* carbohydrate-carbohydrate interactions between apposed membranes. For example, myelin, the membranous sheath which is spirally wound around nerve axons is rich in two glycosphingolipids, GalC and its sulphated form, galactosylceramide 1³-sulfate. These interactions are often mediated by divalent cations. Water disrupts these interactions between single sugar molecules, but the interactions between polymeric multivalent glycoconjugates or membrane lipid domains, that present a multivalent array of carbohydrate head groups, are stronger and persist in water. These can be strong enough to cause specific adhesion of sponge cells into a multicellular organism. In the case of mammalian cells, they may cause transient adhesion of cells allowing up-regulation of proteins such as integrins to maintain adhesion and the interaction can be homotypic, similar to that for the lextanglioside (Boggs et al., 2008).

Immunomodulatory activity of polysaccharides

Polysaccharides isolated from various traditional medicinal plants, especially higher plants (Xu et al., 2009) have been shown to greatly affect the immune system both *in vivo* and *in vitro* by stimulating cytokine and chemokine production, ROS production and cell proliferation. They are reported to be non-toxic, with no significant side effects, which is a major problem associated with many other immunomodulatory agents, such as bacterial polysaccharides and synthetic compounds. These polysaccharides therefore have the potential as immunomodulators (Belska et al., 2010; Schepetkin and Quinn, 2006) as they have been shown to increase macrophage cytotoxic activity against tumour cells and microorganisms, activate phagocytic activity and enhance secretion of cytokines and chemokines (Chen et al., 2010; Schepetkin and Quinn, 2006).

Several studies have shown the immunomodulatory effects of certain polysaccharides and how they activate innate immune cells and T-helper cells leading to the up-regulation of T-lymphocytes, B-lymphocytes, and macrophages and release of different cytokine profiles (Chen et al., 2010; Cheng et al., 2008; Fan et al., 2012; Kouakou et al., 2013; Makino et al., 2006; Wang et al., 2013; Wong et al., 2011; Xia et al., 2012; Xu et al., 2009; Zhang et al., 2013). For example, an *in vivo* immunomodulatory study by Wong et al. (2011) using sclerotial polysaccharides isolated by hot water extraction from *Pleurotus tuber-regium* and *Pheidole rhinocerus* (PTW and PRW, respectively) on both healthy BALB/c mice and healthy athymic nude mice had shown that innate immune cells such as monocyte/macrophages and T-helper cells dendritic cells are activated by the mushroom are activated

by the mushroom sclerotial polysaccharides upon detection of release cytokines like IL-1 β , IFN- γ , TNF- α , IL-2 and IL-6 (Wong et al., 2011).

Also, in a study by Zhang et al. (2013) on hot water extracted polysaccharides from stems of *Taxillus chinensis* and *Uncaria rhyncophylla* in a dose-dependent manner (10 and 100 $\mu\text{g/ml}$) have shown an increase production of TNF and NO, especially on 100 $\mu\text{g/ml}$ concentration in treated macrophages when compared with untreated ones. Hence it can be seen that these polysaccharide fractions display significant immunomodulatory activities as indicated by increased production of TNF- α as well as NO (Zhang et al., 2013).

Another example is on the recent study by Wang et al. (2013) on *Kadsura marmorata*. This plant has been used in traditional Chinese medicine as one of the major ingredients for the treatment of bronchial asthma, eczema and acute and chronic infection due to either viruses or bacteria. The study which was investigated *in vitro* on chicken lymphocytes and macrophages has shown that *K. marmorata* had up-regulate the production of chicken T-lymphocytes, B-lymphocytes and peritoneal macrophages which in turn could stimulate the cell proliferation. It also significantly enhances cytokine secretion. This was ascertained when different concentrations (dose dependent manner) of *K. marmorata* were used. Hence was suggested as having potential effects on regulating the immune system (Wang et al., 2013).

Protein glycation

During the lifetime of the cells, proteins are exposed to a multitude of modification process or non-enzymatic glycosylation reaction between reducing sugars and amine residues in proteins which usually occurred as a result of elevation of glucose concentrations (Shuvaev et al., 2001; Vlassopoulos et al., 2013). Oxidation and glycation are among the major non enzymatic mechanisms. Such kind of alterations can affect tissues and are accumulated in the cells with an indisputable effect on metabolism.

In the reaction known as Maillard, that involves three main stages: early, intermediate and late, glucose or other reducing sugars such as fructose, pentoses, galactose, mannose, ascorbate, xylulose, at the early stage, react with a free amino group of several molecules, including proteins, nucleic acids and lipids, to form an unstable aldimine compound called the Schiff base. However, through rearrangement, this base gives rise to a stable ketoamine, known as Amadori product (Figure 3) (Lapolla et al., 2005).

These molecular and cellular alterations, though not dangerous at the initial stage, can become damaging and pathogenic when sufficiently abundant. Among the various targets that are sensitive to alteration, blood plasma is sustainably and continually exposed to numerous




Figure 3. Maillard reaction showing the three steps of non-enzymatic glycation reaction (Lapolla et al., 2005).

metabolites, which in turn induce an oxidative stress (Rondeau and Bourdon, 2011).

Products of AGE are believed to be risk markers or risk factors of disease (Kasper and Funk, 2001; Vlassopoulos et al., 2013). Soluble AGE compounds in the circulation and extracellular fluids and another glycation product known as glycated hemoglobin (HbA1c) have been demonstrated to be the biomarkers for the diagnosis of diabetes and monitoring of glucose control in diabetic individuals as well as macrovascular diseases (Indurthi et al., 2012; Vlassopoulos et al., 2013).

Cellular receptors for advanced glycation end products

The AGEs that are in the circulation or extracellular space can interact with two types of cell surface receptors. Clearance and scavenger receptors are predominantly involved in AGE capture, removal and degradation. This group of receptors includes p60 (OST-48), p90 (80 K-H), galectin-3, type I and type II macrophage scavenger receptors, CD-36, FEEL-1 and -2, scavenger receptor proteins SR-BI and SR-BII and the lectin-like oxidized low-density lipoprotein receptor 1

(Lox-1). The other types of AGE-receptors initiate specific cellular signalling events in response to AGE exposure. Among these various AGE-receptors, a receptor for advanced glycation end-products (RAGE) is the receptor for AGEs which accumulate in various diseases and inflammations and is suggested to be involved in cellular migration and spreading over binding of amphoterin, a developmental-regulated heparin-binding protein (Indurthi et al., 2012; Kasper and Funk, 2001).

RAGE is a 35 kDa member of the immunoglobulin superfamily of cell surface proteins that interacts with a wide range of ligands, including AGE products, modified low density lipoproteins, amyloid fibrils, amphoterin (HMGB1) and various S100 proteins. It shares structural homology with other immunoglobulin like receptors (Leclerc et al., 2009) and is a specific binding protein for AGE moieties which have been detected on monocytes/macrophages, smooth muscle cells, hepatocytes, neurons, pericytes, T-lymphocytes, mesangial cells, type I pneumocytes, osteoblast-like cells and endothelial cells and its expression can be abnormally up or down-regulated in human disease (Kasper and Funk, 2001; Xie et al., 2013b).

Activation of the RAGE pathway has been shown to be important in wound healing, tumour growth and




Figure 4. A schematic representation of RAGE showing the extracellular part comprising three immunoglobulin like domain, the V, C1 and C2 domain. A single transmembrane helix connects the extracellular domain with the short intracellular domain (ID). RAGE ligands interact with the extracellular domains as pointed out. (*) Indicates that the RAGE domain where S100 binds is not known (Leclerc et al., 2009).

metastasis, as well as in systemic amyloidosis (Germanová et al., 2010; Mahajan et al., 2010).

The mature 382 amino-acid long RAGE is composed of an extracellular part (314 aa), a single transmembrane spanning helix (27 aa) and a short cytosolic domain (41 aa) (Figure 4) (Leclerc et al., 2009). Complete length RAGE contains a V-type domain, which functions for ligand binding, two C-type domains, a transmembrane spanning helix, and a C-terminal cytosolic domain (ctRAGE) which is required for signal transduction. However RAGEs with the deletion of V-type domain (variant 2) are unable to interact with its ligands and therefore irresponsive to any stimuli, whereas RAGEs without ctRAGE (variant 1) have negative actions by competing with endogenous full-length RAGE for ligand binding, thereby blunting the RAGE-dependent signal transduction (Figure 5) (Xie et al., 2013).

RAGE has been documented in promoting inflammation and endothelial activation, which in turn also accelerates coronary atherosclerotic development. Its activation is enhanced by accumulation of its ligands in inflammatory compartments, initiating a vicious cycle causing further up-regulation of the receptor and sustained cell activation. These molecules has also been shown to be highly expressed in human atherosclerotic lesions where it co-localizes with pro-inflammatory and pro-oxidative mediators (Mahajan et al., 2010).

Advanced glycation end products in cellular fluids

Normal tissue remodelling is thought to be contributed by AGE modified proteins. However, under pathological conditions such as diabetes, rheumatoid arthritis (RA) and osteoarthritis, renal insufficiency and in smokers, AGEs can be found elevated in body fluids such as serum, synovial fluid and urine. Among them, serum and synovial fluid pentosidine is the superior indicator for so many diseases status (Kasper and Funk, 2001). Soluble receptor for advanced glycation end products (sRAGE) which is the alternative splicing of RAGE mRNA and the proteolytic cleavage of membrane-bound RAGE has the ability to bind with the same ligands, sRAGE acts as a competitive inhibitor of RAGE, and precludes the cell-bound RAGE signaling. RAGE adversely involves in several pathological processes of human diseases, such as diabetes, Alzheimer's disease and chronic inflammatory diseases such as RA (Chayanupatkul and Honsawek, 2010).

It has been reported recently that another extracellular newly identified RAGE-binding protein, known as EN-RAGE, has been found. It is a member of the S100 protein family, also called S100A12 or CAAF1 or calgranulin C, which is comprised of closely related low molecular weight (10-14 kDa) acidic calcium-binding proteins with EF hands. These proteins were purified




Figure 5. Structures of complete length RAGE and its variants which contains a first 22 amino acid signal recognition sequence, followed by a V-type domain, two C-type domains, a transmembrane domain and a cytosolic signal domain. RAGE variant 1 lacks the cytosolic domain whereas variant 2 lack the V-domain (Xie et al., 2013).

from bovine lung extract and expressed abundantly in the esophageal epithelium, neutrophils and monocytes/macrophages in humans. In the other hand to sRAGE, EN-RAGE is a natural pro-inflammatory ligand for RAGE.

EN-RAGE binding to RAGE activates intracellular signal cascades including MAP-kinase and NKkB which subsequently induces secretion of cytokines such as IL-1 β and TNF- α , expression of adhesion molecules there by activating inflammatory responses and enhancing the activation and migration of monocytes/macrophages to damaged vessels leading to the pathogenesis of inflammatory diseases (Hasegawa et al., 2003; Kalousova et al., 2012; Kim et al., 2012).

Additionally, S100A12 is able to interact with scavenger receptors; it is also chemotactic for neutrophils and monocytes and by binding copper. It is able to generate ROS and so acts as a pro-oxidant agent. Increased level of EN-RAGE are found in various inflammatory compartments like synovial fluid and its serum concentrations are increased in inflammatory diseases and correlated with the disease activity. High concentrations of S100A12 are described in patients with neurodegenerative

diseases and atherosclerosis. EN-RAGE interacting with RAGE could also be implicated in cancerogenesis particularly at sites with chronic and persistent inflammation (Kalousova et al., 2012).

Immunomodulatory effects of advanced glycation end-products

AGE triggers inflammatory response by binding to RAGE there by mediating inflammation known as a key underlying cause in the development of vascular complications leading to an increased expression of cytokines, growth factors and adhesion molecules with mediate an immune response (Choi et al., 2013; Hirose et al., 2011). Intracellularly, AGEs that are formed inside and outside of cells induces a more unspecific chemical reaction, the production of ROS increases which causes cross-linking with other cellular proteins and generation of lipid peroxidation product (Kasper and Funk, 2001). RAGE, found in most tissues, have potent immunomodulatory actions, promoting ROS production and inflammation (Vlassopoulos et al., 2013).




Figure 6. AGE-RAGE interaction and NF- κ B activation leading to oxidant stress, vasoconstriction and a procoagulant state (Lapolla et al., 2005).

AGEs are known to be high in patients with poorly controlled disease, and its accumulation is a fundamental mechanism that causes the complications in diseases like diabetes. T lymphocytes contain a surface receptor for AGE-modified proteins that is inducible by the stimulation of the T-lymphocyte mitogen phytohemagglutinin. AGEs have also been found to act on macrophages through the AGE receptor on the plasma membrane, though it is not known whether the proliferation of activated T lymphocytes would be affected by AGEs (Liu et al., 2005).

Several targets of advanced glycation include structural proteins, such as collagen and aggrecan, plasma proteins, including immunoglobulins and albumin, and intracellular proteins, such as haemoglobin and lens crystallin. Protein glycation in particular has the potential to alter many cellular functions, and as a consequence AGE has been suggested to contribute towards the pathogenesis of many diseases. Moreover, AGE-modified proteins have been shown to generate free radicals such as superoxide and NO, and to up-regulate pro-inflammatory cytokines by monocyte and/or macrophages. Other responses elicited by AGE include induction of IL-1 and expression of tissue factor by monocytes and release of TNF α by THP-1 monocytic cells incubated with AGE-human serum albumin (HSA) (Pertyńska-Marczewska et al., 2004). Other consequences observed in AGE-RAGE interaction are an increase in the procoagulant response to TNF- α , a reduction in thrombomodulin expression, and an increase in endothelin-1 levels (Figure 6) (Lapolla et al., 2005).

CONCLUSION

Carbohydrates are favourable candidates for disease control, because they are present on the cell surface and act as an identification tag, through which they can interact with their surroundings, whether the extracellular matrix or homotypic or heterotypic cells. Interfering with the normal cell recognition phenomenon using a large or a small sugar molecule has been reported to block the progression of tumours by interfering with angiogenesis, cell-cell, cell-matrix interactions, tumour invasion and metastasis. Modification of a natural plant carbohydrate to facilitate its absorption by the body is a more cost-effective and easier way to prevent disease progression than synthesizing a competitive sugar molecule. The generation of AGEs is an inevitable process *in vivo* and their accumulation in tissues has been implicated in the process of aging and also in the pathogenesis of several pathological conditions. A number of *in vitro* and *in vivo* studies have demonstrated that carbohydrates and AGEs modulate various indices of immune function. Further analysis of monosaccharide composition of polysaccharides could be an essential parameter for the evaluation of quality of standard of polysaccharide materials and basic information on it before being applied for any investigation. Methods of isolating a particular polysaccharide is also paramount: their extraction by different methods might yield different chemical components, as such reliable and effective methods should be applied. Although, so far investigations of the receptor-mediated pathways for the immunomodulatory

activity of certain polysaccharides have been reported, a more detailed mechanistic study is required to explain how the innate immune cells such as macrophages are activated by polysaccharides. Future studies could be directed not only to its ability to control cancer which is widely and commonly known it for, but also in other diseases. Further studies should be applied on the most effective compounds that could suppress AGE-mediated inflammatory signalling cascades, which involved the expression of pro-inflammatory intermediates and NF- κ B transcriptional activity. Suppression of intracellular ROS production by scavenging ROS and reducing NADPH oxidase activity was a key regulatory mechanism for inhibiting AGE-mediated inflammatory response. Given the key role of AGE-RAGE signalling in the progression of diabetic complications, the suppressive effects of those compounds on AGE-mediated inflammation could be applied to prevent the development of diabetic complications.

Conflict of Interests

The author(s) have not declared any conflict of interests.

REFERENCES

- Baba E, Erskine R, Boyson JE, Cohen GB, Davis DM, Malik P, Strominger JL (2000). N-linked carbohydrate on human leukocyte antigen-C and recognition by natural killer cell inhibitory receptors. *Hum. Immunol.* 61(12):1202-1218.
- Belska NV, Guriev AM, Danilets MG, Trophimova ES, Uchasova EG, Ligatcheva AA, Belsky YP (2010). Water-soluble polysaccharide obtained from *Acorus calamus* L. classically activates macrophages and stimulates Th1 response. *Int. Immunopharmacol.* 10(8):933-942.
- Boggs JM, Gao W, Hirahara Y (2008). Myelin glycosphingolipids, galactosylceramide and sulfatide, participate in carbohydrate-carbohydrate interactions between apposed membranes and may form glycosynapses between oligodendrocyte and/or myelin membranes. *Biochim. Biophys. Acta* 1780(3):445-455.
- Bowman KG, Bertozzi CR (1999). Carbohydrate sulfotransferases: mediators of extracellular communication. *Chem. Biol.* 6(1):R9-R22.
- Chayanupatkul M, Honsawek S (2010). Soluble receptor for advanced glycation end products (sRAGE) in plasma and synovial fluid is inversely associated with disease severity of knee osteoarthritis. *Clin. Biochem.* 43(13-14):1133-1137.
- Chen J-R, Yang Z-Q, Hu T-J, Yan Z-T, Niu T-X, Wang L, Wang M (2010). Immunomodulatory activity *in vitro* and *in vivo* of polysaccharide from *Potentilla anserina*. *Fitoterapia* 81(8):1117-1124.
- Chen Z, Kwong Huat Tan B, Chan SH (2008). Activation of T lymphocytes by polysaccharide-protein complex from *Lycium barbarum* L. *Int. Immunopharmacol.* 8(12):1663-1671.
- Cheng A, Wan F, Wang J, Jin Z, Xu X (2008). Macrophage immunomodulatory activity of polysaccharides isolated from *Glycyrrhiza uralensis* fish. *Int. Immunopharmacol.* 8(1):43-50.
- Choi H-J, Jang H-J, Chung T-W, Jeong S-I, Cha J, Choi J-Y, Ha K-T (2013). Catalpol suppresses advanced glycation end-products-induced inflammatory responses through inhibition of reactive oxygen species in human monocytic THP-1 cells. *Fitoterapia* 86:19-28.
- Fan L, Ding S, Ai L, Deng K (2012). Antitumor and immunomodulatory activity of water-soluble polysaccharide from *Inonotus obliquus*. *Carbohydr. Polym.* 90(2):870-874.
- Germanová A, Koucký M, Hájek Z, Pařízek A, Zima T, Kalousová M (2010). Soluble receptor for advanced glycation end products in physiological and pathological pregnancy. *Clin. Biochem.* 43(4-5):442-446.
- Ghazarian H, Idoni B, Oppenheimer SB (2011). A glycobiology review: Carbohydrates, lectins and implications in cancer therapeutics. *Acta Histochem.* 113(3):236-247.
- Han E, Ding L, Jin S, Ju H (2011). Electrochemiluminescent biosensing of carbohydrate-functionalized CdS nanocomposites for in situ label-free analysis of cell surface carbohydrate. *Biosensors Bioelectron.* 26(5):2500-2505.
- Hasegawa T, Kosaki A, Kimura T, Matsubara H, Mori Y, Okigaki M, Iwasaka T (2003). The regulation of EN-RAGE (S100A12) gene expression in human THP-1 macrophages. *Atherosclerosis* 171(2):211-218.
- Hirasawa Y, Sakai T, Ito M, Yoshimura H, Feng Y, Nagamatsu T (2011). Advanced-glycation-end-product-cholesterol-aggregated-protein accelerates the proliferation of mesangial cells mediated by transforming-growth-factor-beta 1 receptors and the ERK-MAPK pathway. *Eur. J. Pharmacol.* 672(1-3):159-168.
- Hirose J, Yamabe S, Takada K, Okamoto N, Nagai R, Mizuta H (2011). Immunohistochemical distribution of advanced glycation end products (AGEs) in human osteoarthritic cartilage. *Acta Histochem.* 113(6):613-618.
- Hsieh YSY, Chien C, Liao SKS, Liao S-F, Hung W-T, Yang W-B, Wong C-H (2008). Structure and bioactivity of the polysaccharides in medicinal plant *Dendrobium huoshanense*. *Bioorg. Med. Chem.* 16(11):6054-6068.
- Hu M-X, Xu Z-K (2011). Carbohydrate decoration of microporous polypropylene membranes for lectin affinity adsorption: comparison of mono- and disaccharides. *Colloids Surf. B Biointerfaces* 85(1):19-25.
- Huang Q, Jin Y, Zhang L, Cheung PCK, Kennedy JF (2007). Structure, molecular size and antitumor activities of polysaccharides from *Poria cocos* mycelia produced in fermenter. *Carbohydr. Polym.* 70(3):324-333.
- Hyun S, Kim J, Kwon M, Yu J (2007). Selection and syntheses of tentacle type peptides as 'artificial' lectins against various cell-surface carbohydrates. *Bioorg. Med. Chem.* 15(1):511-517.
- Indurthi VSK, Leclerc E, Vetter SW (2012). Interaction between glycosylated serum albumin and AGE-receptors depends on structural changes and the glycation reagent. *Arch. Biochem. Biophys.* 528(2):185-196.
- Jeurink PV, Noguera CL, Savelkoul HFJ, Wichers HJ (2008). Immunomodulatory capacity of fungal proteins on the cytokine production of human peripheral blood mononuclear cells. *Int. Immunopharmacol.* 8(8):1124-1133.
- Kalousová M, Kuběna AA, Benáková H, Dusilová-Sulková S, Tesář V, Zima T (2012). EN-RAGE (extracellular newly identified receptor for advanced glycation end-products binding protein) and mortality of long-term hemodialysis patients: A prospective observational cohort study. *Clin. Biochem.* 45(7-8):556-560.
- Kasper M, Funk RHW (2001). Age-related changes in cells and tissues due to advanced glycation end products (AGEs). *Arch. Gerontol. Geriatr.* 32(3):233-243.
- Khodse VB, Fernandes L, Gopalkrishna VV, Bhosle NB, Fernandes V, Matondkar SGP, Bhushan R (2007). Distribution and seasonal variation of concentrations of particulate carbohydrates and uronic acids in the northern Indian Ocean. *Mar. Chem.* 103(3-4):327-346.
- Kim J-K, Cho ML, Kamjanapratum S, Shin I-S, You SG (2011). *In vitro* and *in vivo* immunomodulatory activity of sulfated polysaccharides from *Enteromorpha prolifera*. *Int. J. Biol. Macromol.* 49(5):1051-1058.
- Kim J-K, Park S, Lee MJ, Song YR, Han SH, Kim SG, Yoo T-H (2012). Plasma levels of soluble receptor for advanced glycation end products (sRAGE) and proinflammatory ligand for RAGE (EN-RAGE) are associated with carotid atherosclerosis in patients with peritoneal dialysis. *Atherosclerosis* 220(1), 208-214.
- Kouakou K, Schepetkin IA, Yapi A, Kirpotina LN, Jutila MA, Quinn MT (2013). Immunomodulatory activity of polysaccharides isolated from *Alchornea cordifolia*. *J. Ethnopharmacol.* 146(1):232-242.
- Krautwald M, Münch G (2010). Advanced glycation end products as biomarkers and gerontotoxins - A basis to explore methylglyoxal-lowering agents for Alzheimer's disease? *Exp. Gerontol.* 45(10):744-751.
- Lapolla A, Traldi P, Fedele, D. (2005). Importance of measuring products of non-enzymatic glycation of proteins. *Clin. Biochem.* 38(2):103-115.

- Leclerc E, Fritz G, Vetter SW, Heizmann CW (2009). Binding of S100 proteins to RAGE: An update. *Biochim. Biophys. Acta (BBA) Mol. Cell Res.* 1793(6):993-1007.
- Lepenes B, Lee J, Sonkaria S (in press, accessed 2013). Targeting C-type lectin receptors with multivalent carbohydrate ligands. *Adv. Drug Deliv. Rev.* 65(9):1271-1281.
- Liu C-T, Chen K-M, Lee S-H, Tsai L-J (2005). Effect of supplemental L-arginine on the function of T lymphocytes and the formation of advanced glycosylated end products in rats with streptozotocin-induced diabetes. *Nutrition* 21(5):615-623.
- Mahajan N, Bahl A, Dhawan V (2010). C-reactive protein (CRP) up-regulates expression of receptor for advanced glycation end products (RAGE) and its inflammatory ligand EN-RAGE in THP-1 cells: Inhibitory effects of atorvastatin. *Int. J. Cardiol.* 142(3):273-278.
- Makino S, Ikegami S, Kano H, Sashihara T, Sugano H, Horiuchi H, Oda M (2006). Immunomodulatory Effects of Polysaccharides Produced by *Lactobacillus delbrueckii* ssp. *bulgaricus* OLL1073R-1. *J. Dairy Sci.* 89(8):2873-2881.
- Middleton D, Curran M, Maxwell L (2002). Natural killer cells and their receptors. *Transplant Immunol.* 10(2-3):147-164.
- Nakamura M, Tsutsumi M, Ishikawa Y, Umemiya H, Hasegawa T, Izawa K, Hasegawa T (2013). Glycosylated tris-bipyridine ferrous complexes to provide dynamic combinatorial libraries for probing carbohydrate-carbohydrate interactions. *Tetrahedron* 69(14):3019-3026.
- Nangia-Makker P, Conklin J, Hogan V, Raz A (2002). Carbohydrate-binding proteins in cancer, and their ligands as therapeutic agents. *Trends Mol. Med.* 8(4):187-192.
- Pertyńska-Marczewska M, Kiriakidis S, Wait R, Beech J, Feldmann M, Paleolog EM (2004). Advanced glycation end products upregulate angiogenic and pro-inflammatory cytokine production in human monocyte/macrophages. *Cytokine* 28(1):35-47.
- Pesesse X, Backers K, Moreau C, Zhang J, Blero D, Paternotte N, Erneux C (2006). SHIP1/2 interaction with tyrosine phosphorylated peptides mimicking an immunoreceptor signalling motif. *Adv. Enzyme Regul.* 46(1):142-153.
- Ponnusamy S, Zhang H, Kadam P, Lin Q, Lim TK, Sandhu JS, Choolani M (2012). Membrane proteins of human fetal primitive nucleated red blood cells. *J. Proteom.* 75(18):5762-5773.
- Rondeau P, Bourdon E (2011). The glycation of albumin: Structural and functional impacts. *Biochimie* 93(4):645-658.
- Schepetkin IA, Quinn MT (2006). Botanical polysaccharides: Macrophage immunomodulation and therapeutic potential. *Int. Immunopharmacol.* 6(3):317-333.
- Shao B-M, Xu W, Dai H, Tu P, Li Z, Gao X-M (2004). A study on the immune receptors for polysaccharides from the roots of *Astragalus membranaceus*, a Chinese medicinal herb. *Biochem. Biophys. Res. Commun.* 320(4):1103-1111.
- Shuvaev VV, Laffont I, Serot J-M, Fujii J, Taniguchi N, Siest G (2001). Increased protein glycation in cerebrospinal fluid of Alzheimer's disease. *Neurobiol. Aging* 22(3):397-402.
- Sørensen ALT, Clausen H, Wandall HH (2012). Carbohydrate clearance receptors in transfusion medicine. *Biochim. Biophys. Acta* 1820(11):1797-1808.
- Stanilova SA, Dobрева ZG, Slavov ES, Miteva LD (2005). C3 binding glycoprotein from *Cuscuta europea* induced different cytokine profiles from human PBMC compared to other plant and bacterial immunomodulators. *Int. Immunopharmacol.* 5(4):723-734.
- Tang X-H, Yan L-F, Gao J, Yang X-L, Xu Y-X, Ge H-Y, Yang H-D (2012). Antitumor and immunomodulatory activity of polysaccharides from the root of *Limonium sinense* Kuntze. *Int. J. Biol. Macromol.* 51(5):1134-1139.
- Tzimagiorgis G, Michailidou EZ, Kritis A, Markopoulos AK, Kouidou S (2011). Recovering circulating extracellular or cell-free RNA from bodily fluids. *Cancer Epidemiol.* 35(6):580-589.
- Vlassopoulos A, Lean MEJ, Combet E (2013). Role of oxidative stress in physiological albumin glycation: A neglected interaction. *Free Radic. Biol. Med.* 60:318-324.
- Wang H, Deng X, Zhou T, Wang C, Hou Y, Jiang H, Liu G (2013). The in vitro immunomodulatory activity of a polysaccharide isolated from *Kadsura marmorata*. *Carbohydr. Polym.* 97(2):710-715.
- Wang X-M, Sun R-G, Zhang J, Chen Y-Y, Liu N-N (2012). Structure and antioxidant activity of polysaccharide POJ-U1a extracted by ultrasound from *Ophiopogon japonicus*. *Fitoterapia* 83(8):1576-1584.
- Weng BB-C, Lin Y-C, Hu C-W, Kao M-Y, Wang S-H, Lo D-Y, Chiou R Y-Y (2011). Toxicological and immunomodulatory assessments of botryosphaeran (β -glucan) produced by *Botryosphaeria rhodina* RCYU 30101. *Food Chem. Toxicol.* 49(4):910-916.
- Wong K-H, Lai CKM, Cheung PCK (2011). Immunomodulatory activities of mushroom sclerotial polysaccharides. *Food Hydrocoll.* 25(2):150-158.
- Wu JH, Singh T, Herp A, Wu AM (2006). Carbohydrate recognition factors of the lectin domains present in the *Ricinus communis* toxic protein (ricin). *Biochimie* 88(2):201-217.
- Wu L, Ma L, Nicholson LFB, Black PN (2011). Advanced glycation end products and its receptor (RAGE) are increased in patients with COPD. *Respir. Med.* 105(3):329-336.
- Xia L, Liu X, Guo H, Zhang H, Zhu J, Ren F (2012). Partial characterization and immunomodulatory activity of polysaccharides from the stem of *Dendrobium officinale* (Tiepi shihu) *in vitro*. *J. Funct. Foods* 4(1):294-301.
- Xie J, Méndez JD, Méndez-Valenzuela V, Aguilar-Hernández MM (2013a). Cellular signalling of the receptor for advanced glycation end products (RAGE). *Cell. Signal.* 25(11):2185-2197.
- Xie J-H, Shen M-Y, Nie S-P, Liu X, Zhang H, Xie M-Y (2013b). Analysis of monosaccharide composition of *Cyclocarya paliurus* polysaccharide with anion exchange chromatography. *Carbohydr. Polym.* 98(1):976-981.
- Xu H, Yao L, Sun H, Wu Y (2009). Chemical composition and antitumor activity of different polysaccharides from the roots of *Actinidia eriantha*. *Carbohydr. Polym.* 78(2):316-322.
- Zha X-Q, Luo J-P, Luo S-Z, Jiang S-T (2007). Structure identification of a new immunostimulating polysaccharide from the stems of *Dendrobium huoshanense*. *Carbohydr. Polym.* 69(1):86-93.
- Zhang L, Koyyalamudi SR, Jeong SC, Reddy N, Bailey T, Longvah T (2013). Immunomodulatory activities of polysaccharides isolated from *Taxillus chinensis* and *Uncaria rhynchophylla*. *Carbohydr. Polym.* 98(2):1458-1465.

Biotechnology and Molecular Biology Reviews

Related Journals Published by Academic Journals

- *Journal of Cell and Animal Biology*
- *African Journal of Environmental Science and Technology*
- *African Journal of Biochemistry Research*
- *African Journal of Agricultural Research*
- *African Journal of Microbiology Research*
- *African Journal of Biotechnology*
- *African Journal of Pharmacy and Pharmacology*
- *Scientific Research and Essays*

academicJournals