



International Journal for Biotechnology and Molecular Biology Research

Volume 6 Number 5, June 2015
ISSN 2141-2154



*Academic
Journals*

ABOUT IJBMBR

The **International Journal for Biotechnology and Molecular Biology Research (IJBMBR)** (ISSN 2141-2154) is published Monthly (one volume per year) by Academic Journals.

International Journal for Biotechnology and Molecular Biology Research (IJBMBR) provides rapid publication (monthly) of articles in all areas of the subject such as Green energy from chemicals and bio-wastes, Studies in the graft copolymerization of acrylonitrile onto cassava starch by ceric ion induced initiation, Antimutagenic activity of aqueous extract of Momordica charantia, Ethnomedicinal plants and other natural products with anti-HIV active compounds and their putative modes of action etc.

The Journal welcomes the submission of manuscripts that meet the general criteria of significance and scientific excellence. Papers will be published shortly after acceptance. All articles published in IJBMBR are peer-reviewed.

Submission of Manuscript

Please read the **Instructions for Authors** before submitting your manuscript. The manuscript files should be given the last name of the first author

[Click here to Submit manuscripts online](#)

If you have any difficulty using the online submission system, kindly submit via this email ijbmb@academicjournals.org.

With questions or concerns, please contact the Editorial Office at ijbmb@academicjournals.org.

Editors

Prof Atagana, Harrison

*Institute for Science and Technology Education
University of South Africa*

Prof. UC Banerjee

*Department of Pharmaceutical Technology
(Biotechnology)
National Institute of Pharmaceutical Education and
Research
Punjab, INDIA*

Dr. Y. Omid

*Faculty of Pharmacy,
Research Center for Pharmaceutical Nanotechnology,
School of Advanced Biomedical Sciences,
Tabriz University of Medical Sciences,
Tabriz, Iran.*

Prof. Mohamed E. Wagih

*University of New Brunswick (UNB-SJ),
Saint John College, NB,
E2L 4L5, Canada*

Dr. Sripada M. Udupa

*ICARDA-INRA Cooperative Research Project,
International Center for Agricultural Research in the
Dry Areas(ICARDA), B.P. 6299,
Rabat Instituts, Rabat, Morocco*

Dr. Amjad Masood Husaini

*Sher-e-Kashmir University of Agricultural Sciences &
Technology
Bohlochipora, Dr. Ali Jan Road,
Nowshera, Srinagar, J&K-190011, India*

Dr. Om Prakash Gupta

*Directorate of Wheat Research (ICAR)
Post Box-158, A
grasain Marg, Karnal-132001,
Haryana, India*

Editorial Board

Dr. Amro Hanora

*Suez Canal University, Department of Microbiology
and Immunology,
Faculty of Pharmacy, Suez Canal University,
Box 41522 Ismailia, Egypt*

Dr. C. Rajasekaran

*VIT University
School of Bio-Sciences & Technology (SBST)*

Dr. Yasar Karadag

*Gaziosmanpasa University
Faculty of Agriculture,
Department of Field Crops, Tokat-Turkey*

Dr. Ahmet Tutus

*KSU (Kahramanmaras Sutcu Imam University)
Faculty of Forestry,
Department of Forest Industrial Engineering,
Kahramanmaras 46100 Turkey*

Dr. Vinod Joshi

*Desert Medicine Research Centre,
Indian Council of Medical Research
New Pali Road, Jodhpur, India*

Dr. Eshrat Gharaei Fathabad

*K.M.18 Khazarabad road,
Sari, Mazandaran, Iran*

Dr. Shashideep Singhal

*121 Dekalb Ave, Brooklyn,
NY 11201, New York, USA*

Dr Masayoshi Yamaguchi

*101 Woodruff Circle, 1305 WMRB,
Atlanta, Georgia 30322-0001, USA*

Dr. Okonko Iheanyi Omezuruike

*Department of Virology,
Faculty of Basic Medical Sciences,
College of Medicine,
University College Hospital,
Ibadan, Nigeria*

Dr. S. M. Shahid

*University of Karachi,
Karachi-75270, Pakistan*

Prof. Reda Helmy Sammour

*Botany Department,
Faculty of Science,
Tanta University,
Tanta, Egypt*

Dr. Premendra D. Dwivedi

*Food Toxicology Division,
Room No 303,
P.O. Box 80,
M. G. Road, Lucknow-226001, UP, India*

Dr. Amro Hanora

*Microbiology and Immunology department,
Faculty of Pharmacy,
Suez Canal University,
Box 41522 Ismailia, Egypt*

Dr. Tamilnadu

*1501 N. Campbell Ave
Tucson, AZ 85724 India*

Dr. Yadollah Omid

*Faculty of Pharmacy,
Tabriz University of Medical Sciences,
Daneshgah St., Tabriz, Iran*

Dr. Mohsen Selim Asker

*National Research Centre, Dokki, Cairo,
Egypt*

Dr. Fanuel Lampiao

P.O.Box 360, Blantyre, Malawi

Prof. Mohamed E. Wagih

Saint John, NB, E2L 4L5, Canada

Dr. Santosh Kumar Singh

*Centre of Experimental Medicine and Surgery,
Institute of Medical Sciences,
Banaras Hindu University, Varanasi-221005, India*

Dr. Zhanhuan Shang

*No.768, Jiayuguan West Road,
Lanzhou City, Gansu Province, China*

Dr. Worlanyo Eric Gato

*Southern Illinois University – Carbondale,
1245 Lincoln Dr, 144 Neckers, Carbondale IL 62901*

Dr. Chun Chen

*College of Life Sciences,
China Jiliang University
Xueyuan Street, Xiasha, Hangzhou,
Zhejiang Province, PR China*

Dr. Efthimios J.Karadimas

*LGI, Leeds NHS Trust 10th Timoleonos Vassou str,
11521, Athens Greece*

Dr. Samuel Toba Anjorin

University of Abuja, Abuja, Nigeria

Dr. Rupali Agnihotri

*Department of Periodontics,
Manipal college of Dental Sciences,
Manipal,576104. Karnataka. India*

Dr. Mahbuba Begum

*Tuber Crops Research Centre,
Joydebpur, Gajipur-1701, Bangladesh*

Prof. S. Mohan Karuppaiyl

*School of Life Sciences
Srtm University
Nanded. MS. India*

Dr. Neveen Helmy Aboelsoud

*Complementary Medicine Researches and Application
Department
National Research Center, Cairo
Egypt.*

Dr. D.E. Chandrashekar Rao

*National Research Council,
Plant Biotechnology Institute
Canada*

Dr. Nikolaos Papanas

*Democritus University of Thrace
G. Kondyli 22, Alexandroupolis, Greece*

Dr. Sivakumar Swaminathan

*Iowa State University
USA*

Dr. El Sayed Hussein El Sayed Ziedan

*National Research Centre,
Plant Pathology Department
Tahrir St.,Dokki Cairo,
Egypt*

Dr. Chethan Kumar M

*Post Graduate Departments of Bio-technology and Biochemistry,
Ooty Road, Mysore - 570 025,
Karnataka,
India*

Dr. M. Sattari

*Rice Research Ins. of Iran
Iran*

Dr. Zaved Ahmed Khan

*VIT University
India*

Dr. Subbiah Poopathi

*Vector Control Research Centre
Indian Council of Medical Research
(Ministry of Health & Family Welfare,
Govt. of India)
Medical Complex, Indira Nagar
India*

Dr. Reyazul Rouf Mir

*International Crops Research Institute for the Semi-Arid Tropics (ICRISAT),
Patancheru - 502 324, Greater Hyderabad,
India*

Dr. Prasanna Kumar S

*Virginia Commonwealth University,
USA*

Dr. Naseem Ahmad

*Plant Biotechnology Laboratory
Department of Botany
Aligarh Muslim University
Aligarh- 202 002, (UP)
India*

Dr. Zhen-Xing Tang

*Food Bioengineering institute, Hangzhou Wahaha Co.
Ltd, Hangzhou,
Zhejiang,
China*

Dr. Jayanthi Abraham

*VIT (Vellore Institute of Technology) University,
Tamilnadu,
India*

Dr. Gobianand Kuppannan

*National Institute of Animal Science
South Korea*

Dr. R. Harikrishnan

*Jeju National University
South Korea*

Dr. Asit Ranjan Ghosh

*Vellore Institute of Technology (VIT) University,
School of Bio Sciences & Technology,
Medical Biotechnology Division, Vellore-632014,
India*

Dr. Kamal Dev

*Shoolini University of Biotechnology and Management
Sciences (SUBMS)
India*

Dr. Wichian Sittiprapaporn

*Maharakham University
Thailand*

Dr. Vijai Kumar Gupta

*Molecular Glycobiotechnology Group, Department of
Biochemistry,
School of Natural Sciences,
National University of Ireland, Galway,
Ireland*

Dr. Jeffy George

*Department of Microbiology and Immunology
F. Edward Hébert School of Medicine
Uniformed Services University of the Health Sciences
4301 Jones Bridge Road,
Bethesda, MD 20814
USA.*

Dr. Gyanendra Singh

*Stanley S. Scott Cancer Center,
School of Medicine,
Louisiana State University Health Sciences Center
New Orleans, LA 70112,
USA.*

Dr. Anupreet Kour

*1620 Chevy Chase Dr.
Champaign, IL 61821
USA.*

Dr. Arun Sharma

*Institute for Plant Genomics and Biotechnology (IPGB)
Borlaug Center,
TAMU 2123
Texas A&M University
College Station, TX 77843
USA.*

Dr. Mohsen Asker

*Microbial Biotechnology Dept.
National Research Centre
Cairo,
Egypt.*

Dr. Elijah Miinda Ateka

*Department of Horticulture,
Jomo Kenyatta University of Agriculture and
Technology (JKUAT)
Kenya.*

Dr. Jozélio Freire De Carvalho

*Faculdade de Medicina Da USP, Reumatologia
Av. Dr. Arnaldo, 455 - 3º andar – Sala 3133.
São Paulo - SP
Brazil*

Dr. Premendra Dhar Dwivedi

*Food Toxicology Division
Industrial Institute of Toxicology Research,
Post Box No: 80, Mahatma Gandhi Marg,
Lucknow 226001,
India*

Dr. Muhammad Abd El-Moez El-Saadani

*Universities and Research Center District,
New Borg El-Arab,
P.O.Box: 21934 Alexandria,
Egypt.*

Dr. Donald J. Ferguson

*Advanced Orthodontic Training Program,
Nicolas & Asp University College
Dubai,
UAE*

Dr. Kalyan Goswami

*Department of Biochemistry & JB Tropical Disease
Research Centre,
Mahatma Gandhi Institute of Medical Sciences,
Sevagram, Wardha-442102*

Dr. A.K. Handa

*National Research Centre for Agroforestry,
Gwalior Road, JHANSI-284003 UP
India.*

Dr. Amjad M.Husaini

*Metabolic Engineering & Biotechnology Laboratory
Division of Plant Breeding & Genetics
Sher-e-Kashmir University of Agricultural Sciences &
Technology of Kashmir
J&K-191121,
India*

Dr. Vinod Joshi

*Laboratory of Virology & Molecular Biology,
Desert Medicine Research Centre,
Pali Road, Jodhpur-342 005,
India*

Dr. T. Kalaivani

*D/O S. Thiagarajan
B-43, Rajaram Nagar,
Salem - 636 007,
Tamil Nadu, India*

Dr. Priya Kalia

*Orthopaedic Research Unit,
Department of Surgery,
Cambridge University, Cambridge,
UK*

Dr. Patricia Khashayar

*Tehran University of Medical Sciences
Endocrinology and Metabolism Research Center
Shariati Hospital*

Dr. Zaringhalam Moghadam

*Shahid Beheshti Medical University (M.C)
Tehran,
Iran*

Dr. Okeke Ikechukwu Linus

*Department of Surgery, University of Ibadan
Nigeria.*

Dr. Rajesh Kumar Patel

*Centre for Analysis and Learning in Livestock and Food
(CALF)
National Dairy Development Board (NDDB)
Anand- 388 001 (Gujarat)
INDIA*

Dr. Pooja Ralli-Jain

*Department of Pathology and Laboratory Medicine
University of California Irvine,
Irvine, California,
U.S.A.*

Dr. Meltem Sesli

*College of Tobacco Expertise,
Turkish Republic, Celal Bayar University 45210,
Akhisar, Manisa,
Turkey*

Dr. Reda H. Sammour

*Tanta University, Faculty of Science, Tanta,
Egypt*

Dr. Seyed Soheil Saeedi Saravi

*Mazandaran University of Medical sciences, Sari,
Iran*

Dr. R. Senthil Kumar

*St. Matthew's University, School of Medicine
Grand Cayman
Cayman Islands*

Dr. Mohammad Reza Shakibaie

*Kerman University of Medical Sciences, Kerman,
Iran*

Dr. Srividya Shivakumar

*Dept of Microbiology,
CPGS, Jain university,
Bangalore*

Dr. Shashideep Singhal

*The Brooklyn Hospital Center
NewYork-Presbyterian Healthcare System
Brooklyn,
NY.*

Dr. Sripada M. Udupa

*International Center for Agricultural Research in the
Dry Areas (ICARDA),
B.P. 6299, Rabat Instituts, Rabat,
Morocco.*

Dr. Wei Wu

*Institute for Biocomplexity and Informatics
Department of Bio Science
The University of Calgary
Canada*

Dr. Xiao-Bing Zhang

*Molecular Regeneration Laboratory, MC1528B
11234 Anderson Street
Loma Linda, CA 92350*

Prof. Dr. Ozfer Yesilada

*Inonu University
Faculty of Arts and Sciences
Department of Biology
44280 Malatya
Turkey*

Dr. Edson Boasquevisque

*Universidade do Estado do Rio de Janeiro- UERJ
Av 28 de setembro, 87, fundos (LMMC-IBRAG).
Vila Isabel, city: Rio de Janeiro/ RJ
Brasil*

Dr. Abhilash M.

*The Oxford College of Engineering
Hosur Road, Bangalore - 560068*

Dr. Nasar Uddin Ahmed

*Department of Genetics and Plant Breeding
Patuakhali Science and Technology University
Dumki, Patuakhali-8602
Bangladesh*

Dr. Mervat Morsy EL- Gendy

*Chemistry of Natural and Microbial Products
Department,
National Research Center, Dokki, Cairo,
Egypt*

Dr. Gjurmakch Aliev

*Health Science and Healthcare Administration
Program,
University of Atlanta, Atlanta, Georgia,
USA*

Dr. Muhammad Asgher

*Department of Chemistry and Biochemistry,
University of Agriculture,
Faisalabad,
Pakistan*

Dr. Anand Bharatkumar

*Parul Institute of Pharmacy, Limda, Waghodia,
Vadodara*

Dr. Chinmoy Kumar Bose,

*Netaji Subhash Chandra Bose Cancer Research
Institute
16A, Park Lane, Park Street, Kolkata 700 016,
India.*

Dr. Mousumi Debnath

*Jaipur Engineering College and Research Centre
(JECRC) Department of Biotechnology,
Shri Ram ki Nangal, Via Vatika ,Tonk Road , Jaipur-
303905 ,
India*

Dr. Dolan C. Saha

*Dept. of Biochemistry and Molecular Biology,
Faculty of Medicine,
University of Calgary,
Canada*

Dr. Ramasamy Harikrishnan

*Department of Aquatic Biomedical Sciences
School of Marine Biomedical Science
College of Ocean Sciences
Jeju National University
Jeju city, Jeju 690 756,
South Korea*

Dr. Abdul Haque

*Health
Biotechnology division, nibge,
Faisalabad,
Pakistan*

Dr. Kuvalekar Aniket Arun

*Interactive Research School for Health Affairs
(IRHSA),
Bharati Vidyapeeth University, Pune, Maharashtra,
India*

Dr. Asit Ranjan Ghosh

*School of Bio Science & Technology,
Division of Medical Biotechnology,
Vellore Institute of Technology (VIT) University,
Vellore-632014,
India*

Dr. Prasanna Kumar Santhekadur

*Department of Human and Molecular Genetics
Virginia Commonwealth University
Richmond,
VA*

Dr. Majid Sattari

*Rice Research Institute of Iran
Iran*

Dr. Mihael Cristin Ichim

*National Institute Research and Development for
Biological Sciences /
"Stejarul" Research Centre for Biological Sciences
Alexandru cel Bun St., 6, Piatra Neamt, 610004,
Romania*

Dr. Sailas Benjamin

*Enzyme Technology Laboratory
Biotechnology Division
Department of Botany
University of Calicut
Kerala - 673 635
India*

Dr. Sreeramanan Subramaniam

*School of Biological Sciences,
Universiti Sains Malaysia (USM),
Minden Heights, 11800, Penang,
Malaysia*

Dr. Vijai Kumar Gupta,

*Department of Biochemistry, NUI, Galway,
Ireland*

Dr. Vitor Engrácia Valenti

*Universidade Federal de São Paulo
Rua Napoleão de Barros, 715, Térreo
São Paulo, SP
Brazil.*

Dr. Ravindra Pogaku

*School of Engineering and IT
Universiti Malaysia Sabah
88999 Kota Kinabalu
Sabah,
Malaysia*

Dr. Ahmed Eid Abdel-Hamid Eweis Fazary

*School of Pharmacy,
College of Medicine,
National Taiwan University, Taipei 100,
Taiwan.*

Dr. Mohammad Hashemi

*Dept. of Clinical Biochemistry,
School of Medicine,
Zahedan University of Medical Sciences,
Zahedan,
Iran*

Dr. Hesham, Abd El-Latif

*Genetics Department,
Assiut University, Assiut 71516,
Egypt.*

Prof. Jia-ying Xin

*College of Food Engineering
Harbin University of Commerce
138 Tongda Road
Daoli District
Harbin 150076, Heilongjiang
P.R.China*

Dr. Kabir Mohammad Humayun

*Plant Molecular Biotech Lab
Department of Medical Biotechnology
College of Biomedical Science
Kangwon National University
Kangwon-do, Chuncheon, 200-701
South Korea*

Dr. Kalpesh Gaur

*Geetanjali College of Pharmaceutical Studies Manwa
Khera,
Udaipur- 313002. Rajasthan,
India*

Dr. Meganathan, Kannan

*Center for Biologics Evaluation and Research (CBER),
U.S. Food and Drug Administration (FDA),
Bldg. NIH 29A, Room 2C-10,
8800 Rockville Pike,
Bethesda, MD 20892.
USA.*

Assist. Prof. Ali Karadeniz

*Department of Physiology,
Faculty of Veterinary Medicine,
University of Atatürk 25240 ERZURUM
Turkey*

Dr. Matthew Kostek

*Department of Kinesiology
University of Connecticut
Storrs CT*

Dr. Tansu Kucuk

*Gulhane School of Medicine
Department of Obstetrics and Gynecology
Etlik 06018 Ankara,
Turkey*

Dr. Kuo-Sheng Hung

*Department of Neurosurgery
Taipei Medical University - Wan Fang Medical Center
111 Section 3, Hsing-Long Rd,
Taipei 116,
Taiwan*

Dr. V. Manju

*Department of Biochemistry,
Periyar University,
Salem -11.*

Dr. Mbagwu Ferdinand Nkem

*Department of Plant science and Biotechnology,
Faculty of Science,
Imo State University
Nigeria.*

Dr. Anand Pithadia

*Parul Institute of Pharmacy
Vadodara, Gujarat,
India*

Dr. Radhakrishnan Ramaraj

*Department of Internal Medicine
University of Arizona
Tucson 85724
AZ*

Dr. M. Rasool

*School of Bio Sciences and Technology,
VIT University,
Vellore-632104, Tamil Nadu,
India*

Dr. Reda A.I. Abou-Shanab

*Genetic Engineering & Biotechnology Research
Institute (GEBRI)
Mubarak City for Scientific Research and Technology
Applications
New Burg El-Arab City, Universities and Research
Institutes
Zone, P.O. 21934, Alexandria,
Egypt.*

Dr. MR. Pravin Babarao Suruse

*Department of Pharmaceutics
Sharad Pawar College of Pharmacy
Wanadongri, Hingna Road
Nagpur- 441 110. (M. S.)*

Dr. Jan Woraratanadharm

*GenPhar, Inc.,
Mount Pleasant,
SC*

Dr. Serap Yalin

*Mersin University Pharmacy Faculty
Department of Biochemistry, Mersin
Turkey*

Dr. YongYong Shi

*Bio-X Center,
Shanghai Jiao Tong University,
Hao Ran Building, 1954 Hua Shan Road,
Shanghai 200030,
PR China*

Dr. Jyotdeep Kaur

*Department of Biochemistry,
Post Graduate Institute of Medical Education and
Research (PGIMER),
Chandigarh*

Dr. Rajkumar

*Dept. Of Radiation Biosciences,
Institute of Nuclear Medicine and Allied Sciences
Brig. S.K. Mazumdar Road, Timarpur,
Delhi-110054
India*

Dr. Meera Sumanth

*Visveswarapura Institute of Pharmaceutical Sciences,
22nd Main, 24th Cross, B.S.K II stage,
Bangalore-560070
Karnataka,
India.*

Dr, Jai S. Ghosh

*Department of Microbiology,
Shivaji University,
Kolhapur 416004,
India*

Prof. Dr. Alaa H. Al-Charrakh

*Babylon University, College of Medicine.
Dept. of Microbiology
Hilla, Iraq*

Instructions for Author

Electronic submission of manuscripts is strongly encouraged, provided that the text, tables, and figures are included in a single Microsoft Word file (preferably in Arial font).

The **cover letter** should include the corresponding author's full address and telephone/fax numbers and should be in an e-mail message sent to the Editor, with the file, whose name should begin with the first author's surname, as an attachment.

Article Types

Three types of manuscripts may be submitted:

Regular articles: These should describe new and carefully confirmed findings, and experimental procedures should be given in sufficient detail for others to verify the work. The length of a full paper should be the minimum required to describe and interpret the work clearly.

Short Communications: A Short Communication is suitable for recording the results of complete small investigations or giving details of new models or hypotheses, innovative methods, techniques or apparatus. The style of main sections need not conform to that of full-length papers. Short communications are 2 to 4 printed pages (about 6 to 12 manuscript pages) in length.

Reviews: Submissions of reviews and perspectives covering topics of current interest are welcome and encouraged. Reviews should be concise and no longer than 4-6 printed pages (about 12 to 18 manuscript pages). Reviews are also peer-reviewed.

Review Process

All manuscripts are reviewed by an editor and members of the Editorial Board or qualified outside reviewers. Authors cannot nominate reviewers. Only reviewers randomly selected from our database with specialization in the subject area will be contacted to evaluate the manuscripts. The process will be blind review.

Decisions will be made as rapidly as possible, and the Journal strives to return reviewers' comments to authors as fast as possible. The editorial board will re-review manuscripts that are accepted pending revision. It is the goal of the AJMR to publish manuscripts within weeks after submission.

Regular articles

All portions of the manuscript must be typed double-spaced and all pages numbered starting from the title page.

The Title should be a brief phrase describing the contents of the paper. The Title Page should include the authors' full names and affiliations, the name of the corresponding author along with phone, fax and E-mail information. Present addresses of authors should appear as a footnote.

The Abstract should be informative and completely self-explanatory, briefly present the topic, state the scope of the experiments, indicate significant data, and point out major findings and conclusions. The Abstract should be 100 to 200 words in length. Complete sentences, active verbs, and the third person should be used, and the abstract should be written in the past tense. Standard nomenclature should be used and abbreviations should be avoided. No literature should be cited.

Following the abstract, about 3 to 10 key words that will provide indexing references should be listed.

A list of non-standard **Abbreviations** should be added. In general, non-standard abbreviations should be used only when the full term is very long and used often. Each abbreviation should be spelled out and introduced in parentheses the first time it is used in the text. Only recommended SI units should be used. Authors should use the solidus presentation (mg/ml). Standard abbreviations (such as ATP and DNA) need not be defined.

The Introduction should provide a clear statement of the problem, the relevant literature on the subject, and the proposed approach or solution. It should be understandable to colleagues from a broad range of scientific disciplines.

Materials and methods should be complete enough to allow experiments to be reproduced. However, only truly new procedures should be described in detail; previously published procedures should be cited, and important modifications of published procedures should be mentioned briefly. Capitalize trade names and include the manufacturer's name and address. Subheadings should be used. Methods in general use need not be described in detail.

Results should be presented with clarity and precision. The results should be written in the past tense when describing findings in the authors' experiments. Previously published findings should be written in the present tense. Results should be explained, but largely without referring to the literature. Discussion, speculation and detailed interpretation of data should not be included in the Results but should be put into the Discussion section.

The Discussion should interpret the findings in view of the results obtained in this and in past studies on this topic. State the conclusions in a few sentences at the end of the paper. The Results and Discussion sections can include subheadings, and when appropriate, both sections can be combined.

The Acknowledgments of people, grants, funds, etc should be brief.

Tables should be kept to a minimum and be designed to be as simple as possible. Tables are to be typed double-spaced throughout, including headings and footnotes. Each table should be on a separate page, numbered consecutively in Arabic numerals and supplied with a heading and a legend. Tables should be self-explanatory without reference to the text. The details of the methods used in the experiments should preferably be described in the legend instead of in the text. The same data should not be presented in both table and graph form or repeated in the text.

Figure legends should be typed in numerical order on a separate sheet. Graphics should be prepared using applications capable of generating high resolution GIF, TIFF, JPEG or Powerpoint before pasting in the Microsoft Word manuscript file. Tables should be prepared in Microsoft Word. Use Arabic numerals to designate figures and upper case letters for their parts (Figure 1). Begin each legend with a title and include sufficient description so that the figure is understandable without reading the text of the manuscript. Information given in legends should not be repeated in the text.

References: In the text, a reference identified by means of an author's name should be followed by the date of the reference in parentheses. When there are more than two authors, only the first author's name should be mentioned, followed by 'et al'. In the event that an author cited has had two or more works published during the same year, the reference, both in the text and in the reference list, should be identified by a lower case letter like 'a' and 'b' after the date to distinguish the works.

Examples:

Abayomi (2000), Agindotan et al. (2003), (Kelebeni, 1983), (Usman and Smith, 1992), (Chege, 1998;

1987a,b; Tijani, 1993,1995), (Kumasi et al., 2001) References should be listed at the end of the paper in alphabetical order. Articles in preparation or articles submitted for publication, unpublished observations, personal communications, etc. should not be included in the reference list but should only be mentioned in the article text (e.g., A. Kingori, University of Nairobi, Kenya, personal communication). Journal names are abbreviated according to Chemical Abstracts. Authors are fully responsible for the accuracy of the references.

Examples:

Chikere CB, Omoni VT and Chikere BO (2008). Distribution of potential nosocomial pathogens in a hospital environment. *Afr. J. Biotechnol.* 7: 3535-3539.

Moran GJ, Amii RN, Abrahamian FM, Talan DA (2005). Methicillinresistant *Staphylococcus aureus* in community-acquired skin infections. *Emerg. Infect. Dis.* 11: 928-930.

Pitout JDD, Church DL, Gregson DB, Chow BL, McCracken M, Mulvey M, Laupland KB (2007). Molecular epidemiology of CTXM-producing *Escherichia coli* in the Calgary Health Region: emergence of CTX-M-15-producing isolates. *Antimicrob. Agents Chemother.* 51: 1281-1286.

Pelczar JR, Harley JP, Klein DA (1993). *Microbiology: Concepts and Applications.* McGraw-Hill Inc., New York, pp. 591-603.

Short Communications

Short Communications are limited to a maximum of two figures and one table. They should present a complete study that is more limited in scope than is found in full-length papers. The items of manuscript preparation listed above apply to Short Communications with the following differences: (1) Abstracts are limited to 100 words; (2) instead of a separate Materials and Methods section, experimental procedures may be incorporated into Figure Legends and Table footnotes; (3) Results and Discussion should be combined into a single section.

Proofs and Reprints: Electronic proofs will be sent (e-mail attachment) to the corresponding author as a PDF file. Page proofs are considered to be the final version of the manuscript. With the exception of typographical or minor clerical errors, no changes will be made in the manuscript at the proof stage.

Fees and Charges: Authors are required to pay a \$650 handling fee. Publication of an article in the International Journal for Biotechnology and Molecular Biology Research is not contingent upon the author's ability to pay the charges. Neither is acceptance to pay the handling fee a guarantee that the paper will be accepted for publication. Authors may still request (in advance) that the editorial office waive some of the handling fee under special circumstances

Copyright: © 2015, Academic Journals.

All rights Reserved. In accessing this journal, you agree that you will access the contents for your own personal use but not for any commercial use. Any use and or copies of this Journal in whole or in part must include the customary bibliographic citation, including author attribution, date and article title.

Submission of a manuscript implies: that the work described has not been published before (except in the form of an abstract or as part of a published lecture, or thesis) that it is not under consideration for publication elsewhere; that if and when the manuscript is accepted for publication, the authors agree to automatic transfer of the copyright to the publisher.

Disclaimer of Warranties

In no event shall Academic Journals be liable for any special, incidental, indirect, or consequential damages of any kind arising out of or in connection with the use of the articles or other material derived from the IJBMBR, whether or not advised of the possibility of damage, and on any theory of liability.

This publication is provided "as is" without warranty of any kind, either expressed or implied, including, but not limited to, the implied warranties of merchantability, fitness for a particular purpose, or non-infringement. Descriptions of, or references to, products or publications does not imply endorsement of that product or publication. While every effort is made by Academic Journals to see that no inaccurate or misleading data, opinion or statements appear in this publication, they wish to make it clear that the data and opinions appearing in the articles and advertisements herein are the responsibility of the contributor or advertiser concerned. Academic Journals makes no warranty of any kind, either express or implied, regarding the quality, accuracy, availability, or validity of the data or information in this publication or of any other publication to which it may be linked.

ARTICLES

RNA Interference – A fine tuner of gene regulation: A review

S.K. Dash, Sushil K. Mohapatra and H.N. Malik

Review

RNA Interference – A fine tuner of gene regulation: a Review

S.K. Dash^{1*}, Sushil K. Mohapatra² and H.N. Malik²

¹Department of Animal Genetics and Breeding, COVS, GADVASU, India.

²Animal Biotechnology Centre, ICAR-NDRI, India.

Received 9 October, 2014; Accepted 19 May, 2015

The discovery of RNA interference by Fire and Mello has opened up a new arena for research in molecular biology. The interfering RNAs (iRNAs) are 21-23 nucleotide RNA duplexes, which have the ability to reduce or abolish gene activity, and this property is being used in reverse genetics. Dicer and RISC (RNA-induced silencing complex) play important role in mechanism of RNA interference. The specificity, efficiency and potency make RNAi an attractive tool for analyzing the function of genes. Researchers are investigating the possibility of using RNA interference in treatment of various important diseases such as HIV, cancer, neurodegenerative diseases, malaria and various other diseases.

Key words: RNA interference, Dicer, RISC, reverse genetics.

INTRODUCTION

Gene expression has fundamental importance in all living organisms. RNA was supposed to be the key genetic material before its role was taken over by DNA in most of the living organisms. The discovery of catalytic RNA (ribozyme) has evolutionary implications and it is now realized that RNA plays a key role in gene expression. The regulation of gene expression is done at two stages, first is the transcriptional level in which transcriptional factors are involved along with other mechanisms such as histone acetylation and DNA methylation and second is the post transcriptional level that is by mRNA processing, translation of mRNA and RNA interference (RNAi). Mechanism of RNA interference (RNAi) was first given by A. Fire and C. Mello and this is a conserved biological response to double stranded RNA, which

mediates resistance to both endogenous parasitic and exogenous pathogenic nucleic acids, as well as regulates the expression of protein coding genes (Hannon, 2002). Earlier RNAi was called by other names, including post transcriptional gene silencing and transgene silencing, but it was obvious that they were the same phenomenon only after these were characterized at the molecular level. RNAi can be defined as the ability of exogenous or endogenous double stranded RNA to suppress the expression of the gene which corresponds to the sequence of double stranded RNA. The RNAi is a knockdown and not knockout mechanism of gene silencing, with the advancement of genetic engineering (Volpe and Martienssen, 2011). RNAi is proving to be an alternative approach for stable silencing of genes in a

*Corresponding author. E-mail: shaktikant07@gmail.com.

variety of different animal species. RNAi has been linked to various cellular processes, including the formation of centromeric structure (Djupedal and Ekwall, 2009) and gene regulation, through microRNAs and heterochromatin formation (Dehio, 1994; Wassenegger, 2005). *Science* named RNAi as the 'Technology of the Year' in 2002. During the last few years RNAi has moved to the cutting edge of molecular biology.

DISCOVERY OF RNA INTERFERENCE

At first gene silencing in plants drew the interest of researchers. It was reported that introduction of transgenes homologous to endogenous genes resulted in gene suppression of these genes, this process was known as co-suppression, in which there is degradation of the endogenous and the transgene mRNA (Paddison et al., 2002). A similar process of silencing, called Quelling was also reported in fungi (*N. crassa*) (Romano and Macino, 1992). The explanation for the gene silencing studied in plants for several years was provided by A. Fire et al. (1998) through post transcriptional gene silencing (PTGS) called RNA interference (RNAi). They received Nobel Prize in Physiology or Medicine in 2006 for this novel discovery. In their work Fire and Mello studied the phenotypic effect after injection of single stranded and double stranded unc-22 RNA into gonad of three different groups of *C. elegans*, the unc-22 gene encodes a myofilament protein and the decrease in unc-22 activity is known to produce severe twitching movements. They found that the group that was injected with double stranded RNA produced severe twitching movement while the other two groups exhibited normal or wild type movements. Similar process of RNAi has been reported in *Drosophila* (Filippo et al., 2011) and in mammals (Yang et al., 2001).

RNA INTERFERENCE (RNAi)

Double stranded (dsRNAs) are produced in cell either through transcription of microRNA genes, which consists of inverted repeats or through viral replication (dsRNA Rotavirus). In RNAi long (70-90 nts) dsRNA is digested into short interfering or siRNA (21-23 nts). This siRNA silences homologous genes by destruction of its mRNA or through inhibition of translation of its mRNA or by inducing chromatin modifications in its promoter region (Fire et al., 1998). The interfering RNAs (iRNAs) may be classified into two types, short interfering RNA and micro RNA. The siRNA are exogenous and 21-23 nucleotides in length. They can be designed *in vitro* or the single stranded and double stranded RNA virus infecting the cell are other sources of siRNA. The non protein coding microRNA (miRNA) genes are the other source of iRNAs. These miRNA genes produce iRNAs of 21- 22 nucleotides.

length (Zamore et al., 2000).

MECHANISM OF RNAi

Endogenous miRNA

The Primary miRNA produced from the transcription of miRNA genes, undergo first stage processing in the nucleus and are acted upon by the complex consisting of enzymes Drosha and Pasha, to produce short hairpin pre miRNAs or sh- pre miRNAs. These pre miRNAs are exported into the cytoplasm by the enzyme Exportin5. In the cytoplasm the pre miRNA undergo second stage processing by the activity of enzyme Dicer (a type of ribonuclease III), which cleaves the pre miRNA to produce siRNA duplex. The Dicer then loads the guide strand into the RISC (RNA induced silencing complex); the guide strand is the strand of siRNA duplex which participates in silencing of its complementary mRNA sequence and the one that does not is the passenger strand. The RISC after loaded with the guide strand which carries it to the complementary mRNA so that the guide strand can bind to the complementary sequence in mRNA. The mRNA is then cleaved or degraded if there is a complete complementarity between the two strands. When there is lesser complementarity, leads to suppression of synthesis of the protein in which mRNA is involved.

Exogenous miRNA

The mechanism of down regulation of exogenous miRNA is similar to the endogenous one, except that in endogenous iRNA processing pathway, the first phase processing does not occur, so the Drosha Pasha enzymes do not take part in the processing of mRNA.

Non-Stoichiometric effect of RNAi

RNA interference mechanism involves only small amounts of dsRNA that can wipe out excess of mRNA. This enzymatic mechanism involves RNA dependent RNA polymerase (RdRp). The sense strand which does not participate in RNAi is acted upon by RdRp as a result of which a double stranded RNA is produced which in turn again participates in RNAi pathway (Zamore et al., 2000).

Dicer

The dicer is a cytoplasmic RNase III enzyme with endonuclease activity and it functions as a molecular ruler producing dsRNA fragments 21-23 nucleotides in length. Structurally the dicer is composed of three rigid regions

namely the RNase domain, Platform domain and PAZ domain; these domains are connected by flexible hinges (Filipowicz et al., 2005 and Macrae et al., 2006). Dicer recognizes the end of a dsRNA with PAZ domain and then cuts the dsRNA with its RNase III domains. The distance between PAZ and RNase III domains determines the length of small RNAs generated. Homologs of Dicer exist in many organisms including *C. elegans*, *Drosophila*, yeast and humans. After hydrolysis of substrate RNAs, Dicer aids in loading its small dsRNA product into RISC. The thermodynamic asymmetry along the siRNA or miRNA duplex determines “guide” and “passenger” strands. The dicer presents the strand of the siRNA duplex (whose 5' end is thermodynamically less stable) as the guide strand to the RISC (Bernstein et al., 2001).

RISC complex

The RNA induced silencing complex otherwise called RISC is approximately 500-kDa RNA-multiprotein complex, which triggers mRNA degradation in response to siRNA. RISC is involved in three major functions a) *Helicase*-unwinding of double-stranded siRNA b) *Nuclease*-ribonuclease component cleaves mRNA c) *RNA-dependent RNA polymerase RdRp*- amplification of silencing signal (Cogoni et al., 1999). Argonaute protein is the signature component of RISC and it has two characteristic domains PAZ (also found in Dicer) and PIWI. The cleft of the PAZ domain binds to the characteristic two-nucleotide 3' overhang of the guide strand; the RISC then carries the guide strand to the complementary mRNA to participate in the process of silencing (Macrae et al., 2006).

APPLICATIONS OF RNAI

Reverse genetics

The function of a gene can be identified by blocking the translation of mRNA and studying the resultant phenotypic change, this branch of genetics is termed as reverse genetics. This is one of the major area of research in which RNA interference is applied. A number of genes whose function has been identified through RNAi are SRP72 (Signal Recognition Particle) which functions in the cytoplasmic protein translocation pathway; PEPCK- a rate-controlling enzyme in gluconeogenesis; BCL2- a potent inhibitor of cell death; CAMK2G- which plays an important role in mammary gland development of goat, β -lactoglobulin- whey protein gene in swine.

Therapeutic applications

Diseases caused by microbes are the major cause of death

worldwide. Research using RNAi technique has been found to be effective in the treatment of a number of prominent viral diseases such as HIV and influenza, hepatitis and west Nile virus and bacterial infections such as pneumonia and sepsis. RNAi when used in cell culture studies has given promising results in inhibiting the replication or cellular uptake of viruses and other infectious agents. RNAi does not interact with the DNA of an individual rather it interferes with the translation process of gene expression. This would alleviate the patients concern as no changes to ones DNA is made as incase of gene therapy which suggests that this method of treatment would no more be feared than taking any prescription drug. For this reason therapies based on RNAi have attracted much interest of researchers in the pharmaceutical sector.

Transfection of human cells with siRNAs directed against different genes in the poliovirus genome resulted in resistance of the cells to infection with poliovirus (Gitlin et al., 2002). The ability of siRNAs targeting gene encoding the death receptor Fas to protect mice from liver failure and fibrosis in two models of autoimmune hepatitis has been tested. Intravenous injection of Fas siRNA reduced Fas protein levels in the livers of mice during 10-day period. Fas siRNA treatment was found to inhibit hepatocyte necrosis and inflammatory infiltration and markedly reduced serum concentrations of transaminases demonstrating a clear hepatoprotective effect of siRNA therapy (Song et al., 2003).

Potent and specific inhibition of HIV type 1 replication has been reported by RNAi, despite the success of RNAi mediated inhibition of HIV-encoded RNAs in cell culture, targeting the virus directly represents a substantial challenge as the viruses have a high mutation rate. Therefore down regulation (through RNAi) of cellular cofactors required for HIV infection presents an attractive alternative approach; some of the cellular cofactors that have been successfully downregulated by RNAi are NF-B (Surabhi and Gaynor, 2002) and HIV receptor CD4 (Novina et al., 2002).

The cancer cells exhibit two general abnormalities; firstly, dysregulation of cell cycle resulting in uncontrolled growth and secondly loss of apoptosis due to abnormalities in one or more proteins that mediate apoptosis. miRNAs such as let-7 and miR-15/miR-16 block Ras and BCL2 oncogenes and are promising candidates for cancer treatment (Hannon, 2002). Lentiviral vectors were used to produce transgenic porcine cells expressing shRNAs that inhibit FMD viron replication and mutation (Long and Westhusin, 2007). Transgenic *Anopheles* strains carrying miRNA constructs can silence the genes in the immune system of the mosquito that prevent *Plasmodium* recognition by immune system, such strain of *Anopheles* can be left to the field to replace the wild type and thus malaria can be controlled (Malhotra et al., 2002).

The signal recognition particle (SRP) is known to be

involved in the synthesis of membrane and secretory proteins across or into the endoplasmic reticulum (ER) membrane in eukaryotes (Köch et al., 2003). RNAi silencing of SRP72 and SRP68 lead to the depletion of these particles and induced sudden death in *T. brucei*, due to toxicity leading to accumulation of the pre-SRP in the nucleolus (Lustig et al., 2005). Suppression of prion (PrP) protein in transgenic goats was possible by RNAi and thus a significant (>90%) decrease in PrP expression levels was observed (through Western Blotting) in transgenic goats, carrying miRNA constructs that silence the genes coding for prion particles (Golding et al., 2006). Investigations are going to find out whether RNAi can be used as a remedy for treating various neurodegenerative disorders such as Alzheimer's, Huntington's or Parkinson's disease. One of the strategies for treating neurodegenerative disorders is by blocking disease specific events that initiate these disorders. Pro-apoptotic members of the Bcl-2 family were targeted by RNAi and neuronal death was successfully prevented (Colussi et al., 2000).

CHALLENGES FOR RNAi THERAPY

The RNAi can be used as a major tool in therapeutics however various challenges are there in developing RNAi as a therapy, such as i) efficient targeted delivery to specific cell or tissue types is still not a practical reality for oligonucleotide-based therapeutics ii) Systemic delivery of viral vectors is still a major hurdle iii) Off-target effects (OTE) - arise when an introduced RNA can pair with multiple genes at a time. Today the delivery related research in RNAi is receiving much desired attention.

Conflict of Interests

The author(s) have not declared any conflict of interests.

CONCLUSION

Studies started first in plants, flies, and worms paved way for RNAi for being discovered as an accidental observation. RNAi machinery can handle both endogenous and exogenous dsRNA and the faster identification of gene function is possible through RNAi. It is a powerful experimental tool in reverse genetics and this technique has wide spread future applications in medicine. RNA interference (RNAi) is a relatively new technology that is revolutionizing the way that researchers study mammalian gene expression. RNAi has had significant impact on the ease, speed, and specificity with which the loss of gene function analysis can be done in mammalian cells and animal models. This is becoming the method of choice for researchers doing

loss of function studies.

REFERENCES

- Bernstein E, Caudy AA, Hammond SM, Hannon GJ (2001). Role for a bidentate ribonuclease in the initiation step of RNA interference. *Nature*, 409:363-366.
- Cogoni C, Macino G (1999). Gene silencing in *Neurospora crassa* requires a protein homologous to RNA-dependent RNA polymerase. *Nature*, 399:166-169.
- Colussi PA, Quinn LM, Huang DC, Coombe M, Read SH, Richardson H, Kumar S (2000). Debcl, a proapoptotic Bcl-2 homologue, is a component of the *Drosophila melanogaster* cell death machinery. *J. Cell Biol.* 148:703-714.
- Dehio C, Schell J (1994). Identification of plant genetic loci involved in a post transcriptional mechanism for meiotically reversible transgene silencing. *Proceedings of the National Academy of Sciences of the United States of America*, 91(12):5538-5542.
- Djupedal I, Ekwall K (2009). Epigenetics: heterochromatin meets RNAi. *Cell Res.* 19:282-295.
- Filipowicz W (2005). RNAi: The Nuts and Bolts of the RISC Machine. *Cell*, 122:17-20.
- Filippo MC, Onorati MC, Kothe GO, Burroughs AM, Parsi KM, Breiling A, Federica LS, Saxena A, Miyoshi K, Siomi H, Siomi MC, Carninci P, David SG, Corona DFV, Orlando V (2011). Chromatin-associated RNA interference components contribute to transcriptional regulation in *Drosophila*. *Nature*, 480(7377):391-395.
- Fire A, Xu S, Montgomery MK, Kostas SA, Driver SE, Mello CC (1998). Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*. *Nat.* 391:806-811.
- Gitlin L, Karelsky S, Andino R (2002). Short interfering RNA confers intracellular antiviral immunity in human cells. *Nature*, 418:430-434.
- Hannon GJ (2002). RNA interference. *Nat.* 418:244.
- Köch HG, Hengelage T, Neumann-Haefelin C, MacFarlane J, Hoffschulte HK, Schimz KL, Mechler B, Müller M (1999). *In vitro* studies with purified components reveal signal recognition particle (SRP) and SecA/SecB as constituents of two independent protein-targeting pathways of *Escherichia coli*. *Mol. Biol. Cell*, 10:2163-2173.
- Long CR, Westhusin ME (2007). Development of Foot and Mouth Disease Resistant Livestock Using RNA Interference. *Veterinary Physiology & Pharmacology Texas A & M Univ. College Station, Texas* 77843.
- Lustig Y, Hanoch G, Shai U, Shulamit M (2005). The Trypanosoma brucei signal recognition particle lacks the Alu-domain-binding proteins: purification and functional analysis of its binding proteins by RNAi. *J. Cell Sci.* 118(19):4551-4562.
- Macrae IJ, Li F, Zhou K, Cande WZ, Doudna JA (2006). Structure of Dicer and Mechanistic Implications for RNAi. © 2006 Cold Spring Harbor Laboratory Press 978-087969817-1. *Cold Spring Harb. Symp. Quant. Biol.* 71:73-80.
- Malhotra P, Dasaradhi PV, Kumar A, Mohammed A, Agrawal N, Bhatnagar RK, Chauhan VS (2002). Double-stranded RNA-mediated RNAs gene silencing of cysteine proteases (falcipain-1 and -2) of *Plasmodium falciparum*. *Mol. Microbiol.* 45:1245-1254.
- Novina CD, Murray MF, Dykxhoorn DM, Beresford PJ, Riess J, Lee SK, Collman RG, Lieberman J, Shankar P, Sharp PA (2002). siRNA-directed inhibition of HIV-1 infection. *Nat. Med.* 8:681-686.
- Romano N, Macino G (1992). Quelling: Transient inactivation of gene expression in *Neurospora crassa* by transformation with homologous sequences. *Mol. Microbiol.* 6:3343-3353.
- Song E, Lee SK, Wang J, Ince N, Ouyang N, Min J, Chen J, Shankar P, Lieberman J (2003). RNA interference targeting Fas protects mice from fulminant hepatitis. *Nat. Med.* 9:347-351.
- Surabhi RM, Gaynor RB (2002). RNA interference directed against viral and cellular targets inhibits human immunodeficiency virus type-1 replication. *J. Virol.* 76:12963-12973.
- Volpe T, Martienssen RA (2011). RNA Interference and Heterochromatin Assembly. *Cold Spring Harb. Perspect. Biol.* 3:1-11.
- Wassenegger M (2005). The Role of the RNAi Machinery in Heterochromatin Formation: Minireview. *Cell*, 122:13-16.

Yang S, Stephen T, Eric P, Kyonggeun Y (2001). Specific Double-Stranded RNA Interference in Undifferentiated Mouse Embryonic Stem Cells. *Mol. Cell. Biol.* 21(22):7807-7816.

Zamore PD, Tuschl T, Sharp PA, Bartel DP (2000). RNAi: Double-stranded RNA directs the ATP dependent cleavage of mRNA at 21 to 23 nucleotide intervals. *Cell*, 101:25-33.



International Journal for Biotechnology and Molecular Biology Research

Related Journals Published by Academic Journals

- *African Journal of Environmental Science & Technology*
- *Biotechnology & Molecular Biology Reviews*
- *African Journal of Biochemistry Research*
- *African Journal of Microbiology Research*
- *African Journal of Pure & Applied Chemistry*
- *Journal of Cell and Animal Biology*
- *African Journal of Biotechnology*
- *Scientific Research and Essays*

academicJournals