

African Journal of Microbiology Research

Volume 9 Number 26, 1 July, 2015

ISSN 1996-0808



ABOUT AJMR

The **African Journal of Microbiology Research (AJMR)** (ISSN 1996-0808) is published Weekly (one volume per year) by Academic Journals.

African Journal of Microbiology Research (AJMR) provides rapid publication (weekly) of articles in all areas of Microbiology such as: Environmental Microbiology, Clinical Microbiology, Immunology, Virology, Bacteriology, Phycology, Mycology and Parasitology, Protozoology, Microbial Ecology, Probiotics and Prebiotics, Molecular Microbiology, Biotechnology, Food Microbiology, Industrial Microbiology, Cell Physiology, Environmental Biotechnology, Genetics, Enzymology, Molecular and Cellular Biology, Plant Pathology, Entomology, Biomedical Sciences, Botany and Plant Sciences, Soil and Environmental Sciences, Zoology, Endocrinology, Toxicology. The Journal welcomes the submission of manuscripts that meet the general criteria of significance and scientific excellence. Papers will be published shortly after acceptance. All articles are peer-reviewed.

Submission of Manuscript

Please read the **Instructions for Authors** before submitting your manuscript. The manuscript files should be given the last name of the first author

[Click here to Submit manuscripts online](#)

If you have any difficulty using the online submission system, kindly submit via this email ajmr@academicjournals.org.

With questions or concerns, please contact the Editorial Office at ajmr@academicjournals.org.

Editors

Prof. Dr. Stefan Schmidt,
*Applied and Environmental Microbiology
School of Biochemistry, Genetics and Microbiology
University of KwaZulu-Natal
Private Bag X01
Scottsville, Pietermaritzburg 3209
South Africa.*

Prof. Fukai Bao
*Department of Microbiology and Immunology
Kunming Medical University
Kunming 650031,
China*

Dr. Jianfeng Wu
*Dept. of Environmental Health Sciences,
School of Public Health,
University of Michigan
USA*

Dr. Ahmet Yilmaz Coban
*OMU Medical School,
Department of Medical Microbiology,
Samsun,
Turkey*

Dr. Seyed Davar Siadat
*Pasteur Institute of Iran,
Pasteur Square, Pasteur Avenue,
Tehran,
Iran.*

Dr. J. Stefan Rokem
*The Hebrew University of Jerusalem
Department of Microbiology and Molecular Genetics,
P.O.B. 12272, IL-91120 Jerusalem,
Israel*

Prof. Long-Liu Lin
*National Chiayi University
300 Syuefu Road,
Chiayi,
Taiwan*

N. John Tonukari, Ph.D
*Department of Biochemistry
Delta State University
PMB 1
Abraka, Nigeria*

Dr. Thaddeus Ezeji
*Assistant Professor
Fermentation and Biotechnology Unit
Department of Animal Sciences
The Ohio State University
1680 Madison Avenue
USA.*

Associate Editors

Dr. Mamadou Gueye
*MIRCEN/ Laboratoire commun de microbiologie
IRD-ISRA-UCAD, BP 1386,
DAKAR, Senegal.*

Dr. Caroline Mary Knox
*Department of Biochemistry, Microbiology and
Biotechnology
Rhodes University
Grahamstown 6140
South Africa.*

Dr. Hesham Elsayed Mostafa
*Genetic Engineering and Biotechnology Research
Institute (GEBRI)
Mubarak City For Scientific Research,
Research Area, New Borg El-Arab City,
Post Code 21934, Alexandria, Egypt.*

Dr. Wael Abbas El-Naggar
*Head of Microbiology Department,
Faculty of Pharmacy,
Mansoura University,
Mansoura 35516, Egypt.*

Dr. Abdel Nasser A. El-Moghazy
*Microbiology, Molecular Biology, Genetics Engineering
and Biotechnology
Dept of Microbiology and Immunology
Faculty of Pharmacy
Al-Azhar University
Nasr city,
Cairo, Egypt*

Editorial Board

Dr. Barakat S.M. Mahmoud

*Food Safety/Microbiology
Experimental Seafood Processing Laboratory
Costal Research and Extension Center
Mississippi State University
3411 Frederic Street
Pascagoula, MS 39567
USA*

Prof. Mohamed Mahrous Amer

*Poultry Disease (Viral Diseases of poultry)
Faculty of Veterinary Medicine,
Department of Poultry Diseases
Cairo university
Giza, Egypt*

Dr. Xiaohui Zhou

*Molecular Microbiology, Industrial Microbiology,
Environmental Microbiology, Pathogenesis, Antibiotic
resistance, Microbial Ecology
Washington State University
Bustad Hall 402 Department of Veterinary
Microbiology and Pathology, Pullman,
USA*

Dr. R. Balaji Raja

*Department of Biotechnology,
School of Bioengineering,
SRM University,
Chennai
India*

Dr. Aly E Abo-Amer

*Division of Microbiology, Botany Department, Faculty
of Science, Sohag University.
Egypt.*

Dr. Haoyu Mao

*Department of Molecular Genetics and Microbiology
College of Medicine
University of Florida
Florida, Gainesville
USA.*

Dr. Rachna Chandra

*Environmental Impact Assessment Division
Environmental Sciences
Sálim Ali Center for Ornithology and Natural History
(SACON),
Anaikatty (PO), Coimbatore-641108, India*

Dr. Yongxu Sun

*Department of Medicinal Chemistry and
Biomacromolecules
Qiqihar Medical University, Qiqihar 161006
Heilongjiang Province
P.R. China*

Dr. Ramesh Chand Kasana

*Institute of Himalayan Bioresource Technology
Palampur, Distt. Kangra (HP),
India*

Dr. S. Meena Kumari

*Department of Biosciences
Faculty of Science
University of Mauritius
Reduit*

Dr. T. Ramesh

*Assistant Professor
Marine Microbiology
CAS in Marine Biology
Faculty of Marine Sciences
Annamalai University
Parangipettai - 608 502
Cuddalore Dist. Tamilnadu,
India*

Dr. Pagano Marcela Claudia

*Post doctoral fellowship at Department of Biology,
Federal University of Ceará - UFC,
Brazil.*

Dr. EL-Sayed E. Habib

*Associate Professor,
Dept. of Microbiology,
Faculty of Pharmacy,
Mansoura University,
Egypt.*

Dr. Pongsak Rattanachaiakunsopon

*Department of Biological Science,
Faculty of Science,
Ubon Ratchathani University,
Warin Chamrap, Ubon Ratchathani 34190,
Thailand*

Dr. Gokul Shankar Sabesan

*Microbiology Unit, Faculty of Medicine,
AIMST University
Jalan Bedong, Semeling 08100,
Kedah,
Malaysia*

Dr. Kwang Young Song

*Department of Biological Engineering,
School of Biological and Chemical Engineering,
Yanbian University of Science and Technology,
Yanji,
China.*

Dr. Kamel Belhamel

*Faculty of Technology,
University of Bejaia
Algeria*

Dr. Sladjana Jevremovic

*Institute for Biological Research
Sinisa Stankovic,
Belgrade,
Serbia*

Dr. Tamer Edirne

*Dept. of Family Medicine, Univ. of Pamukkale
Turkey*

Dr. R. Balaji Raja M.Tech (Ph.D)

*Assistant Professor,
Department of Biotechnology,
School of Bioengineering,
SRM University,
Chennai.
India*

Dr. Minglei Wang

University of Illinois at Urbana-Champaign, USA

Dr. Mohd Fuat ABD Razak

*Institute for Medical Research
Malaysia*

Dr. Davide Pacifico

*Istituto di Virologia Vegetale – CNR
Italy*

Prof. Dr. Akrum Hamdy

*Faculty of Agriculture, Minia University, Egypt
Egypt*

Dr. Ntobeko A. B. Ntusi

*Cardiac Clinic, Department of Medicine,
University of Cape Town and
Department of Cardiovascular Medicine,
University of Oxford
South Africa and
United Kingdom*

Prof. N. S. Alzoreky

*Food Science & Nutrition Department,
College of Agricultural Sciences & Food,
King Faisal University,
Saudi Arabia*

Dr. Chen Ding

*College of Material Science and Engineering,
Hunan University,
China*

Dr Svetlana Nikolić

*Faculty of Technology and Metallurgy,
University of Belgrade,
Serbia*

Dr. Sivakumar Swaminathan

*Department of Agronomy,
College of Agriculture and Life Sciences,
Iowa State University,
Ames, Iowa 50011
USA*

Dr. Alfredo J. Anceno

*School of Environment, Resources and Development
(SERD),
Asian Institute of Technology,
Thailand*

Dr. Iqbal Ahmad

*Aligarh Muslim University,
Alighrah
India*

Dr. Josephine Nketsia-Tabiri

*Ghana Atomic Energy Commission
Ghana*

Dr. Juliane Elisa Welke

*UFRGS – Universidade Federal do Rio
Grande do Sul
Brazil*

Dr. Mohammad Nazrul Islam

*NIMR; IPH-Bangalore & NIUM
Bangladesh*

Dr. Okonko, Iheanyi Omezuruike

*Department of Virology,
Faculty of Basic Medical Sciences,
College of Medicine,
University of Ibadan,
University College Hospital,
Ibadan,
Nigeria*

Dr. Giuliana Noratto

*Texas A&M University
USA*

Dr. Phanikanth Venkata Turlapati

*Washington State University
USA*

Dr. Khaleel I. Z. Jawasreh

*National Centre for Agricultural Research and
Extension, NCARE
Jordan*

Dr. Babak Mostafazadeh, MD

*Shaheed Beheshti University of Medical Sciences
Iran*

Dr. S. Meena Kumari

*Department of Biosciences
Faculty of Science
University of Mauritius
Reduit
Mauritius*

Dr. S. Anju

*Department of Biotechnology,
SRM University, Chennai-603203
India*

Dr. Mustafa Maroufpor

Iran

Prof. Dong Zhichun

*Professor, Department of Animal Sciences and
Veterinary Medicine,
Yunnan Agriculture University,
China*

Dr. Mehdi Azami

*Parasitology & Mycology Dept,
Baghaeei Lab.,
Shams Abadi St.
Isfahan
Iran*

Dr. Anderson de Souza Sant'Ana

*University of São Paulo.
Brazil.*

Dr. Juliane Elisa Welke

*UFRGS – Universidade Federal do Rio Grande do Sul
Brazil*

Dr. Paul Shapshak

*USF Health,
Depts. Medicine (Div. Infect. Disease & Internat Med)
and Psychiatry & Beh Med.
USA*

Dr. Jorge Reinheimer

*Universidad Nacional del Litoral (Santa Fe)
Argentina*

Dr. Qin Liu

*East China University of Science
and Technology
China*

Dr. Xiao-Qing Hu

*State Key Lab of Food Science and Technology
Jiangnan University
P. R. China*

Prof. Branislava Kocic

*Specialist of Microbiology and Parasitology
University of Nis, School of Medicine Institute
for Public Health Nis, Bul. Z. Djindjica 50, 18000 Nis
Serbia*

Dr. Rafel Socias

*CITA de Aragón,
Spain*

Prof. Kamal I. Mohamed

*State University of New York at Oswego
USA*

Dr. Adriano Cruz

*Faculty of Food Engineering-FEA
University of Campinas (UNICAMP)
Brazil*

Dr. Mike Agenbag (Michael Hermanus Albertus)

*Manager Municipal Health Services,
Joe Gqabi District Municipality
South Africa*

Dr. D. V. L. Sarada

*Department of Biotechnology,
SRM University, Chennai-603203
India.*

Dr. Samuel K Ameyaw

*Civista Medical Center
United States of America*

Prof. Huaizhi Wang

*Institute of Hepatopancreatobiliary
Surgery of PLA Southwest Hospital,
Third Military Medical University
Chongqing400038
P. R. China*

Prof. Bakhiet AO

*College of Veterinary Medicine, Sudan
University of Science and Technology
Sudan*

Dr. Saba F. Hussain

*Community, Orthodontics and Paediatric Dentistry
Department
Faculty of Dentistry
Universiti Teknologi MARA
40450 Shah Alam, Selangor
Malaysia*

Prof. Dr. Zohair I.F.Rahemo

*State Key Lab of Food Science and Technology
Jiangnan University
P. R. China*

Dr. Afework Kassu

*University of Gondar
Ethiopia*

Prof. Isidro A. T. Savillo

*ISCOF
Philippines*

Dr. How-Yee Lai

*Taylor's University College
Malaysia*

Dr. Nidheesh Dadheech

*MS. University of Baroda, Vadodara, Gujarat, India.
India*

Dr. Omitoyin Siyanbola

*Bowen University,
Iwo
Nigeria*

Dr. Franco Mutinelli

*Istituto Zooprofilattico Sperimentale delle Venezie
Italy*

Dr. Chanpen Chanchao

*Department of Biology,
Faculty of Science,
Chulalongkorn University
Thailand*

Dr. Tsuyoshi Kasama

*Division of Rheumatology,
Showa University
Japan*

Dr. Kuender D. Yang, MD.

*Chang Gung Memorial Hospital
Taiwan*

Dr. Liane Raluca Stan

*University Politehnica of Bucharest,
Department of Organic Chemistry "C.Nenitzescu"
Romania*

Dr. Muhamed Osman

*Senior Lecturer of Pathology & Consultant
Immunopathologist
Department of Pathology,
Faculty of Medicine,
Universiti Teknologi MARA,
40450 Shah Alam, Selangor
Malaysia*

Dr. Mohammad Feizabadi

*Tehran University of medical Sciences
Iran*

Prof. Ahmed H Mitwalli

*State Key Lab of Food Science and Technology
Jiangnan University
P. R. China*

Dr. Mazyar Yazdani

*Department of Biology,
University of Oslo,
Blindern,
Oslo,
Norway*

Dr. Ms. Jemimah Gesare Onsare

*Ministry of Higher, Education
Science and Technology
Kenya*

Dr. Babak Khalili Hadad

*Department of Biological Sciences,
Roudehen Branch,
Islamic Azad University,
Roudehen
Iran*

Dr. Ehsan Sari

*Department of Plant Pathology,
Iranian Research Institute of Plant Protection,
Tehran,
Iran.*

Dr. Snjezana Zidovec Lepej

*University Hospital for Infectious Diseases
Zagreb,
Croatia*

Dr. Dilshad Ahmad

*King Saud University
Saudi Arabia*

Dr. Adriano Gomes da Cruz

*University of Campinas (UNICAMP)
Brazil*

Dr. Hsin-Mei Ku

*Agronomy Dept. NCHU 250 Kuo
Kuang Rd, Taichung,
Taiwan*

Dr. Fereshteh Naderi

*Physical chemist,
Islamic Azad University,
Shahre Ghods Branch
Iran*

Dr. Adibe Maxwell Ogochukwu

*Department of Clinical Pharmacy and Pharmacy
Management,
University of Nigeria,
Nsukka.
Nigeria*

Dr. William M. Shafer

*Emory University School of Medicine
USA*

Dr. Michelle Bull

*CSIRO Food and Nutritional Sciences
Australia*

Prof. Dr. Márcio Garcia Ribeiro (DVM, PhD)

*School of Veterinary Medicine and Animal Science-
UNESP,
Dept. Veterinary Hygiene and Public Health,
State of Sao Paulo
Brazil*

Prof. Dr. Sheila Nathan

*National University of Malaysia (UKM)
Malaysia*

Prof. Ebiamadon Andi Brisibe

*University of Calabar,
Calabar,
Nigeria*

Dr. Julie Wang

*Burnet Institute
Australia*

Dr. Jean-Marc Chobert

*INRA- BIA, FIPL
France*

Dr. Zhilong Yang, PhD

*Laboratory of Viral Diseases
National Institute of Allergy and Infectious Diseases,
National Institutes of Health*

Dr. Dele Raheem

*University of Helsinki
Finland*

Dr. Li Sun

*PLA Centre for the treatment of infectious diseases,
Tangdu Hospital,
Fourth Military Medical University
China*

Dr. Biljana Miljkovic-Selimovic

*School of Medicine,
University in Nis,
Serbia; Referent laboratory for Campylobacter and
Helicobacter,
Center for Microbiology,
Institute for Public Health, Nis
Serbia*

Dr. Xinan Jiao

*Yangzhou University
China*

Dr. Endang Sri Lestari, MD.

*Department of Clinical Microbiology,
Medical Faculty,
Diponegoro University/Dr. Kariadi Teaching Hospital,
Semarang
Indonesia*

Dr. Hojin Shin

*Pusan National University Hospital
South Korea*

Dr. Yi Wang

*Center for Vector Biology, 180 Jones Avenue
Rutgers University, New Brunswick, NJ 08901-8536
USA*

Dr. Heping Zhang

*The Key Laboratory of Dairy Biotechnology and
Engineering,
Ministry of Education,
Inner Mongolia Agricultural University.
China*

Prof. Natasha Potgieter

*University of Venda
South Africa*

Dr. Alemzadeh

*Sharif University
Iran*

Dr. Sonia Arriaga

*Instituto Potosino de Investigación Científica y
Tecnológica/División de Ciencias Ambientales
Mexico*

Dr. Armando Gonzalez-Sanchez

*Universidad Autonoma Metropolitana Cuajimalpa
Mexico*

Dr. Pradeep Parihar

*Lovely Professional University, Phagwara, Punjab.
India*

Dr. William H Roldán

*Department of Medical Microbiology,
Faculty of Medicine,
Peru*

Dr. Kanzaki, L I B

*Laboratory of Bioprospection. University of Brasilia
Brazil*

Prof. Philippe Dorchie

*Laboratory of Bioprospection. University of Brasilia
Brazil*

Dr. C. Ganesh Kumar

*Indian Institute of Chemical Technology,
Hyderabad
India*

Dr. Farid Che Ghazali

*Universiti Sains Malaysia (USM)
Malaysia*

Dr. Samira Bouhdid

*Abdelmalek Essaadi University,
Tetouan,
Morocco*

Dr. Zainab Z. Ismail

*Department of Environmental Engineering, University
of Baghdad.
Iraq*

Dr. Ary Fernandes Junior

*Universidade Estadual Paulista (UNESP)
Brasil*

Dr. Papaevangelou Vassiliki

*Athens University Medical School
Greece*

Dr. Fangyou Yu

*The first Affiliated Hospital of Wenzhou Medical
College
China*

Dr. Galba Maria de Campos Takaki

*Catholic University of Pernambuco
Brazil*

Dr. Kwabena Ofori-Kwakye

*Department of Pharmaceutics,
Kwame Nkrumah University of Science & Technology,
KUMASI
Ghana*

Prof. Dr. Liesel Brenda Gende

*Arthropods Laboratory, School of Natural and Exact
Sciences, National University of Mar del Plata
Buenos Aires,
Argentina.*

Dr. Adeshina Gbonjubola

*Ahmadu Bello University,
Zaria.
Nigeria*

Prof. Dr. Stylianos Chatzipanagiotou

*University of Athens – Medical School
Greece*

Dr. Dongqing BAI

*Department of Fishery Science,
Tianjin Agricultural College,
Tianjin 300384
P. R. China*

Dr. Dingqiang Lu

*Nanjing University of Technology
P.R. China*

Dr. L. B. Sukla

*Scientist –G & Head, Biominerals Department,
IMMT, Bhubaneswar
India*

Dr. Hakan Parlakpınar

*MD. Inonu University, Medical Faculty, Department
of Pharmacology, Malatya
Turkey*

Dr Pak-Lam Yu

*Massey University
New Zealand*

Dr Percy Chimwamurombe

*University of Namibia
Namibia*

Dr. Euclésio Simionatto

*State University of Mato Grosso do Sul-UEMS
Brazil*

Dr. Hans-Jürg Monstein

*Clinical Microbiology, Molecular Biology Laboratory,
University Hospital, Faculty of Health Sciences, S-581
85 Linköping
Sweden*

Dr. Ajith, T. A

*Associate Professor Biochemistry, Amala Institute of
Medical Sciences, Amala Nagar, Thrissur, Kerala-680
555
India*

Dr. Feng-Chia Hsieh

*Biopesticides Division, Taiwan Agricultural Chemicals
and Toxic Substances Research Institute, Council of
Agriculture
Taiwan*

Prof. Dra. Suzan Pantaroto de Vasconcellos

*Universidade Federal de São Paulo
Rua Prof. Artur Riedel, 275 Jd. Eldorado, Diadema, SP
CEP 09972-270
Brasil*

Dr. Maria Leonor Ribeiro Casimiro Lopes Assad

*Universidade Federal de São Carlos - Centro de
Ciências Agrárias - CCA/UFSCar
Departamento de Recursos Naturais e Proteção
Ambiental
Rodovia Anhanguera, km 174 - SP-330
Araras - São Paulo
Brasil*

Dr. Pierangeli G. Vital

*Institute of Biology, College of Science, University of
the Philippines
Philippines*

Prof. Roland Ndip

*University of Fort Hare, Alice
South Africa*

Dr. Shawn Carraher

*University of Fort Hare, Alice
South Africa*

Dr. José Eduardo Marques Pessanha

*Observatório de Saúde Urbana de Belo
Horizonte/Faculdade de Medicina da Universidade
Federal de Minas Gerais
Brasil*

Dr. Yuanshu Qian

*Department of Pharmacology, Shantou University
Medical College
China*

Dr. Helen Treichel

*URI-Campus de Erechim
Brazil*

Dr. Xiao-Qing Hu

*State Key Lab of Food Science and Technology
Jiangnan University
P. R. China*

Dr. Olli H. Tuovinen

*Ohio State University, Columbus, Ohio
USA*

Prof. Stoyan Groudev

*University of Mining and Geology "Saint Ivan Rilski"
Sofia
Bulgaria*

Dr. G. Thirumurugan

*Research lab, GIET School of Pharmacy, NH-5,
Chaitanya nagar, Rajahmundry-533294.
India*

Dr. Charu Gomber

*Thapar University
India*

Dr. Jan Kuever

*Bremen Institute for Materials Testing,
Department of Microbiology,
Paul-Feller-Str. 1, 28199 Bremen
Germany*

Dr. Nicola S. Flanagan

*Universidad Javeriana, Cali
Colombia*

Dr. André Luiz C. M. de A. Santiago

*Universidade Federal Rural de Pernambuco
Brazil*

Dr. Dhruva Kumar Jha

*Microbial Ecology Laboratory,
Department of Botany,
Gauhati University,
Guwahati 781 014, Assam
India*

Dr. N Saleem Basha

*M. Pharm (Pharmaceutical Biotechnology)
Eritrea (North East Africa)*

Prof. Dr. João Lúcio de Azevedo

*Dept. Genetics-University of São Paulo-Faculty of
Agriculture- Piracicaba, 13400-970
Brasil*

Dr. Julia Inés Fariña

*PROIMI-CONICET
Argentina*

Dr. Yutaka Ito

*Kyoto University
Japan*

Dr. Cheruiyot K. Ronald

*Biomedical Laboratory Technologist
Kenya*

Prof. Dr. Ata Akcil

*S. D. University
Turkey*

Dr. Adhar Manna

*The University of South Dakota
USA*

Dr. Cícero Flávio Soares Aragão

*Federal University of Rio Grande do Norte
Brazil*

Dr. Gunnar Dahlen

*Institute of odontology, Sahlgrenska Academy at
University of Gothenburg
Sweden*

Dr. Pankaj Kumar Mishra

*Vivekananda Institute of Hill Agriculture, (I.C.A.R.),
ALMORA-263601, Uttarakhand
India*

Dr. Benjamas W. Thanomsub

*Srinakharinwirot University
Thailand*

Dr. Maria José Borrego

*National Institute of Health – Department of Infectious
Diseases
Portugal*

Dr. Catherine Carrillo

*Health Canada, Bureau of Microbial Hazards
Canada*

Dr. Marcotty Tanguy

*Institute of Tropical Medicine
Belgium*

Dr. Han-Bo Zhang

*Laboratory of Conservation and Utilization for Bio-
resources
Key Laboratory for Microbial Resources of the
Ministry of Education,
Yunnan University, Kunming 650091.
School of Life Science,
Yunnan University, Kunming,
Yunnan Province 650091.
China*

Dr. Ali Mohammed Somily

*King Saud University
Saudi Arabia*

Dr. Nicole Wolter

*National Institute for Communicable Diseases and
University of the Witwatersrand,
Johannesburg
South Africa*

Dr. Marco Antonio Nogueira

*Universidade Estadual de Londrina
CCB/Depto. De microbiologia
Laboratório de Microbiologia Ambiental
Caixa Postal 6001
86051-980 Londrina.
Brazil*

Dr. Bruno Pavoni

*Department of Environmental Sciences University of
Venice
Italy*

Dr. Shih-Chieh Lee

*Da-Yeh University
Taiwan*

Dr. Satoru Shimizu

*Horonobe Research Institute for the Subsurface
Environment,
Northern Advancement Center for Science &
Technology
Japan*

Dr. Tang Ming

*College of Forestry, Northwest A&F University,
Yangling
China*

Dr. Olga Gortzi

*Department of Food Technology, T.E.I. of Larissa
Greece*

Dr. Mark Tarnopolsky

*Mcmaster University
Canada*

Dr. Sami A. Zabin

*Al Baha University
Saudi Arabia*

Dr. Julia W. Pridgeon

*Aquatic Animal Health Research Unit, USDA, ARS
USA*

Dr. Lim Yau Yan

*Monash University Sunway Campus
Malaysia*

Prof. Rosemeire C. L. R. Pietro

*Faculdade de Ciências Farmacêuticas de Araraquara,
Univ Estadual Paulista, UNESP
Brazil*

Dr. Nazime Mercan Dogan

*PAU Faculty of Arts and Science, Denizli
Turkey*

Dr Ian Edwin Cock

*Biomolecular and Physical Sciences
Griffith University
Australia*

Prof. N K Dubey

*Banaras Hindu University
India*

Dr. S. Hemalatha

*Department of Pharmaceutics, Institute of
Technology,
Banaras Hindu University, Varanasi. 221005
India*

Dr. J. Santos Garcia A.

*Universidad A. de Nuevo Leon
Mexico India*

Dr. Somboon Tanasupawat

*Department of Biochemistry and Microbiology,
Faculty of Pharmaceutical Sciences,
Chulalongkorn University,
Bangkok 10330
Thailand*

Dr. Vivekananda Mandal

*Post Graduate Department of Botany,
Darjeeling Government College,
Darjeeling – 734101.
India*

Dr. Shihua Wang

*College of Life Sciences,
Fujian Agriculture and Forestry University
China*

Dr. Victor Manuel Fernandes Galhano

*CITAB-Centre for Research and Technology of Agro-
Environment and Biological Sciences, Integrative
Biology and Quality Research Group,
University of Trás-os-Montes and Alto Douro,
Apartado 1013, 5001-801 Vila Real
Portugal*

Dr. Maria Cristina Maldonado

*Instituto de Biotecnología. Universidad Nacional de
Tucuman
Argentina*

Dr. Alex Soltermann

*Institute for Surgical Pathology,
University Hospital Zürich
Switzerland*

Dr. Dagmara Sirova

*Department of Ecosystem Biology, Faculty Of Science,
University of South Bohemia,
Branisovska 37, Ceske Budejovice, 37001
Czech Republic*

Dr. E. O Igbinosa

*Department of Microbiology,
Ambrose Alli University,
Ekpoma, Edo State,
Nigeria.*

Dr. Hodaka Suzuki

*National Institute of Health Sciences
Japan*

Dr. Mick Bosilevac

*US Meat Animal Research Center
USA*

Dr. Nora Lía Padola

*Imunoquímica y Biotecnología- Fac Cs Vet-UNCPBA
Argentina*

Dr. Maria Madalena Vieira-Pinto

*Universidade de Trás-os-Montes e Alto Douro
Portugal*

Dr. Stefano Morandi

*CNR-Istituto di Scienze delle Produzioni Alimentari
(ISPA), Sez. Milano
Italy*

Dr Line Thorsen

*Copenhagen University, Faculty of Life Sciences
Denmark*

Dr. Ana Lucia Falavigna-Guilherme

*Universidade Estadual de Maringá
Brazil*

Dr. Baoqiang Liao

*Dept. of Chem. Eng., Lakehead University, 955 Oliver
Road, Thunder Bay, Ontario
Canada*

Dr. Ouyang Jinping

*Patho-Physiology department,
Faculty of Medicine of Wuhan University
China*

Dr. John Sorensen

*University of Manitoba
Canada*

Dr. Andrew Williams

*University of Oxford
United Kingdom*

Dr. Chi-Chiang Yang

*Chung Shan Medical University
Taiwan, R.O.C.*

Dr. Quanming Zou

*Department of Clinical Microbiology and Immunology,
College of Medical Laboratory,
Third Military Medical University
China*

Prof. Ashok Kumar

*School of Biotechnology,
Banaras Hindu University, Varanasi
India*

Dr. Chung-Ming Chen

*Department of Pediatrics, Taipei Medical University
Hospital, Taipei
Taiwan*

Dr. Jennifer Furin

*Harvard Medical School
USA*

Dr. Julia W. Pridgeon

*Aquatic Animal Health Research Unit, USDA, ARS
USA*

Dr. Alireza Seidavi

*Islamic Azad University, Rasht Branch
Iran*

Dr. Thore Rohwerder

*Helmholtz Centre for Environmental Research UFZ
Germany*

Dr. Daniela Billi

*University of Rome Tor Vergata
Italy*

Dr. Ivana Karabegovic

*Faculty of Technology, Leskovac, University of Nis
Serbia*

Dr. Flaviana Andrade Faria

*IBILCE/UNESP
Brazil*

Prof. Margareth Linde Athayde

*Federal University of Santa Maria
Brazil*

Dr. Guadalupe Virginia Nevarez Moorillon

*Universidad Autonoma de Chihuahua
Mexico*

Dr. Tatiana de Sousa Fiuza

*Federal University of Goias
Brazil*

Dr. Indrani B. Das Sarma

*Jhulelal Institute of Technology, Nagpur
India*

Dr. Guanghua Wang

*Northeast Institute of Geography and Agroecology,
Chinese Academy of Sciences
China*

Dr. Renata Vadkertiova

*Institute of Chemistry, Slovak Academy of Science
Slovakia*

Dr. Charles Hocart

*The Australian National University
Australia*

Dr. Guoqiang Zhu

*University of Yangzhou College of Veterinary Medicine
China*

Dr. Guilherme Augusto Marietto Gonçalves

*São Paulo State University
Brazil*

Dr. Mohammad Ali Faramarzi

*Tehran University of Medical Sciences
Iran*

Dr. Suppasil Maneerat

*Department of Industrial Biotechnology, Faculty of
Agro-Industry, Prince of Songkla University, Hat Yai
90112
Thailand*

Dr. Francisco Javier Las heras Vazquez

*Almeria University
Spain*

Dr. Cheng-Hsun Chiu

*Chang Gung memorial Hospital, Chang Gung
University
Taiwan*

Dr. Ajay Singh

*DDU Gorakhpur University, Gorakhpur-273009 (U.P.)
India*

Dr. Karabo Shale

*Central University of Technology, Free State
South Africa*

Dr. Lourdes Zélia Zanoni

*Department of Pediatrics, School of Medicine, Federal
University of Mato Grosso do Sul, Campo Grande,
Mato Grosso do Sul
Brazil*

Dr. Tulin Askun

*Balikesir University
Turkey*

Dr. Marija Stankovic

*Institute of Molecular Genetics and Genetic
Engineering
Republic of Serbia*

Dr. Scott Weese

*University of Guelph
Dept of Pathobiology, Ontario Veterinary College,
University of Guelph,
Guelph, Ontario, N1G2W1,
Canada*

Dr. Sabiha Essack

*School of Health Sciences
South African Committee of Health Sciences
University of KwaZulu-Natal
Private Bag X54001
Durban 4000
South Africa*

Dr. Hare Krishna

*Central Institute for Arid Horticulture,
Beechwal, Bikaner-334 006, Rajasthan,
India*

Dr. Anna Mensuali

*Dept. of Life Science,
Scuola Superiore
Sant'Anna*

Dr. Ghada Sameh Hafez Hassan

*Pharmaceutical Chemistry Department,
Faculty of Pharmacy, Mansoura University,
Egypt*

Dr. Kátia Flávia Fernandes

*Biochemistry and Molecular Biology
Universidade Federal de Goiás
Brasil*

Dr. Abdel-Hady El-Gilany

*Public Health & Community Medicine
Faculty of Medicine,
Mansoura University
Egypt*

Dr. Hongxiong Guo

*STD and HIV/AIDS Control and Prevention,
Jiangsu provincial CDC,
China*

Dr. Konstantina Tsaousi

*Life and Health Sciences,
School of Biomedical Sciences,
University of Ulster*

Dr. Bhavnaben Gowan Gordhan

*DST/NRF Centre of Excellence for Biomedical TB
Research
University of the Witwatersrand and National Health
Laboratory Service
P.O. Box 1038, Johannesburg 2000,
South Africa*

Dr. Ernest Kuchar

*Pediatric Infectious Diseases,
Wroclaw Medical University,
Wroclaw Teaching Hospital,
Poland*

Dr. Hongxiong Guo

*STD and HIV/AIDS Control and Prevention,
Jiangsu provincial CDC,
China*

Dr. Mar Rodriguez Jovita

*Food Hygiene and Safety, Faculty of Veterinary
Science.
University of Extremadura,
Spain*

Dr. Jes Gitz Holler

*Hospital Pharmacy,
Aalesund. Central Norway Pharmaceutical Trust
Professor Brochs gt. 6. 7030 Trondheim,
Norway*

Prof. Chengxiang FANG

*College of Life Sciences,
Wuhan University
Wuhan 430072, P.R.China*

Dr. Anchalee Tungtrongchitr

*Siriraj Dust Mite Center for Services and Research
Department of Parasitology,
Faculty of Medicine Siriraj Hospital,
Mahidol University
2 Prannok Road, Bangkok Noi,
Bangkok, 10700, Thailand*

Instructions for Author

Electronic submission of manuscripts is strongly encouraged, provided that the text, tables, and figures are included in a single Microsoft Word file (preferably in Arial font).

The **cover letter** should include the corresponding author's full address and telephone/fax numbers and should be in an e-mail message sent to the Editor, with the file, whose name should begin with the first author's surname, as an attachment.

Article Types

Three types of manuscripts may be submitted:

Regular articles: These should describe new and carefully confirmed findings, and experimental procedures should be given in sufficient detail for others to verify the work. The length of a full paper should be the minimum required to describe and interpret the work clearly.

Short Communications: A Short Communication is suitable for recording the results of complete small investigations or giving details of new models or hypotheses, innovative methods, techniques or apparatus. The style of main sections need not conform to that of full-length papers. Short communications are 2 to 4 printed pages (about 6 to 12 manuscript pages) in length.

Reviews: Submissions of reviews and perspectives covering topics of current interest are welcome and encouraged. Reviews should be concise and no longer than 4-6 printed pages (about 12 to 18 manuscript pages). Reviews are also peer-reviewed.

Review Process

All manuscripts are reviewed by an editor and members of the Editorial Board or qualified outside reviewers. Authors cannot nominate reviewers. Only reviewers randomly selected from our database with specialization in the subject area will be contacted to evaluate the manuscripts. The process will be blind review.

Decisions will be made as rapidly as possible, and the Journal strives to return reviewers' comments to authors as fast as possible. The editorial board will re-review manuscripts that are accepted pending revision. It is the goal of the AJMR to publish manuscripts within weeks after submission.

Regular articles

All portions of the manuscript must be typed double-spaced and all pages numbered starting from the title page.

The Title should be a brief phrase describing the contents of the paper. The Title Page should include the authors' full names and affiliations, the name of the corresponding author along with phone, fax and E-mail information. Present addresses of authors should appear as a footnote.

The Abstract should be informative and completely self-explanatory, briefly present the topic, state the scope of the experiments, indicate significant data, and point out major findings and conclusions. The Abstract should be 100 to 200 words in length. Complete sentences, active verbs, and the third person should be used, and the abstract should be written in the past tense. Standard nomenclature should be used and abbreviations should be avoided. No literature should be cited.

Following the abstract, about 3 to 10 key words that will provide indexing references should be listed.

A list of non-standard **Abbreviations** should be added. In general, non-standard abbreviations should be used only when the full term is very long and used often. Each abbreviation should be spelled out and introduced in parentheses the first time it is used in the text. Only recommended SI units should be used. Authors should use the solidus presentation (mg/ml). Standard abbreviations (such as ATP and DNA) need not be defined.

The Introduction should provide a clear statement of the problem, the relevant literature on the subject, and the proposed approach or solution. It should be understandable to colleagues from a broad range of scientific disciplines.

Materials and methods should be complete enough to allow experiments to be reproduced. However, only truly new procedures should be described in detail; previously published procedures should be cited, and important modifications of published procedures should be mentioned briefly. Capitalize trade names and include the manufacturer's name and address. Subheadings should be used. Methods in general use need not be described in detail.

Results should be presented with clarity and precision. The results should be written in the past tense when describing findings in the authors' experiments. Previously published findings should be written in the present tense. Results should be explained, but largely without referring to the literature. Discussion, speculation and detailed interpretation of data should not be included in the Results but should be put into the Discussion section.

The Discussion should interpret the findings in view of the results obtained in this and in past studies on this topic. State the conclusions in a few sentences at the end of the paper. The Results and Discussion sections can include subheadings, and when appropriate, both sections can be combined.

The Acknowledgments of people, grants, funds, etc should be brief.

Tables should be kept to a minimum and be designed to be as simple as possible. Tables are to be typed double-spaced throughout, including headings and footnotes. Each table should be on a separate page, numbered consecutively in Arabic numerals and supplied with a heading and a legend. Tables should be self-explanatory without reference to the text. The details of the methods used in the experiments should preferably be described in the legend instead of in the text. The same data should not be presented in both table and graph form or repeated in the text.

Figure legends should be typed in numerical order on a separate sheet. Graphics should be prepared using applications capable of generating high resolution GIF, TIFF, JPEG or Powerpoint before pasting in the Microsoft Word manuscript file. Tables should be prepared in Microsoft Word. Use Arabic numerals to designate figures and upper case letters for their parts (Figure 1). Begin each legend with a title and include sufficient description so that the figure is understandable without reading the text of the manuscript. Information given in legends should not be repeated in the text.

References: In the text, a reference identified by means of an author's name should be followed by the date of the reference in parentheses. When there are more than two authors, only the first author's name should be mentioned, followed by 'et al'. In the event that an author cited has had two or more works published during the same year, the reference, both in the text and in the reference list, should be identified by a lower case letter like 'a' and 'b' after the date to distinguish the works.

Examples:

Abayomi (2000), Agindotan et al. (2003), (Kelebeni, 1983), (Usman and Smith, 1992), (Chege, 1998;

1987a,b; Tijani, 1993,1995), (Kumasi et al., 2001) References should be listed at the end of the paper in alphabetical order. Articles in preparation or articles submitted for publication, unpublished observations, personal communications, etc. should not be included in the reference list but should only be mentioned in the article text (e.g., A. Kingori, University of Nairobi, Kenya, personal communication). Journal names are abbreviated according to Chemical Abstracts. Authors are fully responsible for the accuracy of the references.

Examples:

Chikere CB, Omoni VT and Chikere BO (2008). Distribution of potential nosocomial pathogens in a hospital environment. *Afr. J. Biotechnol.* 7: 3535-3539.

Moran GJ, Amii RN, Abrahamian FM, Talan DA (2005). Methicillinresistant *Staphylococcus aureus* in community-acquired skin infections. *Emerg. Infect. Dis.* 11: 928-930.

Pitout JDD, Church DL, Gregson DB, Chow BL, McCracken M, Mulvey M, Laupland KB (2007). Molecular epidemiology of CTXM-producing *Escherichia coli* in the Calgary Health Region: emergence of CTX-M-15-producing isolates. *Antimicrob. Agents Chemother.* 51: 1281-1286.

Pelczar JR, Harley JP, Klein DA (1993). *Microbiology: Concepts and Applications.* McGraw-Hill Inc., New York, pp. 591-603.

Short Communications

Short Communications are limited to a maximum of two figures and one table. They should present a complete study that is more limited in scope than is found in full-length papers. The items of manuscript preparation listed above apply to Short Communications with the following differences: (1) Abstracts are limited to 100 words; (2) instead of a separate Materials and Methods section, experimental procedures may be incorporated into Figure Legends and Table footnotes; (3) Results and Discussion should be combined into a single section.

Proofs and Reprints: Electronic proofs will be sent (e-mail attachment) to the corresponding author as a PDF file. Page proofs are considered to be the final version of the manuscript. With the exception of typographical or minor clerical errors, no changes will be made in the manuscript at the proof stage.

Fees and Charges: Authors are required to pay a \$550 handling fee. Publication of an article in the African Journal of Microbiology Research is not contingent upon the author's ability to pay the charges. Neither is acceptance to pay the handling fee a guarantee that the paper will be accepted for publication. Authors may still request (in advance) that the editorial office waive some of the handling fee under special circumstances

Copyright: © 2015, Academic Journals.

All rights Reserved. In accessing this journal, you agree that you will access the contents for your own personal use but not for any commercial use. Any use and or copies of this Journal in whole or in part must include the customary bibliographic citation, including author attribution, date and article title.

Submission of a manuscript implies: that the work described has not been published before (except in the form of an abstract or as part of a published lecture, or thesis) that it is not under consideration for publication elsewhere; that if and when the manuscript is accepted for publication, the authors agree to automatic transfer of the copyright to the publisher.

Disclaimer of Warranties

In no event shall Academic Journals be liable for any special, incidental, indirect, or consequential damages of any kind arising out of or in connection with the use of the articles or other material derived from the AJMR, whether or not advised of the possibility of damage, and on any theory of liability.

This publication is provided "as is" without warranty of any kind, either expressed or implied, including, but not limited to, the implied warranties of merchantability, fitness for a particular purpose, or non-infringement. Descriptions of, or references to, products or publications does not imply endorsement of that product or publication. While every effort is made by Academic Journals to see that no inaccurate or misleading data, opinion or statements appear in this publication, they wish to make it clear that the data and opinions appearing in the articles and advertisements herein are the responsibility of the contributor or advertiser concerned. Academic Journals makes no warranty of any kind, either express or implied, regarding the quality, accuracy, availability, or validity of the data or information in this publication or of any other publication to which it may be linked.

African Journal of Microbiology Research

Table of Content: Volume 9 Number 26, 1 July, 2015

ARTICLES

Use of thermal imaging for the early detection of signs of disease in pigs challenged orally with *Salmonella typhimurium* and *Escherichia coli*

Md. Manirul Islam, Sonia Tabasum Ahmed, Hong-Seok Mun, A. B. M. Rubayet Bostami, Yae-Jin Kim and Chul-Ju Yang

Prevalence of triple viral infections of human immunodeficiency virus (HIV), hepatitis B and C among tuberculosis patients and associated risk factors: The case of West Arsi Zone, Ethiopia

Endale Mengesha, Tekle Airgecho, Edessa Negera and Mulugeta Kebede

Full Length Research Paper

Use of thermal imaging for the early detection of signs of disease in pigs challenged orally with *Salmonella typhimurium* and *Escherichia coli*

Md. Manirul Islam, Sonia Tabasum Ahmed, Hong-Seok Mun, A. B. M. Rubayet Bostami, Yae-Jin Kim and Chul-Ju Yang*

Department of Animal Science and Technology, Suncheon National University 255 Jungangno, Suncheon, Jeonnam 540-950, Republic of Korea.

Received 13 May, 2015; Accepted 22 June, 2015

A total of 27 piglets were randomly assigned to three treatment groups (control, infection with *Salmonella enterica* serovar Typhimurium KCTC 2515 (ST) and infection with *Escherichia coli* KCTC 2571 (EC)) in a completely randomized design to early detection of signs of disease based on body temperature. Totally three inoculations were done consisting of 14 days each and then thermal images were captured to calculate body temperature of pigs at 0, 2, 6, 12, and 24 h and then every 24 h up to 14 days post inoculation. A reduced average daily gain (ADG) was observed in the first week of all three post inoculation ($P < 0.05$), while the gain:feed ratio was decreased at first week during first inoculation in both ST and EC group compared to control ($P < 0.05$). Body temperature was elevated in ST infected piglets at 24 h, peaked at 72 h ($P < 0.05$) and remained elevated, however, the EC induced piglets showed a subnormal body temperature throughout the experimental period relative to the control ($P < 0.05$). Taken together, the results indicate that signs of disease following experimentally induced bacterial infection in pigs can be detected quickly and easily using thermal images.

Key words: Piglets, growth performance, thermal image, body temperature, disease diagnosis.

INTRODUCTION

Infections with enteric pathogens such as *Salmonella* or *Escherichia coli* are common during intensive pig production and are associated with poor performance and animal welfare (Pijpers et al., 1991; Greiner et al., 2000). Such infections may also pose a risk to human health and lead to expensive veterinarian intervention costs. Delays in detection may result in the outbreak of

diseases and even more expenses. Detection of animal illness and providing individual and group-by-group mass therapy is not effective. Since infection is primarily spread via horizontal transmission from carrier animals to others in close contact (Schwartz, 1991), the best way to handle an illness is early detection of diseases followed by application of treatment before the disease can spread.

*Corresponding author. E-mail: yangcj@scnu.kr. Tel: +82-61-750-3235. Fax: +82-61-750-3239

Healthy animals are important to profit and performance, as well as animal welfare. Timely interventions may interrupt the lifetime carriage of *Salmonella* and *E. coli* and improve the health and growth of weaned pigs.

Swine have an unusual means of regulating core temperature. Due to the low number of sweat glands (Moritz and Henriques, 1947; Montagna and Yun, 1964), pigs reduce their core temperature by increasing the rate of peripheral blood flow through the skin, panting, or moistening the skin with water. However, panting and evaporation are less efficient regulation systems under some environmental conditions and are not always possible in confined pig housing. Fever is the earliest and one of the main clinical signs of many diseases (Plonait, 2004). Previous investigations showed that *S. typhimurium* challenged piglets showed increased body temperature (Balaji et al., 2000), while *E. coli* endotoxin inoculation resulted in subnormal body temperature (Halloy et al., 2004) compared to controls. Based on the way heat is transmitted through the skin of swine, infrared images have the potential for use in the detection of body temperature (Ingram and Weaver, 1969; Godynicki et al., 1985).

Thraulsen et al. (2010) concluded that infrared thermography allows routine measurements of body surface temperature that can be used for early disease detection. Röhlinger et al. (1979) reported the possibility of using infrared cameras (IRCs) to measure body surface temperature for early disease detection in various animals, including swine. Scolari et al. (2011) used an IRC to detect the rise and fall of vulvar skin temperature of sows during estrus. Therefore, the present study was conducted to measure the growth performance and potential for early detection of sign of disease based on body temperature patterns measured using a thermal camera as an early disease diagnostic tool in *Salmonella enterica* serover Typhimurium and *E. coli* infected piglets.

MATERIALS AND METHODS

All experimental procedures used in this study were approved by the Animal Care and Use Committee of Suncheon National University, South Korea.

Animals and experimental design

A total of 27 newly weaned piglets (crossbred ((Landrace x Yorkshire) x Duroc), and mean body weight 8 kg and 28 days of age) were used for three consecutive inoculations. The length of each post inoculation period was 14 days and the experiment was conducted for a total of 42 days. At the onset of the study, each animal was tested bacteriologically (fecal) before receiving the *Salmonella* and *E. coli* challenge and were found to be negative for the bacteria. The piglets were assigned to one of the three treatment groups (control, infection with *Salmonella enterica* serover Typhimurium (ST) and infection with *Escherichia coli* (EC)) in each post inoculation period in a completely randomized design based on their initial body weight. Each treatment had three replicate pens with three pigs per pen. The pigs in each group were

reared in three isolated pens (slatted floor and a space allowance of 0.75 m² per piglet) in an environmentally controlled room and allowed a 1-week adaptation period before commencing the study. Pigs in the ST and EC group were then orally infected with 10 mL mixed suspensions (1:1) of *S. enterica* serover Typhimurium KCTC 2515 (5.5 x 10⁶ cfu/ml) and *E. coli* KCTC 2571 (3.7 x 10⁶ cfu/ml). An ambient temperature of 24 ± 1.49°C was maintained in the pig houses via overhanging electric heaters. During the experimental period, pigs were provided with a commercial diet *ad libitum*. The pens were illuminated by artificial light, and ventilation was provided by ten air changes per hour. The body weight and feed intake were recorded on a weekly basis until the end of the experiment.

Infrared thermography of piglets

Thermal images of piglets were captured using an infrared camera (Thermo Tracer, Type- TH5104R CAT I, NEC San-ei Instruments, Ltd., Tokyo, Japan) at a fixed distance of approximately 1 m from the animal. The emissivity value was set to 0.985 and the thermograph resolution was calibrated to ambient temperature and humidity as per the manufacturer's recommendations. The piglets were numbered temporarily and allowed to settle before infrared thermographic (IT) images were captured. Multiple images were captured and the most informative image for an individual piglet was used. Thermal images were captured at 0, 2, 6, 12, and 24 h and then every day for 14 days post inoculation (dpi) without handling the piglets. Images were stored in a memory card, then transferred to a computer for analysis using the InfRec Analyzer Lite software system. The temperatures acquired from the IT images were recorded from three locations in each piglet (head, body and tail region), and the average values were used to calculate the standard body temperature of piglets.

Statistical analysis

All recorded data were statistically analysed using the SAS system v. 9.1 (SAS Institute. Inc., Cary, NC, USA, 2003). The individual pen was considered as the experimental unit for growth performance and body temperature calculation. Treatment means were computed with the LSMEANs and significant treatment x time interactions were observed for body temperature using thermal camera. Longitudinal significant variations in body temperature due to *Salmonella* and *E. coli* infection in different time periods are denoted by abcde, while treatment effects on specific time period denoted by xyz. Duncan's multiple range tests were used to identify significant differences among treatment groups for growth performances and body temperature. A probability level of P < 0.05 was considered statistically significant.

RESULTS

The average daily gain (ADG), average daily feed intake (ADFI) and gain:feed ratio of piglets are shown in Figures 1, 2 and 3. In the present study, we observed a decreased ADG at the first week of each post inoculation period in the ST and EC groups (P < 0.05) (Figure 1). A reduced ADFI was found in the second week of first inoculation and in the first week of second inoculation in the ST and EC infection groups relative to the control (P < 0.05) (Figure 2). A reduced gain:feed ratio was observed in the first week of first inoculation (P < 0.05) (Figure 3). Although the gain:feed ratio was decreased in




Figure 1. Mean average daily gain (ADG) (kg) of piglets in different weeks during the three inoculation periods. Data are presented as the mean \pm SE. Bars at a week without a common letter differed significantly ($P < 0.05$).




Figure 2. Mean average daily feed intake (ADFI) (kg) in different weeks during the three inoculation periods. Data are presented as the mean \pm SE. Bars at a week without a common letter differed significantly ($P < 0.05$).

EC group, it was increased in ST group during second week of first inoculation and second inoculation period compared to control ($P < 0.05$) (Figure 3). Thermal images of piglets for control, infected with *Salmonella enterica* serovar Typhimurium KCTC 2515 and *Escherichia coli* KCTC 257 are shown in Figure 4.

The body temperature patterns of piglets with bacterial infections are shown in Figure 5, 6 and 7. During the first inoculation, no significant variations were observed until 24 h post inoculation, while piglets challenged with ST showed a gradual increase in temperature by 24 h. The maximum temperature increase relative to the control was observed at 72 h ($P < 0.05$), and this value remained

elevated until the end of the study period (Figure 5). However, the EC induced piglets showed a subnormal body temperature pattern throughout the experimental period during all three inoculation periods relative to the control ($P < 0.05$) (Figures 5, 6 and 7). Average daily body temperature was significantly lower in EC group while it was alleviated in ST group compared to control ($P < 0.05$) (Figure 8).

DISCUSSION

In most animals, illness is commonly expressed by their




Figure 3. Mean average gain:feed ratio in different weeks during the three inoculation periods. Data are presented as the mean \pm SE. Bars at a week without a common letter differed significantly ($P < 0.05$).




Figure 4. Thermal images of piglets for control, infected with *Salmonella enterica* serover Typhimurium KCTC 2515 and *Escherichia coli* KCTC 2571.




Figure 5. Body temperature pattern of piglets during the first inoculation period (14 days). Data are presented as the mean \pm SE. Lines at a particular time period without a common letter differed significantly ($P < 0.05$).




Figure 6. Body temperature pattern of piglets during the second inoculation period (14 days). Data are presented as the mean \pm SE. Lines at a particular time period without a common letter differed significantly ($P < 0.05$).




Figure 7. Body temperature pattern of piglets during the third inoculation period (14 days). Data are presented as the mean \pm SE. Lines at a particular time period without a common letter differed significantly ($P < 0.05$).

behaviour (Weary et al., 2009). Farm animals express a wide range of behaviours such as loss of appetite, drinking and social behaviours, many of which may impact their health and welfare directly or indirectly. In the present study, we observed a decreased ADG at the first week of each post inoculation period in the ST and EC groups (Figure 1), which is consistent with the results reported by Van Heugten et al. (1994). Additionally, reduced body weight gain of pigs was observed over 14 days in response to bacterial infection (Balaji et al., 2000), and decreased weight gain, feed intake and efficiency of feed utilization were observed after repeated

challenge with non-infectious agents in broilers (Klasing et al., 1987). The reduced ADG of piglets during post inoculation period of piglets might be due to effect of *Salmonella* and *E. coli* infection (Van Heugten et al., 1994; Balaji et al., 2000). ADFI did not affect in the first week, it was decreased in the second week during first inoculation which also extended in the first week of second inoculation (Figure 2). Feed intake was depressed in ST induced pigs in the 120 h period following the challenge, with the maximum decrease being observed 48 h after challenge. In a previous study, intake was returned to control levels at between 120 and




Figure 8. Average daily body temperature pattern of piglets during the three inoculation periods (42 days). Data are presented as the mean \pm SE. Bars at a particular inoculation period without a common letter differed significantly ($P < 0.05$).

144 h of the challenge and was supported by Balaji et al. (2000). A number of infectious diseases of swine are characterized by fever, cachexia, inactivity and anorexia (Hart, 1988). The reduced gain:feed ratio might have been due to the effect of reduced weight gain in the respective periods in each post inoculation period. The overall decreased piglets growth performance of this study can be explained as the effect of bacterial infection which might be the indication of illness in pig (Klasing et al., 1987; Balaji et al., 2000).

Physiological responses of infection and inflammation in animals are the result of neuroendocrine and immune systems interactions. Adaptive responses include induction of proinflammatory mediators and fever, reduced feed intake and diminished growth performance, as well as development of a regulated, specific, immune response to ward off pathogens. *Salmonella* and *E. coli* are the most successful at colonizing the gastrointestinal tract under stress conditions, when nutrition and immunity are suboptimum, as is the case in newly weaned pigs (Aumaitre et al., 1995). A recent review described changes in peripheral blood flow resulting in alterations of skin temperature of livestock that could be detected by infrared thermography (Stewart et al., 2005). In the present study, piglets challenged with ST and EC showed remarkable temperature variations at different hours and days in all three post inoculation periods. The body temperature patterns of piglets with bacterial infections are shown in Figures 5, 6 and 7. During the first inoculation, no significant variations were observed until 24 h post inoculation, while piglets challenged with ST showed a gradual increase in temperature by 24 h. The

maximum temperature increase relative to the control was observed at 72 h, and this value remained elevated until the end of the study period (Figure 5). However, the EC induced piglets showed a subnormal body temperature pattern throughout the experimental period relative to the control (Figures 5, 6 and 7). The increased body temperature of ST challenged piglets observed in the present study is consistent with the results reported by Walsh et al. (2012), who found a tendency for increasing body temperature with increasing time of bacterial inoculation. The febrile conditions in our present study started at 12 h post inoculation, then gradually increased until becoming significantly higher on day 8. These increased levels then continued throughout the experimental period. Balaji et al. (2000) reported that the fever associated with ST was gradual in onset, and was sustained for 5 days following challenge. The infrared thermographies used to detect bacterial infection in pigs in the present study have also been applied in many other studies (Loughmiller et al., 2001; Schaefer et al., 2004; Johnson et al., 2011). Infrared techniques were used by Loughmiller et al. (2001) to measure the body temperature of pigs, and the results showed that it is possible to detect a febrile response using body-surface temperature. Pigs have few sweat glands and are therefore forced to cool down by increasing the rate of blood flow through their skin (Moritz and Henriques, 1947; Montagna and Yun, 1964); accordingly, infrared technology may be a good method for detection of increases in body temperature. Manno et al. (2006) reported that surface temperature increases with increasing ambient temperature. *E. coli* are commonly

found in human and animal intestinal tracts and can survive outside the gastrointestinal tract for considerable lengths of time in several environments (Ashbolt, 2004). In the present study, EC induced piglets showed a subnormal body temperature throughout the experimental period in all three post inoculation periods, which is normal and occurs due to starvation and dehydration due to diarrhoea. Feed intake of the EC infected group of this experiment was lower during most of the week in each inoculation period. The body temperature was also lower during the first week, indicating that starvation or lower feed intake can be a cause of subnormal body temperature. Additionally, some pigs infected with EC showed diarrheal symptoms, although we did not record the diarrheal score, which may also have led to subnormal body temperature. The detection of subnormal body temperature in piglets of this study is in agreement with the results of studies conducted by Vianna and Carrive (2005), who investigated changes in temperature at the rat tail in connection with a fear reaction and concluded that it was possible to detect decreased temperature via infrared images. The increased body temperature due to *salmonella* and subnormal body temperature due to *E. coli* infection of this present study could be helpful for the pig farmers to early detection of bacterial infection which is important to take measures for further severe damage. Average daily body temperature was significantly lower in EC group while it was alleviated in ST group compared to control (Figure 8). The overall body temperature in ST group was also alleviated while EC group was decreased compared to control which might be due to the effect of weekly body temperature in pigs. Overall, growth performances and body temperature pattern of challenged piglets could be more beneficial for the farmers in pig monitoring system.

Conclusions

Reduced ADG, ADFI and gain:feed ratio, as well as variations in body temperature are general indicators of bacterial infections in pigs. In the present study, thermal image analysis of heat emissions from pig skin made early detection of ST and EC infection in pigs possible. This method provides the opportunity to measure body temperature continuously without contracting or stressing the animals and with a minimal risk of injuries. Thus, it can be an effective tool to minimize costs and save time while ensuring animal welfare during pig farming.

Conflict of interests

The authors did not declare any conflict of interest.

ACKNOWLEDGEMENTS

This research was supported by the Ministry of Science,

ICT and Future Planning (MSIP), South Korea, under the Convergence Information Technology Research Center (CITRC) support program (NIPA-2014-H0401-14-1008) supervised by the NIPA (National IT Industry Promotion Agency).

REFERENCES

- Ashbolt NJ (2004). Microbial contamination of drinking water and disease outcomes in developing regions. *Toxicology* 198:229-238.
- Aumaitre A, Peiniau J, Madec F (1995). Digestive adaptation after weaning and nutritional consequences in the piglet. *Pig News Info.* 16:73N-79N.
- Balaji R, Wright KJ, Hill CM, Dritz SS, Knoppe EL, Minton JE (2000). Acute phase responses of pigs challenged orally with *Salmonella typhimurium*. *J. Anim. Sci.* 78:1885-1891.
- Godynick V, ELBAB M, Schwarz R (1985). The vascular pattern in the skin of the pig at the time of birth. *Zbl. Vet. Med. C* 14: 304-315.
- Greiner LL, Stahly TS, Stabel TJ (2000). Quantitative relationship of systemic virus concentration on growth and immune response in pigs. *J. Anim. Sci.* 78: 2690-2695.
- Halloy DJ, Bouhet S, Oswald IP, Goret-Nicaise M, Kobisch M, Mainil J, Gustin PG (2004). Pathophysiological changes occurring during *Escherichia coli* endotoxin and *Pasteurella multocida* challenge in piglets: relationship with cough and temperature and predictive value for intensity of lesions. *Vet. Res.* 35:309-324.
- Hart BL (1988). Biological basis of the behavior of sick animals. *Neurosci. Biobehav. R.* 12:123-137.
- Ingram D, Weaver M (1969). A quantitative study of the blood vessels of the pig's skin and the influence of environmental temperature. *Anat. Rec.* 163:517-524.
- Johnson SR, Rao S, Hussey SB, Morley PS, Traub-Dargatz JL (2011). Thermographic eye temperature as an index to body temperature in ponies. *J. Equine Vet. Sci.* 31:63-66.
- Klasing KC, Laurin DE, Peng RK, Fry DM (1987). Immunologically mediated growth depression in chicks: influence of feed intake, corticosterone and interleukin-1. *J. Nutr.* 117: 1629-1637.
- Loughmiller JA, Spire MF, Dritz SS, Fenwick BW, Hosni MH, Hogge SB (2001). Relationship between mean body surface temperature measured by use of infrared thermography and ambient temperature in clinically normal pigs and pigs inoculated with *Actinobacillus pleuropneumoniae*. *Am. J. Vet. Res.* 62:676-681.
- Manno MC, Oliveira RFMD, Donzele JL, Oliveira WPD, Vaz RGMV, Silva BAN, Saraiva EP, Lima KRDS (2006). Efeitos da temperatura ambiente sobre o desempenho de suínos dos 30 aos 60 kg. *Rev. Bras. Zootecn.* 35:471-477.
- Montagna W, Yun JS (1964). The skin of the domestic pig 1. *J. Investig. Dermatol.* 43:11-21.
- Moritz AR, Henriques FC (1947). Studies of Thermal Injury: II. The relative importance of time and surface temperature in the causation of cutaneous burns. *Am. J. Pathol.* 23:695-720.
- Pijpers A, Schoevers EJ, Van Gogh H, Van Leengoed LA, Visser IJ, Van Miert AS, Verheijden JH (1991). The influence of disease on feed and water consumption and on pharmacokinetics of orally administered oxytetracycline in pigs. *J. Anim. Sci.* 69:2947-2954.
- Plonait HG, Puerperium und perinatale V (2004). *Textbook of Pig Diseases*, 4th ed. Parey Verlag.
- Röhlinger P, Grunow C, Reichmann A, Zimmerhackel M (1979). Voruntersuchungen zur Ermittlung der Anwendungsgebiete der Infrarotmeßtechnik in der Veterinärmedizin [Preliminary studies to determine the use of infrared technique in veterinary medicine]. *Monatsh. Veterinarmed.* 287-291.
- SAS (2003). *SAS STAT User's Guide*, version 9.1, SAS Institute Inc. Cary, North Carolina, USA.
- Schaefer A, Cook N, Tessaro S, Deregt D, Desroches G, Dubeski P, Tong A, Godson D (2004). Early detection and prediction of infection using infrared thermography. *Can. J. Anim. Sci.* 84:73-80.
- Schwartz K (1991). Salmonellosis in swine. *Comp. Cont. Edu. Pract.* 13:139-147.

- Scolari SC, Clark SG, Knox RV, Tamassia MA (2011). Vulvar skin temperature changes significantly during estrus in swine as determined by digital infrared thermography. *J. Swine Health Prod.* 19:151-155.
- Stewart M, Webster J, Schaefer A, Cook N, Scott S (2005). Infrared thermography as a non-invasive tool to study animal welfare. *Anim. Welf.* 14:319-325.
- Traulsen I, Naunin K, Mueller K, Krieter J (2010). Application of infrared thermography to measure body temperature of sows. *Zuchtungskunde* 82:437-446.
- Van Heugten E, Spears JW, Coffey MT (1994). The effect of dietary protein on performance and immune response in weanling pigs subjected to an inflammatory challenge. *J. Anim. Sci.* 72:2661-2669.
- Vianna DM, Carrive P (2005). Changes in cutaneous and body temperature during and after conditioned fear to context in the rat. *Eur. J. Neurosci.* 21:2505-2512.
- Walsh MC, Rostagno MH, Gardiner GE, Sutton AL, Richert BT, Radcliffe JS (2012). Controlling Salmonella infection in weanling pigs through water delivery of direct-fed microbials or organic acids. Part I: effects on growth performance, microbial populations, and immune status. *J. Anim. Sci.* 90:261-271.
- Weary DM, Huzzey JM, Von Keyserlingk MAG (2009). Board-invited interview: using behavior to predict and identify ill health in animals. *J. Anim. Sci.* 87:770-777.

Full Length Research Paper

Prevalence of triple viral infections of human immunodeficiency virus (HIV), hepatitis B and C among tuberculosis patients and associated risk factors: The case of West Arsi Zone, Ethiopia

Endale Mengesha², Tekle Airgecho^{3*}, Edessa Negera¹ and Mulugeta Kebede¹

¹Hawassa University, Faculty of Natural and computational Sciences, Department of Biology, P.O. Box 05, Hawassa, Ethiopia.

²Federal Ministry of Ethiopia, P. O. Box 1234, Addis Ababa, Ethiopia.

³Haramaya University, College of Health and Medical Sciences, P. O. Box 235, Harar, Ethiopia.

Received 16 May, 2015; Accepted 22 June, 2015

Tuberculosis, human immunodeficiency virus and hepatitis B and C infections are serious public health problems in Ethiopia. This study was conducted to assess the prevalence of triple viral infections and associated factors among tuberculosis patients who were on anti-tuberculosis treatment in the year 2011/12 in West Arsi Zone, Ethiopia. A cross sectional study design was conducted among 374 TB patients who were on first line anti-TB treatment. Simple random sampling technique was employed to draw the study subjects until the calculated sample size was achieved. Blood samples were collected to determine the HIV serostatus by rapid HIV testing kits; whereas the hepatitis B and C sero-status was determined by instant one step HBsAg and Flavicheck-HCV_{WB} kits, respectively. Socio-demographic and risk factors information was collected and analysed using univariate and multivariate logistic regression. Injection drug users were excluded from this study. From a total of 374 tuberculosis patients enrolled, TB-HIV co-infection was found to be 56 (14.97%) among which 5 (8.92%) hepatitis B and 4 (7.14%) were hepatitis C triple infected. The overall TB-HIV-Hepatitis triple infection prevalence was 2.4% (9/374). Though the current level of triple viral infections among TB patients seems to be low, an integrated program for treatment, prevention and control strategy is crucial to be considered in TB patient management.

Key words: Co-infection, HBV, HCV, HIV, TB, triple infection, West Arsi Zone, Ethiopia.

INTRODUCTION

Human immunodeficiency virus (HIV) continues to be a major global public health problem, having claimed more than 39 million lives so far. In 2013, there were 1.5 [1.4–

1.7] million people died from HIV-related causes globally. In the same year there were approximately 35.0 [33.2–37.2] million people living with HIV with 2.1 [1.9–2.4]

*Corresponding author. E-mail: teklemicro@gmail.com.

million people becoming newly infected globally. Sub-Saharan Africa is the most affected region, with 24.7 [23.5-26.1] million people living with HIV that accounts for almost 70% of the global total of new HIV infections (WHO, 2014).

Ethiopia is one of the Sub-Saharan African countries affected by high burden of HIV/AIDS pandemic. Since the first report of HIV case in 1984 in the country, the disease remained as a major public health concern of the country (FMoH, 2010). There is an estimated prevalence of 2.4% with 1.9% and 2.9% in male and female respectively. There is also difference in prevalence among urban (7.7%) and rural (0.9%) communities (Ministry of Health of Ethiopia, 2010).

According to the World Health Organization, Ethiopia is one among 22 TB high burden countries with an estimated 220,000 (258 per 100,000) incident and 200,000 (237 per 100,000) prevalence cases. There were an estimated 15,000 deaths (18 per 100,000) due to TB, excluding HIV related deaths in 2011 (World Health Organization, 2012).

HIV/AIDS pandemic has caused a resurgence of TB, resulting in increased morbidity and mortality worldwide. HIV and TB have a synergistic interaction; each accentuates progression of the other. Unfortunately, Sub-Saharan Africa region is part of the world where TB has been flourishing unhindered reaching the proportion of 82% TB cases co-infected with HIV in 2010. In the same year an estimated 1.1 million deaths occurred among HIV positive cases of TB including 0.32 million deaths among women. In addition, there were an estimated 0.35 million deaths among incident TB cases that were HIV positive (World Health Organization, 2011).

Though the prevalence of hepatitis co-infection with HIV varies widely across different studies mainly due to the variation in the distribution of risk factors, geographic location and other factors specific to the study population; it is one of the leading causes of morbidity and mortality in individuals with HIV (Iser and Sasadeusz, 2008). The co-infection of HIV with hepatitis (B and C) is attributable due to similar routes of transmission with evidence of significant clinical implications of complicated clinical course, case management and may also adversely affect therapy for HIV infection and vice versa (Padmapriyadarsini et al., 2006; D'Souza and Foster, 2004; Hennessey et al., 2009).

The coexistence of TB, HIV and viral hepatitis infections in the same patient poses a unique challenge to the patient and clinicians. Here, the major concern is most commonly available anti-TB and antiretroviral agents have associated risk of liver damage. In situation of concurrent treatment of these infections is inevitable, carefully chosen and tailored therapy is needed to minimize drug-drug interactions and related toxicities. When this condition is superimposed with HBV and/or HCV infection, the treatment and patient management challenge will be further complicated (Marzuki et al., 2008;

Yee et al., 2003).

Studies revealed that TB/HIV co-infected patients demonstrated a doubling level of liver enzymes among 14% of HBV and 12 % HCV infected patients. Moreover, 45% of patients with both HCV and HIV infection were developed drug-induced hepatitis during TB treatment (Wajiso, 2003; Byrd et al., 1979).

HIV-infected TB patients demonstrated a doubling in liver enzyme levels in 14% of patients with HBV and 12% of patients with HCV infection, although few had symptomatic hepatitis. About 5 of 11 (45%) patients with both HCV and HIV infection developed drug-induced hepatitis during TB treatment; the risk of hepatitis was over 3 times greater than in those with HIV infection alone and over 14 times greater than in those with neither HIV nor HCV infection (Wajiso, 2003; Byrd et al., 1979).

There is very limited or no information on the prevalence of HIV and Hepatitis (B and C) among TB patients thereby very little is known about the clinical burden of triple viral infection among TB patients and subsequent impact on TB treatment outcome in the country.

Shashemene, the capital of West Arsi Zone, is a crossroad town with a high prevalence of TB and HIV infection. According to the 2011(2004 E.C) Ethiopian Ministry of Health report the TB-HIV co-infection rate in the age group of 15-49 years was estimated to be 47% in the area. Therefore, this study was conducted to address the prevalence of triple viral infections and associated risk factors among TB patients in Shashemene, West Arsi zone, Ethiopia.

MATERIALS AND METHODS

The study area

West Arsi zone is one of the 14 zones of Oromia Regional State (the largest of 9 regional states of Ethiopia). The estimated population of the zone is 2,286,987. Shashemene is the capital of the zone with a population of 127, 798. Shashemene Referral Hospital is the only referral hospital in the zone. It is situated at 240 km away from capital of Ethiopia, Addis Ababa to the southern part of the country on the way to Moyale-Kenya. The hospital was established by missionaries as TB and Leprosy clinic in 1950. In 1976 it was upgraded to general public hospital and then in 2008, to referral hospital. The Health Centers used in this study were located within 12 to 35 km radius from the hospital with estimated catchment population of 750,000 (Figure 1).

Study design

A health facility based cross-sectional study was conducted to assess the prevalence of triple viral infection of HIV, HBV and HCV among TB patents at one referral hospital (Shashemene Referral Hospital) and three health centers (Shashemene, Aje and Arsi Negele) from September 2011 to June 2012.

Ethical considerations

The protocol was approved by Hawassa University Department of




Figure 1. Map of the study area.

Biology and College of Health Sciences institutional Ethical Review Committee. Support letters were obtained from the Zonal and district health offices. Informed consent was obtained from all adults (> 18 years) and for children (<18 years) informed consent was obtained from their parents or guardians. A verbal assent was obtained from children of 12-17 years. For all the study subjects, laboratory diagnosis and treatment (if needed) were given free of charge and all results were kept confidentially.

Sample size determination

The sample size was determined using a single population formula with P=37.2% from previous study in the area:

$$N = \frac{(Z_{1-\alpha/2})^2 * P(1 - P)}{d^2}$$

$$N = \frac{(1.96)^2 * (0.372) * (0.628)}{(0.05)^2}$$

N = 359 with 5% non-response rate; N = 359+18=377.

We considered the confidence limits = 95% (α = 0.05), Power =80%, P= 37.2 % (Wajiso, 2003).

Source population and study population

The source population was all types of TB cases following anti -TB treatment in TB clinics of the study area. The study subjects were recruited by randomly selecting TB cases from Shashemene Referral Hospital and Shashemene, Arsi Negele and Aje health centers.

Sampling procedure

A simple random sampling technique was used to select the study subjects. All TB patients those put on first line anti-TB treatment in three health centres and Shashemene Referral Hospital were taken as sampling frame. An identification numbers were given for all registered patients and then the study subjects were selected randomly until the calculated sample size is achieved.

Inclusion and exclusion criteria

All types of TB patients who visited the health facilities and registered for anti-TB treatment were included in the study. However, pediatrics and referred out cases were not included in the study. Individuals who were not willing to give their consent for participation and injection drug users were also excluded.

Socio-demographic and patient data collection

Data on socio-demographic and risk factor were collected from each

consented participants in a well-designed and pre-tested questionnaires.

Nurses working at the respective health facilities were assigned to counsel the patients before blood sample collection for HIV testing through PIHCT (Provider Initiated HIV Counselling and Testing) and hepatitis testing.

Clinical sample collection and transportation

Vein blood samples of 2-5 ml was collected aseptically by the assigned laboratory technologist or technician. The collected blood specimens were transported appropriately to Shashemene Referral Hospital Laboratory for HIV and hepatitis testing within 2 h of collection.

HIV testing

HIV testing was done on whole blood sample by rapid testing kits KHB (Shanghai Kehua Bio-Engineering Co., China, 100% sensitivity and 98% specificity), Stat-pack (Chembio Diagnostics Systems, USA, 99.7% sensitivity and 99.9% specificity) and Uni-Gold (Trinity Biotech Plc, Ireland, 100% sensitivity and 100% specificity) following the national HIV testing algorithm.

Hepatitis testing

HBV was tested for HBsAg using Instant One step test kit (SD BIOLINE, Korea, >99% specificity and >99% sensitivity) and HCV was tested by using Flavicheck[®]-HCV WB (Qualpro diagnostics, India, 100% Sensitivity and 99.6% specificity) test kits using the manufacturer's guideline at Shashemene Referral Hospital Laboratory.

Quality assurance of data

To assure the quality of data to the level maximum, the questionnaires were validated by pre-testing on about 10% of sample size on some randomly selected patients who were not included in the study. Beside manufactures' instruction, there was strict adherence to good laboratory practice principles and standard operational procedures during laboratory analysis as.

Study variables

Independent

These included clinical signs and symptoms, risky behaviours for HIV and Hepatitis infection, Socio-economic and socio-demographic variables.

Dependent

HIV, HBV and HCV serostatus, TB-HIV co-infection and TB-HIV-Hepatitis triple infection were considered as outcome variables.

Statistical analysis

All the collected laboratory and socio-demographic data were properly coded and labelled for easy of entry. All data were double entered into an Excel spread sheet, cleaned, verified using STATA version 9.0 and made ready for analysis.

Descriptive analysis

Means, standard deviations, frequencies, standard error of the

means, proportions, percentages as well as descriptive graphs and tables were used with respect to the given variables.

Univariate analysis

Explanatory variables were individually cross tabulated with the outcome variable and statistical significance was assessed using chi-square, Odd ratio (OR) and 95% confidence interval (CI) were calculated to determine the strength of the association.

Multivariate analysis

Explanatory variables significantly associated ($p \leq 0.05$) with the outcome variable in Univariate analysis were included in a multivariate logistic regressions analysis using the program packages SPSS (SPSS Inc., Chicago, Illinois, USA, version 15.0 and STATA version 9.0) to detect confounding effects and to evaluate the relative influence of the different co-variables outcome variables.

RESULTS AND DISCUSSION

Socio-demographic characteristics of study subjects

A total of 374 TB patients were included in the study. Of the total TB patients, 224 (59.9%) were male while 150 (40.1%) of them were female with male to female ratio of 1.5:1. Among the total study participants, 203 (54.3%) of them were urban residents while 171(45.7%) were rural residents. The mean (\pm SE) age of the respondents was 30.12 \pm 0.645 years with a median age of 27 years. The average family size of the study subjects was 5.29+0.157SE with median of 5 (range 1-19) and the mean number of housewife per husband was 1.69+0.037SE with median of 2 and range of 1-5. Concerning the educational status of the study subjects, 143 (38.2%) were illiterate (unable to write and read) and 231 (61.2%) were literate. About 192 (51.3%) of study subjects were married, 148 (39.6%) were single, 12 (3.2%) were divorced, 15 (4.0%) were widowed and 7 (1.9%) were separated from their partner (due to work, education etc.). Occupationally majority of study subjects; 163 (43.6%) were farmers followed by students, 79 (21.1%) and only 3(0.8%) were commercial sex workers (Table 1).

Socio-demographic variables versus seropositivity of study participants

The overall prevalence of TB-HIV co-infection was 56 (15%). Among 224 male and 150 female TB patients tested for HIV, 32 (14.3%) male and 24 (16.0%) female were seropositive respectively.

The prevalence of Hepatitis (HBV and HCV) in TB-HIV co-infected patients was found to be 9 (16.07%). Of the 9 hepatitis positive TB-HIV co-infected patients 5(8.9%) were with HBV and 4 (7.1%) with HCV co-infection. The

Table 1. Socio-demographic characteristics of the study subjects for the study of triple viral infections of human immunodeficiency virus (HIV), Hepatitis B and C (HBV and HCV) among tuberculosis patients and associated risk factors in West Arsi zone, Ethiopia in 2011/12.

Variable	Category	N=374	%
Sex	Female	150	40.1
	Male	224	59.9
	Total	374	100.0
Age	15-24	147	39.3
	25-34	117	31.3
	35-44	69	18.4
	45-54	15	4.0
	55-64	16	4.3
	>64	10	2.7
	Total	374	100.0
Residence	Urban	203	54.3
	Rural	171	45.7
	Total	374	100.0
Education	Illiterate	143	38.2
	1 st cycle	120	32.1
	2 nd cycle	94	25.1
	12 grade complete	12	3.2
	College and above	5	1.3
	Total	374	100.0
Occupation	Farmer	164	43.9
	Student	79	21.1
	Daily laborers	28	7.5
	Merchants	40	10.7
	Commercial sex workers	3	0.8
	Government employee	13	3.5
	House wife	26	7.0
	Others	21	5.6
	Total	374	100.0
Marital status	Single	148	39.6
	Married	192	51.3
	Divorced	12	3.2
	Partner died	15	4.0
	Partner separated	7	1.9
	Total	374	100.0

mean (\pm SE) age of HIV positive TB patients was 30.41(\pm 1.14) and the mean age of Hepatitis positive TB/HIV co-infected patients was 30.56(\pm 0.86SE). There is no statistically significant difference between the age groups of study participants ($P>0.05$) in HIV seropositivity.

Among the total 56 HIV positive TB patients, 32(57.1%) were male while 24 (42.9%) were female and from 9 hepatitis positive TB-HIV co-infected patients, 4 (44.44%) of them were male while 5 (55.56%) were female with the overall TB-HIV-Hepatitis triple infection prevalence of 9(2.4%) observed in the current study area. About 45 (80.4%) of HIV positive TB patients were urban residents while 11 (19.6%) of them were rural residents. The statistical analysis showed that statistical significant difference between the urban and rural residents ($P<0.05$) with regard to HIV seropositivity among TB patients. Similarly, majority 6(66.7%) of triple infected patients were reside in urban areas while 3 (33.3%) of them were reside in rural area. The HIV seropositivity of respondents were statistically significantly different ($P<0.05$) with respect to their educational status, occupation and marital status (Table 2).

Characteristics and clinical features of study subjects

TB patients were categorized as new, relapse, and defaulter cases. Of the total 374 TB patients enrolled in the study 347 (92.8%) of them were new cases, 17 (4.5%) were relapse and 10 (2.7%) were defaulters. The prevalence of HIV seropositivity in relapse TB cases was 5 (29.4%) and 49 (14.5%) in new TB cases (Table 3). The prevalence of triple infection (TB-HIV and Hepatitis) was found to be 8 (16.3%) in new TB cases and 1 (5.9%) in relapse TB cases. No triple infection was detected in defaulters. From the total of 300 TB patients tested for AFB microscopy, 138 (46.0%) of them were positive for the presence of AFB in the sputum samples. Among the 200 (53.5%) TB patients tested for X-ray, lung infiltration reported in 102 (51.1%), whereas, cavities were reported in 51(25.5%) and 49 (24.5%) were with no findings or not reported. The prevalence of HIV seropositivity was 23 (16.7%) and 25 (15.4%) for AFB positive and negative study participants respectively (Table 3). Among the AFB positive and AFB negative TB-HIV patients, 3 (13.04%) and 4 (16.0%) of them were positive for hepatitis test respectively.

Seropositivity versus sampling area

The TB-HIV co-infection prevalence for study subjects from Shashemene Referral Hospital, Shashemene Health Center, Arsi Negele Health Center and Aje Health Center was 26 (13.9%), 14 (21.9%), 11 (8.92%) and 5 (8.9%) respectively. The HIV seropositivity was slightly higher in Shashemene Health center 14 (21.9%) compared to the other centers. The prevalence of triple infection were 4 (15.4%), 3 (21.4%) and 2 (18.2%) Shashemene Referral Hospital, Shashemene Health Center, Arsi Negele Health Center and Aje Health Center respectively. Triple

Table 2. Socio-demographic characteristics of study subjects versus HIV and Hepatitis (HBV and HCV) Serostatus among tuberculosis patients and associated risk factors in West Arsi Zone, Ethiopia in 2011/12.

Socio-demographic characteristics of TB patients of study area	HIV Positive, n= 56				HIV Negative n=318	P-value
	Hepatitis					
	Positive		Negative			
	HBV	HCV	HBV	HCV		
Sex	Female	2	3	22	21	0.649
	Male	3	1	29	31	
	Total	5	4	51	52	
Age	15-24	0	0	11	11	0.709
	25-34	5	2	22	25	
	35-44	0	2	17	15	
	45-54	0	0	0	0	
	55-64	0	0	1	1	
	>64	0	0	0	0	
	Total	5	4	51	52	
Residence	Urban	3	3	42	42	0.000
	Rural	2	1	9	10	
Education	Illiterate	1	0	10	11	0.026
	1 st cycle	2	2	21	21	
	2 nd cycle	3	2	18	18	
	Grade 12 completed	0	0	1	1	
Occupation	College and above	0	0	1	1	0.000
	Farmer	2	2	11	11	
	Student	0	0	2	2	
	Daily laborers	0	0	12	12	
	Merchants	2	2	14	14	
	Commercial sex workers	0	0	1	1	
	Government employee	0	0	4	4	
	House wife	0	0	3	3	
	Other	1	0	4	5	
Marital status	Single	1	1	14	15	0.000
	Married	1	2	19	20	
	Divorced	2	0	7	9	
	Partner died	0	0	5	5	
	Partner separated	1	1	6	7	

infection was not reported from Aje Health center (Table 4).

Risk factor analysis for HIV and hepatitis seropositivity

On Univariate analysis, behavioural factors such as having multiple partner and having unprotected sex with commercial and non-commercial sex workers were found to be risk for HIV seropositivity. From host factors tested in the analysis, marital status (partners separated,

divorced and widowed), being female, residing in urban, having poor HIV knowledge and being illiterate associated with high risk of acquiring HIV. On multivariate analysis, being urban dweller, marital status (partners separated), and being illiterate were associated with increased seropositivity (Table 5).

DISCUSSION

The observed prevalence of TB-HIV co-infection (15%)

Table 3. HIV and Hepatitis Serostatus, Tuberculosis cases category, form of tuberculosis and TB laboratory diagnosis status among tuberculosis patients and associated risk factors in West Arsi Zone, Ethiopia in 2011/12.

Category	Diagnosis	HIV sero-status		Total
		Positive n (%)	Negative n (%)	
HIV serostatus	HIV rapid test	56 (15%)	318 (85%)	374
Hepatitis	HBsAg	5 (8.9%)	51 (91.1%)	56
	HCV antibody	4 (7.1%)	52 (92.9%)	56
TB category	New	49 (14.1)	298 (85.9)	347
	Relapse	5 (29.4)	12 (69.6)	17
	Defaulter	2 (20)	8 (80)	10
Form of TB	PTB	48 (16.2)	248 (83.8)	296
	EPTB	8 (10.5)	68 (89.5)	76
	Disseminated TB	0	2 (100)	2
AFB Result	Positive	23 (16.7)	115 (83.3)	138
	Negative	25 (15.4)	136 (83.9)	162
	Not performed	8 (10.96)	65 (89)	73
X-ray Result	Infiltrations	15 (14.7)	87 (85.3)	102
	Cavities	9 (17.7)	42 (82.3)	51
	No finding	7 (14.3)	42 (85.7)	49
	Not performed	25 (14.5)	147 (85.5)	172

Table 4. Distribution of Hepatitis B, C (HBV and HCV) and HIV seropositivity among tuberculosis patients and associated risk factors in West Arsi Zone, Ethiopia in 2011/12.

Parameter	HIV serostatus		Hepatitis sero-status			
	Positive N (%)	Negative N (%)	Positive N (%)		Negative N (%)	
			HBV	HCV	HBV	HCV
Shashemene Referral Hospital	26 (13.9)	161 (86.1)	2 (7.7)	2 (7.7)	24 (92.3)	24 (92.3)
Shashemene Health center	14 (21.9)	50 (78.1)	2 (14.3)	1 (7.1)	12 (85.7)	13 (92.9)
Arsi Negele Health Center	11 (18.9)	56 (81.1)	1 (9.1)	1 (9.1)	10 (89.9)	10 (89.9)
Aje Health center	5 (8.9)	51 (91.1)	0 (0)	0 (0)	5 (100)	5 (100)
Total	56 (15.0)	318 (85.0)	5 (8.9)	4 (7.1)	51 (91.1)	52 (92.9)

Table 5. Univariate and multivariate logistic regression analyses of risk factors for HIV infections among tuberculosis patients in West Arsi Zone, Ethiopia in 2011/12.

Factor	Univariate analysis			Multivariate analysis		
	Crude OR	CI	P-value	Adjusted OR/ AOR	CI	P-value
Sex						
Male	1			1		
Female	1.14	0.643-2.031	0.065	1.054	0.371-3.003	0.085
Residence						
Rural	1			1		
Urban	4.143	2.068-8.30	0.000*	2.439	1.282-7.246	0.049*
Educational status						
Illiterate	1			1		
Literate	0.344	0.272-0.691	0.000*	0.141	0.096-0.682	0.000*

Table 5. Contd.

Factor	Univariate analysis			Multivariate analysis		
	Crude OR	CI	P-value	Adjusted OR/ AOR	CI	P-value
Marital status						
Married	1			1		
Single	1.631	0.549-4.811	0.06	1.003	0.537-3.897	0.889
Partners separated	10.466	3.568-31.312	0.001*	2.638	1.077-6.466	0.034*
Divorced	4.121	1.532-9.847	0.005*	1.021	0.899-4.0123	0.654
Widowed	2.193	1.685-11.349	0.044*	1.001	0.0123-2.014	0.998
HIV knowledge						
Yes	1			1		
No	2.54	1.581-10.883	0.042*	3.288	0.637-16.982	0.155
Unprotected sex with commercial sex worker (high risky groups)						
No	1			1		
Yes	1.268	0.673-2.392	0.562	2.410	0.657-8.843	0.185
Unprotected sex with noncommercial partner						
No	1			1		
Yes	2.255	1.779-6.528	0.035*	1.217	0.331-1.999	0.653

reported in this study is lower than those studies conducted previously in the study area (37.2%; 25.7%; 20%). This might be attributed to behavioral changes brought by health education on HIV/AIDS prevention and control program. The program has been implemented by the Federal Ministry of Health at national level (Sharma et al., 2005; Wajiso, 2003; Arega, 2007).

The triple infection of TB-HIV-Hepatitis prevalence had little attention or was no longer been addressed in Ethiopia. The TB-HIV-Hepatitis triple infection, HBV (8.9%) and HCV (7.1%), in the present study pinpoints the need to consider triple infection for future prevention and control strategies in the study area in particular. The prevalence of HBV among TB-HIV co-infected patients obtained in this study is comparable with other findings in other countries that reported the prevalence of HBV among TB-HIV co-infected patients: in Thailand was 9% while the prevalence of HCV was 31%. According to these authors the main route of HCV transmission is injection and drug use while sexual intercourse is the main route for HBV transmission. The low HCV prevalence in this study may be accounted for the low prevalence of the main route of transmission. However this needs further investigation. The prevalence of triple infection of TB-HIV-Hepatitis was higher in this study (TB patients) compared to the studies conducted in other populations (blood donors, health care providers etc., data not indicated) in the same area. This higher prevalence may be due to the fact that HIV and Hepatitis shares common routes of transmissions (Yee et al., 2003).

In this study, there was a significant difference in HIV seropositivity between urban and rural residents ($p=0.000$; adjusted OR=2.439). There was also a difference in triple infected between urban (10.7%) and rural (5.4%) dwelling study participants. This might be due to the differences of the available risk factors in both areas. In addition, the current study shows that, a statistically significant difference in HIV seropositivity between different educational levels ($p=0.000$). HIV seropositivity was lower among literate than illiterates (adjusted OR=0.141) and the same for triple infection. This is probably literates can easily understand the ways of transmission and prevention measures from different sources such as newspaper and periodicals. However, the presence of pre-disposing risk factors especially in urban settings where more literate people are found should not be underestimated. In such cases, target oriented education which induces behavioral changes should be tailored to minimize the high prevalence of HIV sero-prevalence in urban settings.

The result of this study showed that a significant difference of HIV seropositivity among different occupations and the same for triple infection. It was indicated that the leading seropositivity was seen in daily laborers, merchants, commercial sex workers and governmental employee with descending order. Similarly, a study conducted in India indicated that unemployed and business professionals took the larger percentage HIV/TB co-infection (Kumar et al., 2002), another study suggested that higher seropositivity in manual laborers and some

studies conducted in Ethiopia indicated that, these groups of patients were seropositive to HIV (Wajiso, 2003; Devi et al., 2005; UNAIDS, 2009; Kumar et al., 2002).

HIV seropositivity was highest among those partners who live separately for various reasons. The high HIV seropositivity in this group is probably due to the fact that partner separation eventually leads to risky behaviors. The other reason might be when partners know their HIV status, they prefer to live separate in order to make ART treatment more confidential.

The major risk factors for triple infection of HIV-TB-Hepatitis were found to be having multiple sexual partners, followed by unsafe sexual intercourse (data not presented). This is probably due to shared routes of transmission of HIV and hepatitis (B&C), and TB is the opportunistic disease which comes after immune suppression like in HIV patients (Sterling and Sulkowski, 2004; Sy and Jamal, 2006; Solomon et al., 2008; Sirinak et al., 2008).

Conclusions

This study reveal relatively high prevalence rate of triple infection even if TB-HIV co-infection is getting declined than previous reports. The HBV-HCV co-infection was not found. Some behavioral and socio-demographic risk factors such as partners being separated, having multiple sexual partners, dwelling area (being urban) and occupation (unemployed) were found to be strongly associated with triple infection. In general the current study showed the need for more information on the triple infection in order to base therapeutic decisions and to set integrated prevention and control program in the country. This underline the need for further epidemiologic and clinical studies to optimize the management of this complicated medical conditions.

Authors' contributions

EM, EN, TA and MK formulated the study questions and designed the study protocol; EM conducted the field and all laboratory activities. EN, TA and MK supervised the field and laboratory works; EN and TA drafted the manuscript; EM, EN, TA and MK contributed to critical review of the manuscript. All authors equally contributed to the interpretation of the data and writing of the manuscript and read and approved the final version.

Conflict of interests

The authors did not declare any conflict of interest.

ACKNOWLEDGEMENTS

Study participants who were volunteer and open for the question raised and giving blood sample for the study deserves our heartfelt gratitude. We would like also to thank Shashemene Referral Hospital for allowing us to

use the laboratory for sample processing.

REFERENCES

- Arega D (2007). Epidemiology and Drug Resistance Pattern of Mycobacterial Isolates among HIV positive and HIV negative TB patients using Conventional and Molecular Methods in South east Ethiopia. Addis Ababa University.
- Byrd RB, Horn BR, Solomon DA, Griggs GA (1979). Toxic effects of isoniazid in tuberculosis chemoprophylaxis: role of biochemical monitoring in 1,000 patients. *JAMA* 241:1239-1241.
- Devi SB, Naorem S, Singh TJ, Singh KB, Prasad L, Devi TS (2005). HIV and TB coinfection. *J. Ind. Acad. Clin. Med.* 6:220-223.
- D'Souza R, Foster GR (2004). Diagnosis and treatment of chronic hepatitis B. *J. R. Soc. Med.* 97:318-321.
- FMoH (2010). Implementation Guidelines for TB/HIV collaborative activities in Ethiopia. National TB/HIV working group, Addis Ababa, Ethiopia.
- Hennessey KA, Kim AA, Griffin V, Collins NT, Weinbaum CM, Sabin K (2009). Prevalence of infection with hepatitis B and C viruses and co-infection with HIV in three jails: a case for viral hepatitis prevention in jails in the United States. *J. Urban Health* 86:93-105.
- Iser DM, Sasadeusz JJ (2008). Current treatment of HIV/hepatitis B virus coinfection. *J. Gastroenterol. Hepatol.* 23:699-706.
- Kumar P, Sharma N, Sharma N, Patnaik S (2002). Clinical profile of tuberculosis in patients with HIV infection/AIDS. *Indian J. Chest Dis. Allied Sci.* 44:159-164.
- Marzuki O, Fauzi A, Ayoub S, Kamarul Imran M (2008). Prevalence and risk factors of anti-tuberculosis drug-induced hepatitis in Malaysia. *Singap. Med. J.* 49:688.
- Padmapriyadarsini C, Chandrabose J, Victor L, Hanna L, Arunkumar N, Swaminathan S (2006). Hepatitis B or hepatitis C co-infection in individuals infected with human immunodeficiency virus and effect of anti-tuberculosis drugs on liver function. *J. Postgrad. Med.* 52:92.
- Sharma S, Mohan A, Kadiravan T (2005). HIV-TB co-infection: epidemiology, diagnosis & management. *Indian J. Med. Res.* 121: 550-567.
- Sirinak C, Kittikraisak W, Pinjeesekikul D, Charusuntonsi P, Luanloed P, Srisuwanvilai LO, Nateniyom S, Akksilp S, Likanonsakul S, Sattayawuthipong W (2008). Viral hepatitis and HIV-associated tuberculosis: risk factors and TB treatment outcomes in Thailand. *BMC Public Health* 8:245.
- Solomon SS, Srikrishnan AK, Mehta SH, Vasudevan CK, Murugavel KG, Thamburaj E, Anand S, Kumar MS, Latkin C, Solomon S (2008). High prevalence of HIV, HIV/hepatitis C virus co-infection and risk behaviors among IDUs in Chennai, India: a cause for concern. *J. Acquir. Immune Defic. Syndr.* 49(3):327-332.
- Sterling RK, Sulkowski MS (2004) Hepatitis C virus in the setting of HIV or hepatitis B virus coinfection. *Seminars in liver disease*, 2004. Copyright© 2004 by Thieme Medical Publishers, Inc., 333 Seventh Avenue, New York, NY 10001, USA., pp.61-68.
- Sy T, Jamal MM (2006). Epidemiology of hepatitis C virus (HCV) infection. *Int. J. Med. Sci.* 3:41.
- UNAIDS (2009). AIDS epidemic update 2009. Geneva: Joint United Nations Programme on HIV/AIDS (UNAIDS) and World Health Organization (WHO).
- Wajiso K (2003). Assessment of Hiv Sero-Prevalence among Registered Tuberculosis Patients in Arsi Zone. Addis Ababa University.
- WHO (World Health Organization) (2011). WHO Report, Global Tuberculosis Control, 2011. Available on: http://whqlibdoc.who.int/publications/2010/978924156406_9_eng.pdf.
- WHO (World Health Organization) (2012). WHO report, Global Tuberculosis control, 2012. Available at www.who.int/iris/bitstream/10665/75938/1/9789241564502_eng.pdf
- WHO Media Center/ HIV/AIDS (2014). Available on <http://www.who.int/mediacentre/factsheets/fs360/en/> June 3, 2015.
- Yee D, Valiquette C, Pelletier M, Parisien I, Rocher I, Menzies D (2003). Incidence of serious side effects from first-line antituberculosis drugs among patients treated for active tuberculosis. *Am. J. Respir. Crit. Care Med.* 167:1472-1477.

African Journal of Microbiology Research

Related Journals Published by Academic Journals

- *African Journal of Biotechnology*
- *African Journal of Biochemistry Research*
- *Journal of Bacteriology Research*
- *Journal of Evolutionary Biology Research*
- *Journal of Yeast and Fungal Research*
- *Journal of Brewing and Distilling*

academicJournals