

A photograph of two giraffes in a savanna landscape. The giraffes are standing in a grassy field with green bushes and trees in the background. The sky is a clear, pale blue. The giraffes have a characteristic brown and white spotted pattern. One giraffe is in the foreground, and another is slightly behind it to the left. The image is framed with rounded corners.

# Journal of Ecology and The Natural Environment

Volume 7 Number 11, November 2015

ISSN 2006-9847



*Academic  
Journals*

## ABOUT JENE

The **Journal of Ecology and the Natural Environment (JENE)** (ISSN 2006-9847) is published Monthly (one volume per year) by AcademicJournals.

**Journal of Ecology and the Natural Environment (JENE)** provides rapid publication (monthly) of articles in all areas of the subject such as biogeochemical cycles, conservation, paleoecology, plant ecology etc.

The Journal welcomes the submission of manuscripts that meet the general criteria of significance and scientific excellence. Papers will be published shortly after acceptance. All articles published in JENE are peer-reviewed.

## Submission of Manuscript

Please read the **Instructions for Authors** before submitting your manuscript. The manuscript files should be given the last name of the first author

[Click here to Submit manuscripts online](#)

If you have any difficulty using the online submission system, kindly submit via this email [jene@academicjournals.org](mailto:jene@academicjournals.org).

With questions or concerns, please contact the Editorial Office at [jene@academicjournals.org](mailto:jene@academicjournals.org).

## Editors

**Dr. Abd El-Latif Hesham**

*Genetics Department, Faculty of Agriculture, Assiut University,  
Assiut 71516,  
Egypt*

**Dr. Ahmed Bybordi**

*East Azarbaijan Research Centre for Agriculture and Natural Resources, Tabriz, Iran*

**Dr. Sunil Kumar**

*Natural Resource Ecology Laboratory,  
Colorado State University  
1499 Campus Delivery, A204 NESB, Fort Collins,  
Colorado-80526,  
USA*

**Prof. Gianfranco Rizzo**

*University of Palermo  
Dipartimento DREAM – Viale delle Scienze - Building  
9. 90128  
Palermo,  
Italy*

**Dr. Bahman Jabbarian Amiri**

*Kiel University, Germany,  
Ökologie-Zentrum der CAU  
Abt. Hydrologie und Wasserwirtschaft  
Olhausen Straße, 75  
Kiel,  
Germany*

**Dr. Bikramjit Sinha**

*National Institute of Science Technology and Development Studies,  
Pusa Gate, Dr. KS Krishnan Marg, New Delhi 110012,  
India*

**Prof. Gianfranco Rizzo**

*University of Palermo  
Dipartimento DREAM – Viale delle Scienze - Building  
9. 90128  
Palermo,  
Italy*

## Associate Editors

**Dr. Marko Sabovljevic**

*Dept. Plant Ecology, Faculty of Biology,  
University of Belgrade  
Takovska 43, 11000 Belgrade,  
Serbia*

**Dr. Sime-Ngando Téléphone**

*CNRS  
LMGE, UMR 6023, Université Blaise Pascal, 63177,  
Aubière Cedex  
France*

**Dr. Bernd Schierwater**

*ITZ, Ecology and Evolution, TiHo Hannover  
Büenteweg 17d, 30559 Hannover,  
Germany*

**Dr. Bhattacharyya Pranab**

*North-East Institute of Science & Technology  
Medicinal, Aromatic & Economic Plant Division,  
North-East Institute of Science & Technology,  
Jorhat-785006, Assam,  
India*

**Prof. Marian Petre**

*University of Pitesti, Faculty of Sciences  
1 Targul din Vale Street, Pitesti, 110040, Arges  
County,  
Romania.*

**Prof. R.C. Sihag**

*CCS Haryana Agricultural University  
Department of Zoology & Aquaculture, Hisar-125004,  
India*

**Prof. Kasim Tatic**

*School of Economics and Business, University of Sarajevo  
Trg oslobođenja 1, 71000 SARAJEVO,  
Bosnia and Herzegovina*

**Dr. Zuo-Fu Xiang**

*Central South University of Forestry & Technology,  
498 Shaoshan Nanlu,  
Changsha, Hunan, China.*

## Editorial Board

**Dr. Zuo-Fu Xiang**

*Central South University of Forestry & Technology,  
498 Shaoshan Nanlu,  
Changsha, Hunan, China.*

**Dr. Pankaj Sah**

*Higher College of Technology, Muscat,  
Department of Applied Sciences,  
(Applied Biology) Higher College of Technology,  
Al-Khuwair, PO Box 74,  
PC 133, Muscat  
(Sultanate of Oman)*

**Dr. Arti Prasad**

*Mohan Lal Sukhadia University,  
Udaipur, Rajasthan, India.  
123, Vidya Nagar, Hiran Magri,  
Sector-4, Udaipur, Rajasthan,  
India*

**Parviz Tarikhi**

*Mahdasht Satellite Receiving Station  
(Postal): No. 80, 14th Street, Saadat Abad Avenue,  
Tehran 1997994313,  
Iran*

**Bharath Prithiviraj**

*Post Doctoral Research Associate  
Knight Lab, Dept. of Chemistry & Biochemistry  
University of Colorado at Boulder  
USA*

**Dr. Melissa Nursey-Bray**

*Australian Maritime College, Tasmania,  
Australia*

**Parvez Rana**

*Department of Forestry and Environmental Science  
Shahjalal University of Science and Technology  
Bangladesh*

**Mirza Hasanuzzaman**

*Faculty of Agriculture, Sher-e-Bangla Agricultural  
University  
Sher-e-Bangla Nagar, Dhaka-1207,  
Bangladesh*

**Dr. Giri Kattel**

*Murray Darling Freshwater Research Centre, La Trobe  
University  
471 Benetook Avenue, Mildura, Victoria 3500,  
Australia*

**Dr. M. Rufus Kitto**

*Faculty of Marine Science-Obhur station,  
King Abdulaziz University,  
Jeddah 21589, Saudi Arabia*

**Dr. Özge Zencir**

*Kemah Vocational Training School,  
Erzincan University, Kemah,  
Erzincan, Turkey.*

**Dr. Sahadev Sharma**

*Laboratory of Ecology and Systematics,  
Graduate School of Engineering and Science,  
University of the Ryukyus, Senbaru 59,  
Nishihara, Okinawa-903-0213 Japan*

**Dr. Hasan Kalyoncu**

*University of Süleyman Demirel,  
Faculty of Art and Science,  
Department of Biology,  
32100 Isparta/Turkey*

**Hammad Khan**

*Department of Zoology and Fisheries,  
University of Agriculture,  
Faisalabad, Pakistan*

**Mirza Hasanuzzaman**

*Faculty of Agriculture,  
Sher-e-Bangla Agricultural University  
Sher-e-Bangla Nagar, Dhaka-1207,  
Bangladesh*

**Abdurrahman Dunder**

*Siirt University, Science and Arts Faculty,  
Department of Biology,  
56000, Siirt, Turkey*

**Meire Cristina Nogueira de Andrade**

*College of Agronomic Sciences,  
São Paulo State University, Brazil.*

**Imran Ahmad Dar**

*Dept. of Industries and Earth Sciences,  
The Tamil University,  
Ocean and Atmospheric Sciences & Technology Cell,  
(A Unit of Ministry of Earth Sciences, Govt. of  
India).*

**S. Jayakumar**

*Department of Ecology and Environmental  
Sciences,  
School of Life Sciences,  
Pondicherry University,  
Puducherry - 605 014, India*

**Umer Farooq**

*University of Veterinary & Animal Sciences  
Lahore, Pakistan*



# Instructions for Author

**Electronic submission** of manuscripts is strongly encouraged, provided that the text, tables, and figures are included in a single Microsoft Word file (preferably in Arial font).

The **cover letter** should include the corresponding author's full address and telephone/fax numbers and should be in an e-mail message sent to the Editor, with the file, whose name should begin with the first author's surname, as an attachment.

## Article Types

Three types of manuscripts may be submitted:

**Regular articles:** These should describe new and carefully confirmed findings, and experimental procedures should be given in sufficient detail for others to verify the work. The length of a full paper should be the minimum required to describe and interpret the work clearly.

**Short Communications:** A Short Communication is suitable for recording the results of complete small investigations or giving details of new models or hypotheses, innovative methods, techniques or apparatus. The style of main sections need not conform to that of full-length papers. Short communications are 2 to 4 printed pages (about 6 to 12 manuscript pages) in length.

**Reviews:** Submissions of reviews and perspectives covering topics of current interest are welcome and encouraged. Reviews should be concise and no longer than 4-6 printed pages (about 12 to 18 manuscript pages). Reviews are also peer-reviewed.

## Review Process

All manuscripts are reviewed by an editor and members of the Editorial Board or qualified outside reviewers. Authors cannot nominate reviewers. Only reviewers randomly selected from our database with specialization in the subject area will be contacted to evaluate the manuscripts. The process will be blind review.

Decisions will be made as rapidly as possible, and the journal strives to return reviewers' comments to authors as fast as possible. The editorial board will re-review manuscripts that are accepted pending revision. It is the goal of the AJFS to publish manuscripts within weeks after submission.

## Regular articles

All portions of the manuscript must be typed double-spaced and all pages numbered starting from the title page.

**The Title** should be a brief phrase describing the contents of the paper. The Title Page should include the authors' full names and affiliations, the name of the corresponding author along with phone, fax and E-mail information. Present addresses of authors should appear as a footnote.

**The Abstract** should be informative and completely self-explanatory, briefly present the topic, state the scope of the experiments, indicate significant data, and point out major findings and conclusions. The Abstract should be 100 to 200 words in length. Complete sentences, active verbs, and the third person should be used, and the abstract should be written in the past tense. Standard nomenclature should be used and abbreviations should be avoided. No literature should be cited.

Following the abstract, about 3 to 10 key words that will provide indexing references should be listed.

A list of non-standard **Abbreviations** should be added. In general, non-standard abbreviations should be used only when the full term is very long and used often. Each abbreviation should be spelled out and introduced in parentheses the first time it is used in the text. Only recommended SI units should be used. Authors should use the solidus presentation (mg/ml). Standard abbreviations (such as ATP and DNA) need not be defined.

**The Introduction** should provide a clear statement of the problem, the relevant literature on the subject, and the proposed approach or solution. It should be understandable to colleagues from a broad range of scientific disciplines.

**Materials and methods** should be complete enough to allow experiments to be reproduced. However, only truly new procedures should be described in detail; previously published procedures should be cited, and important modifications of published procedures should be mentioned briefly. Capitalize trade names and include the manufacturer's name and address. Subheadings should be used. Methods in general use need not be described in detail.

**Results** should be presented with clarity and precision. The results should be written in the past tense when describing findings in the authors' experiments. Previously published findings should be written in the present tense. Results should be explained, but largely without referring to the literature. Discussion, speculation and detailed interpretation of data should not be included in the Results but should be put into the Discussion section.

**The Discussion** should interpret the findings in view of the results obtained in this and in past studies on this topic. State the conclusions in a few sentences at the end of the paper. The Results and Discussion sections can include subheadings, and when appropriate, both sections can be combined.

**The Acknowledgments** of people, grants, funds, etc should be brief.

**Tables** should be kept to a minimum and be designed to be as simple as possible. Tables are to be typed double-spaced throughout, including headings and footnotes. Each table should be on a separate page, numbered consecutively in Arabic numerals and supplied with a heading and a legend. Tables should be self-explanatory without reference to the text. The details of the methods used in the experiments should preferably be described in the legend instead of in the text. The same data should not be presented in both table and graph form or repeated in the text.

**Figure legends** should be typed in numerical order on a separate sheet. Graphics should be prepared using applications capable of generating high resolution GIF, TIFF, JPEG or Powerpoint before pasting in the Microsoft Word manuscript file. Tables should be prepared in Microsoft Word. Use Arabic numerals to designate figures and upper case letters for their parts (Figure 1). Begin each legend with a title and include sufficient description so that the figure is understandable without reading the text of the manuscript. Information given in legends should not be repeated in the text.

**References:** In the text, a reference identified by means of an author's name should be followed by the date of the reference in parentheses. When there are more than two authors, only the first author's name should be mentioned, followed by 'et al'. In the event that an author cited has had two or more works published during the same year, the reference, both in the text and in the reference list, should be identified by a lower case letter like 'a' and 'b' after the date to distinguish the works.

Examples:

Abayomi (2000), Agindotan et al. (2003), (Kelebeni, 1983), (Usman and Smith, 1992), (Chege, 1998;

1987a,b; Tijani, 1993,1995), (Kumasi et al., 2001) References should be listed at the end of the paper in alphabetical order. Articles in preparation or articles submitted for publication, unpublished observations, personal communications, etc. should not be included in the reference list but should only be mentioned in the article text (e.g., A. Kingori, University of Nairobi, Kenya, personal communication). Journal names are abbreviated according to Chemical Abstracts. Authors are fully responsible for the accuracy of the references.

Examples:

Chikere CB, Omoni VT and Chikere BO (2008). Distribution of potential nosocomial pathogens in a hospital environment. *Afr. J. Biotechnol.* 7: 3535-3539.

Moran GJ, Amii RN, Abrahamian FM, Talan DA (2005). Methicillinresistant *Staphylococcus aureus* in community-acquired skin infections. *Emerg. Infect. Dis.* 11: 928-930.

Pitout JDD, Church DL, Gregson DB, Chow BL, McCracken M, Mulvey M, Laupland KB (2007). Molecular epidemiology of CTXM-producing *Escherichia coli* in the Calgary Health Region: emergence of CTX-M-15-producing isolates. *Antimicrob. Agents Chemother.* 51: 1281-1286.

Pelczar JR, Harley JP, Klein DA (1993). *Microbiology: Concepts and Applications.* McGraw-Hill Inc., New York, pp. 591-603.

### Short Communications

Short Communications are limited to a maximum of two figures and one table. They should present a complete study that is more limited in scope than is found in full-length papers. The items of manuscript preparation listed above apply to Short Communications with the following differences: (1) Abstracts are limited to 100 words; (2) instead of a separate Materials and Methods section, experimental procedures may be incorporated into Figure Legends and Table footnotes; (3) Results and Discussion should be combined into a single section.

Proofs and Reprints: Electronic proofs will be sent (e-mail attachment) to the corresponding author as a PDF file. Page proofs are considered to be the final version of the manuscript. With the exception of typographical or minor clerical errors, no changes will be made in the manuscript at the proof stage.

**Fees and Charges:** Authors are required to pay a \$550 handling fee. Publication of an article in the Journal of Ecology and the Natural Environment is not contingent upon the author's ability to pay the charges. Neither is acceptance to pay the handling fee a guarantee that the paper will be accepted for publication. Authors may still request (in advance) that the editorial office waive some of the handling fee under special circumstances

**Copyright: © 2014, Academic Journals.**

All rights Reserved. In accessing this journal, you agree that you will access the contents for your own personal use but not for any commercial use. Any use and or copies of this Journal in whole or in part must include the customary bibliographic citation, including author attribution, date and article title.

Submission of a manuscript implies: that the work described has not been published before (except in the form of an abstract or as part of a published lecture, or thesis) that it is not under consideration for publication elsewhere; that if and when the manuscript is accepted for publication, the authors agree to automatic transfer of the copyright to the publisher.

**Disclaimer of Warranties**

In no event shall Academic Journals be liable for any special, incidental, indirect, or consequential damages of any kind arising out of or in connection with the use of the articles or other material derived from the JENE, whether or not advised of the possibility of damage, and on any theory of liability.

This publication is provided "as is" without warranty of any kind, either expressed or implied, including, but not limited to, the implied warranties of merchantability, fitness for a particular purpose, or non-infringement. Descriptions of, or references to, products or publications does not imply endorsement of that product or publication. While every effort is made by Academic Journals to see that no inaccurate or misleading data, opinion or statements appear in this publication, they wish to make it clear that the data and opinions appearing in the articles and advertisements herein are the responsibility of the contributor or advertiser concerned. Academic Journals makes no warranty of any kind, either express or implied, regarding the quality, accuracy, availability, or validity of the data or information in this publication or of any other publication to which it may be linked.



# Journal of Ecology and the Natural Environment

Table of Contents: Volume 7 Number 11, November 2015

## ARTICLES

**An inventory of plant species found in gravel borrow pit around Gaborone, Botswana**

Israel A. Legwaila, Gaone T. Thebe and Tebogo Selebatso

**Soil organic carbon stock estimation in range lands of Kumrat Dir Kohistan KPK Pakistan**

Suleman Shah, Javed iqbal, Adnan Ahmad and Alamgir Khan

Full Length Research Paper

# An inventory of plant species found in gravel borrow pit around Gaborone, Botswana

Israel A. Legwaila\*, Gaone T. Thebe and Tebogo Selebatso

Botswana College of Agriculture, Department of Crop Science and Production, Private Bag 0027, Gaborone, Botswana.

Received 27 August, 2015; Accepted 28 September, 2015

**This study sought to establish an inventory of plant species established at three gravel borrow pits around Gaborone, Botswana. At each, plant data were collected from randomly selected 10 x 10 m quadrats within and up to 30 m away from the borrow pits. Species of *Acacia* were the most numerous in the pits, while more non-woody than woody species were established within the pits. Most woody species within the pits were not found around the pits, and most of the non-woody species within the pits were found around them, with the exception of the Tlokweg pit.**

**Key words:** Re-vegetation, reclamation, pioneer species, seed bank, borrow pit, *Acacia*.


## INTRODUCTION

Quarrying for gravel has a number of negative impact that affect the integrity of the environment including degradation of ecosystems. The removal of vegetation reduces the ecology of a site and exposes it to other environmental factors such as soil erosion that may exacerbate the negative impacts already caused. However, efforts have been made worldwide to reduce these negative impacts through reclamation by re-contouring and re-vegetation (Cripps et al., 2004). In both cases, natural succession will occur, re-establishing native vegetation on the sites. This happens through species colonization, spreading, displacement and replacement over time until climax species are established (Davis et al., 1985). These processes happen as the quarry soils change overtime due to weathering and other physical, chemical and biological processes fed by pioneer species, (Legwaila, 2012). Under natural succession, establishment of vegetation on disturbed land is dependent on the availability of a seed bank from

vegetation adjacent to a disturbed site. Landform and soil characteristics may also have an impact on establishment of vegetation (Davy, 2008). Landform in disturbed land may experience accelerated erosion and runoff, inhibited infiltration and unfavourable micro climatic conditions, all of which have a negative effect on the natural recovery processes of disturbed land (Whisenant, 2008). Where topsoil has been replaced after decommissioning of a borrow pit, the soil is expected to carry numerous seeds of plant species from the local environment. However, it generally takes longer for environmental impacts to decrease and for desired outcomes to be achieved under natural succession than under technical reclamation and there will always be left over impacts regardless of the type of reclamation interventions (Figure 1). When technical re-vegetation is undertaken, the process may occur more rapidly resulting in more rapid re-vegetation.

Technical re-vegetation however, requires amelioration

\*Corresponding author. E-mail: [ialegwaila@gmail.com](mailto:ialegwaila@gmail.com).



**Figure 1.** An illustration of the extent of environmental impacts over time after different reclamation interventions (Legwaila, 2012).

**Table 1.** Geographic coordinates of the sites.

Corners	Place					
	Bokaa		Tlokweng		Mmokolodi	
A	S 24° 26' 02	E 26° 02' 49.5	S24° 40' 20.52	E26° 02' 14.8648	S24° 28' 09.0	E 26° 58' 14.0
B	S 24° 26' 03	E 26° 02' 58.8	S24° 40' 19.90	E26° 02' 25.7320	S24° 28' 07.0	E 26° 58' 23.0
C	S 24° 25' 52	E 26° 02' 57.7	S 24° 40' 29.39	E26° 02' 24.4906	S 24° 28' 17.0	E 26° 58' 36.0
D	S 24° 25' 53	E 26° 02' 15.7	S 24° 40' 29.13	E 26° 02' 13.3490	S 24° 28' 25.3	E 26° 58' 28.0

of the quarry soils to support plant life as well as proper choice of plants which can survive the local conditions and provide the functions that are intended for the quarry after reclamation. This requires knowledge and selection of native species that are common to establish in disturbed land and be productive under less than normal conditions of quarry sites to ensure sustainability of the vegetation. It has been found in other studies that at times exotic species establish in decommissioned quarry sites, and has been concluded that they may have been introduced by humans (Davis et al., 2000). It has also been concluded that use of non-native species is “the second most important threat to biodiversity after habitat loss”, (Vermeulen and Whitten, 1999).

This study sought to identify and develop an inventory of plant species that established in borrow pits that were quarried for gravel around the city of Gaborone, Botswana.

**METHODOLOGY**

**Study sites**

This study was conducted at three decommissioned borrow pits around Gaborone. The pits were located in the villages of Bokaa, Mmokolodi and Tlokweng. These borrow pits came about as a result of the high infrastructure construction developments that

occurs in the city of Gaborone and the neighbouring areas. The land in these areas was previously used for communal grazing. The borrow pits were selected based on their close proximity to Gaborone. Their coordinates were taken at four corners of the sites (Table 1). The soils around these sites were slightly variable but generally supported a shrub savanna and savanna vegetation structure. The soils around Bokaa and Tlokweng pits were haplic lixisols which are common in the tropics with predominant dry seasons. They form subsequent to leaching of clay. They have very low levels of plant nutrients and are highly erodible. Mmokolodi on the other hand had eutric regosols which are weakly developed soils very common in unstable landforms. They also have low levels of nutrients and nutrient holding capacity.

**Experimental design**

Within each borrow pit site, four (4) quadrants of 10 m by 10 m were randomly selected. Around each borrow pit, eight quadrants of 10 m by 10 m from different sides of the pits were randomly selected to conduct an inventory of plant species around the pits. This procedure was performed to enable comparison of plant species within the borrow pits and those around them.

**Collection and identification of plant species**

All plants seen in the quadrats were recorded to species level. For those that could not be reliably identified in the field, specimens were collected for later identification in the herbarium at the Botswana College of Agriculture. The same procedure was

**Table 2.** Plant species found within the three borrow pits.

Bokaa		Tlokweg		Mmokolodi	
Plants within pit	Plant outside pit	Plants within pit	Plant outside pit	Plants within pit	Plant outside pit
<i>Acacia erubescens</i> *	<i>Acacia tortilis</i> *	<i>Acacia erubescens</i> *	<i>Acacia erubescens</i> *	<i>Acacia tortilis</i> *	<i>Acacia tortilis</i> *
<i>Acacia gerrardii</i> *	<i>Acacia mellifera</i> *	<i>Acacia gerrardii</i> *	<i>Acacia gerrardii</i> *	<i>Acacia erubescens</i> *	<i>Asparagus bechuanicus</i>
<i>Acacia mellifera</i> *	<i>Achyranthesaspera</i>	<i>Acacia mellifera</i> *	<i>Acacia mellifera</i> *	<i>Acacia gerrardii</i> *	<i>Cambretum imbrebe</i> *
<i>Acacia nigrescens</i> *	<i>Acrotome inflata</i>	<i>Acacia tortilis</i> *	<i>Aristida congesta</i>	<i>Acacia mellifera</i> *	<i>Carissa bispinosa</i> *
<i>Acacia tortilis</i> *	<i>Aristida congesta subsp barbicollis</i>	<i>Achyranthesaspera</i>	<i>Aristida congesta subsp barbicollis</i>	<i>Acacia nigrescens</i> *	<i>Combretum apiculatum</i> *
<i>Acrotome hispida</i>	<i>Aristida conjesta</i>	<i>Aristida congesta subsp barbicollis</i>	<i>Asparagus bechuaniscus</i>	<i>Achyranthes aspera</i>	<i>Combretum hereroense</i> *
<i>Aristida congesta</i>	<i>Asparagus bechuaniscus</i>	<i>Asparagus bechuaniscus</i>	<i>Combretum imberbe</i> *	<i>Acrotome hispida</i>	<i>Dichrostachys cinerea</i>
<i>Aristida congesta subsp. barbicollis</i>	<i>Combretum imberbe</i> *	<i>Ceratotheca triloba</i>	<i>Carissa bispinosa</i> *	<i>Aristida congesta Subsp congesta</i>	<i>Euclea undulata</i> *
<i>Asparagus bechuanicus</i>	<i>Carissa bispinosa</i> *	<i>Corchorus olitorius</i>	<i>Ceratotheca triloba</i>	<i>Aristida congesta subsp barbicollis</i>	<i>Evolvulus alsinoides</i>
<i>Ceratotheca triloba</i>	<i>Chloris virgata</i>	<i>Dicerocaryum eriocarpum</i>	<i>Cucumis myriocarpus</i>	<i>Asparagus bechuanicus</i>	<i>Gomphocarpus fruticosus g</i>
<i>Chloris virgata</i>	<i>Combretum apiculatum</i> *	<i>Dicoma tomentosa</i>	<i>Cyperus turrillii</i>	<i>Corchorus olitorius</i>	<i>Gomphrera celosioides</i>
<i>Corchorus olitorius</i>	<i>Combretum hereroense</i> *	<i>Echinochloa holubii</i>	<i>Dichrostachys cinerea</i> *	<i>Dicerocaryum eriocarpum</i>	<i>Grewia flava</i> *
<i>Cucumis myriocarpus</i>	<i>Cucumis myriocarpus</i>	<i>Ehretia rigida</i> *	<i>Gomphrera celosioiides</i>	<i>Dicoma tomentosa</i>	<i>Grewia flavescens</i> *
<i>Dicerocaryum eriocarpum</i>	<i>Cyperus turrillii</i>	<i>Euclea undulate</i> *	<i>Grewia flava</i> *	<i>Echinochloa holubii</i>	<i>Kalanchoe lanceolata</i>
<i>Dicoma tomentosa</i>	<i>Ehretia rigida</i> *	<i>Gomphrena celosioides</i>	<i>Grewia flavescens</i> *	<i>Euclea undulata</i> *	<i>Kyphocarpa angustifolia</i>
<i>Dodonae aviscosa</i> *	<i>Euclea undulate</i> *	<i>Guillemineadensa</i>	<i>Kalanchoe lanceolata</i>	<i>Evolvulus alsinoides</i>	<i>Peltophorum africanum</i> *
<i>Ehretia rigida</i> *	<i>Evolvulus alsinoides</i>	<i>Hermbstaedtiafleckii</i>	<i>Lantana rugosa</i>	<i>Fimbristylis hispidula</i>	<i>Perotis patens</i>
<i>Euclea undulata</i> *	<i>Gomphocarpus fruticosus</i>	<i>Kyphocarpa angustifolia</i>	<i>Ocimum canum</i>	<i>Gomphrena celosioides</i>	<i>Pogonarthria squarrosa</i>
<i>Evolvulus alsinoides</i>	<i>Gomphrera celosioides</i>	<i>Melinis repens</i>	<i>Pappea capensis</i>	<i>Gomphocarpus fruticosus</i>	-
<i>Fimbristylis hispidula</i>	<i>Grewia flava</i> *	<i>Monsonia angustifolia</i>	<i>Peltophorum africanum</i> *	<i>Grewia flava</i> *	-
<i>Gomphocarpus fruticosus</i>	<i>Grewia flavescens</i> *	<i>Sesbania bispinosa</i>	<i>Pogonarthria squarrosa</i>	<i>Guilleminea densa</i>	-

conducted outside the borrow pits from a distance of 30 m to the edge of the pits.

## RESULTS AND DISCUSSION

Overall there were 44 species (11 woody and 33 non-woody) found at all borrow pits (Table 2). The woody species made a total of 25% of all the species found. The majority of the woody species were *Acacia* species making 46% (five different species). All of the other six species contributed 9% each. A total of 5 woody species, four *Acacia*

species and *Euclea undulate* were present at each of the pits, (Table 3). The predominance of the *Acacia* species might be an indication of their greater tolerance to low levels of resources within the pits (Smith and Smith, 2014; Donfack et al., 1995). This enables them to efficiently exploit the little resources in the soil, better than other species. There was also an exotic species (*Dodonaea viscosa*) at Bokaa pit commonly used as an ornamental plant in Botswana. Its existence in the pit could be attributed to the anthropogenic activities that took place at the pit during operation (Davis et al., 2000). Of the 33 non-woody species

found at the different pits, 12 (36%) were common to all pits (Table 3).

There was a high number of non-woody plants that made the understory made up of grasses and small shrubs. The trees that existed with them might have created conducive micro-climatic and soil conditions, facilitating their growth and establishment (Smith and Smith, 2014; Whisenant, 2008).

It was expected that most plants that established within the borrow pits would either have been from the soil seed bank in the surrounding land or as a result of seed dispersion

**Table 2.** Contd.

<i>Gomphrena celosioides</i>	<i>Kalanchoe lanceolata</i>	<i>Streptopetalum serattum</i>	<i>Streptopetalum serattum</i>	<i>Hermbsaediafleckii</i>	-
<i>Grewia flava*</i>	<i>Kyphocarpa augustifolia</i>	<i>Tricholaena monachne</i>	<i>Tricholaena monachne</i>	<i>Hypertelis bowkeriana</i>	-
<i>Guilleminea densa</i>	<i>Lantana rugosa</i>	<i>Urochloa mosambicensis</i>	<i>Urochloa mosambicensis</i>	<i>Kohautia subverticilla</i>	-
<i>Hermbsaediafleckii</i>	<i>Ocimum canum</i>	<i>Ziziphus mucronata*</i>	<i>Zyzybus mucronata*</i>	<i>Kyphocarpa angustifolia</i>	-
<i>Hypertelis bowkeriana</i>	<i>Pappea capensis</i>	-	-	<i>Melinis repens</i>	-
<i>Kohautia subverticillata</i>	<i>Peltophorum africanum*</i>	-	-	<i>Monsonia angustifolia</i>	-
<i>Kyphocarpa angustifolia</i>	<i>Perotis patens</i>	-	-	<i>Pogonarthria squarrosa</i>	-
<i>Melinis repens</i>	<i>Pogonarthria squarrosa</i>	-	-	<i>Sesbania bispinosa</i>	-
<i>Monsonia angustifolia</i>	<i>Solanum sp</i>	-	-	<i>Sida cordifolia</i>	-
<i>Ocimum canum</i>	<i>Streptopetalum serattum</i>	-	-	<i>Tricholaena monachne</i>	-
<i>Rhigozum brevispinosum*</i>	<i>Tarconanthus camphoratus*</i>	-	-	<i>Urochloa mosambicensis</i>	-
<i>Schmidtia pappophoroides</i>	<i>Tricholaena monachne</i>	-	-	-	-
<i>Sesbania bispinosa</i>	<i>Urochloa mosambicensis</i>	-	-	-	-
<i>Sida cordifolia</i>	-	-	-	-	-
<i>Solanum sp.</i>	-	-	-	-	-
<i>Streptopetalum serratum</i>	-	-	-	-	-
<i>Urochloa mosambicensis</i>	-	-	-	-	-

\*Woody species found within the pits.


**Table 3.** Plant species common at all three borrow pits.

<b>Botanical Name</b>	<b>Common Name</b>
<i>Acacia tortilis*</i>	<i>Mosu (Hairy umbrella thorn)</i>
<i>Acacia gerrardii*</i>	<i>Moga, Moki (Red thorn; Grey haired acacia)</i>
<i>Acacia mellifera*</i>	<i>Mongana (Black thorn; Hook thorn)</i>
<i>Acacia nigrescens*</i>	<i>Mokoba ( Knob thorn)</i>
<i>Aristida congesta subsp. barbicollis</i>	<i>Seloka; Bojang-ja-mothaba-jo-bonnye; (Buffalo grass)</i>
<i>Asparagus bechuanicus</i>	<i>Lesitwa, (Wild asparagus)</i>
<i>Dicerocaryum eriocarpum</i>	<i>Tshetho; Legatapitse (Devil's thorn; boot protector plant)</i>
<i>Dicoma tomentosa</i>	<i>Ombahu</i>
<i>Enchinochloa holubii</i>	<i>(Limpopo grass; Antelope grass; Kalahari water grass)</i>
<i>Euclea undulate*</i>	<i>Mothakola (Thicket euclea)</i>
<i>Gomphrena celosioides</i>	<i>Mositanoka (Prostate globe amaranthas; bachelor's button)</i>
<i>Guilleminea densa</i>	<i>Mohulapitse (Small mat weed)</i>
<i>Hypertelis bowkeriana</i>	<i>Mothabana; Munyu-wa-pasi</i>
<i>Kyphocarpa angustifolia</i>	<i>Mosono-wa-mmutla; Silky burweed; Hare's tail bush</i>

**Table 3.** Contd.

<i>Melinis repens</i>	<i>Lenapa; Senyane (Fairy grass)</i>
<i>Monsonia angustifolia</i>	<i>Phusana; Tsatsalopane (Crane bill)</i>
<i>Sesbania bispinosa</i>	<i>Mositanokana; Selaole (Spiny sesbina)</i>

\*Woody species found within the pits.



**Figure 2.** A comparison of woody species within the pits and adjacent to the pits.

from the vegetation stand surrounding the pits. The distance of the different species from the pits and their dispersal characteristics could also have affected what got established in the pits (Makhabu and Marotsi, 2012). It should be noted that vegetation surrounding the pits was only sampled to a distance of 30 m. It was found that the majority of woody species that grew around the pits were not found growing within the pits (Figure 2). With the exception of the Tlokweg pit, majority of non-woody plants that were found

around the pits were also found growing inside the pits (Figure 3).


**Conclusion**

It can be concluded that *Acacia* species have a great tolerance for poor underdeveloped soils found in decommissioned borrow pits. This can make them suitable candidates for pioneer species in technical reclamation, especially when

there is a limitation of topsoil available for re-vegetation. They can be used to provide conducive environmental conditions for other plant species.

It can also be concluded that most non-woody species are easy to establish in disturbed lands. They are useful in the control of negative impacts such as soil erosion as well as improving the capability for water infiltration. This contributes to the success of other processes such as decomposition which can accelerate development





**Figure 3.** A comparison of non-woody species within the pits and those adjacent to the pits.

of better soil that can support re-vegetation.

### Conflict of interests

The author(s) did not declare any conflict of interest.

### REFERENCES

- Cripps JC, Roubos V, Hughes D, Burton M, Crowther H, Nolan A, Tonks D (2007). Reclamation planning in hard rock quarries: A guide to good practice. Mineral Industry Research Organization (MIRO).
- Davis MA, Grime JP, Thompson K (2000). Fluctuating resources in plant communities: a general theory of invisibility. *J. Ecol.* 88:528-534.
- Davis BNK, Lakhani KH, Brown MC, Park DG (1995). Early Seral communities in a limestone quarry- an experimental study of treatment effects on cover and richness of vegetation. *J. Appl. Ecol.* 22:473-490.
- Davy AJ (2008). Establishment and manipulation of plant population and communities in terrestrial systems. In: *Handbook of ecological restoration, Principles of restoration*, Ed. Perrow M.R, Davy, A.J. Cambridge University Press, Cape Town. pp. 223-241.
- Donfack PI, Flor H, Pontanie R (1995). Dynamic vegetation around abandoned disturbed land. *J. Veg. Sci.* 6:499-508
- Legwaila I (2012). The effects of quarry treatment and distance on the attractiveness of reclaimed limestone quarry landscapes in England. Ph.D Thesis, University of Sheffield, England.
- Makhabu SW, Barotsi B (2012). Changes in herbaceous species composition in absence of disturbance in a *Cenchrusbiflorus*Roxb. Invaded area in Central Kalahari Game Reserve, Botswana. *Int. J. Ecol.* 2012:6.
- Smith TM, Smith RL (2014). *Elements of Ecology*. Benjamin Cummings, San Francisco, USA.
- Vermeulen J, Whitten T (1999). *Biodiversity and Cultural Heritage in the Management of Limestone Resources: Lessons from East Asia*. Directions in Development Series, The World Bank, Washington D.C.
- Whisenant SG (2008). Terrestrial systems. In *Handbook of ecological restoration, Principles of restoration*, Ed. Perrow M.R, Davy, A.J. Cambridge University Press, Cape Town pp. 83-105.

*Full Length Research Paper*

## Soil organic carbon stock estimation in range lands of Kumrat Dir Kohistan KPK Pakistan

Suleman Shah<sup>1\*</sup>, Javed iqbal<sup>2</sup>, Adnan Ahmad<sup>2</sup> and Alamgir Khan<sup>1</sup>

<sup>1</sup>Department of Forestry Shaheed Benazir Bhutto University, Sheringal DIR (U) KPK Pakistan.

<sup>2</sup>Shaheed Benazir Bhutto University, Sheringal Dir Upper KPK, Pakistan.

Received 31 July, 2015; Accepted 15 September, 2015

The present study was carried out in two grass land of Kumrat Dir Kohistan, North Hindukush regions of KPK Pakistan. Locally these two grass land are known as Roshi Dab and Bothore Dab. The area of these two Dabs is about 400 hectare. The elevation of these two Dabs ranges from 7665 feet to 7954 feet while the geographic location of the study site was N 35° 33.249' and E 72° 12.258'. Stratified random sampling method was used for data collection. 18 soil samples were taken with the depth of 0-15 cm and 16-30 cm. The average soil pH was 5.3. The mean soil bulk density was 1.107 -g/cm<sup>3</sup>. In present study it was found that the undisturbed range land stored maximum amount of soil carbon (32.69 tons hac<sup>-1</sup>) as compare to range land near to the agriculture land (29.77 tons hac<sup>-1</sup>) Similarly the rangeland near to forest land stored 35.62 t ha<sup>-1</sup> carbon, also stored more carbon as compare to range land near to agriculture land 29.77 t ha<sup>-1</sup>. The results of the present study confirmed that conversion of range land into agriculture land reduced the soil carbon in the study area.

**Key words:** Carbon stocks, climate change, soil bulk density, range lands, forestry.

### INTRODUCTION

The matter of great concern among scientific communities around the globe is the increase level of CO<sub>2</sub> and other green house gases. This increase level of green house gases leads to global warming and climate change. The most contributing agent in global climate change is Carbon Dioxide (IPCC, 2007). The atmospheric concentrations of CO<sub>2</sub>, N<sub>2</sub>O, CH<sub>4</sub> and other greenhouse gases prolong rise, affects the global climate. The Kyoto Protocol to the United Nations Framework Convention on Climate Change (UNFCCC) and succeeding agreements have set accurate targets in terms of levels and dates for sinking overall greenhouse gas emissions to the atmosphere (Watson et al., 2000).

The current concentration of CO<sub>2</sub> in the atmosphere is 402 ppm in 2014 ([www.CO2now.com](http://www.CO2now.com)). The carbon dioxide comes from different sources like fossils fuel burning, cement industries and other anthropogenic activities. Cement industries and fossil fuel burning contributes 91% that equals to 33.4 billion metric tons and the remaining 9% or 3.3 billion metric tons comes from land use changes particularly from the conversion of forest land to other land uses. This humanity's carbon dioxide 50% or 18.4 billion metric tons goes to atmosphere, 26% or 9.5 billion metric tons goes to land and the remaining 24% or 8.8 billion metric tons goes to oceans ([www.CO2now.com](http://www.CO2now.com)). Rangelands signify 24% of

\*Corresponding author. E-mail: [saayim77@gmail.com](mailto:saayim77@gmail.com).

the world's land area (FAO, 2007). Grassland Soil Organic Matter can be strongly influenced by supervision and management (Conant et al., 2001). SOC losses due to exchange of 'native' or freely grazed grassland (FGG) to cropland and poor grazing land management activities (Davidson and Ackerman, 1993; Guo and Gifford, 2002).

Range lands has so many importance like it support our wild life and livestock, it reduces soil erosion, and ensure clean water by reducing siltation (Milchunas, 2004). Apart from the above, range land acts as a sink of carbon and plays a noticeable role in the global climate change mitigation by catching and storing the atmospheric carbon dioxide (IPCC, 2006). Temperate grasslands contain soil organic matter (SOM) that averages 331 Mg ha<sup>-1</sup>, and grasslands contain 12% of the earth's SOM (Schlesinger, 1997). Therefore, enhancing carbon sequestration in terrestrial ecosystems is an important approach for controlling the increase in atmospheric CO<sub>2</sub> concentration. Grazing decreased soil carbon absorption, especially when initial soil carbon concentration was higher than 2%. Compared with no or light grazing, typical soil carbon concentration decreased in reaction to heavy grazing by 30.0% and moderate grazing by 17.0%. It decreased on average by 16.2% in response to heavy grazing and by 8.2% in response to fair grazing (Shiping et al., 2011).

Land use change can go ahead to change in a range of soil properties, including soil carbon (C), nitrogen (N) content, soil bulk density and pH (Callesen et al., 2003). There is significant alarm that land use change can lead to variation of the soil carbon (C) (Houghton et al., 1999; Schlesinger, 1990) though, Lal (2004) reported that more conventional soil properties such as total C is susceptible to land use change as complicated physical fractionation schemes. Soil is the largest organic C reservoir in the terrestrial biosphere; about two times superior than that of vegetation or the atmosphere (Schlesinger, 1997). Even an insignificant change in SOC storage could result in a major variation in atmospheric CO<sub>2</sub> concentration (Callesen et al., 2003; Wynn et al., 2006). Rangelands stored about 30% of the world's soil carbon (White et al., 2000; Grace, 2004; Milne et al., 2007).

In Pakistan about 60% of the total area of the country comprises rangelands. The area fairly supports 93.5 million head of livestock. Even so, there remains an insufficiency of basic ecological information that is needed to get better understanding of why, when and where rangeland ecosystems function as C sinks or sources. Pakistan is signatory to the Kyoto protocol. The protocol stress on the member countries to quantify the carbon, stored in different sinks. In Pakistan study on range land regarding forage production, carrying capacity and biomass estimation has been carried out. Study regarding carbon stocks of range land particularly on soil carbon is scarce. The present study was carried out in Kumrat valley. The study aimed to find out the total carbon stored in the soil of rangelands in Kumrat Dir

Kohistan.

## METHODOLOGY

### Research design

Stratified random sampling design was used. The range land of the area were divided into three strata that were undisturbed range land, range lands near to forest and range lands near to agriculture land. In all these three strata the soil sample was taken. 18 soil samples were randomly taken in each site at the depth of 0-15 cm and from 15-30 cm.

### Soil samples collection

The elevation and coordinates of each unit are from where soil samples were collected and were measured by GPS. Collection of soil samples was done in two depths; 0-15 and 15-30 cm and the weight of each sample was measured in the field and was put in labeled bags and brought to the laboratory for further analysis.

### Calculation of soil bulk density

The soil samples which were collected in rangelands were brought to the laboratory and were kept in the oven for 48 h on 72°C to dry. The soil samples were put in the known volume of a cylinder (H= 5.12 cm and Cross sectional Area= 20.32 cm<sup>2</sup>). After this the samples were weighted (through digital balance) and their volumes were calculated. The volume of soil core was calculated and was 104 cm<sup>3</sup>. Soil bulk density gcm<sup>-3</sup> was calculated by using following formula:

$$B.D = Mass/volume$$

### Soil pH estimation

For the determination of soil pH, 50 g of air- dried soil was taken into a 500 ml glass beaker, and 50 ml of distilled water were added. After titration, the contents were mixed and allowed to stand for an hour. After this, the soil pH was calculated by using soil pH meter (Mc Lean, 1982).

### Soil carbon estimation in Tons ha<sup>-1</sup>

Soil carbon in tons ha<sup>-1</sup> was calculated from the relation of soil organic carbon (SOC %), Soil bulk density (g/cm<sup>3</sup>) and thickness of horizon (cm). The following formula (Persson et al., 2008; Adnan et al., 2014) was used to find out soil carbon in tones ha<sup>-1</sup>:

$$SOC \text{ (tones ha}^{-1}\text{)} = SOC \text{ content \%} * SBD \text{ (g/cm}^3\text{)} * TH \text{ (cm)} * 100$$

Where, SOC = soil organic carbon; SBD = soil bulk density and TH = thickness of horizon.

$$\text{Soil Organic Carbon (tons hac}^{-1}\text{)} = \text{Soil Organic Content \%} \times \text{Soil Bulk Density (g/cm}^3\text{)} \times \text{Thickness of Horizon (cm)} \times 100.$$

### Statistical analysis

For the statistical analysis different software like MS Excel, Sigma plot software, program and PAST was used. Mean Standard deviation, and CV% Standard error were calculated. Regression

**Table 1.** Details of soil pH, SBD gm/cm<sup>3</sup>, SOM%, SOC%, and SOC tons hac<sup>-1</sup> in all sample plots.

Plot number	Elevation ft	Latitude	Longitude	Avg pH	Avg BD (g/cm <sup>3</sup> )	Total SOM%	SOC %	SOC tons hac <sup>-1</sup> (30) cm
1	7665	N 35° 33'.249"	E 72° 12'.258"	5.21	1.03	1.81	1.04	28.59
2	7689	N 35° 33'.262"	E 72° 12'.135"	4.95	1.01	1.62	0.93	32.90
3	7695	N 35° 33'.244"	E 72° 12'.268"	5.34	0.96	1.71	0.99	32.55
4	7697	N 35° 33'.262"	E 72° 12'.292"	4.91	1.14	1.71	0.99	30.92
5	7711	N 35° 34'.768"	E 72° 10'.956"	5.95	1.13	1.71	0.99	33.31
6	7732	N 35° 34'.765"	E 72° 10'.971"	5.19	1.19	1.83	1.06	37.96
7	7761	N 35° 33'.25"	E 72° 12'.093"	5.85	1.24	1.81	1.04	39.06
8	7781	N 35° 34'.79"	E 72° 10'.997"	4.85	1.07	1.61	0.93	39.34
9	7845	N 35° 34'.673"	E 72° 10'.626"	5.29	0.97	1.82	1.05	28.61
10	7852	N 35° 34'.417"	E 72° 10'.762"	4.86	0.99	1.71	0.99	36.76
11	7853	N 35° 34'.438"	E 72° 10'.774"	5.85	1.13	1.71	0.99	33.76
12	7879	N 35° 34'.654"	E 72° 10'.683"	5.37	1.08	1.77	1.02	34.19
13	7879	N 35° 34'.578"	E 72° 10'.729"	5.42	1.10	1.71	0.99	33.90
14	7884	N 35° 34'.639"	E 72° 10'.712"	5.93	1.07	1.65	0.95	29.61
15	7916	N 35° 34'.556"	E 72° 10'.718"	4.94	1.07	1.4	0.81	26.11
16	7921	N 35° 34'.539"	E 72° 10'.721"	5.01	1.08	1.6	0.92	30.11
17	7934	N 35° 34'.785"	E 72° 10'.449"	5.97	1.23	1.71	0.99	29.99
18	7954	N 35° 34'.776"	E 72° 10'.457"	5.95	1.37	1.65	0.95	30.90
Min	7665	-	-	4.85	0.96	1.4	0.81	26.11
Max	7954	-	-	5.97	1.37	1.83	1.06	39.34
Mean	7813.78	-	-	5.38	1.10	1.69	0.98	32.70
SD	97.02	-	-	0.42	0.10	0.10	0.05	3.76
Std. error	22.86	-	-	0.10	0.02	0.02	0.01	0.88
CV %	1.24	-	-	7.96	9.39	6.07	6.07	11.51

models were developed in order to study the relationship between soil organic matter and elevation and soil organic carbon and elevation.

## RESULTS AND DISCUSSION

### Soil pH of each sample plot

Results of soil pH are given in Table 1. Soil pH ranged from 4.5 to 5.9. The average soil pH was 5.3 which showed the acidity in nature. In order to

find out the relationship between soil pH and elevation, regression model was developed (Figure 4). The relation of soil pH and elevation is polynomial cubic. The value of R<sup>2</sup> was 0.13 and that showed that there was weak relation between elevation and soil pH.

### Soil pH of Range land near to agriculture land

These range lands were the disturbed range land.

Majority of these range lands were converted to agriculture lands. Soil samples were collected in these range land from those area which are not still converted to agriculture fields. These unconverted areas are mostly located at relatively high altitude and sloppy area. Soil samples were taken from those areas and there pH was calculated. The results of soil samples are given in Table 2. The average soil pH in this site was calculated as 5.91. These soils are less acidic in nature then those of undisturbed range land and

**Table 2.** Details of soil pH, SBD gm/cm<sup>3</sup>, SOM%, SOC%, and SOC tons hac<sup>-1</sup> in range land near to agriculture land.

Plot number	Elevation (ft)	Avg pH	Avg BD (g/cm <sup>3</sup> )	Total SOM%	SOC %	SOC (ton hac <sup>-1</sup> 30 cm)
1	7711	5.95	1.01	1.62	0.93	28.59
2	7761	5.85	1.14	1.71	0.99	34.19
3	7853	5.85	1.07	1.61	0.93	29.99
4	7884	5.93	0.99	1.71	0.99	29.61
5	7934	5.97	1.07	1.40	0.81	26.11
6	7954	5.95	1.08	1.60	0.92	30.11
Min	7711	5.85	0.99	1.40	0.81	26.11
Max	7954	5.97	1.14	1.71	0.99	34.19
Mean	7849.5	5.91	1.06	1.60	0.93	29.77
SD	96.20	0.05	0.05	0.11	0.06	2.62
Std. error	39.27	0.02	0.02	0.04	0.02	1.07
CV %	1.22	0.89	5.12	7.04	7.04	8.81

**Table 3.** Details of soil pH, SBD gm/cm<sup>3</sup>, SOM%, SOC%, and SOC tons hac<sup>-1</sup> in undisturbed rangeland.

Plot number	Elevation ft	Avg pH	Avg BD g/cm <sup>3</sup>	Total SOM%	SOC %	SOC ton hac <sup>-1</sup> 30 cm
1	7665	5.21	1.03	1.81	1.04	32.55
2	7695	5.34	0.96	1.71	0.99	28.61
3	7732	5.19	1.19	1.83	1.06	37.96
4	7845	5.29	0.97	1.82	1.05	30.90
5	7879	5.37	1.08	1.77	1.02	33.31
6	7879	5.42	1.10	1.71	0.99	32.90
Min	7665	5.19	0.96	1.71	0.99	28.61
Max	7879	5.42	1.19	1.83	1.06	37.96
Mean	7782.50	5.30	1.05	1.77	1.02	32.71
SD	96.48	0.08	0.08	0.05	0.03	3.10
Std. error	39.38	0.03	0.03	0.02	0.01	1.26
CV %	1.23	1.69	8.18	3.05	3.05	9.48

range lands near to forest lands.

### **Soil pH in undisturbed range land**

The result of soil pH in undisturbed range lands is presented in Table 3. In the present study, it was found that soil pH in undisturbed range land ranges from 5.19 to 5.42 while the mean soil pH was 5.30.

### **Soil pH of range land near to forest land**

The results of soils pH of range land near to the forest land is given in Table 4. The minimum soil pH of range land near the forest land was 4.85 while the maximum soil pH was 5.01. The Mean soil pH was 4.92.

### **Soil bulk density (g/cm<sup>3</sup>)**

In the present study the soil particles were examined and soil bulk density was found in each study site. Soil bulk density ranged from 0.96 to 1.37 g/cm<sup>3</sup>. The mean soil bulk density was 1.107 g/cm<sup>3</sup>. Details of soil is given in Table 3 and Figure 7. In order to find out relationship between soil bulk density and elevation, regression model was developed. The relationship between soil bulk density and elevation is polynomial cubic. The value of R<sup>2</sup> was 0.61. This value of R<sup>2</sup> indicated that there is a positive relation between soil bulk density and elevation. The relation of soil bulk density and elevation is best represented in Figure 5.

### **Soil bulk density (g/cm<sup>3</sup>) in undisturbed range land**

Soil SBD (g/cm<sup>3</sup>) in UDRL ranged from is 0.96 to 1.19

**Table 4.** Details of soil pH, SBD gm/cm<sup>3</sup>, SOM%, SOC%, and SOC tons hac<sup>-1</sup> in rangeland near to forest.

Plot no	Elevation ft	Avg pH	Avg BD g/cm <sup>3</sup>	Total SOM%	SOC %	SOC ton hac <sup>-1</sup> 30 cm
1	7689	4.95	1.13	1.71	0.99	33.90
2	7697	4.91	1.24	1.81	1.04	39.06
3	7781	4.85	1.13	1.71	0.99	33.76
4	7852	4.86	1.07	1.65	0.95	30.92
5	7916	4.94	1.23	1.71	0.99	36.76
6	7921	5.01	1.38	1.65	0.95	39.34
Min	7689	4.85	1.07	1.65	0.95	30.92
Max	7921	5.01	1.37	1.81	1.04	39.34
Mean	7809.33	4.92	1.19	1.70	0.98	35.62
SD	103.50	0.06	0.10	0.05	0.03	3.33
Std. error	42.25	0.02	0.04	0.02	0.01	1.36

g/cm<sup>3</sup>. Mean soil bulk density was 1.058 g/cm<sup>3</sup>. Details are given in Table 3.

#### **Soil bulk density (g/cm<sup>3</sup>) of range land near to forest land**

Table 4 showed that among all three study site Soil bulk density (g/cm<sup>3</sup>) of RLNFL was maximum and was 1.199 g/cm<sup>3</sup>. The minimum and maximum Soil bulk density in this site was 1.07 and 1.37 g/cm<sup>3</sup> respectively. Mean soil bulk density was 1.199 g/cm<sup>3</sup>.

#### **Soil bulk density (g/cm<sup>3</sup>) of each sample plot of range land near to agriculture land**

The soil bulk density near to agriculture land is expressed in Table 2. In this study site BD ranges from 0.99 to 1.14 g/cm<sup>3</sup>. Mean soil bulk density was 1.06 g/cm<sup>3</sup>.

#### **Soil organic matter (%)**

The results showed that the sample plots which were taken in undisturbed range land has maximum amount of soil organic matter than the sample plots taken in the range land near to forest land and agriculture land Table 1. The soil organic matter was calculated by Walkley and Black method in the laboratory. Minimum soil organic matter was recorded in Range land near to agriculture land and maximum soil organic carbon of 1.83% was found in undisturbed range land. Mean soil organic matter was 1.69%. In order to study the relation of soil organic matter and elevation, regression model was developed. The relation between soil organic matter and elevation was negative the value of R<sup>2</sup> was 0.19 (Figure 6).

The results of the present study shows great variation in soil organic matter in each study site. Undisturbed range lands has more SOM%, range lands near to forest has moderate SOM% while the range lands near to agriculture lands has in SOM%. The results show that the sample plots which were taken in undisturbed range land has maximum amount of SOM% than the sample plots taken in the range land near to forest land and agriculture land. SOM% of all these sample units ranged from 1.4 to 1.83%. Mean SOM% was 1.69%. In undisturbed range lands SOM% ranged from 1.71 to 1.83% while the mean soil organic matter was 1.775%. Range lands near to forest land SOM ranged from 1.65 to 1.81% while the mean SOM% was 1.70%. SOM% in range lands near to agriculture land ranged from 1.4 to 1.71% while the mean SOM% was 1.60%. These statistics shows that SOM% is more in undisturbed range land while less in range land near to forest land and lesser in range land near to agriculture lands.

#### **Soil organic matter% of sample plots taken in undisturbed range land**

Table 3 shows the soil organic matter (SOM %) in undisturbed range lands. Minimum SOM% was 1.71% and maximum was 1.83% while the mean SOM% was 1.775%.

#### **Soil organic matter% of sample plots taken in range land near to forest**

Details of soil organic matter in this site are presented in Table 4. Soil organic matter% in this site ranged from 1.65 to 1.81% while the mean soil organic matter % was 1.70%.



### **Soil organic matter% of sample plots taken in range land near to agriculture land**

Table 2 showed that minimum of SOM% was 1.4% and maximum was 1.71% while the mean SOM% was 1.60%.

### **Soil organic carbon SOC%**

The SOC% was calculated from the SOM% by using the following formulae:

$SOC\% = SOM\% / 1.724$  (Walkley and Black, Ahmad et al., 2014).

Where, SOC = soil organic carbon; SOM = soil organic matter; 1.72= constant.

### **Soil organic carbon concentration (SOC%) of each plot taken in undisturbed range land**

Table 3 shows the total soil organic carbon concentration of SOC%% in undisturbed range lands. Minimum of SOC% was 0.99% and maximum was 1.05%. Mean SOC% in these range lands were 1.02%.

### **Soil organic carbon concentration (SOC%) of each plot taken in range land near to forest land**

Table 4 shows the SOC% in each sample unit taken in range land near to forest land. Minimum of SOC% was 0.95% and maximum was 1.04%. Mean SOC% was 0.989%.

### **Soil organic carbon concentration (SOC%) of each plot taken in range land near to agriculture land**

Table 2 shows the SOC% in those range lands which are near to agriculture lands. Minimum SOC% was 0.92% and maximum was 0.99% the mean SOC% was 0.93%.

### **Soil organic carbon concentration in tons $ha^{-1}$ of each sample plot**

Table 1 shows that in each study site the soil organic carbon in  $t\ ha^{-1}$  was calculated. For the calculation of soil organic carbon, the following formula was used (Walkley and Black, 1934):

$SOC\ (tons\ ha^{-1}) = SOC\% \times SOM \times TH \times 100$ . Where, SOC = soil organic carbon; SOM = soil organic matter; TH = thickness of the horizon of the soil.

In the present study, soil organic carbon ranged from 26.11 to 39.34  $t\ ha^{-1}$ . The average soil organic carbon was 32.70  $t\ ha^{-1}$ . The results of the present study reveals

that the soil of undisturbed range land stored maximum soil carbon while the soil of the range land near to the agriculture land hold minimum soil carbon. In the present study, relation of soil organic carbon and elevation was determined through regression model (Figure 8). The relation is poly nominal cubic. The value of R<sup>2</sup> was 0.63.

In Undisturbed range land, SOC (tons  $ha^{-1}$ ) ranged from 28.61 tons to 37.96 tons  $ha^{-1}$ . The Mean SOC was 32.69 tons  $ha^{-1}$ . Range lands near to forests stored an average soil carbon of 35.62  $t\ ha^{-1}$ . Range lands near to agriculture lands stored an average soil carbon of 29.77 tons  $ha^{-1}$ . These results show that the undisturbed range lands soils stored maximum amount of SOC while soils of range lands near to forest lands stored less soil carbon  $t\ ha^{-1}$ .

### **Soil organic carbon $t\ ha^{-1}$ in undisturbed range lands**

Table 3 shows the total SOC in tons  $ha^{-1}$  stored by the undisturbed range lands. Minimum amount of SOC stored by these soils was 28.61 tons  $ha^{-1}$  and maximum amount of SOC stored by these soils was 37.96 tons  $ha^{-1}$ . Mean SOC which was stored by these soils was 32.69 tons  $ha^{-1}$ .

### **Soil organic carbon in tons $ha^{-1}$ in range land near to forest land**

Table 4 shows the total SOC tons  $ha^{-1}$  stored by range land near the forest lands. Minimum SOC was 30.92 tons  $ha^{-1}$  and maximum SOC was 39.34 tons  $ha^{-1}$  while mean SOC was 35.62 tons  $ha^{-1}$ .

### **Soil organic carbon in $t\ ha^{-1}$ in range land near to agriculture land**

Table 2 shows the soil organic carbon in  $t\ ha^{-1}$  in range land near the agriculture land. Minimum SOC was 26.11  $t\ ha^{-1}$  and maximum SOC was 34.19  $t\ ha^{-1}$  while mean SOC was 29.77  $t\ ha^{-1}$ . The results of the present study confirms that in undisturbed range land the soil organic carbon was more as compared to range land near the agriculture land. Soil disturbance occurred due to various agriculture practices that lead to soil erosion and loss of soil organic carbon.

## **CONCLUSION**

The study was conducted in Kumrat valley Dir Kohistan. Elevation of the study site ranged from 7665 to 7954 ft, while the geographic location is N 35° 33'.25 to 34'.785' and E 72° 12'.135' to 12'.997' (Figure 1 and 2). GIS image showed location and description of Kumrat Dir Kohistan (Figure 3). The minimum average soil weight was 103.5 g



Figure 1. Image of Roshi Dab.



Figure 2. Image of Bothore Dab.






Figure 3. GIS image of Dir Kohistan.



**Figure 4.** Relation between soil pH and Elevation (ft).




**Figure 5.** Relationship between soil BD (gm cm<sup>3</sup>) and Elevation (ft).

while the maximum soil sample weight was 142.5 g. The average soil sample weight was 116.77 g. Soil pH ranged from 4.5 to 5.9. The average soil pH was 5.3 which showed the acidity in nature. A regression model between soil pH and Elevation (ft) was developed (Figure 4).


Soil bulk density in all sample unit ranged from 0.96 to 1.37 g/cm<sup>3</sup>. The mean soil bulk density was 1.107 g/cm<sup>3</sup>. Soil SBD (g/cm<sup>3</sup>) in UDRL ranges from 0.96 to 1.19 g/cm<sup>3</sup>. Mean soil bulk density was 1.058 g/cm<sup>3</sup>. RLNFL have the minimum and maximum SD in this site was 1.07 g/cm<sup>3</sup> and 1.37 g/cm<sup>3</sup> respectively. Mean soil bulk density

is 1.199 g/cm<sup>3</sup>. The soil near to agriculture land BD ranges from 0.99 g/cm<sup>3</sup> to 1.14 g/cm<sup>3</sup>. Mean soil bulk density was 1.06 g/cm<sup>3</sup>. A regression model was developed between SBD and Elevation (Figure 5)

SOM% of all these sample units ranges from 1.4% to 1.83%. Mean SOM% is 1.69%. Undisturbed range lands have SOM% ranges from 1.71% to maximum of 1.83% while the mean soil organic matter % is 1.775. Range lands near to forest land have SOM ranges from 1.65% to maximum of 1.81% while the mean SOM% is 1.70%. SOM% in range lands near to agriculture land is ranges from 1.4% to maximum of 1.71% while the mean SOM%



**Figure 6.** Relationship between Soil organic matter % and elevation (ft).




**Figure 7.** Relationship between soil organic carbon % and elevation in each study site.

is 1.60%. These statistics shows that SOM% is more in undisturbed range land while less in range land near to forest land and lesser in range land near to agriculture lands. It is due to the severe disturbance of land soil surface which leads to prevent the soil from carbon storage. A regression model was developed between SOM% and elevation (Figure 6).

SOC% ranged from 0.92 to 1.05%. Mean SOC% was 0.98%. SOC% in undisturbed range lands ranged from 0.99to of 1.05%. Mean SOC% 1.02%. SOC in range lands near to forest lands ranges from 0.95 to 1.04%. Mean SOC% was 0.989%. SOC in range lands near to

agriculture lands ranged from 0.92 0.99% while mean SOC% was 0.93%. A regression model was developed between SOC% and elevation (Figure 7). Land use change can go ahead to change in a range of soil properties, including soil carbon (C), nitrogen (N) content, soil bulk density and pH (Callesen et al., 2003). Though, Lal (2004) reported that more conventional soil properties such as total C, it is susceptible to land use change as complicated physical fractionation schemes. Soil is the largest organic C reservoir in the terrestrial biosphere, about two times superior than that of vegetation or the atmosphere (Schlesinger, 1997). These results showed





**Figure 8.** Relationship between soil carbon t ha<sup>-1</sup> and elevation.

that the undisturbed range land has more carbon stored while range lands near to agriculture lands stored less in their soils. In Undisturbed range land SOC (tons hac<sup>-1</sup>) ranges from 28.61 tons hac<sup>-1</sup> to 37.96 tons hac<sup>-1</sup>. The Mean SOC was 32.69 tons hac<sup>-1</sup>. Range lands near to forests stored an average soil carbon of 35.62 t hac<sup>-1</sup>. Range lands near to agriculture lands stored an average soil carbon of 29.77 tons hac<sup>-1</sup>. These results showed that the undisturbed range lands soils stored maximum amount of SOC while soils of range lands near to forest lands stored less soil carbon t hac<sup>-1</sup>. A regression model was developed between SOC tons hac<sup>-1</sup> and elevation (Figure 8).

## RECOMMENDATION

The present study stated that if the range lands were not disturbed, it will bring a lot of rest in global climate change. As shown in Tables 1, 2, 3 and 4 the maximum organic carbon was stored by the range lands near to forest land and range lands which are not disturbed yet. While in other hand the range lands which are disturbed or will be disturbed in near future will lose the ability of storing organic carbon concentration.

The areas were remote and due to raising human and livestock population, the area was affected harshly. So the forest department is directed to raise awareness in local people and give special attention to this new raising problem.

## Conflict of interests

The author(s) did not declare any conflict of interest.

## REFERENCES

- Ahmad A, Mirza SN, Nizami SM (2014). Assessment of biomass and carbon stocks in coniferous forest of Dir Kohistan, KPK. Pak. J. Agric. Sci. 51(2):335-340.
- Callesen I, Liski J, Raulund-Rasmussen K, Olsson M, Tau-Strand L, Vesterdal L, Westman C (2003). Soil carbon stores in Nordic well-drained forest soils—relationships with climate and texture class. Glob. Change Biol. 9:358-370.
- Davidson E, Ackerman I (1993). Changes in soil carbon inventories following cultivation of previously untilled soils. Biogeochemistry 20: 161-193.
- Milne E, Sessay MF, Easter M, Paustian K, Killian K (2007). Sustainable Land Management Through Soil Organic Carbon Management and Sequestration The GEFSOC Modelling System pp. 359-371.
- FAO (2007). FAO Newspaper Rangelands and Climate Change: Mitigation, 2007
- Grace J (2004). Understanding and managing the global carbon cycle. J. Ecol. 92(2):189-202.
- Guo L, Gifford R (2002). Soil carbon stocks and land use change: a meta-analysis. Glob. Change Biol. 8:345-360.
- Houghton R, Hackler J, Lawrence K (1999). The US carbon budget: contributions from land-use change. Science 285:574
- IPCC (2007) Guidelines for National Greenhouse Inventories. Volume 4. Agriculture, Forestry and Other Land Use. Intergovernmental Panel on Climate Change (IPCC), IPCC/IGES, Hayama, Japan pp. 48-49.
- IPCC (2006). Guidelines for National Greenhouse Gas Inventories. Available online with updates. Available at: [www.ipccnggip.iges.or.jp/public/2006gl/index.html](http://www.ipccnggip.iges.or.jp/public/2006gl/index.html)
- Lal R (2004). Offsetting China's CO<sub>2</sub> Emissions by Soil Carbon Sequestration. Clim. Change 65(3):263-275.
- Schlesinger WH (1997). Biogeochemistry: An Analysis of Global Change (Second Edition). Academic Press, San Diego, California.
- Schlesinger WH (1990). Evidence from chronosequence studies for a low carbon-storage potential of soils. Nature 348:232-234.
- Shiping W, Andreas W, Zhicai Z, Xiaofeng C, Rong L, Yanfen W, Haishan N (2011). Management and land use change effects on soil carbon in northern China's grasslands. Agric. Ecosyst. Environ. 142 (3-4):329-340.
- Walkley A, Black IA (1934). An Examination of Degtjareff Method for Determining Soil Organic Matter and a Proposed Modification of the



- Chromic Acid Titration Method. *Soil Sci.* 37:29-37.
- White A, Cannell MGR, Friend AD (2000). The high-latitude terrestrial carbon sink: a model analysis. *Glob. Change Biol.* 6(2):227-245.
- Wynn J, Bird M, Vellen L, Grand-Clement E, Carter J, Berry S (2006). Continental-scale measurement of the soil organic carbon pool with climatic, edaphic, and biotic controls. *Glob. Biogeochem. Cycles* 20, GB1007.

A photograph of two giraffes in a savanna landscape. One giraffe is in the foreground, facing left, and another is behind it, facing right. The background shows green trees and a clear sky. The image is framed with rounded corners.

# Journal of Ecology and The Natural Environment

## *Related Journals Published by Academic Journals*

- *African Journal of Environmental Science and Technology*
- *International Journal of Biodiversity and Conservation*
- *Journal of Yeast and Fungal Research*
- *Journal of Entomology and Nematology*
- *African Journal of Biology and Environment*
- *Journal of Evolutionary Biology Research*

**academicJournals**