

Journal of Pharmacognosy and Phytotherapy

Volume 7 Number 11 December 2015

ISSN 2141-2502



*Academic
Journals*

ABOUT JPP

The **Journal of Pharmacognosy and Phytotherapy (JPP)** is published monthly (one volume per year) by Academic Journals.

The **Journal of Pharmacognosy and Phytotherapy (JPP)** is an open access journal that provides rapid publication (monthly) of articles in all areas of the subject such as ethnobotany, phytochemistry, ethnopharmacology, zoopharmacognosy, medical anthropology etc.

The Journal welcomes the submission of manuscripts that meet the general criteria of significance and scientific excellence. Papers will be published shortly after acceptance. All articles published in JPP are peer-reviewed.

Submission of Manuscript

Submit manuscripts as e-mail attachment to the Editorial Office at: jpp@academicjournals.org. A manuscript number will be mailed to the corresponding author shortly after submission.

The Journal of Pharmacognosy and Phytotherapy (JPP) will only accept manuscripts submitted as e-mail attachments.

Please read the **Instructions for Authors** before submitting your manuscript. The manuscript files should be given the last name of the first author.

Editors

Dr. (Mrs) Banasri Hazra

Research Scientist (U.G.C.)
Department of Pharmaceutical Technology
Jadavpur University
Calcutta - 700032
India

Dr. Yuanxiong Deng

Dept of Pharmaceutical Science
School of Medicine
Hunan Normal University
Tongzipo Road 371, Changsha 410013,
Hunan China

Prof. Maha Aboul Ela

Beirut Arab University, Faculty of Pharmacy, Beirut
Campus

Dr. S. RAJESWARA REDDY

Assistant Professor, Division of Animal Biotechnology
Department of Biotechnology, School of Herbal
Studies and Naturo Sciences,
Dravidian University, Kuppam – 517 425, A.P.
India

Dr. Mekhfi Hassane

University Mohammed the First, Faculty of Sciences,
Department of biology, Oujda, Morocco
Morocco

Dr. Ilkay Erdogan Orhan

Faculty of Pharmacy, Gazi University,
Ankara, Turkey
Turkey

Dr. Arun Kumar Tripathi

Central Institute of Medicinal and Aromatic Plants
P.O. CIMAP, LUCKNOW-226015,
India

Dr. Wesley Lyeverton Correia Ribeiro

Universidade Estadual do Ceará, Faculdade
de Veterinária/Laboratório de Doenças
Parasitárias Av. Paranjana, 1700
Itaperi - Fortaleza
60740-903, CE - Brazil

Dr. Maryam Sarwat

C/O A.M. Khan, House No. 195

Dr. Yong-Jiang Xu

Saw Swee Hock School of Public Health,
National University of Singapore, Singapore.

Prof. Dr. Adeolu Alex Adedapo

Department of Veterinary Physiology,
Biochemistry and Pharmacology
University of Ibadan, Nigeria

Dr. Joana S. Amaral

Campus de Sta Apolónia,
Ap. 1134, 5301-857 Bragança,
Portugal

Dr. Asad Ullah Khan

Interdisciplinary Biotechnology UNIT
Aligarh Muslim University,
India

Dr. Sunday Ene-ojo Atawodi

Biochemistry Department
Ahmadu Bello University
Zaria, Nigeria

Prof. Fukai Bao

Department of Microbiology and Immunology,
Kunming Medical College
China

Dr. Bhaskar C Behera

Agharkar Research Institute
Dept. of Science & Technology,
Plant Science Division
India

Prof. R. Balakrishna Bhat

Walter Sisulu University
Department of Botany
Mthatha, South Africa

Dr. Mohammad Nazrul Islam Bhuiyan

BCSIR Laboratories;
Chittagong cantonment;
Chittagong-4220;
Bangladesh

Dr. Baojun Bruce Xu

Beijing Normal University-Hong Kong Baptist
University United International College Zhuhai,
Guangdong Province,
China

Dr. Hamad H. Issa

Department of Physical Sciences,
School of natural Sciences,
The University of Dodoma,
Tanzania

Dr. Gagan Deep

Department of Pharmaceutical Sciences
School of Pharmacy,
University of Colorado Denver,
Colorado,
USA

Dr. Fengguo Xu

Dept of Epidemiology and Public Health,
Yong Loo Lin School of Medicine,
National University of Singapore,
Singapore

Dr. Haitao Lv

Medicine and Endocrinology,
Albert Einstein College of Medicine,
Yeshiva University,
USA

Hassane MEKHF

University Mohammed the First,
Faculty of Sciences,
Department of biology,
Laboratory of Physiology and Ethnopharmacology,
Morocco

Dr. Subhash C. Mandal

Division of Pharmacognosy
Pharmacognosy and Phytotherapy Research
Laboratory,
Department of Pharmaceutical Technology,
Jadavpur University,
India.

Dr. Adibe Maxwell Ogochukwu

Clinical Pharmacy and Pharmacy Management,
Faculty of Pharmaceutical Sciences,
University of Nigeria, Nsukka
Enugu state,
Nigeria.

Dr. Odukoya, Olukemi Abiodun

Department of Pharmacognosy,
Faculty of Pharmacy
University of Lagos.
Nigeria.

Dr. Qinxue Richard Ding

Medical Center at Stanford University,
Palo Alto,
USA

Dr. Sulejman Redžić

Faculty of Science of the University of Sarajevo
33-35 Zmaja od Bosne St.,
Sarajevo,
Bosnia and Herzegovina

Dr. Michal Tomczyk

Medical University of Bialystok,
Faculty of Pharmacy,
Department of Pharmacognosy,
Poland

Dr. Ugur Çakilcioglu

Firat University,
Faculty of Science and Arts,
Department of Biology,
Elazig
Turkey

Prof. Samson Sibanda

National University of Science and Technology
Cnr Gwanda Road/Cecil Avenue,
Ascot, Bulawayo,
Zimbabwe

Instructions for Author

Electronic submission of manuscripts is strongly encouraged, provided that the text, tables, and figures are included in a single Microsoft Word file (preferably in Arial font).

The **cover letter** should include the corresponding author's full address and telephone/fax numbers and should be in an e-mail message sent to the Editor, with the file, whose name should begin with the first author's surname, as an attachment.

Article Types

Three types of manuscripts may be submitted:

Regular articles: These should describe new and carefully confirmed findings, and experimental procedures should be given in sufficient detail for others to verify the work. The length of a full paper should be the minimum required to describe and interpret the work clearly.

Short Communications: A Short Communication is suitable for recording the results of complete small investigations or giving details of new models or hypotheses, innovative methods, techniques or apparatus. The style of main sections need not conform to that of full-length papers. Short communications are 2 to 4 printed pages (about 6 to 12 manuscript pages) in length.

Reviews: Submissions of reviews and perspectives covering topics of current interest are welcome and encouraged. Reviews should be concise and no longer than 4-6 printed pages (about 12 to 18 manuscript pages). Reviews are also peer-reviewed.

Review Process

All manuscripts are reviewed by an editor and members of the Editorial Board or qualified outside reviewers. Authors cannot nominate reviewers. Only reviewers randomly selected from our database with specialization in the subject area will be contacted to evaluate the manuscripts. The process will be blind review.

Decisions will be made as rapidly as possible, and the journal strives to return reviewers' comments to authors as fast as possible. The editorial board will re-review manuscripts that are accepted pending revision. It is the goal of the JPP to publish manuscripts within weeks after submission.

Regular articles

All portions of the manuscript must be typed double-spaced and all pages numbered starting from the title page.

The **Title** should be a brief phrase describing the contents of the paper. The Title Page should include the authors' full names and affiliations, the name of the corresponding author along with phone, fax and E-mail information. Present addresses of authors should appear as a footnote.

The **Abstract** should be informative and completely self-explanatory, briefly present the topic, state the scope of the experiments, indicate significant data, and point out major findings and conclusions. The Abstract should be 100 to 200 words in length. Complete sentences, active verbs, and the third person should be used, and the abstract should be written in the past tense. Standard nomenclature should be used and abbreviations should be avoided. No literature should be cited.

Following the abstract, about 3 to 10 key words that will provide indexing references should be listed.

A list of non-standard **Abbreviations** should be added. In general, non-standard abbreviations should be used only when the full term is very long and used often. Each abbreviation should be spelled out and introduced in parentheses the first time it is used in the text. Only recommended SI units should be used. Authors should use the solidus presentation (mg/ml). Standard abbreviations (such as ATP and DNA) need not be defined.

The **Introduction** should provide a clear statement of the problem, the relevant literature on the subject, and the proposed approach or solution. It should be understandable to colleagues from a broad range of scientific disciplines.

Materials and methods should be complete enough to allow experiments to be reproduced. However, only truly new procedures should be described in detail; previously published procedures should be cited, and important modifications of published procedures should be mentioned briefly. Capitalize trade names and include the manufacturer's name and address. Subheadings should be used. Methods in general use need not be described in detail.

Results should be presented with clarity and precision. The results should be written in the past tense when describing findings in the authors' experiments. Previously published findings should be written in the present tense. Results should be explained, but largely without referring to the literature. Discussion, speculation and detailed interpretation of data should not be included in the Results but should be put into the Discussion section.

The Discussion should interpret the findings in view of the results obtained in this and in past studies on this topic. State the conclusions in a few sentences at the end of the paper. The Results and Discussion sections can include subheadings, and when appropriate, both sections can be combined.

The Acknowledgments of people, grants, funds, etc should be brief.

Tables should be kept to a minimum and be designed to be as simple as possible. Tables are to be typed double-spaced throughout, including headings and footnotes. Each table should be on a separate page, numbered consecutively in Arabic numerals and supplied with a heading and a legend. Tables should be self-explanatory without reference to the text. The details of the methods used in the experiments should preferably be described in the legend instead of in the text. The same data should not be presented in both table and graph form or repeated in the text.

Figure legends should be typed in numerical order on a separate sheet. Graphics should be prepared using applications capable of generating high resolution GIF, TIFF, JPEG or Powerpoint before pasting in the Microsoft Word manuscript file. Tables should be prepared in Microsoft Word. Use Arabic numerals to designate figures and upper case letters for their parts (Figure 1). Begin each legend with a title and include sufficient description so that the figure is understandable without reading the text of the manuscript. Information given in legends should not be repeated in the text.

References: In the text, a reference identified by means of an author's name should be followed by the date of the reference in parentheses. When there are more than two authors, only the first author's name should be mentioned, followed by 'et al'. In the event that an author cited has had two or more works published during the same year, the reference, both in the text and in the reference list, should be identified by a lower case letter like 'a' and 'b' after the date to distinguish the works.

Examples:

Cole (2000), Steddy et al. (2003), (Kelebeni, 1983), (Bane and Jake, 1992), (Chege, 1998; Cohen, 1987a,b;Tristan, 1993,1995), (Kumasi et al., 2001)

References should be listed at the end of the paper in alphabetical order. Articles in preparation or articles submitted for publication, unpublished observations, personal communications, etc. should not be included in the reference list but should only be mentioned in the article text (e.g., A. Kingori, University of Nairobi, Kenya, personal communication). Journal names are abbreviated according to Chemical Abstracts. Authors are fully responsible for the accuracy of the references.

Examples:

Ansell J, Hirsh J, Poller L (2004). The pharmacology and management of the vitamin K antagonists: the Seventh ACCP Conference on Antithrombotic and Thrombolytic. Therapy. 126:204-233

Ansell JE, Buttaro ML, Thomas VO (1997). Consensus guidelines for coordinated outpatient oral anti coagulation therapy management. Ann. Pharmacother. 31:604-615

Charnley AK (1992). Mechanisms of fungal pathogenesis in insects with particular reference to locusts. In: Lomer CJ, Prior C (eds), Pharmaceutical Controls of Locusts and Grasshoppers: Proceedings of an international workshop held at Cotonou, Benin. Oxford: CAB International. pp 181-190.

Jake OO (2002). Pharmaceutical Interactions between *Striga hermonthica* (Del.) Benth. and fluorescent rhizosphere bacteria Of *Zea mays*, L. and *Sorghum bicolor* L. Moench for *Striga* suicidal germination In *Vigna unguiculata*. PhD dissertation, Tehran University, Iran.

Furmaga EM (1993). Pharmacist management of a hyperlipidemia clinic. Am. J. Hosp. Pharm. 50: 91-95

Short Communications

Short Communications are limited to a maximum of two figures and one table. They should present a complete study that is more limited in scope than is found in full-length papers. The items of manuscript preparation listed above apply to Short Communications with the following differences:

(1) Abstracts are limited to 100 words; (2) instead of a separate Materials and Methods section, experimental procedures may be incorporated into Figure Legends and Table footnotes; (3) Results and Discussion should be combined into a single section.

Proofs and Reprints: Electronic proofs will be sent (e-mail attachment) to the corresponding author as a PDF file. Page proofs are considered to be the final version of the manuscript. With the exception of typographical or minor clerical errors, no changes will be made in the manuscript at the proof stage.

Fees and Charges: Authors are required to pay a \$550 handling fee. Publication of an article in the Journal of Pharmacognosy and Phytotherapy (JPP) is not contingent upon the author's ability to pay the charges. Neither is acceptance to pay the handling fee a guarantee that the paper will be accepted for publication. Authors may still request (in advance) that the editorial office waive some of the handling fee under special circumstances.

Copyright: © 2013, Academic Journals.

All rights Reserved. In accessing this journal, you agree that you will access the contents for your own personal use but not for any commercial use. Any use and or copies of this Journal in whole or in part must include the customary bibliographic citation, including author attribution, date and article title.

Submission of a manuscript implies: that the work described has not been published before (except in the form of an abstract or as part of a published lecture, or thesis) that it is not under consideration for publication elsewhere; that if and when the manuscript is accepted for publication, the authors agree to automatic transfer of the copyright to the publisher.

Disclaimer of Warranties

In no event shall Academic Journals be liable for any special, incidental, indirect, or consequential damages of any kind arising out of or in connection with the use of the articles or other material derived from the JPP, whether or not advised of the possibility of damage, and on any theory of liability.

This publication is provided "as is" without warranty of any kind, either expressed or implied, including, but not limited to, the implied warranties of merchantability, fitness for a particular purpose, or non-infringement. Descriptions of, or references to, products or publications does not imply endorsement of that product or publication. While every effort is made by Academic Journals to see that no inaccurate or misleading data, opinion or statements appear in this publication, they wish to make it clear that the data and opinions appearing in the articles and advertisements herein are the responsibility of the contributor or advertiser concerned. Academic Journals makes no warranty of any kind, either express or implied, regarding the quality, accuracy, availability, or validity of the data or information in this publication or of any other publication to which it may be linked.

ARTICLE

Research Article

Pharmacological studies of methanolic extracts of *Sonchus arvensis* from Kathmandu

263

Bhupendra Kumar Poudel, Jagannath Prasad Sah, Shyam Raj Subedi, Mohan Prasad Amatya, Sadhana Amatya and Tirtha Maiya Shrestha

Full Length Research Paper

Pharmacological studies of methanolic extracts of *Sonchus arvensis* from Kathmandu

Bhupendra Kumar Poudel*, Jagannath Prasad Sah, Shyam Raj Subedi, Mohan Prasad Amatya, Sadhana Amatya and Tirtha Maiya Shrestha

Institute of Medicine, Tribhuvan University, Nepal.

Received 5 August, 2015; Accepted 17 November, 2015

***Sonchus arvensis*, a perennial sowthistle, is a common but underutilized species of Kathmandu, Nepal. Several uses like sedative, antioxidant and kidney stone eradicating properties have been identified till date, but study on other pharmacological activities is not yet explored. Therefore, this plant was collected from Kathmandu; aerial parts of the plant were dried, crushed, and extracted using a Soxhlet apparatus. The methanolic extract was then concentrated for screening pharmacological effects. While comparing with the standards, the plant was found to possess strong anti-inflammatory activity and inhibitory effect in Gastro Intestinal (GI) motility and locomotor activity in a dose-dependent fashion. The plant, however, did not show skeletal muscle relaxant activity as observed in the traction test and inclined plane test. Thus, it is concluded that the plant possess strong phyto-chemicals having anti-inflammatory activity and inhibitory effect in locomotion and GI motility.**

Key words: *Sonchus*, anti-inflammatory, motility, locomotion, extraction.

INTRODUCTION

In Nepal, the concept of ethno-medicine has developed since the late 19th century (1885-1901 A.D). The first book "Chandra-Nighantu regarding medical plants was published by the Royal Nepal Academy in 1969. Majority of the population is still dependent on botanical medicines which indicates the importance of herbal medicines in the primary health care (Rajbhandari et al., 2001; Amatya et al., 2009). *Sonchus arvensis*, a vigorous herbaceous perennial plant with milky sap and creeping roots is abundantly available in 1000 to 4100 m in range and grows in sandy, loamy or clayey soils (Holm et al. 1997). In many areas, this sowthistle is considered a noxious weed, as it grows quickly in a wide range of

conditions and its wind-borne seeds allow them to spread rapidly (APNI, 2012). Since, this plant is least explored and there are only few pharmacological studies, this study is done to lay a strong foundation for the future development of herbal medicines from this plant's methanolic extract.

MATERIALS AND METHODS

Sample preparation

Samples of the plant were collected from Nagarjun Hill, Kathmandu, Nepal; the aerial parts were dried, crushed, and extracted using a

*Corresponding author. E-mail: poudel.bhupen@gmail.com. Tel: 977-9841380139.

Soxhlet apparatus. Twenty grams of the powdered material was extracted with 200 ml volumes of petroleum ether, diethyl ether, methanol, and water, each in a stepwise manner. The extracts were concentrated and stored in refrigerator. Only methanolic extract was used for pharmacological screening. Male albino mice (30 ± 2.12 g) were used for gastrointestinal motility, spontaneous locomotor activity, and skeletal motility activity tests, whereas male wistar rats (280 ± 3.43 g) were used for anti-inflammatory test.

Gastrointestinal motility: Charcoal meal test

Mice were divided into 4 groups of 3 mice each; the first and second group were administered 500 and 100 mg/kg of methanol extract made in normal saline solution intraperitoneally (i.p), respectively; third group were administered atropine sulphate (5 mg/kg, i.p) and fourth group were administered control vehicle i.p. After 30 min, all animals were fed with 1 ml of charcoal meal (animal charcoal 12 g, tragacanth 2 g, and water 130 ml) intragastrically with the aid of feeding needle. All the mice were killed after 30 min by inhalation of chloroform. The abdomen was opened, the intestine quickly isolated and the small intestine from pylorus to caecum was cut by scissor and its length was measured by ruler. The intestinal distance moved by the charcoal meal from pylorus was measured to calculate percent of charcoal movement from pylorus to caecum as (Rouf et al., 2003):

$$\% \text{ of Charcoal movement} = \frac{\text{Distance travelled by the charcoal meal}}{\text{Total length of the small intestine}} \times 100$$

Spontaneous locomotor activity

Mice were divided into 5 groups, each consisting of 3 mice. First group, serving as control, received i.p, the vehicle only and second group received i.p 5 mg/kg of diazepam serving as standard, while the remaining groups were injected i.p with 125, 250 and 500 mg/kg of the methanolic extract i.p. After 30 min of administration of extract, each mouse was placed in an open square field 50×50 cm surrounding and subdivided into 25 squares of 10×10 cm. The percentage inhibition in locomotion was calculated as (Kulkarni 2000):

$$\% \text{ Inhibition} = 100 - \frac{\text{No. of squares crossed in test}}{\text{No. of squares crossed in control}} \times 100$$

Skeletal muscle activity: Traction technique

Mice were divided into five groups of five mice in each. Each mouse of first three groups were given methanolic extract i.p at the dose of 125, 250, and 500 mg/kg respectively. One group received only distilled water as a control and another group received diazepam at the dose of 5 mg/kg as a standard. After 30 min, the capability of the mice to touch the wire with at least one of the hind paws within 5 s was observed and percentage failure was calculated by the formula (Chattopadhyay et al., 2003):

$$\% \text{ Failure} = \frac{\text{No. of mice failed to grasp the wire}}{\text{Total no. of mice}} \times 100$$

Skeletal muscle activity: Inclined plane test

Mice were divided in 5 groups of 4 mice each. First group was given 125mg/kg, methanol extract, second group 250mg/kg methanol

extract, third group 500 mg/kg, fourth group diazepam 5 mg/kg and the fourth group i.e. control group was given distilled water only i.p. After 30 min they were placed in the inclined plane 45° for 2 min and their ability to remain at such inclination was noted.

Percentage failure was calculated by the following formula (Chattopadhyay et al., 2003):

$$\% \text{ failure} = \frac{\text{no of mice failed to remain in the inclined plane}}{\text{total no of mice}} \times 100$$

Anti-inflammatory activity: Carrageenan test

It was studied in male wistar rats (280 ± 3.43 g) as per the method described by Winter et al. (1962). Rats were divided into 5 groups, each consisting of 3 rats. One group serving as control received i.p the vehicle (normal saline) only and the other group received i.p 100 mg/kg of aspirin and 10 mg/kg diclofenac serving as standard, while the remaining groups were injected i.p with 200 and 400 mg/kg of the methanol extract. One hour later each animal was injected with 0.1 ml of 1% carrageenan suspension in normal saline into the sub-plantar region of right hind paw. The diameter of the paw was measured in each rat by vernier caliper before and 3 h after carrageenan injection. The diameter of oedema was recorded as the difference between the two readings.

The percentage inhibition of oedema was calculated as (Arumozhi et al., 2005):

$$\% \text{ Inhibition} = 100 - \frac{\text{Change in diameter of test group}}{\text{Change in diameter of control group}} \times 100$$

RESULTS AND DISCUSSION

The extractive value for methanol extract was higher than for other extracts; and hence was used for pharmacological screening. Similar higher percentage yield was found for methanolic extract of *S. arvensis* than with other fractions (Ali 2012).

In this study, the extract decreased propulsion of the charcoal meal through the gastrointestinal tract of mice dose dependently when compared with the control group. The 500 mg/kg; intraperitoneal (ip) methanol and 5 mg/kg; ip atropine had comparable inhibitory activity in intestinal motility (Figure 1). This reduction in gastrointestinal motility by methanol extract of *S. arvensis* may be due to antisecretory effects. The numerous phytochemicals like tannins, polyphenolic compounds, flavonoids, quercetin and other chemical compounds may be speculated for antimotility effect (Ezekwesili et al., 2010). Hence, this activity of the plant may be useful in treatment of diarrhoea as an antimotility agent.

Similarly, methanol extract significantly inhibited locomotor activity on mice in dose dependent fashion indicating antidepressant property (Figure 2). Therefore, a standardized *S. arvensis* extract or its purified constituents could be of potential interest for the GI motility disorders.

The traction test and inclined plane test revealed that the methanol extract didn't possess skeletal muscle relaxant activity (Table 1 and 2).




Figure 1. Effect of methanol extract on intestinal motility.




Figure 2. Effect of methanol extract on locomotor activity.

Further, methanol extract inhibited carrageenan induced acute paw oedema in dose dependent manner indicating anti-inflammatory activity. 400 mg/kg; ip methanol extract had higher anti-inflammatory activity than 10 mg/kg; ip diclofenac (Figure 3). Carrageenan releases prostaglandins (Winter et al., 1962) and inflammation occurs because of a proteolytic process with formation of kinin-like mediator(s) (Rosa and Sorrentino 1968).

S. arvensis contains various compounds palmitic acid, β -sitosterol, daucosterol, quercetin, apigenin-7-O- β -glucopyranoside, luteolin-7-O- β -D-glucopyranoside, quercetin-3-O- β -D-glucopyranoside and rutin (Jiang et al.

2009) which might have produced above pharmacological effects. However, further studies are required to isolate the major bioactive constituents and to verify the findings.

Conclusions


This study revealed significant anti-inflammatory, inhibitory effect on both gastrointestinal and locomotor activity. The study, however, revealed no skeletal muscle activity. It may, therefore, be concluded from this study that the plant possessed constituents that revealed the

Table 1. Effect of methanol extract of on skeletal muscle of mice by traction test.

| Treatment | Dose (mg/kg; ip) | No. of mice failed to grasp the wire | Failure (%) |
|------------------|------------------|--------------------------------------|-------------|
| Control | - | 0 | 0 |
| Methanol extract | 125 | 0 | 0 |
| Methanol extract | 250 | 0 | 0 |
| Methanol extract | 500 | 0 | 0 |
| Diazepam | 5 | 4 | 100% |

Table 2. Effect of methanol extract on skeletal muscle of mice by inclined plane test.

| Treatment | Dose (mg/kg; ip) | Animal falling after treatment | Failure (%) |
|---------------------|------------------|--------------------------------|-------------|
| Control | - | 0 | 0 |
| Methanol extract | 125 | 0 | 0 |
| Methanol extract | 250 | 0 | 0 |
| M Methanol extract | 500 | 0 | 0 |
| Standard (Diazepam) | 5 | 4 | 100 |

**Figure 3.** Effect of methanol extract on anti-inflammatory activity.

above pharmacological properties which may generate lead molecules for development of newer drugs. However, to reach any conclusive decision, a detailed phytochemical study for isolation, purification, identification, and characterization of the compound and biological studies with exact mechanism of action responsible for the particular activity, is necessary. Hence, further scientific investigation and specific studies are highly recommended for better evaluation of the potential effectiveness of the plant.

ACKNOWLEDGEMENT

The authors are thankful to Department of Pharmacy,

Institute of Medicine, Tribhuvan University for providing laboratory and laboratory materials to conduct experiments.

Conflict of Interests

The authors have not declared any conflict of interest.

REFERENCES

- Ali Khan R (2012). Evaluation of flavonoids and diverse antioxidant activities of *Sonchus arvensis*. *Chem. Central J.* 6:126.
- Amatya MP, Srivastava DL, Shakya KS, Pradhan NB, Karmacharya K, Shrestha UK, Ranjit R (2009). Phytochemical and Pharmacological studies of *Osbeckiastellata* Buch.-Ham. ex D. Don. *Plant Res. Bull. Dept.* 31:56-63.

- Australian Plant Name Index (APNI) (2012). IBIS database. Centre for Australian National Biodiversity Research. Australian Government, Canberra. Available at: <http://www.cpbr.gov.au/cgi-bin/apni>
- Arumozhi DK, Veeranjanyulu A, Badhankar SL, Arora SK (2005). Pharmacological investigations of *sapindustrifoliatus* in various *in vitro* and *in vivo* models of inflammation. *Indian J. Pharmacol.* 37:96-102.
- Chattopadhyay DP, Arunachalam G, Mandal SC, Bhadra R, Mandal AB (2003). CNS activity of the methanol extracts of *Mallotuspeltatus* (geist) Muell Arg. leaf: An ethnomedicine of onge. *J. Ethnopharmacol.* 85:99-105.
- Ezekwesili JO, Nkemdilim UU, Okeke CU (2010). Mechanism of antidiarrhoeal effect of ethanolic extract of *Psidium guajava* leaves. *Biokemistri* 22(2):201.
- Holm L, Doll J, Holm E, Pancho J, Herberger J (1997). *World weeds: natural histories and distribution.* John Wiley & Sons. P. 1129.
- Jiang L, QingQiang Y, YanYing X (2009). Study on chemical constituents of *Sonchus arvensis* L. *Food Drug* 11(3):27-29.
- Kulkarni SK (2000). *Hand Book of Experimental Pharmacology*, 2nd ed India: VallabhPrakashan. pp. 43-56.
- Rajbhandari KR (2001). *Ethnobotany of Nepal.* Kathmandu: Ethnobotanical Society of Nepal (ESON). P. 189.
- Rosa MD, Sorrentino L (1968). The mechanism of the inflammatory effect of carrageenin. *Eur. J. Pharmacol.* pp. 340-342.
- Rouf ASS, Islam MS, Rahman MT (2003). Evaluation of antidiarrhoeal activity *Rumexmaritimus* root. *J. Ethnopharmacol.* 84:307-310.
- Winter CA, Risley EA, Nuss GW (1962). Carrageenin-induced edema in hind paw of the rat as an assay for antiinflammatory drugs. *Proc. Soc. Exp. Biol. Med.* 111:544-7.



Related Journals Published by Academic Journals

- African Journal of Pharmacy and Pharmacology
- Research in Pharmaceutical Biotechnology
- Medical Practice and Reviews
- Journal of Clinical Pathology and Forensic Medicine
- Journal of Medicinal Plant Research
- Journal of Drug Discovery and Development
- Journal of Clinical Virology Research

academicJournals