

A photograph of a zebra in a green field, used as a background for the journal cover. The zebra is in profile, facing left, with its head slightly lowered. The background is a soft-focus green field.

Journal of Ecology and The Natural Environment

Volume 8 Number 1, January 2016

ISSN 2006-9847



*Academic
Journals*

ABOUT JENE

The **Journal of Ecology and the Natural Environment (JENE)** (ISSN 2006-9847) is published Monthly (one volume per year) by Academic Journals.

Journal of Ecology and the Natural Environment (JENE) provides rapid publication (monthly) of articles in all areas of the subject such as biogeochemical cycles, conservation, paleoecology, plant ecology etc.

The Journal welcomes the submission of manuscripts that meet the general criteria of significance and scientific excellence. Papers will be published shortly after acceptance. All articles published in JENE are peer-reviewed.

Contact Us

Editorial Office: jene@academicjournals.org

Help Desk: helpdesk@academicjournals.org

Website: <http://www.academicjournals.org/journal/JENE>

Submit manuscript online <http://ms.academicjournals.me/>

Editors

Dr. Abd El-Latif Hesham

*Genetics Department, Faculty of Agriculture, Assiut University,
Assiut 71516,
Egypt*

Dr. Ahmed Bybordi

East Azarbaijan Research Centre for Agriculture and Natural Resources, Tabriz, Iran

Dr. Sunil Kumar

*Natural Resource Ecology Laboratory,
Colorado State University
1499 Campus Delivery, A204 NESB, Fort Collins,
Colorado-80526,
USA*

Prof. Gianfranco Rizzo

*University of Palermo
Dipartimento DREAM – Viale delle Scienze - Building
9. 90128
Palermo,
Italy*

Dr. Bahman Jabbarian Amiri

*Kiel University, Germany,
Ökologie-Zentrum der CAU
Abt. Hydrologie und Wasserwirtschaft
Olhausen Straße, 75
Kiel,
Germany*

Dr. Bikramjit Sinha

*National Institute of Science Technology and Development Studies,
Pusa Gate, Dr. KS Krishnan Marg, New Delhi 110012,
India*

Prof. Gianfranco Rizzo

*University of Palermo
Dipartimento DREAM – Viale delle Scienze - Building
9. 90128
Palermo,
Italy*

Associate Editors

Dr. Marko Sabovljevic

*Dept. Plant Ecology, Faculty of Biology,
University of Belgrade
Takovska 43, 11000 Belgrade,
Serbia*

Dr. Sime-Ngando Téléphone

*CNRS
LMGE, UMR 6023, Université Blaise Pascal, 63177,
Aubière Cedex
France*

Dr. Bernd Schierwater

*ITZ, Ecology and Evolution, TiHo Hannover
Büenteweg 17d, 30559 Hannover,
Germany*

Dr. Bhattacharyya Pranab

*North-East Institute of Science & Technology
Medicinal, Aromatic & Economic Plant Division,
North-East Institute of Science & Technology,
Jorhat-785006, Assam,
India*

Prof. Marian Petre

*University of Pitesti, Faculty of Sciences
1 Targul din Vale Street, Pitesti, 110040, Arges
County,
Romania.*

Prof. R.C. Sihag

*CCS Haryana Agricultural University
Department of Zoology & Aquaculture, Hisar-125004,
India*

Prof. Kasim Tatic

*School of Economics and Business, University of Sarajevo
Trg oslobođenja 1, 71000 SARAJEVO,
Bosnia and Herzegovina*

Dr. Zuo-Fu Xiang

*Central South University of Forestry & Technology,
498 Shaoshan Nanlu,
Changsha, Hunan, China.*

Editorial Board

Dr. Zuo-Fu Xiang

*Central South University of Forestry & Technology,
498 Shaoshan Nanlu,
Changsha, Hunan, China.*

Dr. Pankaj Sah

*Higher College of Technology, Muscat,
Department of Applied Sciences,
(Applied Biology) Higher College of Technology,
Al-Khuwair, PO Box 74,
PC 133, Muscat
(Sultanate of Oman)*

Dr. Arti Prasad

*Mohan Lal Sukhadia University,
Udaipur, Rajasthan, India.
123, Vidya Nagar, Hiran Magri,
Sector-4, Udaipur, Rajasthan,
India*

Parviz Tarikhi

*Mahdasht Satellite Receiving Station
(Postal): No. 80, 14th Street, Saadat Abad Avenue,
Tehran 1997994313,
Iran*

Bharath Prithiviraj

*Post Doctoral Research Associate
Knight Lab, Dept. of Chemistry & Biochemistry
University of Colorado at Boulder
USA*

Dr. Melissa Nursey-Bray

*Australian Maritime College, Tasmania,
Australia*

Parvez Rana

*Department of Forestry and Environmental Science
Shahjalal University of Science and Technology
Bangladesh*

Mirza Hasanuzzaman

*Faculty of Agriculture, Sher-e-Bangla Agricultural
University
Sher-e-Bangla Nagar, Dhaka-1207,
Bangladesh*

Dr. Giri Kattel

*Murray Darling Freshwater Research Centre, La Trobe
University
471 Benetook Avenue, Mildura, Victoria 3500,
Australia*

Dr. M. Rufus Kitto

*Faculty of Marine Science-Obhur station,
King Abdulaziz University,
Jeddah 21589, Saudi Arabia*

Dr. Özge Zencir

*Kemah Vocational Training School,
Erzincan University, Kemah,
Erzincan, Turkey.*

Dr. Sahadev Sharma

*Laboratory of Ecology and Systematics,
Graduate School of Engineering and Science,
University of the Ryukyus, Senbaru 59,
Nishihara, Okinawa-903-0213 Japan*

Dr. Hasan Kalyoncu

*University of Süleyman Demirel,
Faculty of Art and Science,
Department of Biology,
32100 Isparta/Turkey*

Hammad Khan

*Department of Zoology and Fisheries,
University of Agriculture,
Faisalaad, Pakistan*

Mirza Hasanuzzaman

*Faculty of Agriculture,
Sher-e-Bangla Agricultural University
Sher-e-Bangla Nagar, Dhaka-1207,
Bangladesh*

Abdurrahman Dundar

*Siirt University, Science and Arts Faculty,
Department of Biology,
56000, Siirt, Turkey*

Meire Cristina Nogueira de Andrade

*College of Agronomic Sciences,
São Paulo State University, Brazil.*

Imran Ahmad Dar

*Dept. of Industries and Earth Sciences,
The Tamil University,
Ocean and Atmospheric Sciences & Technology Cell,
(A Unit of Ministry of Earth Sciences, Govt. of
India).*

S. Jayakumar

*Department of Ecology and Environmental
Sciences,
School of Life Sciences,
Pondicherry University,
Puducherry - 605 014, India*

Umer Farooq

*University of Veterinary & Animal Sciences
Lahore, Pakistan*

Journal of Ecology and the Natural Environment

Table of Contents: Volume 8 Number 1, January 2016

ARTICLE

Using ecological niche models to plan conservation in a changing environment: A case for the plant *Chasmanthera dependens* Hochst (Menispermaceae) in West Africa

1

Andrew Chibuzor Iloh and Oluwatoyin Temitope Ogundipe

Full Length Research Paper

Using ecological niche models to plan conservation in a changing environment: A case for the plant *Chasmanthera dependens* Hochst (Menispermaceae) in West Africa

Andrew Chibuzor Iloh^{1*} and Oluwatoyin Temitope Ogundipe²

¹Biodiversity and Climate Research Group, Biotechnology Advanced Laboratory, Sheda Science and Technology Complex, Abuja, Nigeria.

²Molecular Systematics Laboratory, Department of Botany, University of Lagos, Lagos, Nigeria.

Received 13 May, 2015; Accepted 6 October, 2015

Climatic envelope modeling techniques implemented in two algorithms, Genetic Algorithm for Rule-set Production (GARP) and bioclimatic variables (BIOCLIM) were used to assess effects of climatic conditions on distributions of plants and anticipate how climate would have delimited their distribution under future conditions using a liana species *Chasmanthera dependens* as a case example. In all, 120 geo-referenced plant records generated from fieldwork and drawn from data served by the Global Biodiversity Information Facility (GBIF). Environmental variables were derived from monthly temperature and rainfall data from WorldClim; eliminating environmental variables with correlations of 0.75 and left eight (8) variables for analysis. Results show that the current suitable range (ecological niche) of the model plant was broad across the tropical rain forest regions. Predictions to future climate scenarios (2050), predicted a significant reduction of suitable distributional areas for the species suggesting possible loss of plant species. Indeed, *ex-situ* conservation may be the most appropriate conservation tool for this species and others in similar situations.

Key words: Bioclimatic variables (BIOCLIM), climate change, conservation, ecological niche models, genetic algorithm for rule-set production (GARP).

INTRODUCTION

Viéet al. (2008) have reported that the numbers of threatened species are on the increase every year; and more attention needs to be given to them in the light of emerging issues of climatic change. According to McCarty (2001) and Hughes (2000), climate change at global and regional scales is predicted to alter species distributions, life histories, community composition and

ecosystem function.

In particular, population losses caused by climate change threaten both species diversity and the delivery of critical ecosystem services (Hughes, 2000). Climate change is now seen to be one of the major causes of biodiversity loss; these impacts are further weakening already fragile ecosystems, for example, the continuous

*Corresponding author. E-mail: chibaziloh@gmail.com.

expansion of deserts in the northern Nigeria (CBD, 2007). The effects of climate change on biodiversity require and demand detailed assessment especially in developing countries like Nigeria and across West Africa more generally.

Lianas (woody vines) according to Schnitzer and Bongers (2002), are non-self-supporting epiphytic plants that uses the architecture of trees to ascend to the forest canopies. They are particularly abundant and diverse in lowland tropical forests, where they constitute up to 40% of the woody biomass representing more than 25% of the woody species (Schnitzer and Bongers, 2002), and contribute substantially to forest leaf area and biomass (Gerwing and Farias, 2000; Chave et al., 2001).

Interestingly, Martin et al. (2004) reports that they are more prevalent in areas of secondary forest succession and are often able to compete effectively against tree and shrub species in specialized environmental conditions such as areas of both acute and chronic disturbance. Furthermore, with increases in global temperatures due to climate change, it has been projected that vine growth rates will further increase (Phillips et al., 2002).

Chasmanthera dependens liana can provide information on liana roleas important structural components and key species richness indicators of tropical forests; where they are important physiognomic and structural components (Gentry, 1991; Schnitzer and Bongers, 2002). It belongs to the family Menispermaceae and is reported as the only *Chasmanthera* species in West Africa. Ortiz et al. (2007) reported that *Chasmanthera* and other Menispermaceae are poorly known for lack of a comprehensive phylogenetic hypothesis. Since *C. dependens* is mainly wild, genetic erosion owing to over-exploitation could occur.

This risk may increase owing to effects of climate change, resulting in population decline. Moritz et al. (2008) have reported the impacts of a century of climate change on small-mammal communities in Yosemite National in the USA while Rubidge et al. (2012) has reported that climate-induced range contraction drives genetic erosion in alpine mammals. These changes in environmental conditions can rapidly shift allele frequencies in populations of species with relatively short generation times thus causing a decrease in their populations (Hoffmann and Willi, 2008). Therefore, an understanding of how plants will fare in the present changing climate would provide information essential to effectively protecting the species.

The most important tools in conservation planning with respect to climate change according to Elith and Leathwick (2009) should be the deployment of species distribution models which will evaluate present and potential future species ranges in relation to climate, soils and other predictive variables. These models according to Thuiller et al. (2005), can highlight individual species that may be at risk because of climate change, and geographic areas that may face substantial shifts in

diversity and species composition (Williams et al., 2005; Loarie et al., 2009). In some cases, models suggest that protected areas may no longer maintain populations of key species, possibly the very ones that the reserves were created to protect (Araújo et al., 2004).

Thorn et al. (2009) and Marini et al. (2009b) have also reported its usefulness in targeting conservation efforts for threatened species. Conversely, shrinking distributions and range shifts could create new refugia - areas where threatened species would be concentrated in the future (Loarie et al., 2009). This according to Marini et al. (2009a) will provide frameworks for identifying spatially optimal conservation strategies which includes designing networks for these priority areas.

Using model plant *C. dependens* Hochst (Menispermaceae) in West Africa, we present ecological niche models (ENM) which were used to determine the potential distribution of the species across West Africa, predict future distribution patterns and thus use these models to develop conservation planning priorities.

MATERIALS AND METHODS

Input data

Two types of data were used as the basis for this study: geographic location records of occurrence of the species and geospatial data sets describing environmental variation across the region. According to Peterson et al. (2012), occurrence records should be from sites spread over the range such that a species' occurrence patterns can be assessed in the light of its influenced major environmental dimensions of variation. Such broad sampling of occurrence is known to increase the accuracy of ecological niche model results (Peterson et al., 2012).

Hence, the entire African range of the liana was taken into account in model calibration, but results were presented only for West Africa. Occurrence data were obtained from field collections across five West African countries; Ghana; Togo, Benin, Nigeria and Cameroun. Additional occurrence data were acquired from data mediated by the Global Biodiversity Information Facility (GBIF), from the Census of the flora of Benin, Naturalis Biodiversity Centre (NL) - Botany Wageningen, Missouri Botanical Garden, National Herbarium Nederland, University of Ghana - Ghana Herbarium, Real Jardín Botánico (Madrid), Vascular Plant Herbarium (MA), Botany (UPS), Vascular Plant Collection at the Botanische Staatssammlung München, Herbarium Togoense, Herbarium du Bénin, Herbarium Senckenbergianum (FR) and TAIIF (Taiwan e-Learning and Digital Archives Program, TELDAP). In order to have single occurrence information from the various herbaria data collected, duplicate records were removed from the data set.

All data were then displayed in relation to country boundaries and examined visually to detect and remove records in incorrect locations that are outliers that did not fall within the country boundaries which may be due to improper geo-referencing from the herbarium records.

Environmental data

In order to generate more biologically meaningful variables, bioclimatic variables (BIOCLIM) derived from the monthly temperature and rainfall values were used as climatic data from

these experiments. Current and future climate data in the form of 19 “bioclimatic” variables were obtained (<http://www.worldclim.org>) at a 2.5’ spatial resolutions. These variables represent annual trends (for example, mean annual temperature, annual precipitation) seasonality (for example, annual range in temperature and precipitation) and extreme or limiting environmental factors (for example, temperature of the coldest and warmest month, and precipitation of the wet and dry quarters).

Current data were from the WorldClim data set (Hijmans et al., 2005) while the 2050, were from the General Circulation Model (GCM): CCCMA climate model, A2 scenario generated by the Intergovernmental Panel on Climate Change (IPCC, 2007).

GCMs depict the climate using a three dimensional grid over the globe, typically having a horizontal resolution of between 250 and 600 km, 10 to 20 vertical layers in the atmosphere and sometimes as many as 30 layers in the oceans. Their resolution is thus quite coarse relative to the scale of exposure units in most impact assessments. Moreover, many physical processes, such as those related to clouds, also occur at smaller scales and cannot be properly modelled. Instead, their known properties must be averaged over the larger scale in a technique known as parameterization. This is one source of uncertainty in GCM-based simulations of future climate. Others relate to the simulation of various feedback mechanisms in models concerning, for example, water vapour and warming, clouds and radiation, ocean circulation and ice and snow albedo. For this reason, GCMs may simulate quite different responses to the same forcing; simply because of the way certain processes and feedbacks are modelled. However, these differences in response are usually consistent with the climate sensitivity range.

The A2 marker scenario (A2-ASF) was developed using Atmospheric Stabilisation Framework Model (ASF), an integrated set of modelling tools that was also used to generate the first and the second sets of IPCC emission scenarios (SA90 and IS92).

Overall, the A2-ASF quantification is based on the following assumptions (Sankovski et al., 2000): Relatively slow demographic transition and relatively slow convergence in regional fertility patterns; Relatively slow convergence in inter-regional GDP per capita differences; Relatively slow end-use and supply-side energy efficiency improvements (compared to other storylines); Delayed development of renewable energy as well as no barriers to the use of nuclear energy.

ENM approaches

Several studies have advised consensus approaches in ENM development (Pearson et al., 2007). We applied two algorithms GARP (Stockwell, 1999) and BioCLIM (Busby, 1991) to construct ENMs. Present day ENMs were developed based on occurrences within the mask appropriate to the model species; in this case, the mask area was the 5 West African countries from where we sampled its populations.

The climate envelope techniques BioCLIM is a classic bioclimatic modelling approach. It fits a minimal envelope in a multidimensional climate space and use presence-only instead of presence/absence data, which can be highly advantageous: many data sets provide presence-only data, and even if absence points are available, they are not always reliable, especially for areas that are not thoroughly inventoried or for species that are difficult to detect. On the other hand, if information on absence points is available and reliable, it is to a model’s disadvantage not to employ it (Jeschke and Strayer, 2008).

GARP (Genetic Algorithm for Rule-set Production) does not need presence/absence data for its application, but presence-only data are sufficient. GARP includes multiple, non-deterministic iterative procedures that incorporate various model distribution methods such as logistic regression and range envelopes, producing with

Table 1. Bioclimatic variables, derived from temperature and precipitation, used in the analysis.

S/N	Parameter	Meaning
1	Bio 1	Annual mean Temperature
2	Bio 2	Mean monthly Temperature
3	Bio 5	Max temp of warmest month.
4	Bio 6	Minimum temp of coldest month
5	Bio 12	Annual precipitation
6	Bio 13	Precipitation of the wettest month
7	Bio 14	Precipitation of the driest month
8	Bio 15	Precipitation seasonality

each run predicted binary maps of presences and absences. Multiple optimal models are produced for each data set, which can be converted into presence likelihoods. However, BioCLIM (envelop algorithm) is less complex than GARP, and often perform poorly in simulations (Wisiz et al., 2008). Different algorithms produce different outputs, but in general convey presence probabilities, or some arbitrary value that could be interpreted in similar fashion.

For Desktop GARP (version 1.1.6) to construct ENMs, 100 random replicate models were developed using the default parameters of convergence limit (0.01) and maximum iterations (1000). The GARP was run on the OpenModeller platform (1.10) (Muñoz et al., 2011). Also using the OpenModeller software, the BIOCLIM was calculated by disregarding the lower and higher 5% to reduce the impact of outliers (Muñoz et al., 2011).

Prior to this, we performed a two-tailed spearman correlation analysis between all pairwise combinations of 19 variables based on their values at 1000 points selected randomly from across West Africa. Eliminating environmental variables with correlations of 0.75 and less, that is, variables that least likely limited geographic distributions, for example, removing mean values rather than maxima or minima (Wiens et al., 2006; Paz, et al., 2015), we were left with eight (8) bioclimatic variables used to construct the models (Table 1).

The Niche A software was also used to provide an improved method to calculate the Area Under Curve (AUC) by calculating the AUC value of the partial area of the Receiver Operating Characteristic (ROC). This was run on the E value of 0.5.

RESULTS

Results from the model construct are presented in Figures 1A and 1B for ENM of *C. dependens* using BIOCLIM and Figures 2A and 2B for ENM of *C. dependens* using GARP.

Figures 1A (BIOCLIM) and 2A (GARP) predicted current highly suitable area for the species across the tropical forest belt of West Africa, this is in line that the model plant is widely distributed in the tropical forest regions of West Africa. Models do not clearly show accurately the distribution from the Camerouns, this was understandable as the presence-only data used from the Camerouns was limited. However, both models showed a possibility of plant survival and distribution in the tropical region of the Camerouns. The models also showed some marginally suitable areas on the maps of Nigera, Benin, Togo and Ghana which were likely to be the sudan




Figure 1A-B. Ecological niche models for *Chasmanthera dependens* using BIOCLIM and showing West African potential distribution at present and projected to 2050.




Figure 2A-B. Ecological niche models for *C. dependens* using GARP and showing West African potential distribution at present and projected to 2050.

regions. This could be defined as the plant's fundamental niche that is; the range of environmental conditions in which the species survives. Areas above the sudan (sahel) as well as the magrove regions were not suitable for plant growth as analysed by the models. Models showed a pseudo-probability surface of the test plant distribution.

Figures 1B and 2B presents future distribution areas of the species across the tropical belt of West Africa. Models were able to show a contraction in niche from the present distribution to the future distribution. This contraction in niche could clearly be seen in Nigeria, Benin, Togo and the Ghana. Maximum suitable areas on the 2050 maps were mainly in the South-Western parts of Nigeria and majorly in the Camerouns. Interestingly, there was little or no contraction in this area.

DISCUSSION

In this study, a total of 120 geo-reference plant records were used in the model calibration. According to Hernandez et al.(2006), the ability to estimate ecological

niches effectively is influenced strongly by species' ecological characteristics independent of sample size and models built with few points, while not as accurate as those based on large data sets and potentially not appropriate for all applications, can still be useful (Hernandez et al., 2006). The Bioclimatic variables used to run the models had correlations below 0.75 according to Peterson et al. (2011) variables, a correlation below 0.75 give best model results.

Both algorithms used had an area under the curve (AUC) of the partial area of the receiver operating characteristic (ROC) of BIOCLIM0.94 and GARP0.95, respectively. The use of partial-area ROC approaches to provide a firmer foundation for evaluation of predictions from ecological niche models (Peterson et al., 2008). This is because models evaluated favorably by traditional ROC, AUCs are not necessarily the best when niche modeling considerations are incorporated into the design of the test thus a recommendation that a modification of ROC that remedies these problems is used (Lobo et al., 2008; Peterson et al., 2008).

The model predicted current suitable area for the species across the tropical forest belt of West Africa. Our

present model results showed an ecological niche representation which coincides with the geographic distribution based on herbarium specimens and field work of the plant species corroborating that niche conditions of the species are represented in the known geographic range, that is; a realized niche that is the range of environmental conditions in which a species is really found. This result is in line with many studies that indicate that a species' fundamental niche is almost always larger than its realized niche in terms of geographic footprint (Pearson and Dawson, 2003; Peterson, 2003a). Cuni-Sanchez et al. (2010), for example, used ecological niche modeling to estimate the fundamental niche of the tree *Adansonia digitata* (baobab) and have considered this area to reflect the global cultivation potential or invasive potential of the species.

As these models have proven to identify potential distribution areas of species it is important to note that geographical distribution of species are influenced by biological interactions and historical factors which are not taken into account in the modeling algorithms (Illoldi-Rangel and Escalante, 2008). Hence, ENMs are ecological niche modeling interpreted as the geographical representation of environmental conditions required by the species (Martínez-Meyer, 2005).

As reported in Rojas-Soto et al.(2003), Sánchez-Cordero et al.(2005), predicted distributions of species from ENM approaches provides clear gains as well as offer a number of opportunities over raw existence data. Whereas, as raw occurrence data may be bias in detectability and sampling efforts, Soberón and Peterson (2004) reports that modeled distributions as input information for these models can overcome these biases to an impressive degree. These methods according to Pressey and Cowling, (2001) help to compensate for the lack of comprehensive distributional data in regional conservation planning.

Pittock (2009) has reported that estimated variables and process applied in a model construction are likely to influence the outcome of the model hence increasing the range of uncertainty in climate projection. Therefore, rather than projecting what will happen really in the future, climate projection is a procedure for providing insights into what may happen. As such, reports will enable a robust decision-making in the face of uncertainty as well as how much to offer conservation planning (Araújo and New, 2007).

Our ENMs were able to demonstrate the likely climate change effects on species' potential distributions across West Africa, these effects of climate change agree well with reports from parallel studies in other regions (Peterson, 2003b; Peterson, 2003c; Peterson et al., 2002, Thuiller, 2004; Araújo and New, 2007; Christensen et al., 2007). Again, Williams et al. (2003) has reported that these findings are in contrast with the severe losses predicted for more diverse environments, such as tropical

rainforest.

Models were also able to show massive loss of potential habitat in Benin Republic. Gouwakinnou (2013) has also reported this in the tree species *Sclerocarya birrea* in Benin Republic. Interestingly, in the Camerouns, there was little or no contraction in this area, is there a possibility that the refugia hypothesis which suggests that the current diversity in the tropics could be attributable to successive isolation and subsequent expansion of populations in response to frequent oscillations of the climate during the quaternary period (Mittelbach et al., 2007). Since fundamental niche might be lost, there is a possibility that areas in the Cameroun might just be point of refugia for West African plants where conditions might be favorable for their existence.

In presenting the results of this study, it should be taken into account that changes in the environment brought about by the effects of climatic change under emission scenarios are only hypotheses of the changes in the environmental requirements (niche) of the species. This according to Pearson and Dawson (2003) takes into account the reactions individual species exhibits to climatic change and therefore, these possibilities may not be considered in the bioclimatic models (Pearson and Dawson, 2003).

The ability for species to adapt to new conditions or move to new places where conditions are favorable is crucial survival (Pearson and Dawson, 2003). However, it is also possible that species do not adapt to new changes, and that the range lost automatically represents the total disappearance, minimally or in part, of its geographical range. The reduced potential habitat in this study may poses these challenges hence, if plant species is not able to adapt to the new changes, it might automatically lose their current distribution.

Conservation concerns

Our understanding of plant distributions is precursors to effective protection of our forest especially in Africa. However, these practices will be greatly set aback with the issues of climate change. The ENMs of this study were able to demonstrate the likely climate change effects on species' potential distributions across West Africa and thus maintaining diversity under the present climatic scenarios is likely not to be sustained due to the possible reduction in ecological niche in future. Indeed, *ex-situ* conservation may be the most appropriate conservation tool for this species and others in similar situations across West Africa.

Conflict of interests

The author(s) did not declare any conflict of interest.

ACKNOWLEDGMENTS

Special thanks to Prof. A. Townsend Peterson for granting the first author the opportunity to attend the ENM course organized by the Biodiversity Informatics Training Curriculum, University of Kansas, USA. We also thank him for all the discussions and useful comments regarding this manuscript. I also thank the reviewers of this manuscript, they provided a lot of questions and suggestions that have made this work a better one. Thank you so much.

REFERENCES

- Araújo MB, New M (2007). Ensemble forecasting of species distribution. *Trends Ecol. Evol.* 22:42-47.
- Araújo MB, Cabeza M, Thuiller W, Hannah L, Williams PH (2004). Would climate change drive species out of reserves? An assessment of existing reserve-selection methods. *Glob. Change Biol.* 10:1618-1626.
- Busby JR (1991). BIOCLIM - a bioclimatic analysis and predictive system. In: Margules, C.R. and Austin, M.P. (Eds.) *Nature Conservation: Cost effective biological surveys and data analysis*. CSIRO, Canberra, pp. 64-68.
- Chave J, Riéra B, Dubois M (2001). Estimation of biomass in a Neotropical forest in French Guiana: spatial and temporal variability. *J. Trop. Ecol.* 17:79-96.
- Christensen JH, Hweitson B, Busuioac A, Chen A, Gao X, Held R, Magana R, Rueda V, Mearns L, Menendez CG, Räisänen J, Rinke A, Sarr A, Whetton P, Arritt R, Benestad R, Beniston M, Bromwich D, Caya D, Comiso J, de-Elia R, Dethloff K (2007). Regional climate predictions. In: Solomon S, Qin D, Manning Z, Chen M, Marquis KB, Averyt KB, Tignor M and Miller H (Eds.). *Climate change, 2007: the physical science basis. Contribution of the working group 1 to the fourth Assessment report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, UK. pp. 847-940.
- Convention on Biological Diversity (CBD) (2007). *Emerging issues for biodiversity conservation in a changing climate*. CBD, Montreal, Canada. 600p.
- Elith J, Leathwick JR (2009). Species distribution models: ecological explanation and prediction across space and time. *Annu. Rev. Ecol. Syst.* 40: 677-697.
- Gentry AH (1991). The distribution and evolution of climbing plants. In: Putz, FE. and Mooney, HA (Eds.). *The biology of Vines* Cambridge University Press, Cambridge, UK. pp. 3-51.
- Gerwing JJ, Farias DL (2000). Integrating liana abundance and forest stature into an estimate of total aboveground biomass for an eastern Amazonian forest. *J. Trop. Ecol.* 16:327-335.
- Gouwakinnou GN (2013). Using niche modeling to plan conservation of an indigenous tree species under changing climate: example of *Sclerocarya birrea* in Benin. *West Afr. Res. Develop. Sub-Saharan Afr.* 6:1-8.
- Hernandez PA, Grahm CH, Master LL, Albert DL (2006). The effect of sample size and species characteristics on performance of different species distribution modeling methods. *Ecography* 29:773-785.
- Hijmans RJ, Cameron SE, Parra JL, Jarvis A (2005). Very high resolution interpolated climatic surfaces for global land areas. *Int. J. Climatol.* 24:1965-1978.
- Hoffmann AA, Willi Y (2008). Detecting genetic responses to environmental change. *Nat. Rev. Genet.* 9:421-432.
- Hughes L (2000). Biological consequences of global warming: is the signal already. *Trends Ecol. Evol.* 15(2):56-61.
- Illoldi-Rangel P, Escalante T (2008). De los modelos de nicho ecológico a las áreas de distribución geográfica. *Biogeografía* 3:7-12.
- IPCC (Intergovernmental Panel on Climate Change) (2007). *Climate Change 2007: the scientific basis. Contribution of Working Groups I, II and III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change Core Writing Team, Pachauri, R.K. and Reisinger, A. (Eds.) IPCC, Geneva, Switzerland. 104p.*
- Jeschke JM, Strayer DL (2008). Usefulness of Bioclimatic Models for Studying Climate Change and Invasive Species. *Ann. N.Y. Acad. Sci.* 1134:1-24
- Loarie SR, Duffy PB, Hamilton H, Asner GP, Field CB, Ackerly DD (2009). The velocity of climate change. *Nature* 462:1052-1055.
- Lobo JM, Jiménez-Valverde A, Real R (2008). AUC: a misleading measure of the performance of predictive distribution models. *Glob. Ecol. Biogeogr.* 17:145-151.
- Marini MA, Barbet-Massin M, Lopes ME, Jiguet F (2009a). Major current and future gap of Brazilian reserves to protect Neotropical savanna birds. *Biol. Conserv.* 142:3039-3050.
- Marini MA, Barbet-Massin M, Lopes ME, Jiguet F (2009b). Predicted climate-driven bird distribution changes and forecasted conservation conflicts in a Neotropical savanna. *Conserv. Biol.* 23:1558-1567.
- Martin PH, Sherman RE, Fahey TJ (2004). Forty years of tropical forest recovery from agriculture: structure and floristic of secondary and old-growth riparian forest in the Dominican Republic. *Biotropica* 36:297-317.
- Martinez-Meyer E (2005). Climate change and biodiversity: some considerations in forecasting shifts in species' potential distributions. *Biodivers. Inform.* 2:42-55.
- McCarty JP (2001). Ecological consequences of recent climate change. *Conserv. Biol.* 15: 320-331.
- Mittelbach GG, Schemske DW, Cornell HV, Allen AP, Brown JM, Bush MB, Harrison SP, Hurlbert AH, Knowlton N, Lessios H., McCain CM, McCune AR, McDade LA, McPeck MA, Near TJ, Price TD, Ricklefs RE, Roy K, Sax DF, Schluter D, Sobel JM, Turelli M (2007). Evolution and the latitudinal diversity gradient: speciation, extinction and biogeography. *Ecol. Lett.* 10:315-331.
- Moritz C, Patton JL, Conroy CJ, Parra JL, White GC, Beissinger SR (2008). Impact of a century of climate change on small-mammal communities in Yosemite National Park, USA. *Science* 322:261-264.
- Muñoz MES, Giovanni R, Siqueira MF, Sutton T, Brewer P, Pereira RS, Canhos DAL, Canhos VP (2011). Open Modeller: a generic approach to species' potential distribution modelling. *Geoinformatica* 15(1):111-135.
- Ortiz R, Kellogg EA, Van der Werff H (2007). Molecular phylogeny of the moonseed family (Menispermaceae): implications for morphological diversification. *Am. J. Bot.* 94(8):1425-1438.
- Paz A, Ibáñez R, Lips KR, Crawford AJ (2015). Testing the role of ecology and life history in structuring genetic variation across a landscape: a trait-based phylogeographic approach. *Mol. Ecol.* 24:3723-3737.
- Pearson RG, Dawson TP (2003). Predicting the impacts of climate change on the distribution of species: are bioclimatic envelopes models useful? *Glob. Ecol. Biogeogr.* 12:361-371.
- Pearson RG, Raxworthy C, Nakamura M, Peterson AT (2007). Predicting species' distributions from small numbers of occurrence records: a test case using cryptic geckos in Madagascar. *J. Biogeogr.* 34:102-117.
- Peterson AT (2003a). Predicting the geography of species' invasions via ecological niche modelling. *Q. Rev. Biol.* 78(4):419-433.
- Peterson AT (2003b). Projected climate change effects on Rocky Mountain and Great Plains birds: generalities of biodiversity consequences. *Glob. Change Biol.* 9:647-655.
- Peterson AT (2003c). Subtle recent distributional shifts in Great Plains bird species. *Southwest. Nat.* 48:289-292.
- Peterson AT, Ortega-Huerta MA, Bartley J, Sánchez-Cordero V, Soberón J, Buddemeier RH, Stockwell DRB (2002). Future projections for Mexican faunas under global climate change scenarios. *Nature* 416:626-629.
- Peterson AT, Papeş M, Soberón J (2008). Rethinking receiver operating characteristic analysis applications in ecological niche modeling. *Ecol. Model.* 213:63-72.
- Peterson AT, Radocoy T, Hall E, Kerbis JP, Celesia GG (2012). The potential distribution of the African lion in the face of changing global climate. *Oryx* 48(4):555-564.

- Peterson AT, Soberon J, Pearson RG, Anderson RP, Martinez-Meyer E, Nakamura M, Araújo MB (2011). *Ecological Niches and Geographic Distributions*. Princeton University Press, 328p.
- Phillips V, Martinez L, Arroyo TRB, Killeen T, Lewis SL, Malhi Y, Mendoza AM, Neill A, Vargas PN, Alexiades M, Ceron C, Fiore AD, Erwin T, Jardim A, Palacios W, Saldias M, Vinceti B (2002). Increasing dominance of large lianas in Amazonian Forest. *Nature* 418:770-774.
- Pittock BA (2009). *Climate change: the science, impacts and solutions*. 2nd edition CSIRO Publishing, Australia. 200p.
- Pressey RL, Cowling RM (2001). Reserve selection algorithms and the real world. *Conserv. Biol.* 15:275-277.
- Rojas-Soto OR, Alcantara-Ayala O, Navarro AG (2003). Regionalization of the avifauna of the Baja California Peninsula, Mexico: A parsimony analysis of endemism and distributional modelling approach. *J. Biogeogr.* 30:449-461.
- Rubidge EM, Patton JL, Lim M, Burton AC, Brashares JS, Moritz C (2012). Climate-induced range contraction drives genetic erosion in an alpine mammal. *Nat. Clim. Change* 2:285-288.
- Sanchez AC, Osborne PE, Haq N (2010). Identifying global potential for baobab tree cultivation using ecological niche modelling. *Agroforest. Syst.* 80:191-201.
- Sánchez-Cordero V, Cirelli V, Munguia M, Sarkar S (2005). Place prioritization for biodiversity representation using species' ecological niche modeling. *Biodivers. Inform.* 2:11-23.
- Sankovski A, Barbour W, Pepper WJ (2000). Quantification of the IS99 emission scenario storylines using the atmospheric stabilization framework (ASF). *Technol. Forecast. Soc. Change* 63:263-287.
- Schnitzer SA, Bongers F (2002). The ecology of lianas and their role in forests. *Trends Ecol. Evol.* 17:223-230.
- Soberón J, Peterson AT (2004). Biodiversity informatics: Managing and applying primary biodiversity data. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 359:689-698.
- Stockwell DRB (1999). The GARP modelling system: problems and solutions to automated spatial prediction. *Int. J. Geogr. Inf. Sci.* 13: 143-158.
- Thorn JS, Nijman V, Smith D, Nekaris KAI (2009). Ecological niche modelling as a technique for assessing threats and setting conservation priorities For Asian slowlorises (Primates: Nycticebus). *Divers. Distrib.* 15:289-298.
- Thuiller W (2004). Patterns and uncertainties of species range shifts under climate change. *Glob. Change Biol.* 10:2020-2027.
- Thuiller W, Lavorel S, Araújo MB, Sykes MT, Prentice IC (2005). Climate change threats to plant diversity in Europe. *Proc. Nat. Acad. Sci. USA* 102:8245-8250.
- Vié JC, Hilton-Taylor C, Pollock C, Ragle J, Smart J, Stuart SN, Tong R (2008). The IUCN Red List: a key conservation tool. In: Vié, J.C., Hilton-Taylor, C. and S.N. Stuart (Eds). *The 2008 Review of the IUCN Red List of Threatened Species*. IUCN Gland, Switzerland 16p.
- Wiens J, Graham C, Moen D, Smith S, Reeder T (2006). Evolutionary and ecological causes of the latitudinal diversity gradient in hylid frogs: tree frog trees unearth the roots of high tropical diversity. *Am. Nat.* 168:579-596.
- Williams P, Hannah L, Andelman S, Midgley GF, Araújo M, Hughes G, Manne L, Martinez-Meyer E, Pearson R (2005). Planning for climate change: identifying minimum dispersal corridors for the Cape Proteaceae. *Conserv. Biol.* 19:1063-1074.
- Williams SE, Bolitho EE, Fox S (2003). Climate change in Australian tropical rainforests: an impending environmental catastrophe. *Proc. R. Soc. Lond. B Biol. Sci.* 270:1887-1892.
- Wisz MS, Hijmans RJ, Li J, Peterson AT, Graham CH, Guisan A, NCEAS Predicting Species Distributions Working Group (2008). Effects of sample size on the performance of species distribution models. *Divers. Distrib.* 14:763-773.

