



Journal of  
**Medical Laboratory and  
Diagnosis**

Volume 7 Number 6 December 2016  
ISSN 2141-2618



*Academic  
Journals*

## ABOUT JMLD

**Journal of Medical Laboratory and Diagnosis (JMLD)** is published monthly (one volume per year) by Academic Journals.

**Journal of Medical Laboratory and Diagnosis (JMLD)** provides rapid publication (monthly) of articles in all areas of the subject such as Parasitism, Helminthology, Cloning vector, retroviral integration, Genetic markers etc.

### Contact Us

Editorial Office: [jml@academicjournals.org](mailto:jml@academicjournals.org)

Help Desk: [helpdesk@academicjournals.org](mailto:helpdesk@academicjournals.org)

Website: <http://www.academicjournals.org/journal/JMLD>

Submit manuscript online <http://ms.academicjournals.me/>

## Editors

### **Dr. Ratna Chakrabarti**

*Department of Molecular Biology and Microbiology,  
University of Central Florida,  
Biomolecular Research Annex,  
12722 Research Parkway,  
Orlando,  
USA.*

### **Dr. Rajni Kant**

*Scientist D (ADG),  
(P&I Division) Indian Council of Medical Research  
Post Box 4911, Ansari Nagar,  
New Delhi-110029  
India.*

### **Dr. Ramasamy Harikrishnan**

*Faculty of Marine Science, College of Ocean  
Sciences  
Jeju National University  
Jeju city, Jeju 690 756  
South Korea.*

### **Dr. Rokkam Madhavi**

*Andhra University  
Visakhapatnam - 530003  
Andhra Pradesh  
India.*

### **Dr. Mukabana Wolfgang Richard**

*School of Biological Sciences  
University of Nairobi  
P.O. Box 30197 - 00100 GPO  
Nairobi,  
Kenya.*

### **Dr. Lachhman Das Singla**

*College of Veterinary Science  
Guru Angad Dev Veterinary and Animal Sciences  
University  
Ludhiana-141004  
Punjab  
India.*

## Editorial Board

**Dr. Imna Issa Malele**

*Tsetse & Trypanosomiasis Research Institute  
Tanzania.*

**Dr. Mausumi Bharadwaj**

*Institute of Cytology & Preventive Oncology,  
(Indian Council of Medical Research)  
I-7, Sector - 39  
Post Box No. 544  
Noida - 201 301  
India.*

**Dr. James Culvin Morris**

*Clemson University  
214 Biosystems Research Complex  
Clemson SC 29634  
USA.*

# Journal of Medical Laboratory and Diagnosis

Table of Content: Volume 7 Number 6 December 2016

## ARTICLE

- Assessment of antimicrobial susceptibility testing techniques in some laboratories in Cameroon** 36  
Tchoula Mamiafo Corinne, Gonsu Kanga Hortense , Toukam Michel, Emilia Enjema Lyonga-Mbamyah, Andreumont Antoine.

*Full Length Research Paper*

## Assessment of antimicrobial susceptibility testing techniques in some laboratories in Cameroon

Tchoula Mamiafo Corinne<sup>1\*</sup>, Gonsu Kamga Hortense<sup>2,3</sup>, Toukam Michel<sup>2</sup>, Emilia Enjema Lyonga-Mbamyah<sup>2</sup>, Andreumont Antoine<sup>4</sup>.

<sup>1</sup>Department of Microbiology, School of Health Sciences, Catholic University of Central Africa, Yaoundé, Cameroon.

<sup>2</sup>Department of Microbiology, Haematology, Parasitology and Infectious Diseases, Faculty of Medicine and Biomedical Sciences, University of Yaounde I, Yaounde, Cameroon.

<sup>3</sup>Bacteriology Laboratory, Yaoundé University Teaching Hospital, Yaoundé, Cameroon.

<sup>4</sup>Department of Microbiology, Paris-Diderot Faculty of medicine, Paris, France.

Received 25 September, 2016; Accepted 24 November, 2016

The increasing bacterial resistance to antimicrobial agents has rendered susceptibility testing an indispensable tool for appropriate antibiotic selection. This study is aimed at evaluating the technical methods of antibiogram in some medical laboratories in Cameroon. A descriptive cross-sectional study was carried out. The data collection was done with two material, the questionnaire and the observation sheet. We enrolled 13 laboratories through a non-probabilistic technique (Quota sampling). Quality control of media and antibiotic discs as well as their conservation did not comply with the standards. Over 76.9% of laboratories did not have the Mac Farland standard range. One hundred percent of the laboratories used the 90 mm diameter petri dishes. Five to fourteen discs were deposited with a mean of 08 discs per Petri dish. The reading of inhibition zones was done by visual estimation in 56.8% of the laboratories. Only 38.5% of the laboratories had reference strains. The interpretation of the susceptibility testing's crude results were not made by 92.3% of the laboratories. Most laboratories do not have a standard of antimicrobial susceptibility testing. The importance of the strict execution of susceptibility testing procedure should be integrated by the technicians.

**Key words:** Antimicrobial susceptibility testing, laboratories, standards.

### INTRODUCTION

A limited number of methods for antimicrobial susceptibility testing (AST) of medically important microorganisms have survived the maturation of modern diagnostic clinical microbiology (Van Belkum and Dunne, 2013). Antimicrobial susceptibility testing is a technique used to measure the ability of an antibiotic to inhibit

bacterial growth in vivo. This technique is universal and should not undergo any changes except updates. There are few activities in the clinical microbiology laboratory that utilize more technologist time and laboratory resources than AST. It has been suggested that, at least in terms of direct relevance to the care of patients with

\*Corresponding author E-mail: [tchoulacorinne@yahoo.fr](mailto:tchoulacorinne@yahoo.fr).

infection, AST may be the single most important activity performed in the clinical microbiology laboratory. AST results are often used to dictate specific management for individual patients, summary AST data is used to drive empiric antimicrobial therapy, and, finally, formulary decisions in some cases are made based on AST results from the laboratory. (Doern, 2011). The interpretative reading of the susceptibility testing has many parameters which must be taken into consideration before prescribing an antibiotic. With the growing increase in bacteria resistance to antimicrobial agents, susceptibility testing has become an indispensable tool in the judicious choice of antibiotics (Genne and Hans, 2003). There is natural resistance and acquired resistance to antibiotics, for example *Escherichia coli* is naturally resistant to penicillin while ampicillin resistance can be acquired.

Cameroon has many laboratories of different categories. The Bacteriology Unit is found in both public and private laboratories. Nowadays, antimicrobial susceptibility testing is a common practice in these laboratories, but many of them seem not to use standard methods for achieving the susceptibility testing. This study is aimed at assessing the technical methods of antimicrobial susceptibility testing in some medical laboratories in Cameroon and evaluating the gap between these practices and those of the benchmarks.

## METHODOLOGY

During a period of six months we carried out a cross-sectional study during which we enrolled Public health institutions and Private laboratories by a non-probabilistic technic. After the identification of the target laboratories, we asked for their consents; then during a period of one week per laboratory, we observed every single steps of the realization of the antimicrobial susceptibility testing. Finally we compared our results to those required by the standard.

### Participating laboratories

This cross-sectional survey was carried out at thirteen laboratories. These centers are located in 08 regions of Cameroon, and receive a large number of specimens from patients living in surrounding towns. Public and private medical laboratories which carry out antimicrobial susceptibility testing and agreed to participate were included in the study. Each laboratory was assigned a code for privacy reasons. For quality control purpose, our collection tools were based on the standards currently used. All the procedures used in this study were in reference to the current revision of the Helsinki Declaration.

An authorization to carry out this research was provided by the laboratory Heads. The survey protocol was approved by the Cameroon National Ethics Committee (Reference number: 2014/03/434/L/CNERSH/SP).

### Interview data

At enrollment, standardized data collection forms were filled, including identification of each laboratory, the procedure for acquisition, transport and storage of materials, quality controls and

the qualifications of the laboratory personnel.

### Data collection

After obtaining an authorization from the administration, data were collected through an observation sheet. Information on the techniques used for the antimicrobial susceptibility testing from media preparation to the interpretation of the crude results were also collected and their practices compared to those of the standards.

### Quality control

Our collection tools were based on the standards currently used in other countries and those used in the laboratories in Cameroon. The standards mostly found were those of the « Comité de l'Antibiogramme de la Société Française de Microbiologie », CA-SFM (2013) and of the Clinical and Laboratory Standards Institute (CLSI, 2010).

### Statistical analysis

Data analysis was done using CSPRO 4.1. Quantitative variables were expressed as mean  $\pm$  standard deviation, and categorical variables as percentages. Descriptive analysis tools were used including those of the univariate descriptive statistics.

## RESULTS

### Characteristics of the laboratories

Of the 18 laboratories solicited, 13 agreed to participate, and gave a participation rate of 72.2%. Most of the laboratories (76.9%) were public institutions. The repartition of the laboratories according to the qualification of the laboratory Head showed that 69.2% of the laboratories were headed by clinical biologist and the others by pharmacist biologist, biomedical scientists and laboratory technicians.

### Appreciation of the culture media utilization

The different media used by the laboratories for the susceptibility testing of fastidious bacteria are shown in Table 1. We found out that 100% of the media used by the laboratories were homemade. They rigorously follow the instructions on the manufacturers' manual. One of the laboratories did not autoclave media during preparation but heated on a Bunsen burner for 30 min before the spreading on petri dishes. One laboratory also used Columbia agar instead of Mueller Hinton (MH) for susceptibility testing of non-fastidious bacteria. None of the laboratories used MH-F (Mueller Hinton+5% horse blood) agar.

All the laboratories used 90 mm diameter Petri dishes. Only 2 laboratories used square Petri dishes. The media were on average 4.5 mm thick and they were stored in a

**Table 1.** Media used for fastidious bacteria.

Medium	Number
Columbia agar with fresh human blood (GS) and Columbia agar with cooked human blood and Polyvitex (CHOC+P)	2
Columbia agar with cooked human blood (CHOC)	1
CHOC+P	5
Columbia agar with fresh human blood	1
Columbia agar with fresh sheep blood	1
Mueller Hinton (MH)	2
MH with fresh sheep blood 8%	1
<b>Total</b>	<b>13</b>

**Table 2.** Appreciation of culture media utilization.

Medium utilization	Yes	No	NA
Drying of agar before use	3	10	0
Expiration of dehydrated media	1	12	0
Conservation to protect from light	13	0	0
Conservation in dry area	13	0	0
Compliance of the appearance of media (Color, quantity, horizontal surface)	13	0	0
Adequacy of additives concentration	8	3	2

refrigerator after preparation at an average temperature of 7.8°C. The duration of storage time varied from 02 weeks to 02 months with a mean of 03 weeks. Generally, the preservation of dehydrated MH medium was satisfactory. However, additives such as blood, multivitamin supplement was not always in the prescribed concentrations. The media were not dried before use in 10 laboratories.

We have also evaluated the storage and use of antibiotics discs. The discs were kept at an average temperature of 8.9°C (the ones in use and the stock). Only 2/13 laboratories kept their discs in closed containers with desiccants. In addition, 3/13 laboratories were using expired discs. Three quarters of laboratories (76.9%) did not have a Mc Farland standard. In one case, the Mc Farland was available but not used while in another the Mc Farland was used but not as recommended by the standards. We found only the 0.5 McFarland standard.

### Appreciation of the general procedure

The flooding technique was the most commonly used. The colonies used were pure in most laboratories, however, only one laboratory used the densitometer to standardize their inoculum (Table 3).

Visual estimation of the diameter of the inhibition zone was a common practice. The choice of culture media was appropriate in 84.6% of the laboratories, however 76.92% of the laboratories did not use the recommended media

for fastidious bacteria (Table 4). The law of 15-15-15 meaning it is necessary to use the bacterial inoculum within 15 min of preparation, placing antibiotic discs 15 min after swabbing, and incubating 15 min after the deposition of the discs was not observed in any of the laboratories.

The number of discs placed on the agar varied between 5 and 14, with an average of 08 discs per Petri dish of 90 mm diameter (Table 5). The plates were incubated for 24 h at a temperature of 37 ± 1°C and fastidious bacteria were incubated in a CO<sup>2</sup>-enriched environment using a lit candle in a jar. The reading of the incubated Petri dishes was visual for 11 laboratories (Table 6).

### Interpretation of the results

For all the laboratories, the reading of the inhibition zones were made on the back of the agar but not on a black background as recommended by the standards of the European Committee on Antimicrobial Susceptibility Testing (EUCAST) (Table 7). Over 53.8% of laboratories estimated the diameters of the inhibition zones with the naked eye while 46.2% used a ruler and/or calipers. Interpretative reading was not done by 12 of the laboratories. The standard reference most mentioned by the laboratory staff was the "CA-SFM". Only 30.8% of laboratories used CA-SFM to categorize strains as sensitive (S), resistant (R) or intermediate (I). The other laboratories used the antibiotics manufacturer's manual



**Table 3.** Division of the laboratories with respect to the culturing technique used.

Technics	Number	Percentages (%)
Swabbing	05	38.46
Flooding	06	46.15
Not recommended	02	15.39

**Table 4.** Adequation between the type of media and the type of bacteria.

Gender	Percentages (%)
<i>Enterobacteria</i>	84.6
<i>Pseudomonas</i> spp.	84.6
<i>Stenotrophomonas maltophilia</i>	84.6
<i>Acinetobacter</i> spp.	84.6
<i>Staphylococcus</i> spp.	84.6
<i>Enterococcus</i> spp.	84.6
<i>Streptococcus pneumoniae</i>	53.8
<i>Streptococcus</i> (group A, B, C and G)	53.8
<i>Streptococcus viridans</i> group	53.8
<i>Haemophilus</i> spp.	53.8
<i>Moraxella catarrhalis</i>	53.8
<i>Listeria monocytogenes</i>	53.8

**Table 5.** Number of antibiotic discs per Petri dish.

Diameters of the Petri dishes (mm)	Number of discs		
	Minimum	Maximum	Mean
90	5	14	8.2
150	16	16	16.0

written according to the CLSI standards.

Concerning the results issued, all laboratories mentioned the identified bacteria, the isolates were categorized sensitive, resistant or intermediate according to their reaction to the different antibiotics tested. However, the method used, the manufacturer of media or discs, and the origin of the critical points were not included on the results.

### Quality control of the procedure

The search for resistant phenotypes was not performed by any of the laboratories. Although 92.3% of the laboratories controlled the quality of the media prepared, this quality control was limited to sterility testing. Out of 69.2% of the laboratories did not control the quality of their antibiotic discs. For the other laboratories, the control was generally monthly, trimestrial and even semestrial. In some laboratories, this check was done

when a new packet was opened. Only 38.5% of laboratories used reference strains to assess the overall performance of the test and the strains were those of *Escherichia coli*, *Pseudomonas aeruginosa* and *Staphylococcus aureus*. The strains were stored either on nutrient agar at room temperature and away from light or low-temperature in brain-heart infusion (BHI) with glycerol. Finally, only 15.4% of the laboratories were part of an external program of quality assessment in Bacteriology.

### DISCUSSION

In 84.61% of the laboratory, the antimicrobial susceptibility testing was not done according to the standards nor were they updated. This is contrary to the review of Patel in 2012 who suggests standard reference methods should be used for identification so that the subject bacteria are consistently and correctly identified to the genus

**Table 6.** Duration of the incubation according to the gender of bacteria.

Gender	Duration (hours)		
	Minimum	Maximum	Mean
<i>Enterobacteria</i>	24	26	24,7
<i>Pseudomonas</i> spp.	24	26	24,7
<i>S. maltophilia</i>	24	26	24,7
<i>Acinetobacter</i> spp.	24	26	24,7
<i>Staphylococcus</i> spp.	24	26	24,7
<i>Enterococcus</i> spp.	24	26	24,7
<i>S. pneumoniae</i>	24	26	24,7
<i>Streptococcus</i> (group A, B, C, G)	24	26	24,7
<i>S. viridans</i> group	24	26	24,7
<i>Haemophilus</i> spp.	24	26	24,7
<i>M. catarrhalis</i>	24	26	24,7
<i>L. monocytogenes</i>	24	26	24,7

**Table 7.** Measurement of the inhibition diameters.

Measurement	Number	Percentage (%)
With a ruler	03	23,1
With a caliper	03	23,1
Visual estimation	07	53,8
<b>Total</b>	13	100,0

and/or species level. It could be due to either ignorance of the existence of references, or the difficulty to obtain these controls that need to be paid for.

Of the 13 laboratories that agreed to participate, 69.2% were headed by a clinical biologist. This is similar to the work of Nke et al. (2013), who found in their study in Yaounde, that all laboratory Heads were clinical biologists. This shows that our laboratories are headed by qualified specialists.

All the media used for susceptibility testing were homemade, but the type of media was not suitable with the type of bacteria in 11/13 of the laboratories. This is different from a study suggesting that the composition and preparation of the agar and broth media used should be determined, optimised, and documented in a detailed standard operating procedure (Patel, 2012). This shows that their choice was not based on the reference committee on susceptibility testing guidelines.

The storage temperature of the plates was generally respected, while their conservation, the thickness of the agar and the number of discs per petrie dish were not adequate; It is different from a review proposing that growth and incubation conditions (time, temperature, atmosphere e.g. CO<sub>2</sub>), agar depth, number of concentrations tested per broth and agar dilution should be determined, optimised, and documented in

a detailed standard operating procedure (Patel, 2012). This could be explained by the fact that some of these laboratories do not follow the expert rules proposed by the standards..

Only 03 laboratories had a McFarland standard. In one of them the Mc Farland was available but not used and in the other two the use of this standard was not correct. This does not meet the requirements of the review of Patel in 2012 suggesting that the optimum concentration of the inocula must be determined to obtain accurate susceptibility results. This can be explained by the ignorance of the technicians concerning the standards

Eleven laboratories used the flooding technique for economy reasons. Most often, the users of this method did not bother to dry the agar before placing the discs. This was contrary to the "EUCAST" practice, which advises avoiding sub-inoculated agar. We found out that 16.7% of laboratories did not use any of the recommended techniques, but added a colony directly on the agar using a swab.

The temperature and especially the incubation period was not met. References recommended minimum 16 h and maximum 20 to 24 h (EUCAST, 2015, CA-SFM, 2013). Also, these standards recommend a temperature of 35 ± 1°C. This could be due to the fact that this duration was not actually measured but most of the time

it is estimated.

Interpretive reading was not done by 12 of the laboratories. This was contrary to the work of Leclercq et al. (2013) also Reller et al. (2009) who said the results of a susceptibility test must be interpreted by the laboratory prior to communicating a report to a patient's physician. This could be explained by the fact that the biologists are not in charge of only one bench and therefore have too much work.

In regards to the result sheet there were no mention of the method used, type of media or discs, nor the origin of the critical points. Quality control of media prepared was limited to sterility testing. This does not meet the requirements of the WHO and EUCAST that recommend checking the pH, the concentration of divalent cations, thymine and of thymidine, however it is always recommended to do a fertility test and / or specificity. The notion of reference strains is still not well understood in our laboratories; which explains why internal and external quality controls were not done regularly in ten of the laboratories.

## Conclusion

These results suggest that most laboratories do not have a standard of antimicrobial susceptibility testing, and whenever they have one, they do not strictly comply with the guidelines. Moreover, the importance of the strict and correct execution of susceptibility testing procedure has not been well integrated by the technicians. Interpretative reading of the antibiogram should be learned by the head of laboratories in order to bring out the different phenotypes.

## Conflict of interest

There is no conflict of interest regarding any of the authors.

## REFERENCES

- CA-SFM. (2013). Comité de l'antibiogramme de la société française de microbiologie.
- Clinical Laboratory Standard Institute (CLSI) (2010). Performance standards for Antimicrobial Susceptibility testing Sixteenth informational supplement. 2006; 26(25):M100-S16 Wayne, Pennsylvania, USA.
- Doern GV (2011). Antimicrobial susceptibility testing. *J. Clin. Microbiol.* 49 (9 Supplement):S4-S4.
- EUCAST, The European Committee on Antimicrobial Susceptibility Testing (2015). Breakpoint tables for interpretation of MICs and zone diameters. Version 5.0, 2015. <http://www.eucast.org>
- Genné D, Hans HS (2003). De l'antibiogramme à la prescription d'un antibiotique. *Forum Med Suisse* N° 20.
- Leclercq R, Cantón R, Brown DF, Giske CG, Heisig P, MacGowan AP, Mouton JW, Nordmann P, Rodloff AC, Rossolini GM, Soussy CJ, Steinbakk M, Winstanley TG, Kahlmeter G (2013). EUCAST expert rules in antimicrobial susceptibility testing. *Clin. Microbiol. Infect.* 19(2):141-60.
- Nke A (2013). Evaluation des pratiques pré-analytiques dans quelques laboratoires d'analyses médicales de la ville de Yaoundé. Thèse de spécialisation en biologie clinique. <http://www.hsd-fmsb.org/index.php/hsd/article/view/327>
- Patel RM (2012). The guiding principles on antimicrobial susceptibility testing. *Bull. Pharm. Res.* 2(3):146-53.
- Reller LB, Weinstein M, Jorgensen JH, Ferraro MJ (2009). Antimicrobial susceptibility testing: a review of general principles and contemporary practices. *Clin. Infect. Dis.* 49(11):1749-1755
- van Belkum A, Dunne WM (2013). Next-generation antimicrobial susceptibility testing. *J. Clin. Microbiol.* 51(7):2018-2024.

The background of the entire page is a laboratory scene. A pipette is shown in the upper left, dispensing a yellowish liquid into a glass beaker in the center. The scene is lit with a strong blue light, creating a professional and scientific atmosphere. The text is overlaid on semi-transparent dark blue and light blue rectangular areas.

# Journal of Medical Laboratory and Diagnosis

## *Related Journals Published by Academic Journals*

- *Journal of Diabetes and Endocrinology*
- *Journal of Veterinary Medicine and Animal Health*
- *Research in Pharmaceutical Biotechnology*
- *Journal of Physiology and Pathophysiology*
- *Journal of Infectious Diseases and Immunity*
- *Journal of Public Health and Epidemiology*

**academic**Journals