

The background of the cover is a photograph of a person's hands holding a large, spotted fish, likely a walleye, against a bright blue sky. The fish is held vertically, with its head pointing upwards. In the background, there is a blurred landscape featuring a body of water, a line of tall grasses, and a row of trees.

# International Journal of Fisheries and Aquaculture

Volume 9 Number 4 April 2017

ISSN 2006-9839



*Academic  
Journals*

## ABOUT IJFA

**The International Journal of Fisheries and Aquaculture (IJFA)** (ISSN: 2006-9839) is an open access journal that provides rapid publication (monthly) of articles in all areas of the subject such as algaculture, Mariculture, fishery in terms of ecosystem health, Fisheries acoustics etc.

The Journal welcomes the submission of manuscripts that meet the general criteria of significance and scientific excellence. Papers will be published shortly after acceptance. All articles published in the IJFA are peer-reviewed.

### Contact Us

**Editorial Office:** [ijfa@academicjournals.org](mailto:ijfa@academicjournals.org)

**Help Desk:** [helpdesk@academicjournals.org](mailto:helpdesk@academicjournals.org)

**Website:** <http://www.academicjournals.org/journal/IJFA>

**Submit manuscript online** <http://ms.academicjournals.me/>

## Editors

**Dr. V.S. Chandrasekaran**

*Central Institute of Brackishwater Aquaculture  
Indian Council of Agricultural Research (ICAR)  
Chennai,  
India.*

**Prof. Nihar Rajan Chattopadhyay**

*Department of Aquaculture  
Faculty of Fishery Sciences  
West Bengal University of Animal & Fishery Sciences  
West Bengal,  
India.*

**Dr. Lourdes Jimenez-Badillo**

*Ecology and Fisheries Centre  
Universidad Veracruzana  
Veracruz,  
México.*

**Dr. Kostas Kapiris**

*Institute of Marine Biological  
Resources of H.C.M.R.  
Athens,  
Greece.*

**Dr. Masoud Hedayatifard**

*Department of Fisheries Sciences and Aquaculture  
College of Agriculture and Natural Resources  
Advanced Education Center  
Islamic Azad University  
Ghaemshahr,  
Iran.*

**Dr. Zhang Xiaoshuan**

*China Agricultural University  
Beijing,  
China.*

**Dr Joseph Selvin**

*Marine Bioprospecting Lab  
Dept of Microbiology  
Bharathidasan University  
Tiruchirappalli,  
India.*

## Editorial Board

**Dr. Dada Adekunle Ayokanmi**  
*Department of Fisheries and Aquaculture  
Federal University of Technology  
Akure,  
Nigeria.*

**Dr. Ramasamy Harikrishnan**  
*Faculty of Marine Science  
College of Ocean Sciences  
Jeju National University  
Jeju City,  
South Korea.*

**Dr. Kawser Ahmed**  
*Lab. of Ecology, Environment and Climate Change  
Department of Fisheries  
University of Dhaka  
Dhaka,  
Bangladesh.*

**Dr. Maxwell Barson**  
*Biological Sciences Department  
University of Zimbabwe  
Harare,  
Zimbabwe.*

**Dr. Christopher Marlowe Arandela Caipang**  
*Faculty of Biosciences and Aquaculture  
Bodø University College  
Bodø,  
Norway.*

**Dr. William S. Davidson**  
*Department of Molecular Biology and Biochemistry  
Simon Fraser University  
Burnaby,  
Canada.*

**Dr. Babak Ghaednia**  
*Iran Shrimp Research Center (ISRC)  
Bushehr,  
Iran.*

**Prof. Ratha Braja Kishore**  
*Department of Zoology  
Biochemical Adaptation Laboratory  
Banaras Hindu University  
Varanasi,  
India.*

**Prof B. Sharma**  
*Department of Biochemistry  
Centre for Biotechnology  
University of Allahabad  
Allahabad,  
India.*

**Dr. Sebastián Villasante**  
*Fisheries Economics and Natural Resources Research Unit  
University of Santiago de Compostela  
A Coruña,  
Spain.*

**Dr. Mohamed Hamed Yassien**  
*National Institute of Oceanography and Fisheries  
Suez,  
Egypt.*

**Dr. Riaz Ahmad**  
*Department of Zoology  
Aligarh Muslim University  
Aligarh,  
India.*

**Dr. SM Nurul Amin**  
*Lab. Marine Science & Aquaculture  
Institute of Bioscience  
Universiti Putra Malaysia  
Selangor,  
Malaysia.*

**Dr. Melcher Andreas**  
*University of Natural Resources and Applied Life Sciences  
Vienna,  
Austria.*

**Dr. Ayotunde Ezekiel Olatunji**  
*Department of Fisheries and Aquatic Science  
Cross River University of Technology  
Calabar,  
Nigeria.*

**Dr. Ramachandra Bhatta**  
*Animal and Fisheries Sciences University  
College of Fisheries  
Kankanady,  
India.*

**Dr. Subha Ganguly**  
*Department of Fish Processing Technology  
Faculty of Fishery Sciences  
West Bengal University of Animal and Fishery Sciences  
Chakgaria,  
India.*

**Dr. Yusuf Bozkurt**  
*Iskenderun Technical University  
Faculty of Marine Sciences and Technology  
Iskenderun Hatay,  
Turkey.*

**Prof. Rina Chakrabarti**  
*Aqua Research Lab  
Department of Zoology  
University of Delhi  
Delhi,  
India.*

**Prof. Nihar Rajan Chattopadhyay**  
*Faculty of Fishery Sciences*  
*University of Animal and Fishery Sciences*  
*Chakgaria,*  
*India.*

**Dr. Sullip Kumar Majhi**  
*Division of Fisheries*  
*Indian Council of Agricultural Research Complex for Neh*  
*Region*  
*Meghalaya,*  
*India.*

**Dr. Hooman Rahmati-Holasoo**  
*Department of Aquatic Animal Health*  
*Faculty of Veterinary Medicine*  
*University of Tehran*  
*Tehran,*  
*Iran.*

**Dr. Liberty Napilan-Espectato**  
*Institute of Fisheries Policy and Development Studies*  
*College of Fisheries and Ocean Sciences*  
*Iloilo,*  
*Philippines.*

**Dr. Zeinab Abdel-Baki El-Greisy**  
*National Institute of Oceanography and Fisheries (NIOF),*  
*KayetBey,*  
*Egypt.*

**Dr. Thanitha Thapanand-Chaidee**  
*Department of Fishery Biology*  
*Faculty of Fisheries*  
*Kasetsart University*  
*Chatujak,*  
*Thailand.*

**Dr. Gabriel Ugwemorubong Ujagwung**  
*Dept. of Fisheries and Aquatic Environment*  
*Faculty of Agriculture*  
*Rivers State University of Science and Technology*  
*Port Harcourt,*  
*Nigeria.*

**Dr. Margherita Gloria**  
*School of Agriculture*  
*Food Science and Veterinary Medicine*  
*University College Dublin*  
*Belfield,*  
*Ireland.*

**Dr. Derya Guroy**  
*Canakkale Onsekiz Mart University*  
*Faculty of Fisheries*  
*Aquaculture Department*  
*Canakkale,*  
*Turkey.*

**Dr. Daniela Giannetto**  
*Department of Biology*  
*Faculty of Sciences*  
*Muğla Sıtkı Koçman University*  
*Turkey.*

**Dr. Terngu Akaahan**  
*Biological Sciences Department*  
*Federal University of Agriculture Makurdi*  
*Nigeria.*

**ARTICLE**

- Dietary effect of varying linseed oil compositions on growth response, survival and polyunsaturated fatty acid levels in Tilapia (*Oreochromis niloticus*)** 31  
Omolo K. M., Onyango A., Magoma G., Munguti J. and Ogila K.

## Full Length Research Paper

# Dietary effect of varying linseed oil compositions on growth response, survival and polyunsaturated fatty acid levels in Tilapia (*Oreochromis niloticus*)

Omolo K. M.<sup>1\*</sup>, Onyango A.<sup>2</sup>, Magoma G.<sup>1,5</sup>, Munguti J.<sup>3</sup> and Ogila K.<sup>4</sup>

<sup>1</sup>Department of Biochemistry, Jomo Kenyatta University of Agriculture and Technology, P. O. Box 62000-00200 Kenya.

<sup>2</sup>Department of Food Science and Technology, Jomo Kenyatta University of Agriculture and Technology, P. O. Box 62000-00200 Kenya.

<sup>3</sup>Kenya Marine and Fisheries Research Institute, National Aquaculture Research, Development and Training Center, Sagana. Kenya.

<sup>4</sup>Department of Zoology, Jomo Kenyatta University of Agriculture and Technology, P. O. Box 62000-00200 Kenya.

<sup>5</sup>Pan-African University, Institute for Basic Sciences, Technology and Innovation, P. O. Box 62000 00200 Kenya.

Received 28 February, 2017; Accepted 18 April, 2017

The present study evaluated the impact of varying dietary linseed oil composition on growth, survival and tissue polyunsaturated fatty acids (PUFA) profiles in Nile tilapia (*Oreochromis niloticus*). Five iso-nitrogenous diets with varying linseed and sunflower oil concentrations were formulated and fish fed twice daily for 3 months. Commercial diet was used as a control in triplicate tanks set for each diet treatment. Growth parameters were measured from changes in body weight and length. A 75:25 ratio of sunflower oil to linseed oil gave a better survival and specific growth rate than 100% linseed oil or 100% sunflower oil. Tissue PUFA composition were determined using gas chromatography. High dietary linseed oil composition (100%) resulted into significantly high ( $P < 0.05$ ) total n3 fatty acids (9.9-25%) and DHA (1.8-7.9%) in muscles whereas liver n3 fatty acids and DHA composition ranged between 9.3-25.5 and 0.7-2.6%, respectively. Muscle and liver n3/n6 ratio ranged between 0.7-2.2 and 0.7-2.6 while tissue arachidonic acid (ARA 20.4 n6) content ranged between 2.6-3.5% in muscles and 3.4-4.5% in the liver. ARA 20.4 n6 values were low relative to the dietary precursor, linoleic acid, LA, 18.2 n6. Fatty acid deposition in the tissues increased with the feeding period with the third feeding month recording significantly higher DHA, total n3 and n3/n6 ratio. Based on the result, dietary linseed oil > 50% reduced growth and survival rate in tilapia, however, it increased tissue accumulation of essential fatty acids which also increased with the length of feeding period.

**Key words:** Growth, Tilapia, n3 fatty acids, linseed oil, polyunsaturated fatty acids (PUFA).

## INTRODUCTION

Agricultural output originating from fisheries and aquaculture must increase by over 60% to feed the

world in 2050 considering the perceived benefits of fish oil (Leaf and Weber, 1988; Bonaa et al., 1990) and the

\*Corresponding author. E-mail: komolo@jkuat.ac.ke. Tel: +254723623791

Author(s) agree that this article remain permanently open access under the terms of the [Creative Commons Attribution License 4.0 International License](https://creativecommons.org/licenses/by/4.0/)



alarming number of people, mostly in developing countries, suffering from hunger and poverty (FAO, 2014). With the proven benefits of omega 3 highly unsaturated fatty acids (n3-HUFA) (Kris-Etherton et al., 2000; Gebauer et al., 2006; Kusunto et al., 2007), numerous governmental and non-governmental organizations across the world currently advise increased fish intake as a means of improving the health of their citizens (Munguti et al., 2014).

To meet the demand, quality and affordable feeds are integral component of sustainable intensive and semi-intensive aquaculture systems (Munguti et al., 2014). Recent studies have reported the existence of a correlation between the nutritional value of fish and the dietary composition of feeds (Župan et al., 2016; Ljubojević et al., 2015; Ljubojević et al., 2013). Fish oils (FO) are regarded as good lipid sources of aqua-feed formulations (National Research Council, NRC, 2011), however, the global fish oil production has reached a plateau and is not expected to raise much beyond the current level of production (Ng, 2002). Also, it is predicted that within a decade or so, fish oil production may not be sufficient to meet the demand of aquaculture (FAO, 2007). Therefore, the introduction of alternative lipid sources is necessary to enable sustainable aquaculture development (Jordal et al., 2007; Bouraoui et al., 2011).

Reducing the utilization of FO in aqua-feeds formulations while ensuring that appropriate n-3 long chain polyunsaturated fatty acids (n3 LCPUFA) proportion are available in the final product is a challenge (Turchini et al., 2009). However, vegetable oils which are rich in C<sub>18</sub> polyunsaturated fatty acids have been considered the most sustainable alternatives to fish oil (Montero et al., 2003; Lee, 2001). In addition, studies have reported considerable success in partial or total replacement of FO with VO in many fish species (Ng, 2005; Turchini et al., 2009, 2010). Moreover, recent studies have also evaluated the effects of replacing high-quality FO with VO sources on growth performance, fatty acid profile or health parameters of fishes, such as gilthead seabream (*Sparus aurata*) (Menoyo et al., 2004) and European sea bass (*Dicentrarchus labrax L.*) (Mourente et al., 2007). However, the extent of the effect is determined by the vegetable oil source and level of dietary replacement. Linseed oil (LO) is distinguished by the highest content of α-linolenic acid (18:3n-3, ALA) compared to other vegetable oils (Popa et al., 2012). Alpha -linolenic acid is the metabolic precursor of n-3 long-chain polyunsaturated fatty acids (LC-PUFA) such as eicosapentaenoic acid (20:5n-3, EPA) and docosahexaenoic acid (22:6n-3, DHA) (Brenna, 2002).

Linseed oil has replaced dietary FO without affecting growth performance in various fish species (Menoyo et al., 2005; Francis et al., 2006). However, replacement of FO with LO has also led to decreased concentration of n-3 highly unsaturated fatty acids in fish flesh especially when higher dietary linseed oil were fed (Menoyo et al.,

2005; Turchini et al., 2011). On the contrary, it has been shown that when substituted with sunflower oil, dietary linseed oil can increase the EPA/DHA content of Nile tilapia fillets (Justi et al., 2003; Visentainer et al., 2005). The effect of dietary linseed oil on immunity thus survival of various fish species has also been reported (Montero et al., 2010; Kiron et al., 2011).

In the recent past, fish production in Kenya has risen steadily from 1,012 metric tons produced in 2003 to the present production of 21,487 metric tons courtesy of Kenya government fish farming program (Charo-Karissa et al., 2010). However, the major challenges to Kenyan aquaculture sector are the unavailability of quality, efficient and inexpensive farm-made feeds (Munguti et al., 2014). The current research explored the possible use of linseed oil as a dietary replacement of FO in farm-made tilapia feeds and the impact on survival, growth parameters and fillet PUFA profile.

## MATERIALS AND METHODS

### Study site, experimental set up and sampling

Proximate composition analysis of experimental diets and fatty acid composition of tilapia muscles and liver were done in Food Biochemistry laboratory, Food Science department-Jomo Kenyatta University of Agriculture and Technology (JKUAT). Tilapias which were 4 weeks old were set up in a 1000-liters experimental tank for a period of 3 months. The fingerlings were obtained from Kenya Marine and Fisheries Research Institute, KEMFRI, Sagana fish hatcheries. Three tanks containing 40 tilapia were set for each experimental diet. A separate tank was used as a holding tank for two weeks prior to the feeding experiments for fish acclimatization. Feeding was done twice/day at 9.00 am and 4.00 pm.

Continuous water circulation was maintained using a water pump with water conditions and quality checked and maintained regularly for optimum water quality. After the feeding period, 5 tilapia were randomly sampled after 24 h fasting period, anaesthetized and dissected for liver and muscles.

### Diet ingredients

Experimental diets used in this study includes freshwater shrimps (*Caridina nilotica*) as the main protein source, rice bran, wheat flour, popcorn maize flour, vegetable oil blends and vitamin and mineral premixes. Fresh water shrimps were obtained from Wich lum beach along Lake Victoria in Siaya County, Kenya.

Vegetable oil blend comprised of linseed oil, olive oil and sunflower oil. Popcorn maize, wheat flour, olive oil and sunflower oil were obtained from local supermarkets vitamin and mineral premixes obtained from Tam feeds and linseed oil was extracted from linseeds using extruder machine at the Biomechanical engineering workshop, Jomo Kenyatta University of Agriculture and Technology (JKUAT).

### Diet formulation

Powdered experimental ingredients were weighed and pre-mixed prior to the addition of vegetable oil blends. They were then dry mixed thoroughly for 2 min in a bench top food mixer before addition of distilled water and vegetable oil blends and mixing



**Table 1.** Macronutrient content of experimental diets.

Macronutrients (g/100 g)	Washout	Diet 1	Diet 2	Diet 3	Diet 4	Diet 5
Fish shrimps	51.7	51.7	51.7	51.7	51.7	51.7
Rice bran	10.3	10.3	10.3	10.3	10.3	10.3
Wheat flour	10	10	10	10	10	10
Popcorn flour	10	10	10	10	10	10
Yeast	1	1	1	1	1	1
Vegetable oil	14	14	14	14	14	14
Oil proportions (% w/w in 14% oil)						
Sunflower <sup>1</sup>	0	0	25	50	75	100
Olive oil <sup>1</sup>	100	0	0	0	0	0
Linseed oil <sup>2</sup>	0	100	75	50	25	0
Premixes <sup>3</sup>	2	2	2	2	2	2

<sup>1</sup>Obtained from local supermarket; <sup>2</sup>Extracted from linseeds using oil extruder machine, BEED Department, JKUAT-Kenya; <sup>3</sup>Obtained from Tam feeds, Nairobi, Kenya.

continued for further 10 min. All the experimental diets were made on grade 12 meat mincer as an extruder fitted with a die plate with 2 mm diameter holes. The soft feed dough was then cold extruded into the 2 mm die-size strand, pelleted and dried at ambient temperature for 3 h. The feeds were then placed on a sieve and oven dried at 40°C for approximately 24 h until the moisture content was 10% (w/w). The dried feeds were then broken into 2 to 3 mm pellets, sealed in plastic bags and stored at -20°C until commencement of feeding trials.

All equipment used for making up feeds were washed and dried before the next diet was produced to avoid cross-contamination. Pearson's square method was used in feed formulation to determine the proper dietary proportions of high and low protein feed stuffs to add to a feed to meet the dietary requirements (Table 1).

### Proximate analysis

Moisture content, crude protein, crude fat, crude fiber and ash for diet ingredients and experimental diets were determined according to AOAC methods specification 950.46 (AOAC, 1995) (Table 1). In brief, moisture content was determined by weighing 2 g of sample into a moisture dish and transferred to an oven previously heated to temperatures of 105°C and drying done for 1 h.

The final weight of the sample was taken after the drying period and cooling in a desiccator. The flour residue was then reported as total solids and loss in weight as moisture by following formula (AOAC, 1995, method 925. 10):

$$\text{Moisture (\%)} = \frac{W_1 - W_2}{W_1} \times 100$$

$W_1$  = Weight of sample before drying and  $W_2$  = Weight of sample after drying.

Crude protein was determined by Semi-Micro Kjeldahl method where about 1 g of sample was weighed into a digestion flask together with a catalyst composed of 5 g of  $K_2SO_4$  and 0.5 g of  $CuSO_4$  and 15 ml of concentrated  $H_2SO_4$ . The mixture was heated in a fume hood till the digest color turned blue signifying the end of the digestion process. The digest was cooled, transferred to a 100 ml volumetric flask and topped up to the mark with distilled water. A blank digestion with the catalysts and acid was also made. Ten milliliter of diluted digest was transferred into a distilling flask and

washed with about 2 ml distilled water. 15 ml of 40% NaOH was added and this was also washed with about 2 ml distilled water. Distillation was done to a volume of about 60 ml distillate.

The distillate was titrated using 0.02 N-HCL to an orange colour of the mixed indicator which signified the end point (AOAC, 1995; Method 20.87-32.1.22). Calculations were done using the following formula:

$$\text{Nitrogen\%} = (V_1 - V_2) \times N \times f \times 0.014 \times \frac{100}{V} \times \frac{100}{S}$$

$V_1$  = Titer for the sample (ml);  $V_2$  = Titer for blank (ml);  $N$  = Normality of standard HCL solution (0.002);  $F$  = Factor of standard HCL solution;  $V$  = Volume of diluted digest taken for distillation (10ml);  $S$  = Weight of sample taken (g).

Crude protein % = Nitrogen × protein factor

Crude fat was determined through soxhlet extraction method which gives intermittent extraction of oil with excess of fresh organic solvent used. About 5 g of samples were weighed into extraction thimbles and the initial weights of the extraction flasks taken. Fat extraction was done using petroleum ether in soxhlet extraction apparatus for 8 h. The extraction solvents were evaporated and the extracted fat dried in an oven for about 15 min before the final weights of the flasks with extracted fat were taken (AOAC, 1995; Method 920.85-32.1.13). Calculations were done using the following formula:

$$\text{Crude fat (\%)} = \frac{W_1 - W_2}{W_1} \times 100$$

$W_1$  = Weight of sample before extraction;  $W_2$  = Weight of sample after extraction.

Crude fiber was determined by approximately weighing 2 g of the sample was weighed into a 500 ml conical flask. About 200 ml of boiling 1.25%  $H_2SO_4$  was added and boiling done for 30 min under reflux condenser. Filtration was done under slight vacuum with Pyrex glass filter and the residue washed to completely remove the acid with boiling water. Approximately 200 ml of boiling 1.25% NaOH was added to the washed residue and boiling done under reflux for another 30 min. Filtration was done using the same glass filter previously used with the acid. The residue was rinsed with boiling water followed by 1% HCL and again washed with boiling

water to rinse the acid from the residue.

The residue was washed twice with alcohol and thrice with ether. It was then dried in an oven at 105°C in a porcelain dish to a constant weight ( $W_1$ ). Incineration was done in a muffle furnace at 550°C for 3 h, the dish was then cooled in a desiccator and the final weight ( $W_2$ ) taken (AOAC, 1995, Method 920.86-32.1.15). Calculations were done as follows:

$$\text{Crude fiber (\%)} = \frac{W_1 - W_2}{W} \times 100$$

Where,  $W_1$  = Weight of acid and alkali digested sample;  $W_2$  = Weight of incinerated sample after acid and alkali digestion;  $W$  = Weight of sample.

Ash was determined from Sample weights of between 2 and 5 g were weighed in pre-conditioned crucibles. The samples were first charred by flame to eliminate smoking before being incinerated at 550°C in a muffle furnace to the point of white ash. The residues were cooled in desiccators and the weights taken (AOAC, 1995, Method 925.03-32.1.05).

Calculations were done as shown below:

$$\text{Crude ash (\%)} = \frac{\text{Weight of Ash (g)}}{\text{Weight of sample (g)}} \times 100$$

#### Extraction of total lipids and preparation of fatty acid methyl esters

Lipids extraction was done according to the procedure by Bligh and Dyer (1959). Lipids in experimental diets, tilapia liver and muscles were extracted by homogenization of finely ground 0.5 g of samples in chloroform-methanol (2:1, v/v) containing 0.01% butylated hydroxytoluene (BHT) as antioxidant and cold isotonic saline, 0.9% sodium chloride. This was mixed vigorously and allowed to stand for 20 min. The mixture was then centrifuged at 3000 rpm for 10 min and the aqueous layer was then separated from organic layer using a micropipette. The bottom layer, chloroform, was then transferred to 100 ml reflux flask, quick fit, and evaporated to dryness under vacuum evaporator. Fatty acid methyl esters (FAME) were prepared from vegetable oils and total lipid extracted by acid-catalyzed trans-esterification and addition of 5 ml of 1%  $H_2SO_4$  (v/v) in methanol at 70°C, for 3 h. FAME were then extracted into 750 ml of distilled water and 10 ml of hexane, dehydrated using anhydrous sodium sulphate,  $Na_2SO_4$  and concentrated to 0.5 ml under vacuum evaporator. The concentrated FAME were stored in GC vials for later GC analysis

#### GC-analysis

FAME were quantified using gas chromatography with on-column injection, equipped with a fused silica capillary column (SUPELCO Column Omegawax™530, 30 m × 0.5 mm × 0.5 μm) with nitrogen as carrier gas and temperature programming from 170 to 220°C for 18 min<sup>-1</sup> and final time of 47 min totaling to a run time of 75 min. Injection and detection temperatures were 240 and 260°C respectively. The programmer rate for both GC and decoder were set at 5 min<sup>-1</sup> with an attenuation of 3.

All the GC analyses were done under same conditions. Individual methyl esters in the sample were identified by comparison with known FAME standards obtained from Kobian chemicals.

#### Growth performance of tilapia

Growth performance was measure through the following ways:

Initial body weight (IBW) (g): Measured before transferring to experimental tanks

After washout body weight (g) (WBW): Measured at the 3-weeks washout period

Final Body weight (g) (FBW): Measured at the end of the experimental feeding period

Length (cm): Measured from the tip of the snout to the end of the tail

Weight gain (%): Calculated as [(FBW-WBW)/WBW] × 100

Specific growth rate (SGR, %): Calculated based on [(log of FBW - log of WBW)/ feeding days] × 100

Survival (%): Calculated as [(initial number of fish before feeding - dead fish number during feeding/initial number of fish before feeding) × 100]

#### Statistical analyses

Performed using GenStat version 41.0. Significance in polyunsaturated fatty acid compositions of liver and muscles of tilapia fed different experimental diets was determined by analysis of variance (One-way ANOVA). When significant differences were discerned, treatment means were compared using Duncan's Multiple Range Test (DMRT) (Duncan DB. 1955). Values throughout the text are expressed as means ± standard error. In all the analysis, treatment significance was accepted at  $P < 0.05$ .

## RESULTS

### Growth and survival

Commercial diet (control) and diet 4 had significantly high ( $P < 0.05$ ) final body (147, 97) and final length (21.44, 17.4) respectively (Table 5). Weight gain (%) and specific growth rate were also significantly higher ( $P < 0.05$ ) in both diet 4 (1689%, 1.3) and commercial diet (2625%, 1.5) respectively (Table 5). Significantly high survival rate was observed in commercial diet (98.3%) with relatively low survival rate observed in diet 1 (Table 5). Survival and specific growth rate significantly increased with reduction of dietary linseed oil. Body weight and length before and after washout were 2.6/5.4 and 1.6/ 2.7 respectively (Table 4)

### Proximate and fatty acid composition of diet

There was no significant difference ( $P < 0.05$ ) in the experimental dietary proximate composition of protein, crude fats, ash and fiber except for commercial diet which had significantly lower crude protein and fiber (Table 2). Fatty acid composition of diets varied significantly ( $p < 0.05$ ) among the diets with significantly high linolenic acid (C18:3), oleic acid (C18:1) and linoleic acid (C18:2) composition in diet1, washout diet and diet 5 respectively (Table 3). The dietary fatty acid composition variation was attributed to the vegetable oils supplemented at different proportions in the diets (Table 1) and difference in fatty acid composition of linseed oil, olive oil and sunflower oil (Figure 1). Polyunsaturated fatty acid compositions of tilapia muscle and liver following 3-

**Table 2.** Proximate composition of feed ingredients and diets.

Feed ingredients	Proximate compositions				
	Protein	Crude fat	Ash	Moisture	Fibre
Diet 1	45.9±0.21 <sup>d</sup>	23.2±0.09 <sup>ab</sup>	13.6±1.2 <sup>ab</sup>	9.1±0.16 <sup>a</sup>	3.3 ±0.23 <sup>c</sup>
Diet 2	45.4±0.9 <sup>d</sup>	25.0±6.46 <sup>ab</sup>	13.2±1.02 <sup>ab</sup>	8.9±0.16 <sup>a</sup>	2.9±0.03b <sup>c</sup>
Diet 3	44.6±0.4 <sup>d</sup>	25.1±1.5 <sup>ab</sup>	11.7±0.36 <sup>ab</sup>	10.1±0.22 <sup>a</sup>	3.3±0.25 <sup>c</sup>
Diet 4	45.5±0.8 <sup>d</sup>	25.0 ±3.3 <sup>ab</sup>	12.7±0.73 <sup>ab</sup>	10.3±0.35 <sup>a</sup>	3.3±0.1 <sup>c</sup>
Diet 5	44.2±0.4 <sup>d</sup>	24.9±1.2 <sup>ab</sup>	12.8±1.05 <sup>ab</sup>	9.8±0.32 <sup>a</sup>	3.4±0.2 <sup>c</sup>
Washout	45.6±0.6 <sup>d</sup>	25.4±2.3 <sup>ab</sup>	14.9±0.33 <sup>b</sup>	9.4±0.16 <sup>a</sup>	3.3±0.2 <sup>c</sup>
Commercial <sup>1</sup>	36.0±3.97 <sup>c</sup>	16.3±0.4 <sup>ab</sup>	13.5±0.06 <sup>ab</sup>	7.7±0.12 <sup>a</sup>	2.2±0.13 <sup>b</sup>
Caridina <sup>2</sup>	61.1±0.14 <sup>e</sup>	18.2±2.67 <sup>ab</sup>	15.6±3.71 <sup>b</sup>	9.7±0.11 <sup>a</sup>	0.6±0.26 <sup>a</sup>
Popcorn flour <sup>3</sup>	10.8±0.21 <sup>b</sup>	7.5±0.45 <sup>a</sup>	6.4±4.2 <sup>a</sup>	9.3±0.95 <sup>a</sup>	3.3±0.18 <sup>c</sup>
Wheat flour <sup>3</sup>	10.9±0.31 <sup>b</sup>	7.9±0.98 <sup>a</sup>	8.7±2.03 <sup>ab</sup>	8.7±1.17 <sup>a</sup>	1.9±0.04 <sup>b</sup>
Rice Bran <sup>4</sup>	8.7±0.35 <sup>a</sup>	9.9±5.96 <sup>a</sup>	16.2±2.1 <sup>b</sup>	8.7±0.69 <sup>a</sup>	6.1±0.17 <sup>d</sup>

<sup>1</sup>Skretting, Fontaine-Les-Vervins-France obtained from Jambo fish farm, Kiambu, Kenya; <sup>2</sup>Obtained from Wich lum beach of Lake Victoria, Kenya; <sup>3</sup>Obtained from local supermarket; <sup>4</sup>Obtained from Mwea rice mills.

**Table 3.** Fatty acid composition (%) of diet.

Acid	Diet 1	Diet 2	Diet 3	Diet 4	Diet 5	Washout	Commercial
C10:0	1.4 <sup>a</sup> ±0.21	1.8 <sup>ab</sup> ±0.61	1.8 <sup>ab</sup> ±0.17	2.1 <sup>b</sup> ±0.19	3.2 <sup>c</sup> ±0.23	3.5 <sup>c</sup> ±0.5	1.5 <sup>ab</sup> ±0.37
C12:0	1.9 <sup>ab</sup> ±0.35	2.2 <sup>b</sup> ±0.41	2.3 <sup>b</sup> ±0.27	2.9 <sup>c</sup> ±0.34	3.4 <sup>c</sup> ±0.41	3.3 <sup>c</sup> ±0.22	1.4 <sup>a</sup> ±0.33
C14:0	2.4 <sup>b</sup> ±0.31	3.2 <sup>c</sup> ±0.4	3.7 <sup>c</sup> ±0.24	3.5 <sup>c</sup> ±0.16	2.6 <sup>b</sup> ±0.2	2.2 <sup>ab</sup> ±0.13	1.8 <sup>a</sup> ±0.36
C16:0	11.7 <sup>a</sup> ±0.19	12.4 <sup>b</sup> ±0.17	13.7 <sup>c</sup> ±0.11	14.5 <sup>d</sup> ±0.31	15.5 <sup>e</sup> ±0.45	15.3 <sup>e</sup> ±0.34	12.8 <sup>b</sup> ±0.45
C18:0	3.3 <sup>a</sup> ±0.36	4.4 <sup>b</sup> ±0.23	4.9 <sup>bc</sup> ±0.26	5.1 <sup>cd</sup> ±0.17	5.3 <sup>cd</sup> ±0.18	5.5 <sup>d</sup> ±0.15	3.4 <sup>a</sup> ±0.12
C20:0	3.1 <sup>c</sup> ±0.21	2.9 <sup>c</sup> ±0.21	2.6 <sup>bc</sup> ±0.42	2.3 <sup>ab</sup> ±0.51	2.1a±0.37	1.8 <sup>a</sup> ±0.17	2.6 <sup>bc</sup> ±0.24
<sup>a</sup> ∑SFAs	23.3 <sup>a</sup> ±0.41	26.4 <sup>b</sup> ±0.51	28.5 <sup>c</sup> ±0.23	29.7 <sup>d</sup> ±0.19	31.6 <sup>e</sup> ±0.14	31.1 <sup>e</sup> ±0.1	23.1 <sup>a</sup> ±0.13
C16:1	3.9 <sup>ab</sup> ±0.16	4.2 <sup>abc</sup> ±0.28	4.4 <sup>bcd</sup> ±0.37	4.7 <sup>cd</sup> ±0.26	4.9 <sup>d</sup> ±0.19	4.9 <sup>d</sup> ±0.2	3.7 <sup>a</sup> ±0.18
C18:1	18.1 <sup>a</sup> ±0.13	20.0 <sup>b</sup> ±0.31	20.3 <sup>b</sup> ±0.23	20.4 <sup>b</sup> ±0.26	22.4 <sup>d</sup> ±0.15	29.8 <sup>e</sup> ±2.3	21.6 <sup>c</sup> ±0.15
<sup>b</sup> ∑MUFAS	21.8 <sup>a</sup> ±0.63	24.1 <sup>b</sup> ±0.25	24.6 <sup>bc</sup> ±0.13	25.1 <sup>cd</sup> ±0.38	27.3 <sup>e</sup> ±1.4	34.6 <sup>f</sup> ±2.1	25.2 <sup>d</sup> ±0.13
C18:2	16.3 <sup>a</sup> ±0.17	19.5 <sup>b</sup> ±0.81	22.3 <sup>e</sup> ±0.63	24.7 <sup>f</sup> ±0.34	28.5 <sup>g</sup> ±1.5	20.5 <sup>c</sup> ±0.22	21.4 <sup>d</sup> ±0.41
C18:3	18.7 <sup>f</sup> ±1.1	16.5 <sup>e</sup> ±1.2	14.5 <sup>d</sup> ±0.31	12.8 <sup>c</sup> ±0.51	4.5 <sup>b</sup> ±1.3	3.2 <sup>a</sup> ±0.36	14.5 <sup>d</sup> ±0.24
C20:5	1.2 <sup>a</sup> ±0.18	1.3 <sup>a</sup> ±0.16	1.1 <sup>a</sup> ±0.12	1.1 <sup>a</sup> ±0.14	1.1 <sup>a</sup> ±0.03	1.2 <sup>a</sup> ±0.31	1.5 <sup>a</sup> ±0.18
C22:6	2.4 <sup>a</sup> ±0.21	2.3 <sup>a</sup> ±0.19	2.3 <sup>a</sup> ±0.22	2.1 <sup>a</sup> ±0.17	2.3 <sup>a</sup> ±0.25	2.2 <sup>a</sup> ±0.32	3.21 <sup>b</sup> ±1.56
<sup>c</sup> ∑PUFAS	38.3 <sup>c</sup> ±0.61	39.3 <sup>d</sup> ±0.31	39.9 <sup>e</sup> ±0.21	40.2 <sup>e</sup> ±0.35	36.0 <sup>b</sup> ±0.5	26.8 <sup>a</sup> ±2.3	40.3 <sup>e</sup> ±0.43
<sup>d</sup> ∑n3	22.2 <sup>g</sup> ±0.38	19.9 <sup>f</sup> ±0.7	17.6 <sup>d</sup> ±0.8	15.6 <sup>c</sup> ±0.83	7.6 <sup>b</sup> ±0.73	6.4 <sup>a</sup> ±0.51	19.1 <sup>e</sup> ±0.26
∑n6	16.3 <sup>a</sup> ±0.21	19.5 <sup>b</sup> ±0.43	22.3 <sup>e</sup> ±0.19	24.7 <sup>f</sup> ±0.33	28.5 <sup>g</sup> ±0.27	20.5 <sup>c</sup> ±0.17	21.4 <sup>d</sup> ±0.52
∑n3/n6	1.5 <sup>c</sup> ±0.59	1.1 <sup>bc</sup> ±0.22	0.9 <sup>ab</sup> ±0.28	0.7 <sup>ab</sup> ±0.18	0.4 <sup>a</sup> ±0.05	0.4 <sup>a</sup> ±0.05	1.1 <sup>bc</sup> ±0.17

Values reported are means± standard error (n=3) as determined using Duncan's multiple range test. Means within the same row with different superscripts varied significantly (p<0.05). Fatty acids: C10:0 Capric Acid, C12:0 Lauric acid, C14:0 Myristic acid, C14:1 Myristoleic acid, C16:0 palmitic acid, C16:1 palmitoleic acid, C18:0 Stearic acid, C18:1 Oleic acid C18:2 Linoleic acid, C18:3 α-Linolenic acid, C20:0 Arachidic acid, C20:5 Eicosapentaenoic acid, C22:6 docosahexaenoic acid <sup>a</sup>∑SFAs: Total saturated fatty acids <sup>b</sup>∑MUFAS: Total monounsaturated fatty acids <sup>c</sup>∑PUFAS: Total Polyunsaturated fatty acids. <sup>d</sup>∑n3: Total omega-3 fatty acids ∑n6: Total omega-6 fatty acids.

months feeding period are presented in Tables 6 and 7. In all the experimental diets, there was significant (P<0.05) changes in fatty acid composition in tilapia tissues with feeding period. The tilapia liver and muscles fatty acid composition trends were also related to the dietary fatty acid composition.

Even though there was no particular trend in linoleic

acid composition in the feeding period in all experimental diets, significant increase in the tissues composition of linolenic acid, arachidonic acid, eicosapentaenoic acid (EPA), docosahexaenoic acid (DHA), omega -3 and omega-3/omega-6 ratio was observed (Tables 6 and 7). Tissue composition of linolenic acid increased with increased inclusion of linseed oil in the diet with relatively


**Table 4.** Mean body weight and length before and after washout in tilapia.

Body weight (g)		Body length (cm)	
Before washout	After washout	Before washout	After washout
2.6±0.2	5.4±0.4	1.6±0.1	2.7±0.1

**Table 5.** Different growth parameters for tilapia.

	Diet 1	Diet 2	Diet 3	Diet 4	Diet 5	Washout	Commercial
FBW <sup>1</sup>	46.8±6.3 <sup>a</sup>	55±4.0 <sup>ab</sup>	64.4 <sup>b</sup> ±6.4	97.4±4.9 <sup>c</sup>	69.2±6.8 <sup>b</sup>	66.4±8.8 <sup>b</sup>	147.6±2.9 <sup>d</sup>
FBL <sup>2</sup>	13.5±1.2 <sup>a</sup>	14.9±0.7 <sup>ab</sup>	16.5±0.3 <sup>bc</sup>	17.4±0.3 <sup>c</sup>	16.5±0.4 <sup>bc</sup>	16.5±0.3 <sup>bc</sup>	21.44±0.4 <sup>d</sup>
Weight gain (%)	750 <sup>a</sup>	937 <sup>ab</sup>	1099 <sup>ab</sup>	1689 <sup>c</sup>	1216 <sup>b</sup>	1175 <sup>ab</sup>	2625 <sup>d</sup>
SGR <sup>3</sup> (%)	0.9±0.05 <sup>a</sup>	1.1±0.04 <sup>ab</sup>	1.16±0.01 <sup>b</sup>	1.3±0.02 <sup>c</sup>	1.2±0.06 <sup>bc</sup>	1.2±0.08 <sup>b</sup>	1.5±0.01 <sup>d</sup>
Survival (%)	94 <sup>a</sup>	95.67 <sup>a</sup>	96 <sup>b</sup>	97.67 <sup>c</sup>	97 <sup>c</sup>	97.33 <sup>c</sup>	98.33 <sup>d</sup>

<sup>1</sup>FBW, Final body weight (grams); <sup>2</sup>FBL-Final body length (cm); <sup>3</sup>SGR, Specific growth rate.



**Figure 1.** Compositions (%) of C18 fatty acids in vegetable oils used for experimental diets formulation. \*C18:0; Stearic acid, C18:1; Oleic acid, C18:2; Linoleic acid, C18:3; α-Linolenic acid.

high tissue composition of linolenic acid observed in diet 1. In addition, the accumulation of linolenic acid increased with feeding period in both tilapia tissues with significantly high composition observed in month 3 (Tables 6 and 7). The compositions of DHA, EPA, arachidonic acid,  $\sum n3$ , and  $n3/n6$  also increased with feeding period in both tilapia tissues with significantly high composition observed in diet 1 (Tables 6 and 7).

## DISCUSSION

The possible use of VO as a replacement for dietary FO in fish diets has been a subject of study in the recent

past. The data from these studies indicate that FO can partially or totally be replaced with VO, however, with varying outcomes on growth performance and fatty acid profile for different fish species. Results from this study show that > 50% dietary linseed oil lowered growth parameters and survival ability of tilapia (*Oreochromis niloticus*).

Present finding is consistent with earlier reports that high dietary linseed oil lowered the growth performance of tilapia (Li et al., 2016; Francis et al., 2006). Moreover, earlier reports indicate that high dietary n-3 polyunsaturated fatty acids (PUFAs) depressed growth in hybrid tilapia and tilapia zillii (Huang et al., 1998; Kanazwa et al., 1980). In contrast, replacing FO with

**Table 6.** Monthly changes of muscles polyunsaturated fatty acid composition in tilapia fed diets containing varying composition of linseed, sunflower and olive oil.

Acid	Month	Treatment						
		Diet 1	Diet 2	Diet 3	Diet 4	Diet 5	Washout	Commercial
C18:2	1	12.0 <sup>ab</sup> ±0.14	13.7 <sup>gh</sup> ±0.9	14.4 <sup>gh</sup> ±0.83	15.7 <sup>i</sup> ±0.62	18.8 <sup>j</sup> ±0.63	15.7 <sup>i</sup> ±0.47	16.5 <sup>i</sup> ±0.82
	2	10.2 <sup>a</sup> ±0.82	11.6 <sup>abc</sup> ±0.2	14.6 <sup>h</sup> ±0.48	18.3 <sup>j</sup> ±0.5	19.4 <sup>k</sup> ±0.5	18.7 <sup>j</sup> ±0.4	16.2 <sup>i</sup> ±0.12
	3	11.9 <sup>bcd</sup> ±0.38	12.9 <sup>def</sup> ±0.15	13.2 <sup>efg</sup> ±0.45	14.3 <sup>gh</sup> ±0.5	16.3 <sup>i</sup> ±0.36	14.3 <sup>gh</sup> ±0.31	12.5 <sup>ede</sup> ±0.14
C18:3	1	7.8 <sup>i</sup> ±0.27	4.8 <sup>cde</sup> ±0.73	4.3 <sup>bcd</sup> ±0.19	3.7 <sup>bc</sup> ±0.26	3.3 <sup>b</sup> ±0.15	1.3 <sup>a</sup> ±0.31	5.3 <sup>ef</sup> ±0.41
	2	10.8 <sup>kl</sup> ±0.23	9.3 <sup>ij</sup> ±0.45	8.9 <sup>i</sup> ±0.61	7.4 <sup>gh</sup> ±0.21	5.5 <sup>ef</sup> ±0.31	4.8 <sup>cde</sup> ±0.13	8.5 <sup>hi</sup> ±0.63
	3	11.9 <sup>kl</sup> ±0.1	10.3 <sup>l</sup> ±0.63	10.2 <sup>jk</sup> ±0.42	8.4 <sup>hi</sup> ±0.71	6.2 <sup>fg</sup> ±0.37	5.3 <sup>ef</sup> ±0.34	11.7 <sup>i</sup> ±0.06
C20:4	1	1.6 <sup>a</sup> ±0.11	1.9 <sup>abc</sup> ±0.37	2.3 <sup>bcd</sup> ±0.24	2.5 <sup>cdef</sup> ±0.34	2.7 <sup>efgh</sup> ±0.31	2.8 <sup>efgh</sup> ±0.21	1.8 <sup>ab</sup> ±0.13
	2	2.0 <sup>abcd</sup> ±0.23	2.5 <sup>cdef</sup> ±0.42	2.7 <sup>efgh</sup> ±0.31	2.9 <sup>efghi</sup> ±0.27	3.2 <sup>ghij</sup> ±0.21	3.3 <sup>hi</sup> ±0.43	2.4 <sup>bcd</sup> ±0.27
	3	2.6 <sup>defg</sup> ±0.31	2.9 <sup>efghi</sup> ±0.33	3.1 <sup>ghij</sup> ±0.23	3.3 <sup>hi</sup> ±0.14	3.5 <sup>i</sup> ±0.41	3.5 <sup>i</sup> ±0.22	2.8 <sup>efgh</sup> ±0.25
C20:5	1	3.5 <sup>hi</sup> ±0.68	1.2 <sup>bc</sup> ±0.73	0.8 <sup>b</sup> ±0.41	0.3 <sup>a</sup> ±0.21	0.3 <sup>a</sup> ±0.16	1.6 <sup>c</sup> ±0.48	2.7 <sup>efg</sup> ±0.93
	2	3.6 <sup>hij</sup> ±0.36	2.9 <sup>fg</sup> ±0.51	2.7 <sup>efg</sup> ±0.21	2.2 <sup>d</sup> ±0.23	3.2 <sup>gh</sup> ±0.14	1.1 <sup>b</sup> ±0.47	2.7 <sup>efg</sup> ±0.74
	3	5.9 <sup>kl</sup> ±0.73	5.3 <sup>k</sup> ±0.33	4.1 <sup>j</sup> ±0.38	3.7 <sup>ij</sup> ±0.62	2.4 <sup>de</sup> ±0.13	1.3 <sup>bc</sup> ±0.41	2.4 <sup>def</sup> ±0.41
C22:6	1	4.3 <sup>g</sup> ±0.5	2.2 <sup>de</sup> ±0.57	1.8 <sup>cd</sup> ±0.19	1 <sup>abc</sup> ±0.23	0.8 <sup>ab</sup> ±0.14	0.4 <sup>a</sup> ±0.11	3.1 <sup>ef</sup> ±0.51
	2	6.5 <sup>hi</sup> ±0.37	4.8 <sup>g</sup> ±0.5	3.2 <sup>f</sup> ±0.91	2.3 <sup>def</sup> ±0.71	2.3 <sup>def</sup> ±0.17	1.5 <sup>bcd</sup> ±0.72	5.7 <sup>gh</sup> ±0.17
	3	7.9 <sup>ij</sup> ±0.52	7.4 <sup>i</sup> ±0.21	7.3 <sup>ij</sup> ±0.13	6.6 <sup>hi</sup> ±0.5	2.5 <sup>def</sup> ±0.13	1.8 <sup>cd</sup> ±0.29	5.9 <sup>h</sup> ±0.18
Σn3	1	17.2 <sup>g</sup> ±0.34	8.2 <sup>bc</sup> ±0.11	6.8 <sup>b</sup> ±0.23	4.8 <sup>a</sup> ±0.31	4.3 <sup>a</sup> ±0.61	3.2 <sup>a</sup> ±0.41	10.9 <sup>de</sup> ±0.56
	2	20.6 <sup>hij</sup> ±0.9	16.8 <sup>g</sup> ±0.2	14.6 <sup>f</sup> ±0.31	11.6 <sup>e</sup> ±0.71	10.7 <sup>de</sup> ±0.71	8.2 <sup>bc</sup> ±0.72	16.6 <sup>g</sup> ±0.93
	3	25.3 <sup>k</sup> ±0.37	22.6 <sup>j</sup> ±0.17	21.2 <sup>ij</sup> ±0.36	18.2 <sup>gh</sup> ±0.92	10.6 <sup>de</sup> ±0.76	9.0 <sup>cd</sup> ±0.43	19.5 <sup>hi</sup> ±0.68
ΣPUFA	1	28.4 <sup>gh</sup> ±0.31	21.7 <sup>cd</sup> ±0.18	20.9 <sup>bc</sup> ±0.71	20.2 <sup>b</sup> ±0.76	22.9 <sup>de</sup> ±0.47	18.7 <sup>a</sup> ±0.64	27.2 <sup>fg</sup> ±0.88
	2	30.6 <sup>jk</sup> ±0.43	28.1 <sup>gh</sup> ±0.37	29.0 <sup>hi</sup> ±0.27	29.8 <sup>ij</sup> ±0.16	32.9 <sup>l</sup> ±0.18	26.6 <sup>f</sup> ±0.94	32.5 <sup>±0.93</sup>
	3	37.0 <sup>no</sup> ±0.25	35.3 <sup>o</sup> ±1.2	34.2 <sup>mn</sup> ±0.39	32.3 <sup>m</sup> ±0.92	26.7 <sup>kl</sup> ±0.62	23.1 <sup>e</sup> ±0.23	31.8 <sup>kl</sup> ±0.31
Σn6	1	12.0 <sup>ab</sup> ±0.41	13.7 <sup>gh</sup> ±0.93	14.3 <sup>gh</sup> ±0.93	15.7 <sup>i</sup> ±0.26	18.8 <sup>j</sup> ±0.63	15.7 <sup>i</sup> ±0.74	16.5 <sup>i</sup> ±0.82
	2	10.2 <sup>a</sup> ±0.82	11.6 <sup>abc</sup> ±0.45	14.6 <sup>h</sup> ±0.48	18.3 <sup>j</sup> ±0.9	22.4 <sup>k</sup> ±0.5	18.7 <sup>j</sup> ±0.54	16.2 <sup>i</sup> ±0.23
	3	11.9 <sup>bcd</sup> ±0.38	12.9 <sup>def</sup> ±0.51	13.2 <sup>efg</sup> ±0.54	14.3 <sup>gh</sup> ±0.23	16.3 <sup>i</sup> ±0.36	14.3 <sup>gh</sup> ±0.24	12.5 <sup>cde</sup> ±0.14
n3/n6	1	1.5 <sup>def</sup> ±0.64	0.8 <sup>bc</sup> ±0.2	0.7 <sup>abc</sup> ±0.25	0.5 <sup>ab</sup> ±0.14	0.4 <sup>a</sup> ±0.12	0.4 <sup>a</sup> ±0.12	0.9 <sup>c</sup> ±0.15
	2	2.2 <sup>g</sup> ±0.4	1.7 <sup>f</sup> ±0.9	1.2 <sup>d</sup> ±0.14	0.8 <sup>bc</sup> ±0.18	0.7 <sup>abc</sup> ±0.1	0.6 <sup>abc</sup> ±0.1	1.2 <sup>d</sup> ±0.4
	3	2.2 <sup>g</sup> ±0.46	1.8 <sup>f</sup> ±0.61	1.7 <sup>f</sup> ±0.14	1.3 <sup>de</sup> ±0.12	0.7 <sup>abc</sup> ±0.14	0.7 <sup>abc</sup> ±0.25	1.6 <sup>ef</sup> ±0.15

\*Values reported are means± standard error (n=3) as determined using Duncan's multiple range test. Means within the same row with different superscripts varied significantly (p<0.05). C18:2: Linoleic acid, C18:3: α-Linolenic acid, C20:5: Eicosapentaenoic acid, EPA, C22:6: Docosahexaenoic acid, DHA, Σω3: Total omega-3 fatty acids, ΣPUFA: Total Polyunsaturated fatty acids, Σω6: Total omega 6 fatty acids.

linseed oil at higher levels had no effect in the growth of Atlantic salmon and Murray cod (Menoyo et al., 2005; Turchini et al., 2011). In addition, there was no reduction in growth rates of European sea bass, *Dicentrarchus labrax* when FO was replaced with rapeseed oil, linseed oil and olive oil at 60%. Furthermore, growth performance in Atlantic salmon, *Salmo salar*, was also not affected when FO was replaced with VO at 50% (Storebakken, 2002).

It has been established that Nile tilapia and hybrid tilapia require both n-3 and n-6 polyunsaturated fatty acids as essential fatty acids for optimal growth (Chen et al., 2013; FAO, 2014). In addition, dietary n-3 and n-6 fatty acids requirements in tilapia are in the form of α-

linolenic and linoleic acids (Izquierdo et al., 2003; Tocher, 2003), therefore, dietary VO should not alter growth in fish because the needs of essential fatty acids (EFAs) are covered within the VO (Corraze and Kaushik, 2009). Dietary linoleic (C18:2) and α-linolenic (C18:3) acids were provided in the present study through sunflower and linseed oil respectively.

High dietary linseed oil (>50%) in the diets 1 to 3 (Table 1) might have resulted into n3/n6 imbalance creating oxidative stress as n-3 fatty acids physiological tolerance for tilapia might have been exceeded thus possible compromised immune system in the study subjects (Li et al., 2016) reduced survival rate in this study. In addition, it has been established that n-6 fatty acids have superior

**Table 7.** Monthly changes of liver polyunsaturated fatty acid composition in tilapia fed diets containing varying composition of linseed and sunflower.

Acid	Month	Treatments						
		Diet1	Diet 2	Diet 3	Diet 4	Diet 5	Washout	Commercial
C18:2	1	11.9 <sup>bc</sup> ±0.52	13.5 <sup>ef</sup> ±0.4	14.1 <sup>fg</sup> ±0.64	15.7 <sup>h</sup> ±0.97	18.7 <sup>j</sup> ±0.92	15.6 <sup>h</sup> ±0.34	16.8 <sup>i</sup> ±0.41
	2	10.2 <sup>a</sup> ±0.82	11.6 <sup>b</sup> ±0.24	14.6 <sup>g</sup> ±0.4	18.3 <sup>l</sup> ±0.91	22.4 <sup>k</sup> ±0.51	18.7 <sup>i</sup> ±0.53	16.1 <sup>hi</sup> ±0.45
	3	10.3 <sup>a</sup> ±0.72	12.5 <sup>cd</sup> ±0.61	13.1 <sup>de</sup> ±0.31	13.9 <sup>fg</sup> ±0.82	16.2 <sup>hi</sup> ±0.64	14.4 <sup>g</sup> ±0.13	11.7 <sup>b</sup> ±0.24
C18:3	1	7.32 <sup>h</sup> ±0.23	4.9 <sup>cd</sup> ±0.52	4.5 <sup>c</sup> ±0.46	3.8 <sup>b</sup> ±0.41	3.5 <sup>b</sup> ±0.29	1.6 <sup>a</sup> ±0.21	6.4 <sup>fg</sup> ±0.23
	2	10.8 <sup>k</sup> ±0.32	9.3 <sup>j</sup> ±0.54	8.9 <sup>ij</sup> ±0.69	7.4 <sup>h</sup> ±0.53	5.5 <sup>de</sup> ±0.51	1.8 <sup>a</sup> ±0.13	8.5 <sup>i</sup> ±0.7
	3	12.2 <sup>m</sup> ±0.86	11.3 <sup>l</sup> ±0.55	10.3 <sup>k</sup> ±0.42	9.4 <sup>i</sup> ±0.6	6.5 <sup>g</sup> ±0.51	2.5 <sup>ab</sup> ±0.13	10.9 <sup>kl</sup> ±0.94
C20:4	1	2.1 <sup>a</sup> ±0.51	2.6 <sup>abc</sup> ±0.41	2.9 <sup>bcd</sup> ±0.12	3.1 <sup>cdef</sup> ±0.22	3.3 <sup>cdefg</sup> ±0.27	3.8 <sup>ghij</sup> ±0.13	2.3 <sup>ab</sup> ±0.16
	2	3.0 <sup>cde</sup> ±0.81	3.3 <sup>cdefg</sup> ±0.24	3.5 <sup>defgh</sup> ±0.34	3.7 <sup>efghi</sup> ±0.35	4.0 <sup>ghij</sup> ±0.43	4.2 <sup>hij</sup> ±0.41	3.1 <sup>cdef</sup> ±0.24
	3	3.4 <sup>defg</sup> ±0.23	3.6 <sup>defghi</sup> ±0.33	3.8 <sup>ghij</sup> ±0.16	4.0 <sup>ghij</sup> ±0.32	4.3 <sup>ij</sup> ±0.31	4.5 <sup>i</sup> ±0.23	3.5 <sup>defgh</sup> ±0.21
C20:5	1	3.8 <sup>i</sup> ±0.12	1.8 <sup>e</sup> ±0.68	1.0 <sup>c</sup> ±0.24	0.2 <sup>a</sup> ±0.01	0.5 <sup>b</sup> ±0.17	2.0 <sup>ef</sup> ±0.27	2.3 <sup>f</sup> ±0.86
	2	3.6 <sup>i</sup> ±0.36	2.9 <sup>gh</sup> ±0.51	2.7 <sup>g</sup> ±0.21	2.2 <sup>f</sup> ±0.31	2.2 <sup>f</sup> ±0.14	1.1 <sup>c</sup> ±0.47	2.7 <sup>g</sup> ±0.74
	3	6.3 <sup>k</sup> ±0.52	5.2 <sup>j</sup> ±0.59	5.1 <sup>i</sup> ±0.22	3.7 <sup>i</sup> ±0.24	2.1 <sup>ef</sup> ±0.27	1.5 <sup>d</sup> ±0.29	3.8 <sup>i</sup> ±0.17
C22:6	1	5.4 <sup>h</sup> ±0.43	3.0 <sup>f</sup> ±0.12	1.9 <sup>cd</sup> ±0.31	1.4 <sup>b</sup> ±0.12	1.2 <sup>b</sup> ±0.19	0.6 <sup>a</sup> ±0.23	3.1 <sup>f</sup> ±0.61
	2	6.5 <sup>i</sup> ±0.14	4.8 <sup>g</sup> ±0.45	2.4 <sup>de</sup> ±0.71	2.3 <sup>de</sup> ±0.21	1.5 <sup>bc</sup> ±0.73	1.2 <sup>b</sup> ±0.24	3.3 <sup>f</sup> ±0.91
	3	7.4 <sup>l</sup> ±0.16	7.4 <sup>i</sup> ±0.91	7.3 <sup>i</sup> ±0.17	7.0 <sup>m</sup> ±0.51	2.8 <sup>ef</sup> ±0.31	1.6 <sup>bc</sup> ±0.21	7.8 <sup>kl</sup> ±0.95
Σn3	1	16.3 <sup>i</sup> ±0.31	9.6 <sup>e</sup> ±0.43	7.3 <sup>c</sup> ±0.21	5.2 <sup>b</sup> ±0.17	5.0 <sup>b</sup> ±0.62	4.1 <sup>a</sup> ±0.72	11.6 <sup>g</sup> ±0.18
	2	20.6 <sup>k</sup> ±0.15	16.8 <sup>i</sup> ±0.43	14.6 <sup>h</sup> ±0.32	11.6 <sup>g</sup> ±0.71	10.7 <sup>f</sup> ±0.24	8.2 <sup>d</sup> ±0.92	16.6 <sup>i</sup> ±0.91
	3	25.5 <sup>n</sup> ±0.51	23.8 <sup>m</sup> ±0.51	22.4 <sup>l</sup> ±0.34	19.8 <sup>i</sup> ±0.62	11.1 <sup>f</sup> ±0.24	9.3 <sup>e</sup> ±0.23	22.3 <sup>l</sup> ±0.61
ΣPUFA	1	28.0 <sup>f</sup> ±0.31	22.9 <sup>c</sup> ±0.26	21.1 <sup>b</sup> ±0.16	20.7 <sup>b</sup> ±0.52	23.5 <sup>d</sup> ±0.82	19.4 <sup>a</sup> ±0.53	28.2 <sup>f</sup> ±0.92
	2	30.6 <sup>i</sup> ±0.41	28.1 <sup>f</sup> ±0.71	29.0 <sup>g</sup> ±0.36	29.8 <sup>h</sup> ±0.23	32.9 <sup>i</sup> ±0.41	26.6 <sup>e</sup> ±0.91	32.5 <sup>j</sup> ±0.9
	3	36.6 <sup>n</sup> ±0.83	36.1 <sup>m</sup> ±0.71	35.3 <sup>l</sup> ±0.51	33.5 <sup>k</sup> ±0.94	27.1 <sup>e</sup> ±0.17	23.4 <sup>d</sup> ±0.62	33.8 <sup>k</sup> ±0.34
Σn6	1	11.7 <sup>b</sup> ±0.52	13.3 <sup>de</sup> ±0.92	13.9 <sup>ef</sup> ±0.32	15.6 <sup>hi</sup> ±0.43	18.6 <sup>k</sup> ±0.41	15.4 <sup>h</sup> ±0.85	16.6 <sup>j</sup> ±0.94
	2	10.2 <sup>a</sup> ±0.81	11.6 <sup>b</sup> ±0.21	14.6 <sup>g</sup> ±0.52	18.3 <sup>k</sup> ±0.91	22.4 <sup>l</sup> ±0.51	18.7 <sup>k</sup> ±0.53	16.1 <sup>ij</sup> ±0.56
	3	10.3 <sup>a</sup> ±0.72	12.5 <sup>c</sup> ±0.62	13.1 <sup>cd</sup> ±0.31	13.9 <sup>ef</sup> ±0.81	16.2 <sup>ij</sup> ±0.61	14.4 <sup>fg</sup> ±0.15	11.7 <sup>b</sup> ±0.25
n3/n6	1	1.4 <sup>d</sup> ±0.96	0.8 <sup>abc</sup> ±0.17	0.6 <sup>a</sup> ±0.22	0.4 <sup>a</sup> ±0.26	0.3 <sup>a</sup> ±0.17	0.3 <sup>a</sup> ±0.18	0.7 <sup>abc</sup> ±0.14
	2	2.3 <sup>g</sup> ±0.94	1.7 <sup>def</sup> ±0.19	1.2 <sup>bcd</sup> ±0.59	0.8 <sup>abc</sup> ±0.12	0.7 <sup>ab</sup> ±0.12	0.6 <sup>a</sup> ±0.17	1.3 <sup>cde</sup> ±0.94
	3	2.6 <sup>h</sup> ±0.81	2.0 <sup>g</sup> ±0.17	1.8 <sup>efg</sup> ±0.22	1.5 <sup>def</sup> ±0.29	0.7 <sup>abc</sup> ±0.18	0.7 <sup>ab</sup> ±0.4	2.0 <sup>fg</sup> ±0.27

\*Values reported are means± standard error (n=3) as determined using Duncan's multiple range test. Means within the same row with different superscripts varied significantly (p<0.05). C18.2: Linoleic acid, C18.3: α-Linolenic acid, C20.5: Eicosapentaenoic acid, EPA, C22.6: Docosahexaenoic acid, DHA, Σω3: Total omega-3 fatty acids, ΣPUFA: Total Polyunsaturated fatty acids, Σω6: Total omega 6 fatty acids.

growth-promoting effects than n-3 fatty acids in red belly tilapia (Lim et al., 2009) confirming a relatively better growth rates at high (>50%) sunflower oil content in the present study.

Studies have shown that fish species, environmental factors, fish size and age as well as diets affect fatty acid composition of fish tissues (Saito et al., 1999; Kiessling et al., 2001). In this study, tilapia were raised under same exogenous conditions therefore culturing period and diet were the possible factors that could affect fatty acid profile of fish under study. From our findings, DHA, total n3 and n3/n6 ratio increased significantly (P<0.05) from month 1 to month 3 in both tissues (Table 6 and 7), an observation which relates with previous finding that fatty

acid profile in fish is dependent on fish age (Parlov et al., 2009; Nemova et al., 2015a; Nemova et al., 2015b; Denis and Nina, 2016). A study on Atlantic salmon indicated that in two year feeding period, the levels of n3 fatty acids increased from 24 to 29.5%, DHA increased from 5.4 to 8.6% and n3/n6 ratio increased from 2.5 to 3 (Svetlana et al., 2003). In the finding, DHA, n3 fatty acids and n3/n6 ratios increased as follows DHA (4.3 to 7.1%, diet 1 muscles; 5.4 to 7.4% diet 1 liver), n3 fatty acids (17 to 25% in diet 1 muscle; 5.4 to 7.4% in diet 1 liver) and n3/n6 ratios (1.5 to 2.2 in diet 1 muscles; 1.4 to 2.6 in diet 1 liver) (Tables 6 and 7).

The increase in these values correlated with dietary increase in linseed oil proportions suggests active

metabolism of fatty acid from dietary lipid source in tilapia.

Provision of dietary fatty acid in excess promotes intensive beta-oxidation for energy production (Stubhaug et al., 2006, 2007). The present findings is consistent with a study by Stubhaug et al. (2007), that there is preference in fatty acids beta-oxidation in which some fatty acids, specifically omega 3 long chain polyunsaturated fatty acids, LC-PUFAS such as DHA, are spared from catabolism. This explains the tissue accumulation of omega 3 fatty acid particularly, DHA, in our study. This accumulation correlates with the levels of linseed oil inclusion in the diet. Specifically, diet 1 and 2 with > 50% linseed oil inclusion had significantly high liver DHA and omega 3 fatty acids (7.4, 7.4 and 25.5,23.8%) respectively (Table 7) and muscle DHA and omega 3 fatty acids (7.9, 7.4 and 25.3, 22.6) (Table 6).

Tissue polyunsaturated fatty acid values in our study increased with increased dietary proportion of linseed oil, finding which is consistent with previous study indicating significant increase in total n3 fatty acids and DHA content as dietary linseed oil increases (Li et al., 2015). Tonial et al. (2009) observed that Nile tilapia fed dietary flaxseed oil had significant increase in muscle total n3 fatty acids and DHA content. The finding supports suggestion that tilapia is capable of converting dietary alpha- linolenic acid to LC-PUFA (Sargent et al., 2002) and that they can elongate and desaturate dietary alpha-linolenic acids to tissue DHA. However, the level of conversion of C18 PUFA to highly polyunsaturated fatty acids (HUFA) varies among species (Sargent et al., 2002). This variation reflects different rates of assimilation and catabolism of dietary fatty acids once consumed (Yones et al., 2013). The extent of interactions between dietary fatty acids may also determine final tissue lipids composition (Fonseca-Madrigal et al., 2005; Matsushita et al., 2006).

Arachidonic acid (C20:4) levels reported in the present study ranged between 3.4 and 4.5% in liver and 2.6 and 3.5% in tilapia muscles. These values were lower than values observed by Feirreira et al. (2011), 5.8 to 8.12% and Ribeiro et al (2008), 5.8 to 9.2% in tilapia. In addition, values reported in our study were low relative to the dietary linoleic acid (C18:2) (Table 3). These low values might have been due to varied roles played by arachidonic acid which includes being a source for eicosanoids formation and roles it play in resisting stressors prevalent under intensive culture system (Bell and Sargent, 2003; Li et al., 2016). In addition, previous studies indicate that excessive dietary supply of C18 polyunsaturated fatty acids may create selective competition disrupting bioconversion thus low levels of arachidonic acid (Ruyter et al., 2006). This is because alpha linolenic acid and linoleic acids use almost same enzymes for their metabolism (Visentainer, 2007).

The activities and expression of these enzymes are also affected by dietary lipids (Visentainer, 2007). In

conclusion, dietary fish oil can successfully be replaced with linseed oil for adequate essential fatty acids in tilapia muscles, however, high dietary linseed oil (>50%) lowers survival and growth rate of tilapia. From our data, a ratio of 25:75 linseed to sunflower oil gave a relatively better survival and growth levels, however, studies are needed to establish the optimal dietary linseed oil that promote growth, survival and tissue deposition of essential fatty acids.

## CONFLICT OF INTERESTS

The authors have not declared any conflict of interests.

## ACKNOWLEDGMENTS

The author extends their thanks to Research Production and Extension Division, Jomo Kenyatta University of Agriculture and Technology (JKUAT-RPE) and Africa-ai-Japan Project Office for financial support as well as Food Science and Technology Department, JKUAT for laboratory space and instrumentation aid to carry out analyses.

## REFERENCES

- AOAC (1995). Official methods of analysis, 16th edition. Association of Official Analytical Chemists, Inc., Arlington, Virginia, USA.
- Bell JG, Sargent JR (2003). Arachidonic acid in aquaculture feeds: Current status and future opportunities. *Aquaculture* 218:491-499.
- Bligh EC, Dyer WJ (1959). A rapid method of total lipid extraction and purification. *Can. J. Biochem. Physiol.* 37:913-917.
- Bonaa KH, Bjerve KS, Straume B, Gram IT, Thelle D (1990). Effect of eicosapentaenoic and docosahexaenoic acids on blood pressure in hypertension. A population-based intervention trial from the Tromso study. *N. Engl. J. Med.* 322:795-801.
- Bouraoui L, Sánchez-Gurmaches J, Cruz-García L, Gutiérrez J, Benedito-Palos L, Pérez-Sánchez J, Navarro I (2011). Effect of dietary fish meal and fish oil replacement on lipogenic and lipoprotein lipase activities and plasma insulin in gilthead sea bream (*Sparus aurata*). *Aquac. Nutr.* 17: 54-63.
- Brenna JT (2002). Efficiency of conversion of alpha-linolenic acid to long chain n-3 fatty acids in man. *Curr. Opin. Clin. Nutr. Metab. Care.* 5(2):127-32.
- Charo-Karisa H, Gachupin M (2010). Overview of the Fish Farming Enterprise Productivity Program. In: End of Year Report Fish Farming Enterprise Productivity Program Phase I, Aquaculture Development Working Group Ministry of Fisheries Development, Kenya.
- Chen CY, Sun BL, Li XX, Li PY, Guan WT, Bi YZ, Pan Q (2013). N-3 essential fatty acids in Nile tilapia, *Oreochromis niloticus*: quantification of optimum requirement of dietary linolenic acid in juvenile fish. *Aquaculture* 416-417:99-104.
- Corraze G, Kaushik S (2009). Alimentation lipidique et remplacement des huiles de poisson par des huiles végétales en pisciculture. *Cahiers Agric.* 18:112-118.
- Denis AE, Nina NN (2016). Age-Specific Lipid and Fatty Acid Profiles of Atlantic Salmon Juveniles in the Varzuga River. *Int. J. Mol. Sci.* 17:1050.
- Duncan DB (1955). Multiple-range and multiple F tests. *Biometrics* 11:1-42.
- FAO (2007). Aquaculture: status and prospects.
- FAO - Food and Agricultural Organization of the United Nations (2014).



- Species Profiles. Website. Available at: <http://www.fao.org/fishery/affris/species-profiles/nile-tilapia/nutritional-requirements/en/>.
- Ferreira MW, Araujo FG, Costa DV, Rosa PV, Figueiredo HCP, Murgas LDS (2011) Influence of dietary oil sources on muscle composition and plasma lipoprotein concentration in Nile tilapia (*Oreochromis niloticus*). *J World Aquac. Soc.* 42:24-33.
- Fonseca-Madrigal J, Karlazos V, Campbell PJ, Bell JG, Tocher DR (2005). Influence of dietary palm oil on growth tissue fatty acid composition and fatty acid metabolism in liver and intestine in rainbow trout (*Oncorhynchus mykiss*). *Aquac. Nutr.* 11:241-250.
- Francis DS, Turchini GM, Jones PL, De Silva SS (2006). Effects of dietary oil source on growth and fillet fatty acid composition of Murray cod, *Maccullochella peelii peelii*. *Aquaculture* 253:547-556.
- Gebauer SK, Posta TL, Harris WS, Kris-Etherton PM (2006). N-3 fatty acid dietary recommendations and food sources to achieve essentiality and cardiovascular benefits. *Ame. J. Clin. Nutr.* 83(Suppl. 6):1526-1535.
- Huang CH, Huang MC, Hou PC (1998). Effect of dietary lipids on fatty acid composition and lipid peroxidation in sarcoplasmic reticulum of hybrid tilapia, *Oreochromis niloticus* × *O. aureus*. *Comp. Biochem. Physiol. Part A*, 120:331-336.
- Izquierdo MS, Obach A, Aranzamendi L, Montero D, Robania L, Rosenlund G (2003). Dietary lipid sources for seabream and sea bass: growth performance, tissue composition and flesh quality. *Aquac. Nutr.* 9:397-407.
- Justi KC, Hayashi C, Visentainer JV, de Souza NE, Matsushita M (2003). The influence of feed supply time on the fatty acid profile of Nile tilapia (*Oreochromis niloticus*) fed on a diet enriched with n-3 fatty acids. *Food Chem.* 80(4):489-493.
- Kanazawa A, Teshima S, Sakamoto M, Awal A (1980). Requirements of Tilapia zilli for essential fatty acids. *Bull. Jap. Soc. Sci. Fish.* 46:1353-1356.
- Kiessling A, Pickova J, Johansso L, Asgard T, Storebakken T, Kiessling KH (2001). Changes in fatty acid composition in muscle and adipose tissue of farmed rainbow trout (*Oncorhynchus mykiss*) in relation to ration and age. *Food Chem.* 73:271-284.
- Kiron V, Thawonsuwan J, Panigrahi A, Scharsack JP, Satoh S (2011). Antioxidant and immune defences of rainbow trout (*Oncorhynchus mykiss*) offered plant oils differing in fatty acid profiles from early stages. *Aquac. Nutr.* 17:130-140.
- Kris-Etherton PM, Taylor DS, Yu-Poth S (2000). Polyunsaturated fatty acids in the food chain in the United States. *Am. J. Clin. Nutr.* 71:179-188.
- Kusunto A, Ishikura Y, Kawashima H (2007). Effects of archidonate-enriched triacylglycerol supplemented on serum fatty acids and platelet aggregation in healthy male subjects with a fish diet. *Br. J. Nutr.* 98:626-633.
- Leaf A, Weber PC (1988). Cardiovascular effects of n-3 fatty acids. *Engl. J. Med.* 318:549-557.
- Lee SM (2001). Review of the lipid and essential fatty acid requirements of rockfish (*Sebastes schlegelii*). *Aquac. Res.* 32:8-17.
- Li FJ, Lin X, Lin SM, Chen YW, Guan T (2016). Effects of dietary fish oil substitution with linseed oil on growth, muscle fatty acid and metabolism of tilapia (*Oreochromis niloticus*). *Aquac. Nutr.* 22:499-508.
- Lim C, Aksoy M, Klesius P (2009). Lipids and fatty acid requirements of tilapia – Dietary supplementation essential for health, reproduction. *Global Aquac. Advocate* 12(5):86-87.
- Ljubojević D, Ćirković M, Novakov N, Jovanović R, Janković S, Đorđević, Mašić Z (2013). Productivity and meat nutrient in fish: the diet effect. *Kafkas universiteti veteriner fakultesi dergisi*, 19(1):43-49.
- Ljubojević D, Radosavljević V, Puvača N, Baloš MŽ, Đorđević V, Jovanović R, Ćirković M (2015). Interactive effects of dietary protein level and oil source on proximate composition and fatty acid composition in common carp (*Cyprinus carpio* L.). *J. Food Comp. Anal.* 37:44-50.
- Matsushita M, Justi KC, Padre RG, Milinsk MC, Hayashi C, Gomes STM, Visentainer JV, Souza NE (2006). Influence of diets enriched with different vegetable oils on the performance and fatty acid profile of Nile tilapia (*Oreochromis niloticus*) fingerlings. *Acta Sci. Technol.* 28:125-131.
- Menoyo DL, Opez-Bote CJ, Obach A, Bautista JM (2005). Effect of dietary fish oil substitution with linseed oil on the performance, tissue fatty acid profile, metabolism, and oxidative stability of Atlantic salmon. *J. Anim. Sci.* 83:2853-2862.
- Menoyo, D, Izquierdo MS, Robaina L, Gines R, Lopez-Bote CJ, Bautista JM (2004). Adaptation of lipid metabolism, tissue composition and flesh quality in gilthead seabream (*Sparus aurata*) to the replacement of dietary fish oil by linseed and soyabean oils. *Br. J. Nutr.* 92:41-52.
- Montero D, Mathlouthi F, Tort L, Afonso JM, Torrecillas S, Fernández-Vaquero A, Negrin D, Izquierdo MS (2010). Replacement of dietary fish oil by vegetable oils affects humoral immunity and expression of proinflammatory cytokines genes in gilthead sea bream *Sparus aurata* Fish Shellfish Immunol. 29(6):1073-1081.
- Montero D, Kalinowski T, Obach A, Robaina L, Tort L, Caballero MJ and Izquierdo MS. (2003). Vegetable lipid sources for gilthead seabream (*Sparus aurata*): effects on fish health. *Aquaculture* 225:353-370.
- Mourente G, Good JE, Thompson KD, Bell JG (2007). Effects of partial substitution of dietary fish oil with blends of vegetable oils, on blood leucocyte fatty acid compositions, immune function and histology in European sea bass (*Dicentrarchus labrax* L.). *Br. J. Nutr.* 98:770-779.
- Munguti JM, Safina M, Paul SO, Domitila NK, Mary O, Harrison C, Erick OO (2014). An overview of current status of Kenyan fish feed industry and feed management practices, challenges and Opportunities *Int. J. Fish. Aqua. Stud.* (6):128-137.
- National Research Council (NRC) (2011). Nutrient Requirements of fish and shrimp. National Academy Press, Washington, D.C., USA
- Nemova NN, Murzina SA, Nefedova ZA, Veselov AE (2015a). Features in the lipid status of two generations of fingerlings (0+) of Atlantic salmon (*Salmo salar* L.) inhabiting the Arenga River (Kola Peninsula). *Int. J. Mol. Sci.* 16:17535-17545.
- Nemova NN, Nefedova ZA, Murzina SA, Veselov AE, Ripatti PO, Pavlov DS (2015b). The effect of environmental conditions on the dynamics of fatty acids in juveniles of the Atlantic salmon (*Salmo salar* L.). *Russ. J. Ecol.* 46:267-271.
- Ng WK (2002). Potential of palm oil utilisation in aquaculture feeds. *Asia Pac. J. Clin. Nutr.* 11(Suppl 7):S473-476.
- Ng WK (2005). Lipid nutrition of farmed tilapia. *Global Aquac. Advocate* October, 60–61.
- Parlov DS, Nefedova ZA, Veselov AE, Nemova NN, Ruokolainen TR, Vasil'eva OB, Ripatti PO (2009). Age dynamics of lipid status of juveniles of Atlantic salmon (*Salmo salar* L.) from the Varzuga River. *J. Ichthyol.* 49:1073-1080.
- Popa VM, Gruia AR, Dumbrava DN, Moldovan D, Bordean D, Mateescu C (2012). Fatty acid composition and oil characteristics of linseed from Romania J. *Agroalignment. Process Technol.* 18:136-140.
- Ribeiro PAP, Logato PVR, Paula DAJ, Costa AC, Murgas DS, Freitas RTF (2008). Effect of different oils in the diet on lipogenesis and the lipid profile of Nile tilapia. *Rev. Bras. Zootecnia* 37:1331-1337.
- Saito H, Yamashiro R, Alasalvar C, Konno T (1999). Influence of diet on fatty acids of three subtropical fish, subfamily caesioninae (*Caesio diagramma* and *C. tile*) and family siganidae (*Siganus canaliculatus*). *Lipids* 34:1073-1082.
- Sargent JR, Tocher DR, Bell JG (2002). The lipids. In: *Fish Nutrition* (Halver, J.E. & Hardy, R.W. eds), pp. 181–257. Academic Press, San Diego, CA.
- Storebakken T (2002). Nutrient requirements and feeding of finfish for aquaculture Atlantic salmon. Pages 79–102 in C. D. Webster and C. Lim, editors. CABI Publishing, Wallingford, UK.
- Stubhaug L, Li O, Torstensen BE (2006). Beta-oxidation capacity in liver increase during part-smolt transformation of Atlantic salmon (*Salmo salar* L.) fed vegetable oil and fish oil. *J. Fish Biol.* 69:504-517.
- Stubhaug L, Lio O, Torstensen BE (2007). Fatty acid productive value and beta-oxidation capacity in Atlantic salmon (*Salmo salar* L.) fed on different lipid sources along the whole growth period. *Aquac. Nutr.* 13:145-155.
- Svetlana AM, Zinaida A N, Svetlana NP, Alexey EV, Tocher DR (2003). Metabolism and functions of lipids and fatty acids in teleost fish. *Rev Fish Sci* 11:107-184.
- Turchini GM, Francis DS, Senadheera, SPSD, Thanuthong T, De Silva SS (2011). Fish oil replacement with different vegetable oils in Murray cod: evidence of an 'omega-3 sparing effect' by other dietary fatty

- acids. *Aquaculture* 315:250-259.
- Turchini GM, Francs DS (2009). Fatty acid metabolism (desaturase, elongation and beta-oxidation) in rainbow trout fed fish oil- or linseed oil based diets. *Br. J. Nutr.* 102:69-81.
- Visentainer JV, De Souza NE, Makota M, Hayashi C, Franco MRB (2005). Influence of diets enriched with flaxseed oil on the alpha-linolenic, eicosapentaenoic and docosahexaenoic fatty acid in Nile tilapia (*Oreochromis niloticus*). *Food Chem.* 90:557-560.
- Yones AM, El-Saidy DMSD, Abdel-Hakim NF (2013). Effects of fish oil substitution with vegetable oils in diets of juvenile Nile tilapia, *Oreochromis niloticus* (L.) on growth performance, nutrients utilization and muscle fatty acids contents, *Merit Res. J. Food Sci. Technol.* 1(1):009-018.
- Župan B, Ljubojević D, Pelić M, Ćirković M, Đorđević V, Bogut I (2016). Common carp response to the different concentration of linseed oil in diet. *Slovenian Vet. Res.* 53(1).

A hand is shown holding a large, silvery fish, possibly a bass, against a clear blue sky. The fish is held vertically, with its head pointing upwards. The background is a soft-focus landscape with greenery and water.

# International Journal of Fisheries and Aquaculture

## Related Journals Published by Academic Journals

- *Journal of Plant Breeding and Crop Science*
- *African Journal of Agricultural Research*
- *Journal of Horticulture and Forestry*
- *International Journal of Livestock Production*
- *International Journal of Fisheries and Aquaculture*
- *Journal of Cereals and Oilseeds*
- *Journal of Soil Science and Environmental Management*
- *Journal of Stored Products and Postharvest Research*

**academicJournals**