



International Journal of Medicine and Medical Sciences

Volume 9 Number 6 June, 2017

ISSN 2006-9723



*Academic
Journals*

ABOUT IJMMS

The **International Journal of Medicine and Medical Sciences** is published monthly (one volume per year) by Academic Journals.

The **International Journal of Medicine and Medical Sciences (IJMMS)** provides rapid publication (monthly) of articles in all areas of Medicine and Medical Sciences such as:

Clinical Medicine: Internal Medicine, Surgery, Clinical Cancer Research, Clinical Pharmacology, Dermatology, Gynaecology, Paediatrics, Neurology, Psychiatry, Otorhinolaryngology, Ophthalmology, Dentistry, Tropical Medicine, Biomedical Engineering, Clinical Cardiovascular Research, Clinical Endocrinology, Clinical Pathophysiology, Clinical Immunology and Immunopathology, Clinical Nutritional Research, Geriatrics and Sport Medicine

Basic Medical Sciences: Biochemistry, Molecular Biology, Cellular Biology, Cytology, Genetics, Embryology, Developmental Biology, Radiobiology, Experimental Microbiology, Biophysics, Structural Research, Neurophysiology and Brain Research, Cardiovascular Research, Endocrinology, Physiology, Medical Microbiology

Experimental Medicine: Experimental Cancer Research, Pathophysiology, Immunology, Immunopathology, Nutritional Research, Vitaminology and Etiology

Preventive Medicine: Congenital Disorders, Mental Disorders, Psychosomatic Diseases, Addictive Diseases, Accidents, Cancer, Cardiovascular Diseases, Metabolic Disorders, Infectious Diseases, Diseases of Bones and Joints, Oral Preventive Medicine, Respiratory Diseases, Methods of Epidemiology and Other Preventive Medicine

Social Medicine: Group Medicine, Social Paediatrics, Medico-Social Problems of the Youth, Medico-Social Problems of the Elderly, Rehabilitation, Human Ecology, Environmental Toxicology, Dietetics, Occupational Medicine, Pharmacology, Ergonomy, Health Education, Public Health and Health Services and Medical Statistics The Journal welcomes the submission of manuscripts that meet the general criteria of significance and scientific excellence. Papers will be published approximately one month after acceptance. All articles published in IJMMS are peer-reviewed.

Contact

Editorial Office: ijmms@academicjournals.org

Desk: helpdesk@academicjournals.org

Website: <http://www.academicjournals.org/journal/IJMMS>

Submit manuscript online <http://ms.academicjournals.me/>

Editors

Dr. J. Ibekwe

*Acting Editor-in-chief,
International Journal of Medicine and Medical
Sciences Academic Journals
E-mail: ijmms.journals@gmail.com
<http://www.academicjournals.org/ijmms>*

Afrozul Haq

*Editor, Laboratory Medicine
Department of Laboratory Medicine
Sheikh Khalifa Medical City
P.O. Box 51900, ABU DHABI
United Arab Emirates*

Editorial Board

Chandrashekhar T. Sreeramareddy

*Department of Community Medicine,
P O Box No 155, Deep Heights
Manipal College of Medical Sciences,
Pokhara,
Nepal*

Sisira Hemananda Siribaddana

*259, Temple Road, Thalpathpitiya,
Nugegoda, 10250
Sri Lanka*

Dr. santi M. Mandal

*Internal Medicine
UTMB, Galveston, TX,
USA*

Konstantinos Tziomalos

*Department of Clinical Biochemistry
(Vascular Prevention Clinic),
Royal Free Hospital Campus,
University College Medical School, University College
London, London,
United Kingdom*

Cyril Chukwudi Dim

*Department of Obstetrics & Gynaecology
University of Nigeria Teaching Hospital (UNTH)
P.M.B. 01129, Enugu. 400001,
Nigeria*

Mojtaba Salouti

*School of Medical and Basic Sciences,
Islamic Azad University- Zanjan,
Iran*

Imtiaz Ahmed Wani

*Srinagar Kashmir, 190009,
India*

Professor Viroj Wiwanitkit

*Wiwanitkit House, Bangkhae,
Bangkok
Thailand 10160*

Dr. Srinivas Koduru

*Dept of Clinical Sciences
Collage of Health Sciences
University of Kentucky
Lexington USA*

Weiping Zhang

*Department of Oral Biology
Indiana University School of Dentistry
1121 West Michigan Street, DS 271
Indianapolis, IN 46202
USA*

Lisheng XU

*Ho Sin Hang Engineering Building
Department of Electronic Engineering
The Chinese University of Hong Kong
Shatin, N.T. Hong Kong,
China*

Dr. Mustafa Sahin

*Department of Endocrinology and Metabolism
Baskent University,
Ankara,
Turkey*

Dr. Harshdeep Joshi

*Maharishi Markandeshwar
Institute of Medical Sciences and Research
Ambala, (Haryana).
India.*

ARTICLES

| | |
|---|-----------|
| Prevalence and associated factors of female genital mutilation among high school students in Dale Wabera Woreda, Oromia Regional State, Ethiopia | 72 |
| Desalegn Shiferaw, Negussie Deyessa, Gamachu Fufa, Tadele Kinati and Markos Desalegn | |
| Clinical outcomes of patients admitted in intensive care units of Nigist Eleni Mohammed Memorial Hospital of Hosanna, Southern Ethiopia | 79 |
| Mohammed Suleiman Obsa, Abdi Oumer Adem and Getish Bezabih Gete | |

Full Length Research Paper

Prevalence and associated factors of female genital mutilation among high school students in Dale Wabera Woreda, Oromia Regional State, Ethiopia

Desalegn Shiferaw^{1*}, Negussie Deyessa², Gamachu Fufa³, Tadele Kinati⁴ and Markos Desalegn⁵

¹Epidemiology and Biostatistics Unit, Department of Public Health, College of Medicine and Health Sciences, Jigjiga University, Ethiopia.

²School of Public Health, College of Health Sciences, Addis Ababa University, Ethiopia.

³Department of Sociology, College of Social Sciences and Humanities, Jigjiga University, Ethiopia.

⁴Departments of Nursing, College of Medicine and Health Sciences, Jigjiga University, Ethiopia.

⁵Reproductive Health and Nutrition Unit, Department of Public Health, College of Medicine and Health Sciences, Jigjiga University, Ethiopia.

Received 3 April, 2017; Accepted 3 May, 2017

Female genital mutilation is still a serious health problem in Ethiopia. Female genital mutilation conceptualizes harmful traditional practice and is among existing traditional attitudes and practices on practice in different corners of the world. Other than demographic and health survey data, studies showing the prevalence of female genital mutilation is scarce and in some areas nonexistent. Its existence is only, time and again recognized as the practice has various health, social and psychological adversities. This study team found it imperative as the problem warranty study with the aim to assess prevalence of female genital mutilation (FGM) as well as identifying its associated factors among high school students. Institution based cross sectional study design was employed. 95% confidence level and 3% margin of error were used to study the magnitude and associated factors of FGM among 798 high school and preparatory students. A multi stage sampling procedure was employed to select female students in the schools. This research showed percentages and association among the variables by univariate, bivariate and multivariate analysis. Data were collected from 769 study participants with a response rate of 96.4%. About 78%, 95% CI (74.9%, 80.7%) of the study participants underwent the procedure. Age ≥ 17 [AOR=1.8, 95% CI (1.28, 2.61)], grades 11 and 12 [AOR =2.98; 95% CI (1.45, 6.12)] and [AOR=6.52, 95% CI (1.85, 22.94)], respectively, being from rural area [AOR= 1.6, 95% CI (1.01, 2.52)], and house wife and merchant mother occupation [AOR= 2.38, 95% CI (1.07, 5.29)] and [AOR= 2.72, 95% CI (1.04, 7.12)], respectively were independently associated to higher prevalence of female genital mutilation. Irrespective of different forms of interventions in the country, the prevalence of female genital mutilation is still very high in the study area. This study thus revealed that palatable and very convenient intervention strategies need to be crafted to curb the problem at large and for those rural villagers who are at higher risk in particular.

Key words: Female genital mutilation (FGM), high school students, Ethiopia.

INTRODUCTION

Female genital cutting, also termed female genital mutilation (FGM), pertains to any interventions that may involve injury or removal of the female external genitalia either partial or total for any reasons other than therapeutic (World Health Organization, 2006, 2008). The practice was first known as female circumcision, but since the late 1970s this was replaced by the term female genital mutilation (FGM) to give a better reflection of the violation of the woman or girl's basic human rights (World Health Organization, 2008; UNICEF, 2005a; Emam et al., 2011). The practice is widespread in 28 African countries, concentrated around the belt of the Sahel (Wagner, 2011; World Health Organization, 2011; USAID, 2012). Besides, it transcends the aforementioned geographical patterns and similarly practiced in some parts of Middle East and Asia. The practice is also found among immigrants from these areas to Europe, North America, and Australia. Although the practice has been exercised in most developing countries of various cultures, there is no definitive evidence documenting when or why this ritual begun. Some theories suggest that FGM might have been practiced in ancient Greece, Rome, Pre-Islamic Arabia and the Russian Federation (Konjit, 2002).

According to World Health Organization (WHO), about 130 million women and girls in the world have been victims of some kind of FGM. It has also been estimated that each year about three million girls are at risk or are subjected to some kind of ablation, essentially in the 28 countries in sub-Saharan Africa, where this type of ritual has strong, ancestral roots (Mitike and Deressa, 2009; Kaplan et al., 2009). During 2005/2006, the prevalence of FGM/FGC in 27 African countries ranges from 97.9% in Somalia to 0.6% in Uganda and according to EDHS 2005, this value was 74.3% in Ethiopia and 87.2% in Oromia National Regional State (Population Reference Bureau, 2010).

In Ethiopia, if not all or completely holistic for the whole society there are traditional practices that are mostly indigenous in almost all ethnic cultures, which affect the healthy functioning of people, goals, of equality and hinder developmental strategies proposed in different walks of life time. Women and children are more prone to the problem resulted from traditional practices and as a result suffer the negative impacts in which FGM particularly is among the violent traditional action against women. Female genital mutilation (FGM) or mutilating genitals of female is not a new phenomenon and it was in existence long before the beginning of Christianity and Islam as described earlier in this article. It is a national problem because not only it does and influence the

physical, mental, and social life of more than half of Ethiopian population that is women or concerned sex, it also has its own impacts on the socio economic development of the country. Study of this type targeting on prevalence and factors associated is of remarkable input for designing very sound evidence based intervention with the desire to bring sound achievements in the area.

MATERIALS AND METHODS

Study design and population

A descriptive cross-sectional study was utilized in conducting this study during January, 2012. The study was conducted in Dale Wabera Woreda of Kelem Wollega zone, Oromia regional state. The source population were all female students who were enrolled in the two high schools and one preparatory school of the study area in the academic year 2011/2012.

Sample size and sampling procedures

The sample size for the quantitative study was estimated using the single population proportion formula with the assumption of 3% margin of error (d), 95% confidence level ($z_{\alpha/2} = 1.96$) and 87% proportion of FGM of Oromia region taken from Ethiopian Demographic and Health Survey (EDHS) 2005 (Central Statistical Agency, 2005) and using a design effect of 1.5, the calculated sample size with 10% contingency for non-response was 798.

A multi stage sampling procedure was applied to select sample of female students in the schools (stratification in to grades and simple random sampling to select sections and again simple random sampling to select respondents in respective sections).

The calculated sample size was proportionally allocated to the three schools based on the student population they have. First, each school was stratified by grades (9th, 10th, 11th, and 12th). Proportional allocation of sample was done to the respective grades. Secondly, sections were randomly selected from each grade and study subjects were selected by simple random sampling technique.

Data collection procedures

Instrument

Self-administered structured questionnaire which was adopted from Ethiopian Demographic and Health Survey (Central Statistical Agency, 2005) were used to collect data on socio-demographic, status of FGM and other important data related to the objective of the study.

Pre-test was done before the actual data collection started on 30 (4%) female students in the Kebe High School which is found in the Gawo Kebe wereda.

*Corresponding author. E-mail: latisenako@gmail.com.

Author(s) agree that this article remain permanently open access under the terms of the [Creative Commons Attribution License 4.0 International License](https://creativecommons.org/licenses/by/4.0/)

Data collection

Data collection facilitators were experienced individuals and training was given by the principal investigator for two days specially how to create conducive environment for the respondents during data collection, how to give clarity, if there is any inconvenience.

Following an orientation, respondents filled the questionnaire in private by arranging their seat far apart from one another and not allowing their teachers to enter the room.

Data analysis

Data were entered using epi data version 3.1 and exported to Statistical Package for Social Science (SPSS) version 16 for further analysis. Data cleaning took place, during data entry by use of double entry and after data entry by simple frequency and cross tabulation and their consistency was checked. After data cleaning was completed, data were exported to SPSS version 16 for analysis. Data analyses were done through Univariate analysis to see the frequencies of the categories under each variable and the corresponding percentage for each category. We also did bivariable analysis to see association between the dependent variable (in our case female genital mutilation) and independent variables like socio-demographic variables and finally logistic regression was conducted to see the independent effect of the selected independent variables on the status of FGM using odds ratio.

Operational definitions

Female genital mutilation/cutting: This is partial or total removal of the external female genitalia or other injury to the female genital organ whether for cultural or non-therapeutic reasons (World Health Organization, 2006, 2008).

Clitoridectomy: Partial or total removal of the clitoris, with or without excision of part or all of prepuce (World Health Organization, 2006, 2008).

Infibulations: Narrowing the vaginal opening through the creation of a covering seal by cutting and repositioning the labia minora and/or the labia majora, and with or without removal of the clitoris (World Health Organization, 2006, 2008).

Defibulation: Defibulation is a surgical procedure wherein a vertical incision is made on the scar to expose the introitus and create new labia majora.

RESULTS

Socio-demographic characteristics

More than half of the participants (54.2%) were seventeen years old and above with a mean and SD of 16.93 and ± 1.32 respectively. Three hundred and forty-two of the study participants, 44.5% (342) were from grade nine, 31.6% (243) were grade ten, and 14.4% (111) were from grade eleven. More than two-third of the study participants (77.1%) were from rural area and almost all of the study participants (97.9%) were Oromo in terms of their ethnic background, while protestant was the dominant religion in the area (71.3%) (Table 1).

Ninety-five of the practice was done by traditional

circumcisers and the rest was done by health professionals. Among 704 (91.5%) of the study subjects who reported that they have one or more sister(s), 63.6% of the sisters have FGM. From the total number of study subjects who reported mutilation and having sister (s), 56.4% of them have been mutilated with one or more of their sister (s) at the same time. The majority of procedure (64%) was done during night time which was followed by early morning (27.5%) and only 8.5% of it was performed at day time. About 24% of the study participants reported that FGM is required by their religion.

The majority of the respondents (77.7%) agreed with the idea of terminating the practice while 22.3% of them supported to perform FGM in the future.

Prevalence of FGM among the high school and preparatory female students, Dale Wabera, January 2012

The overall prevalence of FGM among the study participants was 77.8%; 95% CI (74.9%, 80.7%). As it is shown below the prevalence increased from 72.5%; 95% CI (69.4%, 75.6%) in grade 9 to 89%; 95% CI (87.9%, 90.1%) in grade twelve (Figure 1).

Associated factors of female genital mutilation

From those factors significantly associated with dependent variable (age, grade, residence, religion, father education, mother education, father occupation, mother occupation, and perceived monthly income) during crude analysis, four of them (age, grade, residence, and mother occupation) remained independently associated with the dependent variable on multivariate analysis.

Among the age groups, those ≥ 17 years were at higher odds of practicing FGM compared to the other age group [AOR 1.8; 95% CI (1.28, 2.6)]. The odds of FGM was higher in grades eleven and twelve students compared to grade nine [AOR=2.98, 95% CI (1.45, 6.12)] and [AOR=6.52, 95%CI (1.85, 22.94)], respectively.

Residence was one of the factors independently associated with magnitude of FGM given that the chance of being mutilated was higher in students from rural area compared to those urban residents [AOR=1.6, 95% CI (1.01, 2.52)].

The other factor independently associated with FGM was mother occupation. The odds of practicing FGM was higher among female students whose mothers were housewives and merchants compared to those with government employee mothers after adjusting for age, grade, and residence [AOR 2.38; 95% CI (1.07, 5.29)] and [AOR 2.72; 95% CI (1.04, 7.12)], respectively (Table 2).

Table 1. Socio-demographic characteristics of high school and preparatory female students of Dale Wabera woreda, January 2012, n=769.

| Variables | Number | % |
|------------------|--------------|------|
| Age | | |
| <17 | 352 | 45.8 |
| ≥17 | 417 | 54.2 |
| Mean ± SD | 16.93 ± 1.32 | |
| Grade | | |
| 9 th | 342 | 44.5 |
| 10 th | 243 | 31.6 |
| 11 th | 111 | 14.4 |
| 12 th | 73 | 9.5 |
| Residence | | |
| Rural | 593 | 77.1 |
| Urban | 176 | 22.9 |
| Religion | | |
| Protestant | 548 | 71.3 |
| Orthodox | 130 | 16.9 |
| Muslim | 91 | 11.8 |
| Ethnicity | | |
| Oromo | 753 | 97.9 |
| Non-Oromo | 16 | 2.1 |

DISCUSSION

As indicated in the study, based on data revealed from the field 77.8% (74.9%, 80.7%) of the students that underwent the procedure which is higher than the Demographic and Health Survey (DHS) (2005) country prevalence (74.3%) but lower than the Oromia regional prevalence (87.2%) (Central Statistical Agency, 2005), this lower prevalence in this study may be due to their age that in DHS those in age group of 15-49 were included and in this current study the maximum age was 23. The finding of this study is also lower than the finding of the base line survey which was 89.4% in Kelem Wollega zone, but higher than that of follow up survey which was 61.3% (EGLDAM, 2011) in the same zone. It is also lower than the finding of the study conducted in West Showa zone which was 96.4%, and the difference may be due to inclusion of higher age group (24-33) years in the former study (Oduro et al., 2006). So, if we would include higher age of the study participants the current prevalence might be higher. The prevalence is lower than the finding of the study conducted among high school female students in south Ethiopia which was 82.2% (Tamire and Molla, 2011) and the possible explanation is the difference in age at which FGM is

performed. When we compare finding of our study with some other countries, the finding of this study is higher than the finding of the study conducted among high school students in Egypt which was 50.3%(17). It is also much higher than that of primary school girls in Addis Ababa which was 25.8% (Tamire and Molla, 2011; Zewde et al., 2009) and might be due to the difference in age of the study participants included and the other may be those of Addis Ababa with more educated parents and has higher access to information.

The age at which FGM is performed on girls varies between countries and even from area to area within the same country. In most countries it is practiced on girls between four and twelve years but it can be done as early as before one year of age and in some areas it can be delayed up to just before their marriage (Population Reference Bureau, 2010; Tag-Eldin et al., 2008; Kerubo, 2010; Olenja and Kamau, 2001; Boyden et al., 2011; Kaplan et al., 2011).

According to the research conducted in Egypt, the average age of mutilation was 10.1±2.3 years (Afifi and Bothmer, 2007), while the figure was 2.9±0.65 years from the study conducted in the primary school girls in Addis Ababa (Zewde et al., 2009). The finding of this study showed that the average age of FGM is 12.95±2.23 years which is much higher than that of the study conducted in Addis Ababa and also slightly higher than the value from the Egypt. As confirmed by different studies, this age varies based on the culture of the specific population (Kaplan et al., 2011).

More than 95% of the procedure was done by traditional practitioners and the rest was done by health professionals. This finding agrees with results of most studies that FGM is mostly done by traditional practitioners commonly known by the name of the village women (World Health Organization, 2008).

Educational level of the study participants were independently associated with the prevalence of FGM especially in those grades eleven and twelve students. The odds of the practice was about three times higher in grade eleven and about 6.5 times higher in grade twelve students when compared with grade nine female students. This finding should be cautiously interpreted because one could be confused with the idea of decreasing trend of FGM as a result of increased community awareness, but the reality may be those who did not undergo FGM this year are candidates for the coming year as the community is still highly in favour of its continuation. This finding is in line with that of study conducted in Egypt and south Ethiopia which showed increased magnitude of the practice along with their grades (Njue and Askew, 2004; Stewart et al., 2002; Tag-Eldin et al., 2008).

The age of the study participants was the other variable independently associated with FGM that those seventeen years and above were with a higher odds of the practice; however, like that of the grades of the study participants,




Figure 1. Prevalence of FGM among high school female students in Dale Wabera woreda, January 2012.

the association with the age needs cautious interpretation. This also agrees with the findings of other studies that the prevalence of FGM increased with the age of the participants (Mitike and Deressa, 2009; Tamire and Molla, 2011).

Even though it is marginal, residence of the study participants was the other factor independently associated to the practice of FGM as those who were from rural areas were at higher odds of the practice and this may be due to the discrepancy in access to information about harmful effects of FGM. This finding agrees with that of high school based study in Egypt and EDHS (2005) findings in which the residence of students was independently associated with the prevalence of FGM (Central Statistical Agency, 2005; Tamire and Molla, 2011; Tag-Eldin et al., 2008).

Mothers' occupation was the other factor significantly associated to FGM. Female students whose mothers were housewives and merchants were at higher odds of practicing FGM compared to those students whose mothers were government employees even though it had no significant association in previous researches with increased prevalence of FGM (Tamire and Molla, 2011). This may be because of those mothers who were government employees that have had better access to information about harmful effects of the practice when compared to housewives mothers who spent most of their time at home and merchants who are mostly busy with their work. Although this study showed no association between magnitude of FGM and education of the parents, other studies showed the presence of significant association with parental educational status (Central Statistical Agency, 2005; Tamire and Molla, 2011; Zewde et al., 2009; Tag-Eldin et al., 2008;

UNICEF, 2005b). This may be due to fear of the challenges they would face from the community even if they have information about harmful consequences of FGM.

As a limitation, the research was conducted among high school students and is not community based and it may not reveal realistic and very precise image of the issue figure in the general community.

Conclusions

Currently most people assume that the prevalence of FGM is decreasing in most parts of the country, but this study showed that this figure is very high (77.8%) in the current study area.

Study participants with age ≥ 17 , grades 11 and 12, residing in rural area, and having housewives and merchant mothers' occupation were at higher odds of FGM. Female genital mutilation is one among traditional practices that is still ongoing across society either in severe or less form in both urban and rural areas. The practice is harmful and cause serious problem on the circumcised women and girls that may be categorized as both short term and long term pain. Community sensitization about the negative impacts of FGM should be designed through different mechanisms such as media, health personnel, community leader, religious leader, and seminar presentation, community education to increase awareness resulting in behavioral change over time. The governmental and nongovernmental organizations working towards building community or girls free from such harmful traditional practice especially in remote areas should rely on evidence based

Table 2. The association of female genital mutilation with selected variables in Dale Wabera woreda, January 2012.

| Variable | COR (95% CI) | AOR (95% CI) |
|--------------------------------|--------------------|---------------------|
| Age | | |
| <17 | 1 | 1 |
| ≥17 | 1.80 (1.29, 3.56) | 1.80 (1.28, 2.60)* |
| Grade | | |
| 9 th | 1 | 1 |
| 10 th | 1.39 (0.95, 2.05) | 1.41 (0.88, 2.26) |
| 11 th | 2.10 (1.19, 3.70) | 2.98 (1.45, 6.12)* |
| 12 th | 3.08 (1.42, 6.66) | 6.52 (1.85, 22.94)* |
| Residence | | |
| Urban | 1 | 1 |
| Rural | 1.74 (1.19, 2.54) | 1.60 (1.01, 2.52)* |
| Religion | | |
| Orthodox | 1 | 1 |
| Protestant | 1.27 (0.82, 1.97) | 1.36 (0.78, 2.37) |
| Muslim | 2.21 (1.09, 4.47) | 1.26 (0.50, 3.19) |
| Father education | | |
| Illiterate | 2.72 (1.32, 5.61) | 1.97 (0.89, 4.38) |
| Primary | 2.37 (1.18, 4.75) | 1.71 (0.79, 3.65) |
| Secondary | 1.77 (0.85, 3.69) | 1.41 (0.65, 3.05) |
| College | 1 | 1 |
| Mothers education | | |
| Illiterate | 4.05 (1.27, 12.91) | 2.96 (0.87, 10.11) |
| Primary | 3.40 (1.06, 10.90) | 2.50 (0.73, 8.49) |
| Secondary | 2.48 (0.72, 8.56) | 2.21 (0.61, 7.96) |
| College | 1 | 1 |
| Fathers' occupation | | |
| Farmer | 2.38 (1.43, 3.96) | 1.53 (0.68, 3.43) |
| Merchant | 2.02 (0.97, 4.19) | 1.56 (0.61, 3.99) |
| Government employee | 1 | 1 |
| Mothers occupation | | |
| House wife | 3.22 (1.50, 6.93) | 2.38 (1.07, 5.29)* |
| Farmer | 2.97 (1.41, 6.26) | 1.95 (0.88, 4.33) |
| Merchant | 2.85 (1.12, 7.29) | 2.72 (1.03, 7.12)* |
| Government employee | 1 | 1 |
| Perceived family income | | |
| Low | 2.29 (1.14, 4.61) | 1.94 (0.92, 4.09) |
| Medium | 1.67 (0.83, 3.34) | 1.56 (0.74, 3.30) |
| High | 1 | 1 |

Adjusted for, age, grade, residence, and religion. *= significant at p< 0.05.

interventions. Women's affairs bureaus and different gender based association should actively work starting

from regional to international level collaboratively with both governments based and non – governmental

organizations to curb the practice from the root. A community based study with stronger study design is also recommended.

CONFLICT OF INTERESTS

The authors have not declared any conflict of interests.

ACKNOWLEDGEMENTS

Authors are grateful to Addis Ababa University who funded this paper and the study participants, data collectors and supervisors for their unreserved participation.

REFERENCES

- Affif M, Bothmer M (2007). Egyptian women's attitudes and beliefs about female genital cutting and its association with childhood maltreatment. *Nurs. Health Sci.* 9:270-276.
- Boyden J, Pankhurst A, Tafere Y (2011). Harmful Traditional Practices and Child Protection: Contested Understandings and Customs of Female Early Marriage and Genital Cutting in Ethiopia. Available at: <http://www.ohchr.org/Documents/HRBodies/CEDAW/HarmfulPractices/JoBoyden-AlulaPankhurst-YisakTafere.pdf>
- Central Statistical Agency (2005). Ethiopian Demographic and Health survey. Available at: [http://www.dhsprogram.com/pubs/pdf/FR179/FR179\[23June2011\].pdf](http://www.dhsprogram.com/pubs/pdf/FR179/FR179[23June2011].pdf)
- EGLDAM (2011). Survey on Harmful Traditional practices in Ethiopia, Oromiya Regional Report. Available at: <http://www.odwace.org/research/46-harmful-traditional-practices-in-ethiopia-oromia-regional-report-august-2011.html>
- Emam EA, EL-Maghawri AM, Labeeb SA (2011). Cultural Awareness about Female Genital Mutilation among Female Employees of Minia University. *J. Am. Sci.* 7 (4):611-617.
- Kaplan A, Hechavarría S, Martín M, Bonhoure I (2011). Health consequences of female genital mutilation/cutting in the Gambia, evidence into action. *BMC Reprod. Health* 8(1):26.
- Kaplan MA, Torán MP, Moreno NJ, Fàbregas MJC, Muñoz OL (2009). Perception of primary health professionals about Female Genital Mutilation: from healthcare to intercultural competence. *BMC Health Serv. Res.* 9:11.
- Kerubo KR (2010). Female Genital Mutilation -effects on women and young girls P 31.
- Konjit K (2002). Gender based violence-an impediment to sexual and reproductive health of women. *Timar Bull.* 3:14.
- Mitike G, Deressa W (2009). Prevalence and associated factors of female genital mutilation among Somali refugees in eastern Ethiopia: a cross-sectional study. *BMC Public Health* 9:264.
- Njue C, Askew I (2004). Medicalization of Female Genital Cutting among the Abagusi in Nyanza Province, Kenya.
- Oduro AR, Ansah P, Hodgson A, Aful TM, Baiden F, Adongo P, Koram KA (2006). Trends in the prevalence of female genital mutilation and its effect on delivery outcomes in the kassena-nankana district of northern Ghana. *Ghana Med. J.* 40(3).
- Olenja J, Kamau J (2001). Olenja, J., & Kamau, J. (2001). Baseline Survey on Female Genital Mutilation Practices in Koibatek District, Rift Valley Province. Ministry of Health/GTZ FGM Project.
- Population Reference Bureau (2010). Female Genital Mutilation/Cutting: Data and Trends. Available at: <http://www.prb.org/Publications/Datasheets/2010/fgm2010.aspx>
- Stewart H, Morison L, White R (2002). Determinants Of Coital Frequency Among Married Women In Central African Republic: The Role Of Female Genital Cutting. *J. Biosoc. Sci.* 34:525-539.
- Tag-Eldin MA, Gadallah MA, Al-Tayeb MN, Abdel-Aty M, Mansour E, Sallem M (2008). Prevalence of female genital cutting among Egyptian girls. *Bull. World Health Org.* 86(4):269-274.
- Tamire M, Molla M (2011). Prevalence and believe in the continuation of Female Genital Cutting among high school girls in Hadiya zone Southern Ethiopia. *BMC Public Health* 13(1):1120.
- UNICEF (2005). Innocent Digest Web site. Changing a harmful social convention: Female genital mutilation/ cutting. Available at: https://www.unicef-irc.org/publications/pdf/fgm_eng.pdf
- UNICEF (2005b). Female Genital Mutilation/Cutting. P 40.
- USAID (2012). Gender Equality and Female Empowering Policy. P 7.
- Wagner N (2011). Violence against women: Female genital cutting a practice impossible to abandon. Available at: http://www.natascha-wagner.com/uploads/9/0/1/5/9015445/violence_against_women_fg.pdf
- World Health Organization (WHO) (2006). Department of Gender and Women's Health, Female genital mutilation: Integrating the prevention and the management of the health complications into the curricula of nursing and midwifery. *Tech. Rep.* WHO/RHR/01.17.144.
- World Health Organization (WHO) (2008). Eliminating Female Genital Mutilation: an interagency statement UNAIDS, UNDP, UNECA, UNISCO, UNFPA, UNHCHR, UNICEF, UNIFEM, WHO. Geneva, Switzerland. *Tech. Rep.* 48.
- World health organization (WHO) (2011). An update on WHO's work on female genital mutilation (FGM).
- Zewde T, Haidar J, Kogi W (2009). Magnitude of and driving factors for female genital cutting in schoolgirls in Addis Ababa, Ethiopia: A cross sectional study. *South Afr. J. Child Health* 4(3):78-82.

Full Length Research Paper

Clinical outcomes of patients admitted in intensive care units of Nigist Eleni Mohammed Memorial Hospital of Hosanna, Southern Ethiopia

Mohammed Suleiman Obsa*, Abdi Oumer Adem and Getish Bezabih Gete

Department of Anesthesia, Wolaita Sodo University School of Medicine, Wolaita, Ethiopia.

Received 25 February, 2017; Accepted 23 March, 2017

An intensive care unit is a special department of hospital or health care facility that provides intensive treatment medicine. Critically ill patients are admitted to the intensive care unit to reduce morbidity and mortality associated with acute illness, trauma or surgical procedures. This study aimed to assess clinical outcomes of patients admitted in intensive care units of Nigist Eleni Mohammed Memorial Hospital of Hosanna from January 2015 to January 2016. Institutional based cross sectional study design was conducted. All the study participants admitted to intensive care unit were included. Regular supervision and follow up was made. Data was entered into Epi info version 7 by investigators and was transported to SPSS version 20 for analysis. Bivariate and multivariate analysis was used to identify factors associated with patient's outcome. A total of 280 clients were enrolled into the study of which 46.42% died. About 26% of patients were found in the age group of 20 to 24 years and majority of them were male (58.2%). Patient with head injury is about six times more likely to die in the intensive care unit than patients with small bowel obstruction (AOR 6.620 (95% CI ((468-93.584)). There were poor outcomes of patients admitted to intensive care unit so that it was strongly recommended to improve quality of care.

Key word: Ethiopia, intensive care unit, cause of death, hospital discharge.

INTRODUCTION

Intensive care unit is a fused area of hospital where patients with acutely life threatening illness/injuries receive a specialized medical and nursing care, such as mechanical ventilation and invasive cardiac monitoring (Donaldson et al, 2000; Winter, 2013). The modern concept of intensive care was founded by anesthetist in Denmark during the polio pandemic (Berthelsen and Cronqvist, 2003). Since then, intensive care units (ICUs)

have significantly improved the quality of care and outcomes of critically ill and injured patients, mainly in high-resource settings (Calvin et al., 1997; Bleck, 2009; Grenvik and Pinsky, 2009). In recent decades, intensive care medicine has developed into highly specialized discipline covering numerous fields of medicine (Berthelsen and Cronqvist, 2003).

Admission into ICU may be required if the patient

*Corresponding author. E-mail: msuleiman43@yahoo.com.

experiences hemodynamic instability requiring frequent monitoring of vital signs, invasive hemodynamic monitoring, rapid titration of intravenous medication with concurrent monitoring. Apart from causing death, type and severity of illness can directly affect the length of ICU stay (De Lange et al., 2009). In sub-Saharan Africa, ICUs have varying qualities and quantities of infrastructure necessary for the provision of proper critical care services (Baelani et al., 2011; Dünser et al., 2006). The reported disease characteristics and mortality rates of patients admitted to ICUs in sub-Saharan Africa vary widely from one population to another (Oke, 2001; Okafor, 2009; Merah et al., 2006).

Critically ill patients are admitted to the intensive care units to reduce morbidity and mortality associated with acute illness, trauma or surgical procedures (Braber and van Zanten, 2010). Up to fifth of patients will die in the ICU (Cook, 2006). Patients that survived in the ICU were discharged to the ward environment. Although, some patients died soon after ICU discharge, such deaths are predictable and cannot be avoided (Campbell et al., 2008; Campos et al., 2011). However, another study revealed that deaths occurring after ICU discharge are unexpected and may be prevented with better standards of care (Chaboyer et al., 2008; McLaughlin et al., 2007). Despite widely analyzed pathophysiological processes and new treatment methods in laboratory and clinical research, less data are available on the causes of death, short- and long-term outcomes of critically ill patients, and associated risk factors. Frequently, data on specific predictive criteria for single diseases have been identified (Benoit et al., 2003; Wunsch et al., 2008; Afessa et al., 2002; Arabi et al., 2004). But, little is known on the exact causes of death and the influence of general risk factors that may consistently complicate the course of critically ill patients regardless of the underlying disease (Estenssoro et al., 2002; Khouli et al., 2005; Chang et al., 2006). Thus, the aim of the study was to determine clinical outcomes of patients admitted in intensive care units of Nigist Eleni Mohammed Memorial Hospital of Hosanna.

MATERIALS AND METHODS

Study setting

The study employed a cross-sectional study design at Nigist Eleni Mohammed Memorial Hospital (NEMMH) from January 2015 to January 2016. NEMMH is one of the governmental hospitals in Ethiopia which is found in Hosanna town. Hosanna town is an administrative city of Hadiya zone and it is located at a distance of 230 km to the south of Addis Ababa, capital city of Ethiopia. The hospital has 9 wards. Adult intensive care unit was considered as it has been providing services for all critically ill patients admitted from different departments excluding neonatal unit. The hospital has a total of 220 beds including 8 and 7 beds in neonatal and adult ICU, respectively.

Source population

All charts of the patients admitted in intensive care unit of Nigist

Eleni Mohammed Memorial Hospital.

Study population

Selected charts of patients admitted in intensive care unit of Nigist Eleni Mohammed Memorial Hospital from January 2015 to January 2016.

Sample size determination and sampling technique

All consecutive patients admitted in intensive care unit of Nigist Eleni Mohammed Memorial Hospital from January 2015 to January 2016.

Data collection tools and procedure

Data was collected using pretested structured questionnaires by two BSc anesthetists and supervised by one MSc holder anesthetist. Patient's charts were reviewed. At the end of data collection, patient's charts were replaced with its original place properly.

Data quality assurance

The structured questionnaire was prepared in English first and translated to the local language, Amharic and again translated back to English to ensure consistency of the questionnaire. Pretest was done on 5% of the sample population. Data collectors and supervisors were trained on each item included in the study tools, objective, relevant to study, right of respondents. During data collection, regular supervision and follow up was made. Investigator cross checked for completeness and consistency of data on a daily basis.

Data analyzing and processing

The data was entered into epi info version 7 and was exported to SPSS version 20 computer program for analysis. Descriptive statistics were used to summarize data, tables and figures for display results. Bivariate and multivariate analyses were used to see the effect of independent variable on outcome variable. Variables which were significant on bivariate analysis at p-value less than 0.2 were taken to multivariate analysis. In multivariate analysis, p-value of less than 0.05 was used as a cut-off point for presence of association. Strength of association was measured by 95% confidence interval and/or odd ratio.

Operational definitions

Clinical outcome

In this research, clinical outcome indicated either patients survived or died at the time of discharge.

Post-ICU patients

Patients who are transferred from ICU to inpatient environments.

Survived

Patients who are alive at the time of discharge.

Table 1. Distribution of sex and age group of patients admitted to intensive care unit at Nigst Eleni Mohammed Memorial Hospital, Hosanna, from January 2015 to January 2016.

| Variables | Category | Frequency | Percentage |
|-----------|-----------------|-----------|------------|
| Sex | Male | 163 | 58.2 |
| | Female | 117 | 41.8 |
| Age group | 15-19 | 61 | 21.8 |
| | 20-24 | 73 | 26.1 |
| | 25-29 | 28 | 10.0 |
| | 30-34 | 29 | 10.4 |
| | 35-39 | 24 | 8.6 |
| | 40-44 | 20 | 7.1 |
| | 45-49 | 16 | 5.7 |
| | 50-54 | 19 | 6.8 |
| | greater than 60 | 10 | 3.6 |
| Religion | Orthodox | 176 | 62.88 |
| | Protestant | 51 | 18.21 |
| | Muslim | 41 | 14.64 |
| | Missing | 7 | 2.50 |
| | Others* | 5 | 1.77 |
| Ethnicity | Hadiya | 132 | 47.14 |
| | Kembata | 63 | 22.50 |
| | Silte | 48 | 17.14 |
| | Gurage | 7 | 2.50 |
| | Missing | 18 | 6.44 |
| | Others** | 12 | 4.28 |

Others* Jehovah witness and wakefata; Others** Wolaita,Oromo and Amhara.

Died

Patients who are not alive at the time of discharge.

Ethical consideration

Ethical clearance and approval was obtained from ethical review committee, Anesthesia Department, Wolaita Soddo University. Permission to conduct was obtained from the hospital. Informed verbal consent was secured from every study participants. The obtained data was only used for study purpose. Confidentiality and anonymity were ensured.

RESULTS

A total of 280 clients were enrolled in the study of which 46.42% died in the ICU. Majority of the patients, 26.1% were between the age group of 20 and 24 years. The mean age of respondents was $31.27 \pm SD (14.019)$, (minimum 15 and maximum 67). Regarding the sex composition of the sampled respondents, about 58.2%

were male, while the remaining 41.8% were female. This means that majority of sampled respondents were male (Table 1).

Admission diagnosis to intensive care unit

Patients with different diagnosis were admitted in intensive care unit of which small bowel obstruction (SBO) was 15.4% and followed by head injury (13.9%), shock (10.4%) and CHF (10.4%). The study also found that about 4.6% of all patients were admitted with acutely exacerbated bronchial asthma and for post-surgical observation (Table 2).

Cross tabulation of level of consciousness, condition of admission and length of ICU stay over patient's outcome

The relationship between time of ICU admission and level of consciousness on survival condition showed that late admitted patients (28.2%) and unconscious patients (25.36%) died in the ICU. This showed that early admitted and conscious patients are more likely to survive than the others (Table 3).

Factors affecting clinical outcome of patients admitted in ICU

The association between factors affecting outcome of patient admitted to ICU revealed that level of consciousness, sex and length of ICU stay for more than 14 days were strongly associated with clinical outcome of patients at p-value less than 0.05. The result of the study also showed that patients with head injury is about six times more likely to die than a patients with small bowel obstruction (Table 4).

DISCUSSION

This study attempted to determine clinical outcome of patients admitted in ICU of Nigst Eleni Mohammed Memorial Hospital. According to findings of this study, relatively higher proportions of patients were found between the age group of 20 and 24 years of age. It was also found that there was male predominance which was consistent with other study (Sawe et al., 2014). A similar finding was reported by other studies of ICUs in sub Saharan Africa as there was a predominantly young-age ICU population (Okafor, 2009; Ohaegbulam et al., 2007; Mhando et al., 2008; Jamison et al., 2006). Similarly, the patient population in this study is younger when compared with patients admitted to ICUs in most developed world (Moran et al., 2008; Towey and Ojara, 2007). The overall young population and male

Table 2. Admission diagnosis of patients admitted to intensive care unit at Nigst Eleni Mohammed Memorial Hospital, Hosanna, from January 2015 to January 2016.

| Admission diagnosis | Frequency | Percentage |
|------------------------------------|-----------|------------|
| Small bowel obstruction (SBO) | 43 | 15.4 |
| Sock | 29 | 10.4 |
| Congestive heart failure (CHF) | 29 | 10.4 |
| Community acquired pneumonia (CAP) | 16 | 5.7 |
| Eclampsia | 28 | 10.0 |
| Head injury | 39 | 13.9 |
| bronchial asthma | 13 | 4.6 |
| postsurgical observation | 13 | 4.6 |
| uterine rupture | 28 | 10.0 |
| Diabetic ketoacidosis (DKA) | 28 | 10.0 |
| Postpartum hemorrhage (PPH) | 14 | 5.0 |

Table 3. Relationship between time of ICU admission, length of ICU stay and level of consciousness on survival condition of patients admitted to intensive care unit at Nigst Eleni Mohammed Memorial Hospital, Hosanna, from January 2015 to January 2016.

| Variables | Category | Died | Survived | Total |
|------------------------|-----------------------|------|----------|-------|
| Time of admission | Early admission | 51 | 45 | 96 |
| | Late admission | 79 | 105 | 184 |
| Level of consciousness | Unconscious | 71 | 96 | 167 |
| | Conscious | 59 | 54 | 113 |
| | Less than a week | 75 | 75 | 150 |
| Length of ICU stay | One week to two weeks | 35 | 51 | 86 |
| | More than two weeks | 20 | 24 | 44 |

predominance in this study may be because of high prevalence of trauma which likely occurred due to the nature of work exposing majority of males on urban streets or the increased level of participation in high-risk activities among male individuals and higher working age male predominance (Towey and Ojara, 2007).

This study also revealed that about 46.42% of patients admitted to ICU died which is higher than a study conducted in University of Nigeria Teaching Hospital on neurological and obstetric patients of which the mortality rates were 43.5 and 33%, respectively (Okafor and Onwuekwe, 2004a; Okafor and Aniebue, 2004b). But, it is lower than the study conducted at National Hospital Abuja in Nigeria on severe head injury patients in the ICU which showed the mortality rate of 68.4% (Ohaegbulam et al., 2007). Another study on critical care obstetric patients in Burkina Faso revealed a mortality rate of 60% (Dao et al., 2003), while the mortality rate in the general ICU population in Uganda was found to be 25% (Gomersall, 2010).

These differences are most likely because the present

study includes all departments of patients admitted to ICU while the others include only specific department or specific diagnosis of patients in the ICU. These differences may also be due to differences in safety and quality of health care services among different health care facilities.

Survival is the main endpoint that is considered important for patients and society (Fernandez et al., 2010). Mortality is a clinical outcome which is easy to define and measure using empirical methods and mortality following ICU discharge is a quality indicator and frequently a predicted event (Fernandez et al., 2006; Frick et al., 2003). The sudden death of post-ICU patients who are expected to survive represents a waste of valuable healthcare resources and a missed opportunity to save life. The result of present study also revealed that about 26% of patients discharged from the ICU died in the ward which is similar to some other studies (Moreno et al., 2001; Green and Edmonds, 2004). This may be due to the fact that post-ICU patients may frequently need a complex care. But, it may be difficult to provide a

Table 4. Factors affecting clinical outcome of patients admitted in ICU of Nigst Eleni Mohammed Memorial Hospital, Hosanna, from January 2015 to January 2016.

| Variables | Category | Df. | Sig. | Exp(B) | 95% C.I. for EXP(B) | |
|---------------------|--------------------------|-----------------|-------|--------|---------------------|--------|
| | | | | | Lower | Upper |
| Age category | 15-19 | 8 | 0.092 | | | |
| | 20-24 | 1 | 0.824 | 1.203 | 0.237 | 6.117 |
| | 25-29 | 1 | 0.879 | .883 | 0.177 | 4.403 |
| | 30-34 | 1 | 0.654 | 1.528 | 0.240 | 9.742 |
| | 35-39 | 1 | 0.689 | 1.483 | 0.215 | 10.220 |
| | 40-44 | 1 | 0.204 | 3.754 | 0.487 | 28.933 |
| | 45-49 | 1 | 0.610 | 2.171 | 0.110 | 42.775 |
| | 50-54 | 1 | 0.359 | 2.579 | 0.341 | 19.487 |
| | greater than 60 | 1 | 0.236 | 0.206 | 0.015 | 2.809 |
| Consciousness level | Unconscious | 1 | 0.000 | 0.279 | 0.137 | 0.567 |
| Sex | Male | 1 | 0.012 | 2.525 | 1.226 | 5.201 |
| Causes of admission | SBO | 10 | 0.710 | | | |
| | Sock | 1 | 0.911 | 1.120 | 0.154 | 8.130 |
| | CHF | 1 | 0.911 | 0.899 | 0.139 | 5.806 |
| | CAP | 1 | 0.930 | 0.903 | 0.095 | 8.604 |
| | Eclampsia | 1 | 0.614 | 0.558 | 0.058 | 5.385 |
| | Head injury | 1 | 0.162 | 6.620 | 0.468 | 93.584 |
| | Bronchial asthma | 1 | 0.351 | 0.371 | 0.046 | 2.973 |
| | Postsurgical observation | 1 | 0.946 | 1.067 | 0.164 | 6.941 |
| | Uterine rupture | 1 | 0.222 | 0.276 | 0.035 | 2.182 |
| | DKA | 1 | 0.487 | 0.536 | 0.093 | 3.104 |
| | PPH | 1 | 0.735 | 0.709 | 0.097 | 5.178 |
| | Time of admission | Early admission | 1 | 0.429 | 1.425 | 0.592 |
| Length of ICU stay | less than 7days | 2 | 0.104 | | | |
| | 7-14days | 1 | 0.277 | 2.084 | .555 | 7.828 |
| | more than 14days | 1 | 0.039 | 4.113 | 1.074 | 15.761 |

complex care in the ward environment. (NICE, 2007). This may also be because inexperienced nurses and doctors struggle to provide the necessary complex care (Endacott et al., 2007). This may also be due to intensive care staff that did not closely follow patients' progress on the ward for a few days to monitor recovery of multisystem disease and assure good continuity of care. Limitations of this study were relative scarcity of clinical data available on the patient's chart and lack of some socio demographic variables as the data was collected retrospectively.

Conclusion

Intensive care units provides patients with severe and life-threatening illnesses and injuries, which require

constant, close monitoring and support from specialist equipment and medications in order to ensure normal bodily functions. Majority of ICU patients in Nigst Eleni Mohammed Memorial Hospital were male. Early admitted and conscious patients are more likely survive than the others. Over two fifth of the patients admitted to ICU died. There may be a limited infrastructure, personnel, and resource which likely contribute to high mortality rates. Therefore, it is very important to have well organized and adequate infrastructures, personnel and resources to provide optimal care for critically ill patients.

ABBREVIATIONS

ICU, Intensive care unit; **JUSH**, Jimma University Hospital; **NEMMH**, Nigst Eleni Mohammed Memorial

Hospital; SSA, sub-Saharan Africa.

CONFLICT OF INTERESTS

The authors have not declared any conflict of interests.

ACKNOWLEDGEMENTS

The authors thank all the staff in the ICUs and record keeper for their support during data collection. They also thank the authors of articles, and on-line information for the valuable works cited in this paper.

REFERENCES

- Afessa B, Morales IJ, Scanlon PD, Peters SG (2002). Prognostic factors, clinical course, and hospital outcome of patients with chronic obstructive pulmonary disease admitted to an intensive care unit for acute respiratory failure. *Crit. Care Med.* 30:1610-1615.
- Arabi Y, Ahmed QA, Haddad S, Aljumah A, Al-shimemeri A (2004). Outcome predictors of cirrhosis patients admitted to the intensive care unit. *European J. Gastroenterol. Hepatol.* 16:333-339.
- Baelani I, Jochberger S, Laimer T, Otieno D, Kabutu J, Wilson I, Baker T, Dünser MW (2011). Availability of critical care resources to treat patients with severe sepsis or septic shock in Africa: a self-reported, continent-wide survey of anaesthesia providers. *Critical Care*, 15, R10.
- Benoit DD, Vandewoude KH, Decruyenaere JM, Hoste EA, Colardyn FA (2003). Outcome and early prognostic indicators in patients with a hematologic malignancy admitted to the intensive care unit for a life-threatening complication. *Critical care medicine*, 31:104-112.
- Berthelsen P, Cronqvist M (2003). The first intensive care unit in the world: Copenhagen 1953. *Acta Anaesthesiologica Scandinavica*, 47:1190-1195.
- Bleck TP (2009). Historical aspects of critical care and the nervous system. *Crit. Care Clin.* 25:153-164.
- Braber A, Van zanten AR (2010). Unravelling post-ICU mortality: predictors and causes of death. *Euro. J. Anaesthesiol.* 27:486-490.
- Calvin JE, Habet K, Parrillo JE (1997). *Critical care in the United States*. *Crit. Care Clin.* 13:363-376.
- Campbell AJ, Cook JA, Adey G, Cuthbertson B (2008). Predicting death and readmission after intensive care discharge.
- Campos RA, Tena SA, Muncharaz AB, Font SM, Sellés AF, Campos LM, Deleón belmar J (2011). Study of post-ICU mortality during 4 years (2006–2009). Analysis of the factors related to death in the ward after discharge from the ICU. *Medicina Intensiva (English Edition)*, 35:150-156.
- Chaboyer W, Thalib L, Foster M, Ball C, Richards B (2008). Predictors of adverse events in patients after discharge from the intensive care unit. *Am. J. Crit. Care* 17:255-263.
- Chang L, Horng CF, Huang YCT, Hsieh YY (2006). Prognostic accuracy of Acute Physiology and Chronic Health Evaluation II scores in critically ill cancer patients. *American Journal of Critical Care*, 15, 47-53.
- Cook D (2006). Methods to assess performance of models estimating risk of death in intensive care patients: a review. *Anaesthesia Intensive Care* 34:164.
- Dao, B, Rouamba A, Ouedraogo D, Kambou T, Bazié A (2003). Transfer of obstetric patients in a pregnant and postpartum condition to an intensive care unit: eighty-two cases in Burkina Faso. *Gynecologie Obstetrique Fertilité* 31:123-126.
- De lange D, Dusseljee J, Brinkman S, Van berkel, G, Van maanen, R, Bosman, R, Joore H, DE Keizer, N, Van der Voort P, Dewaal R (2009). Severity of illness and outcome in ICU patients in the Netherlands: results from the NICE registry 2006-2007. *Neth. J. Crit. Care* 13:16-22.
- Donaldson MS, Corrigan JM, Kohn LT (2000). *To err is human: building a safer health system*, National Academies Press.
- Dünser MW, Baelani I, Ganbold L (2006). A review and analysis of intensive care medicine in the least developed countries. *Crit.Care Med.* 34:1234-1242.
- Endacott R, Kidd T, Chaboyer W, Edington J (2007). Recognition and communication of patient deterioration in a regional hospital: a multi-methods study. *Australian Crit. Care* 20:100-105.
- Estenssoro E, Dubin A, Laffaire E, Canales H, Sáenz G, Moseinco M, Pozo M, Gómez A, Baredes N, Jannello G (2002). Incidence, clinical course, and outcome in 217 patients with acute respiratory distress syndrome. *Crit. Care Med.* 30:2450-2456.
- Fernandez R, Baigorri F, Navarro G, Artigas A (2006). A modified McCabe score for stratification of patients after intensive care unit discharge: the Sabadell score. *Crit. Care*10:R179.
- Fernandez R, Serrano JM, Umaram I, Abizanda R, Carrillo A, Lopez-pueyo MJ, Rascado P, Balerdi B, Suberviola B, Hernandez G (2010). Ward mortality after ICU discharge: a multicenter validation of the Sabadell score. *Intensive Care Med.* 36:1196-1201.
- Frick S, Uehlinger DE, Zenklusen RMZ (2003). Medical futility: predicting outcome of intensive care unit patients by nurses and doctors-a prospective comparative study. *Crit. Care Med.* 31:456-461.
- Gomersall CD (2010). Critical care in the developing world-a challenge for us all. *Crit. Care* 14:131.
- Green A, Edmonds L (2004). Bridging the gap between the intensive care unit and general wards-the ICU Liaison Nurse. *Intensive Crit. Care Nurs.* 20:133-143.
- Grenvik A, Pinsky MR (2009). Evolution of the intensive care unit as a clinical center and critical care medicine as a discipline. *Crit. Care Clin.* 25:239-250.
- Jamison DT, Feachem RG, Makgoba MW, Bos ER, Baingana FK, Hofman KJ, Rogo KO (2006). *Disease and mortality in sub-Saharan Africa*, World Bank Washington, DC.
- Khouli H, Afrasiabi A, Shibli M, Hajal R, Barrett CR, Homel P (2005). Outcome of critically ill human immunodeficiency virus-infected patients in the era of highly active antiretroviral therapy. *J. Intensive Care Med.* 20:279-285.
- Mclaughlin N, Leslie G, Williams T, Dobb G (2007). Examining the occurrence of adverse events within 72 hours of discharge from the intensive care unit. *Anaesth. Intensive Care* 35:486.
- Merah N, Okeke C, Olatosi J (2006). An audit of surgical admissions to the intensive care unit of the Lagos University Teaching Hospital (1997-2002). *Niger. Postgraduate Med. J.* 13:153-156.
- Mhando S, Young B, Lakhoo K, (2008). The scope of emergency paediatric surgery in Tanzania. *Pediatr. Surg. Int.* 24:219-222.
- Moran JL, Bristow P, Solomon PJ, George C, Hart GK, Australian & Committee, NZICSDM(2008). Mortality and length-of-stay outcomes, 1993–2003, in the binational Australian and New Zealand intensive care adult patient database. *Crit. Care Med.* 36:46-61.
- Moreno R, Miranda D, Matos R, Fevereiro, T (2001). Mortality after discharge from intensive care: the impact of organ system failure and nursing workload use at discharge. *Int. Care Med.* 27:999-1004.
- NICE, CF CPA (2007). Acutely ill patients in hospital: Recognition of and response to acute illness in adults in hospital.
- Ohaegbulam S, Okafor U, Ihekire O, Elumelu E (2007). Using the Revised Trauma Score to Predict Outcome in Severely Head Injured Patients in a Developing Nation—A Pilot Study. *Int. J. Med. Health Dev.* 12:51-55.
- Okafor U (2009) Challenges in critical care services in Sub-Saharan Africa: perspectives from Nigeria. *Indian J. Crit. Care Med.* 13:25.
- Okafor U, Aniebue U (2004a). Admission pattern and outcome in critical care obstetric patients. *Int. J. Obstetr. Anesth.* 13:164-166.
- Okafor U, Onwuekwe I (2004b). Disease patterns and outcome for medical neurological patients admitted to a multi-disciplinary intensive care unit. *Int. J. Med. Health Dev.* 9:113-115.
- Oke D (2001). Medical admission into the intensive care unit (ICU) of the Lagos University Teaching Hospital. *Niger. Postgraduate Med. J.* 8:179-182.
- Sawe HR, Mfinanga JA, Lidenge SJ, Mpondo BC, Msangi S, Lugazia E, Mwafongo V, Runyon MS, Reynolds TA (2014). Disease patterns and

- clinical outcomes of patients admitted in intensive care units of tertiary referral hospitals of Tanzania. *BMC Int. Health Human Rights* 14:26.
- Towey R, Ojara S (2007). Intensive care in the developing world. *Anaesthesia* 62:32-37.
- Winter RE (2013). *Unraveling US Health Care: A Personal Guide*, Rowman & Littlefield Publishers.
- Wunsch H, Angus DC, Harrison DA, Collange O, Fowler R, Hoste, EA, Dekeizer NF, Kersten A, Linde-zwirble WT, Sandiumenge A (2008). Variation in critical care services across North America and Western Europe. *Crit. Care Med.* 36:2787-e8.



International Journal of Medicine and Medical Sciences

Related Journals Published by Academic Journals

- *Journal of Medicinal Plant Research*
- *African Journal of Pharmacy and Pharmacology*
- *Journal of Dentistry and Oral Hygiene*
- *International Journal of Nursing and Midwifery*
- *Journal of Parasitology and Vector Biology*
- *Journal of Pharmacognosy and Phytotherapy*
- *Journal of Toxicology and Environmental Health Sciences*

academicJournals