

A close-up photograph of a microscope with a blue and purple color scheme. The background is a blurred bokeh of purple and blue light. The microscope's objective lenses and eyepiece are visible, with a dark semi-transparent overlay at the top containing text.

Scientific Research and Essays

Volume 12 Number 8 30 April, 2017

ISSN 1992-2248



*Academic
Journals*

ABOUT SRE

The Scientific Research and Essays (SRE) is published twice monthly (one volume per year) by Academic Journals.

Scientific Research and Essays (SRE) is an open access journal with the objective of publishing quality research articles in science, medicine, agriculture and engineering such as Nanotechnology, Climate Change and Global Warming, Air Pollution Management and Electronics etc. All papers published by SRE are blind peer reviewed.

Contact Us

Editorial Office: sre@academicjournals.org

Help Desk: helpdesk@academicjournals.org

Website: <http://www.academicjournals.org/journal/SRE>

Submit manuscript online <http://ms.academicjournals.me/>.

Editors

Dr. NJ Tonukari

Editor-in-Chief

Scientific Research and Essays

Academic Journals

E-mail:sre.research.journal@gmail.com

Dr. M. Sivakumar Ph.D. (Tech).

Associate Professor

School of Chemical & Environmental Engineering

Faculty of Engineering

University of Nottingham

JalanBraga, 43500 Semenyih

SelangorDarul Ehsan Malaysia.

Prof. N. Mohamed ElSawi Mahmoud

Department of Biochemistry, Faculty of science, King

Abdul Aziz university,

Saudi Arabia.

Prof. Ali Delice

Science and Mathematics Education Department, Atatürk

Faculty of Education,

Marmara University,Turkey.

Prof. Mira Grdisa

RudjerBoskovicInstitute,BijenickaCesta54,

Croatia.

Prof. Emmanuel HalaKwon-Ndung

Nasarawa State University Keffi Nigeria

PMB1022 Keffi,

Nasarawa State.

Nigeria.

Dr. Cyrus Azimi

Department of Genetics, Cancer Research Center,

CancerInstitute,Tehran University of Medical Sciences,

Keshavarz Blvd.,

Tehran,Iran.

Dr. Gomez, Nidia Noemi

National University of San Luis,

Faculty of Chemistry, Biochemistry and Pharmacy,

Laboratory of Molecular Biochemistry

EjercitodelosAndes950-5700 SanLuis

Argentina.

Prof.M.Nageeb Rashed

Chemistry Department-Faculty of Science,

Aswan South Valley University,

Egypt.

Dr. John W. Gichuki

KenyaMarine& FisheriesResearchInstitute,

Kenya.

Dr. Wong Leong Sing

Department of Civil Engineering,

College of Engineering,

Universiti Tenaga Nasional,

Km7, JalanKajang-Puchong,

43009Kajang, SelangorDarulEhsan, Malaysia.

Prof. Xianyi Li

College of Mathematics and Computational Science

Shenzhen University Guangdong,518060

P.R.China.

Prof. Mevlut Dogan

Kocatepe University, Science Faculty, Physics Dept.

Afyon/Turkey.

Turkey.

Prof. Kwai-Lin Thong

Microbiology Division, Institute of Biological Science,

Faculty of Science, University of Malaya, 50603,

KualaLumpur,

Malaysia.

Prof. Xiacong He

Faculty of Mechanical and Electrical Engineering, Kunming

University of Science and Technology, 253 XueFu Road,

Kunming,

P.R.China.

Prof. Sanjay Misra

Department of Computer Engineering

School of Information and Communication Technology

Federal University of Technology, Minna,

Nigeria.

Prof. Burtram C. Fielding Pr. Sci. Nat.

Department of Medical BioSciences

University of the Western Cape Private Bag X17

Modderdam Road

Bellville,7535,SouthAfrica.

Prof.Naqib Ullah Khan

Department of Plant Breeding and Genetics

NWFP Agricultural University Peshawar 25130,

Pakistan

Editorial Board

Prof. Ahmed M. Soliman

*20MansourMohamedSt.,Apt51,Zamalek,
Cairo,
Egypt.*

Prof. JuanJosé Kasper Zubillaga

*Av.Universidad1953Ed.13depto304,
MéxicoD.F.04340,
México.*

Prof. ChauKwok-wing

*University of Queensland Instituto
Mexicanodel Petroleo, Eje Central
Lazaro Cardenas Mexico D.F.,
Mexico.*

Prof. Raj Senani

*Netaji Subhas Institute of Technology,
Azad Hind Fauj Marg, Sector3,
Dwarka, New Delhi 110075, India.*

Prof. RobinJ Law

*CefasBurnham Laboratory,
Remembrance Avenue Burnhamon Crouch, Essex
CM08HA,
UK.*

Prof. V. Sundarapandian

*IndianInstitute of Information Technologyand
Management-Kerala
Park Centre,
Technopark Campus,Kariavattom P.O.,
Thiruvananthapuram-695581,Kerala,India.*

Prof. Tzung-PeiHong

*Department of Electrical Engineering,
Andat the Department of Computer Science and
Information Engineering
NationalUniversity ofKaohsiung.*

Prof.Zulfiqar Ahmed

*Department of Earth Sciences, box5070,
Kfupm, dhahran-31261, SaudiArabia.*

Prof. Khalifa Saif Al-Jabri

*Department of Civil and Architectural Engineering
College of Engineering, Sultan
Qaboos University
P.O.Box33,Al-Khod123,Muscat.*

Prof. V.Sundarapandian

*Indian Institute of Information Technology &
Management-Kerala
Park Centre,
Technopark,Kariavattom P.O.
Thiruvananthapuram-
695581,KeralaIndia.*

Prof. Thangavelu Perianan

*Department of Mathematics,
Aditanar College,
Tiruchendur-628216India.*

Prof. Yan-zePeng

*Department of Mathematics,
Huazhong University of Science and
Technology,Wuhan430074,P.R.
China.*

Prof. KonstantinosD.Karamanos

*Universite Librede Bruxelles,
CP231 Centre of Nonlinear
Phenomena And Complexsystems,
CENOLIBoulevarddeTriomphe
B-1050,
Brussels,
Belgium.*

Prof. XianyiLi

*School of Mathematics and Physics,
Nanhua University, Hengyang City,
Hunan Province,
P.R.China.*

Dr. K.W.Chau

*HongKong Polytechnic University
Department of Civil & Structural
Engineering, HongKong Polytechnic
University, Hungghom, Kowloon,
HongKong,
China.*

Dr. AmadouGaye

*LPAO-SF/ESPPoBox5085Dakar-FannSENEGAL
University Cheikh Anta Diop
Dakar SENEGAL.*

Prof. MasnoGinting

*P2F-LIPI,Puspiptek-Serpong,
15310 Indonesian Institute of Sciences,
Banten-Indonesia.*

Dr. Ezekiel Olukayode Idowu

*Department of Agricultural Economics,
Obafemi Awolowo University, Ife-Ife,
Nigeria.*

Scientific Research and Essays

Table of Contents: Volume 12 Number 8 30 April, 2017

ARTICLE

- Cytotoxicity of aerial parts of *Indigofera heterantha*** 77
Taj Ur Rahman, Wajiha Liaqat, Khanzadi Fatima Khattak,
Muhammad Iqbal Choudhary, Atif Kamil and
Muhammad Aurang Zeb

Full Length Research Paper

Cytotoxicity of aerial parts of *Indigofera heterantha***Taj Ur Rahman^{1*}, Wajiha Liaqat², Khanzadi Fatima Khattak³, Muhammad Iqbal Choudhary⁴, Atif Kamil⁵ and Muhammad Aurang Zeb⁶**¹Department of Chemistry, Mohi-Ud-Din Islamic University, AJ&K, Pakistan.²Institute of Chemical Sciences, University of Peshawar, Peshawar 25120, K. P. K, Pakistan.³Women University, Swabi, Pakistan.⁴International Center for Chemical and Biological Sciences, H. E. J. Research Institute of Chemistry, University of Karachi, Karachi-75270, Pakistan.⁵Department of Biotechnology Abdul Wali Khan University Mardan, Pakistan.⁶Department of Biochemistry Hazara University, Mansehra, Pakistan.

Received 22 January, 2014; Accepted 5 May, 2016

In the ongoing phytochemical study, an effort was made to investigate cytotoxicity of various crude fractions of aerial parts of *Indigofera heterantha*. The results obtained revealed that all the fractions including *n*-hexane, ethyl acetate, methanol and residue showed brine shrimp (*Artemia salina* Leach) cytotoxicity activity. The data obtained revealed the medicinal importance of the plant and will help the researchers to exploit the phytochemicals for biological activities (cytotoxicity).

Key words: *Indigofera heterantha*, cytotoxicity, aerial parts.

INTRODUCTION

The genus *Indigofera* consists of about 700 species. All species are herbs or shrubs distributed throughout the tropical regions of the globe. In Pakistan, this genus is represented by 24 species. *Indigofera heterantha*, commonly known as ghorega belongs to family leguminosae (Fabeaceae) (Hamayun et al., 2003). This plant is abundantly available in the northern regions of Pakistan. The plant is widely used as traditional medicine for treatment against abdominal pain, spastic pain, skin problems and infectious diseases (Nasir et al., 1997). Some species of genus *Indigofera* such *Indigofera dalecoides* Benth is used for the treatment of diarrhea and pathogenic bacterial infections (Mathabe et al., 2006). *Indigofera oblogifolia* possess antimicrobial (Dahot, 1999), hepatoprotective (Shahjahan et al., 2005) and

lipoxygenase inhibitory activity (Sharif et al., 2005). The species *Indigofera pulchra* has shown snake-venom neutralizing activity (Abubakar et al., 2006). *Indigofera tinctoria* showed antioxidant, free radical scavenging activity and anti-dyslipidemic activity (Prakash et al., 2007; Waako et al., 2007; Puri et al., 2007). *Indigofera mysorensis* has shown anti-diabetic activity. *Indigofera emarginella* have *in-vitro* antimalarial activity against *Plasmodium falciparum* (Chakrabarti et al., 2006). The leaves and flowers of *Indigofera aspalathoides* have cooling effect and used to treat leprosy and cancerous diseases. The leaves of this plant are also applied to abscesses and roots are used to treat dentifrice and mouth ulcers. The roots are chewed in for toothache and apathies. The oil extracted from whole plant, is used for

*Corresponding author. E-mail: taj_urrehman81@yahoo.co.uk.

Table 1. Brine shrimp cytotoxic activities of *n*-hexane fraction of aerial parts of *I. heterantha*.

Dose ($\mu\text{g/ml}$)	No. of shrimps	No. of survivors	LD50 ($\mu\text{g/ml}$)	Std. drug	LD50 ($\mu\text{g/ml}$)
1000	30	16	1362.178	Etoposide	7.4625
100	30	24			
10	30	28			

dandruff (Nadkarni, 1996), syphilis and other skin infectious diseases. The plant is widely prescribed for psoriasis, eczema, burns, boils, ulcers, wounds and used also as antidote to snake venom (Kirtikar and Basu 1975) in Siddha system of medicine. Water soluble fraction of alcoholic extract of dried tender shoots of *I. aspalathoides* showed significant anti-inflammatory effect in experimental albino mice. Some new compounds which includes new ester, new indigoferamide-A and new isoflavone along with three new source compounds dotriacontanoic acid, quercetin and formononetin "4-hydroxy-4-methyl-2-pentanone" were isolated from the seeds of *I. heterantha* Wall exhibiting (Rahman et al., 2014, 2015). The aim of this study was to explore the cytotoxicity of the aerial parts of *I. heterantha* to determine the scientific basis for its use as folk medicine to treat various diseases. Brine shrimps (*A. salina* Leach) lethality bioassay was done so as to assess its safety.

MATERIALS AND METHODS

Plant

The aerial parts of *I. heterantha* were collected during the month of May 2009 from Dir Lower, northern parts of Pakistan. The plant was identified and authenticated by Mr. Samin Jan, Associate Professor, Department of Botany, Islamia College, Peshawar, KPK province, Pakistan. A voucher specimen (No.Sj-36) was deposited in the herbarium of Botany Department, Islamia College, Peshawar Pakistan.

Extraction and isolation

The powdered aerial parts (60 kg) were soaked (cold extraction) in water-methanol (1:19) for seven days. The crude water-methanol extract was filtered and concentrated at reduce pressure using rotary evaporator at 50°C, afforded a crude semi solid mass of 4.7 kg F1. It was then dissolved in chloroform resulting into 87 g soluble fraction F2 and remaining insoluble fraction F3 (370 g). The chloroform soluble fraction F2 was further fractionated with *n*-hexane and methanol afforded 9 g F2-X and 73 g F2-Y crude extracts, respectively using Soxhlet extractor for one day. While the insoluble fraction F3 was further dissolved in ethyl acetate and concentration afforded 270 g crude fraction F4. The ethyl acetate soluble fraction F4 was further partitioned between diethyl ether and water gave 75 g ethereal crude fraction F4-Z, then water fraction (103 g) F4-W and insoluble residue (70 g) F4-I.

Cytotoxic assay

In this bioassay, a shallow rectangular plastic dish (22 x 32 cm)

was filled with artificial sea water. The sea water was prepared by taking commercial salt mixture with double distilled water (3.8 g) in 1:1 ratio. The brine shrimp (*A. salina*) eggs were hatched in the dish. The dish was made unequally partitioned by using an artificial perforated device. About 50 mg of the eggs were sprinkled into large compartment which becomes darken. The minor compartment was exposed to the ordinary light. After two days, nauplii were collected and removed by a pipette from lighted side. A sample of the compounds to be tested was prepared by dissolving 20 mg of each compound in dimethylformamide (DMF) an organic compound with the formula $(\text{CH}_3)_2\text{NCH}$ (2 ml) manufactured by china. Three different stock solution, that is, 550, 50, and 5 mg/ml were transferred to 9 vials (three for every dilution were used for each test sample and LD₅₀ is the average of the three values) with one vial containing DMF was reserved as a control. The solvent was allowed to evaporate by keeping it overnight. Two days, later when the shrimp larvae were ready, 1 ml of sea water and 10 shrimp were added to each vial (30 shrimps/dilution) with a volume adjusted with sea water to 5 ml per vial. After 24 h, the numbers of survivors were counted using standard procedure (McLaughlin et al., 1991; Meyer et al., 1982). The data was analyzed by the use of finny computer program to determine LD₅₀ values (Finney, 1971; McLaughlin et al., 1991; Meyer et al., 1982; Uddin et al., 2011).

RESULTS AND DISCUSSION

Cytotoxicity activity

The result obtained indicated that *n*-hexane fraction showed high activities at 10 $\mu\text{g/ml}$, moderate activities at 100 $\mu\text{g/ml}$ and low activities at 1000 $\mu\text{g/ml}$. Similarly, the ethyl acetate fraction showed activities in the same pattern as mentioned earlier that from high to low at 10, 100 and 1000 $\mu\text{g/ml}$, respectively. Methanol's fractions F1 and F2-X revealed high activities at 10 $\mu\text{g/ml}$, moderate activities at 100 $\mu\text{g/ml}$ and low activities at 1000 $\mu\text{g/ml}$. The residue fractions showed low cytotoxic activities at 1000 $\mu\text{g/ml}$, moderate at 100 $\mu\text{g/ml}$ and high at 10 $\mu\text{g/ml}$.

Tables 1 to 5 show that the active component display mostly high activity at low concentration and low activity at high concentration. The current investigation strongly supported the traditional use of *I. heterantha* for various pathological conditions. The current study proved that the extract fractions of *I. heterantha* have good cytotoxic activity. These activities may be accredited to the existence of alkaloids, polyphenols, phenols, tannins, saponins, anthraquinones, steroids and especially the diterpenes, found in the crude extract and the fractions thereof (Araruna and Carlos, 2010). The plant is commonly used for the treatment of various diseases.

Table 2. Brine shrimp cytotoxic activities of ethyl acetate fraction of aerial parts of *I. heterantha*.

Dose ($\mu\text{g/ml}$)	No. of shrimps	No. of survivors	LD50 ($\mu\text{g/ml}$)	Std. drug	LD50 ($\mu\text{g/ml}$)
1000	30	16	1500.66	Etoposide	7.4625
100	30	22			
10	30	26			

Table 3. Brine shrimp cytotoxic activities of water methanol fraction F1 of aerial parts of *I. heterantha*.

Dose ($\mu\text{g/ml}$)	No. of shrimps	No. of survivors	LD50 ($\mu\text{g/ml}$)	Std. drug	LD50 ($\mu\text{g/ml}$)
1000	30	16	2225.34	Etoposide	7.4625
100	30	22			
10	30	27			

Table 4. Brine shrimp cytotoxic activities of methanol fraction F2-X of aerial parts of *I. heterantha*.

Dose ($\mu\text{g/ml}$)	No. of shrimps	No. of survivors	LD50 ($\mu\text{g/ml}$)	Std. drug	LD50 ($\mu\text{g/ml}$)
1000	30	17	2225.63	Etoposide	7.4625
100	30	22			
10	30	26			

Table 5. Brine shrimp cytotoxic activities of residue fraction of aerial parts of *I. heterantha*.

Dose ($\mu\text{g/ml}$)	No. of shrimps	No. of survivors	LD50 ($\mu\text{g/ml}$)	Std. drug	LD50 ($\mu\text{g/ml}$)
1000	30	18	61	Etoposide	7.4625
100	30	23			
10	30	25			

This data strongly supports that widespread research should be steered to isolate and identify phytochemical constituents responsible for cytotoxic activities to intricate the hidden medicinal potential of the plant (Rahman et al., 2016).

Conclusion

In this study, an effort was made to scientifically authenticate cytotoxic bioassay of the crude extract fractions of aerial parts of *I. heterantha* to elaborate its medicinal importance. The results obtained discloses that it contains a variety of therapeutic agents, which need to be further explored phytochemically. The current study revealed that the aerial parts of the plant have strong cytotoxic activity. Further scientific investigation is needed to explore phytochemicals of this plant, which possess the activity.

CONFLICT OF INTERESTS

The authors have not declared any conflict of interests.

REFERENCES

- Abubakar MS, Balogun E, Abdurrahman EM, Nok AJ, Shok M, Mohammed MG (2006). Ethnomedical treatment of poisonous snakebites: Plant extract neutralized Najanigracollis Venom. *Pharm. Biol.* 9:348-353.
- Araruna K, Carlos B (2010). Anti-inflammatory activities of triterpene lactones from *Lactuca sativa*. *Phytopharmacol.* 1:1-6.
- Chakrabarti R, Damarla RKB, Mullangi R, Sharma VM, Vikramadithyan RK, Rajagopalan R. (2006) Insulin Sensitizing property of *Indigofera mysorensis* extract. *J. Ethnopharmacol.* 10:102-107.
- Dahot MU (1999). Antibacterial and antifungal activity of small protein of *Indigofera oblongifolia* Leaves. *J. EthnoPharmacol.* 14:277-281.
- Finney DJ (1971). *Probit analysis* 3rd ed. Cambridge: Cambridge University Press.
- Hamayun M, Khan A, Khan MA (2003). Common medicinal folk recipes of District Buner, NWFP, Pakistan. *Ethnobot. Leaflets* 1:14.
- Kirtikar KR, Basu BD (1975). *Indian Medicinal plants*, International Book Distributors, Dehra Dun, India. 6:710-713.
- Mathabe MC, Nikolova RV, Lall N, Nyazema NZ (2006). Antibacterial activities of medicinal plants Used for the Treatment of diarrhoea in Limpopo Province, South Afr. *J. Ethno. Pharmacol.* 7:286-289.
- McLaughlin JL, Chang CJ, Smith DL (1991). Bench-top bioassays for the discovery of bioactive natural products: an update. *Studies in natural products chemistry* 9:383-409.
- Meyer BN, Ferrigni NR, Putnam JE, Jacobsen LB, Nichols DE, McLaughlin JL. Brine shrimp (1982). A convenient General bioassay for active plant constituents. *PlantaMedica.* 7:34-39.

- Nasir E, Ali SI (1997). Flora of West Pakistan, Rawalpindi: National Herbarium Agricultural Research Council. 8:83-87.
- Nadkarni KM (1996). Indian Materia Medica, Mumbai: Popular Prakashan Private Limited 3rd edition. 3:1677-1681.
- Prakash D, Suri S, Upadhyay G, Singh BN (2007). Total phenol, antioxidant and free radical scavenging activities of some medicinal plants. Int. J. Food Sci. Nutri. 15:28-33.
- Puri A, Khaliq T, Rajendran SM, Bhatia G, Chandra R, Narender T (2007). Antidyslipidemic activity of *Indigofera tinctoria*. J. Herb Pharmacother. 5:64-70.
- Shahjahan M, Vani G, Devi CS (2005). Protective effect of *Indigofera oblongifolia* in CCl₄-Induced Hepatotoxicity. J. Med. Food. 8:261-266.
- Sharif A, Ahmed E, Malik A, Riaz N, Afza N, Nawaz SA, Arshad M, Choudhary MI (2005). Lipoxygenase Inhibitory constituents from *Indigofera oblongifolia*. Arch. Pharmacol Res. 7:764-770.
- Rahman TU, Arfan M, Liaqat W, Uddin G, Choudhary MI (2014). Rahman TU, Arfan M, Liaqat W, Uddin G, Choudhary MI (2014). Isolation of a Novel Indigoferamide-A from Seeds of *Indigofera heterantha* Wall and its Antibacterial Activity. Rec. Nat. Prod. 8(4):412-416.
- Rahman TU, Khattak KF, Wajaha L, Choudhary MI (2016). Antifungal, insecticidal, cytotoxic and phytotoxic activities of the crude extracts of *Taxus wallichiana* Zucc twigs J. Chem. Pharm. Res. 8:398-402.
- Rahman TU, Uddin G, Khattak KF, Liaqat W, Mohammad G, Choudhary MI, Wadood A, Ahmad A (2014). Isolation and characterization of a novel ester from seeds of *Indigofera heterantha* (Wall). J. Nat. Prod. 7:104-112.
- Rahman TU, Uddin G, Nisa RU, Ludwig R, Liaqat W, Mahmood T, Mohammad G, Choudhary MI, Ayub K (2015). Spectroscopic and density functional theory studies of 7-hydroxy-30-methoxyisoflavone, A new isoflavone from the seeds of *Indigofera heterantha* (Wall), Spectrochim Acta A: Mol. Biomol. Spectrosc. 148:375-381.
- Rahman TU, Khattak KF, Liaqat W, Choudhary MI (2016). Antibacterial, antifungal, insecticidal, cytotoxic and phytotoxic activities of the crude extracts of *Taxus wallichiana* Zucc twigs. J. Chem. Pharm. Res. 8:398-402.
- Uddin G, Rehman TU, Arfan M, Liaquat W (2011). Phytochemical and Biological screening of the seeds of *Indigofera heterantha* wall. Middle-East J. Sci. Res. 8(11):186-190.
- Waako PJ, Katuura E, Smith P, Folb P (2007). East African medicinal plants as a source of lead compounds for the development of new antimalarial drugs. Afr. J. Ecol. 13:106-110.

A close-up, artistic photograph of a microscope with a blue and purple color palette. The microscope is the central focus, with its various lenses and mechanical parts visible. The background is a soft, out-of-focus glow of blue and purple light.

Scientific Research and Essays

Related Journals Published by Academic Journals

- African Journal of Mathematics and Computer Science Research
- International Journal of Physical Sciences
- Journal of Oceanography and Marine Science
- International Journal of Peace and Development Studies
- International NGO Journal

academicJournals