

A close-up photograph of a hand gently holding a baby's foot. The hand is positioned on the left, with fingers wrapped around the foot. The foot is the central focus, showing the texture of the skin and the shape of the toes. The background is a soft, out-of-focus purple or magenta color. The entire image is framed with rounded corners.

International Journal of Nursing and Midwifery

Volume 9 Number 8, August 2017

ISSN 2141-2456



*Academic
Journals*

ABOUT IJNM

The **International Journal of Nursing and Midwifery (IJNM)** is published monthly (one volume per year) by Academic Journals.

International Journal of Nursing and Midwifery (IJNM) is an open access journal that provides rapid publication (monthly) of articles in all areas of the subject such as family practice, women's health care, emergency nursing, psychiatry, geriatrics, pediatrics etc. The Journal welcomes the submission of manuscripts that meet the general criteria of significance and scientific excellence. Papers will be published shortly after acceptance. All articles published in IJNM are peer-reviewed.

Contact Us

Editorial Office: ijnm@academicjournals.org

Help Desk: helpdesk@academicjournals.org

Website: <http://www.academicjournals.org/journal/IJNM>

Submit manuscript online <http://ms.academicjournals.me/>

Editors

Dr. Alleene M. Ferguson Pingnot

*California State University, Stanislaus
One University Circle DBH 260, Turlock, CA 95382
USA*

Dr. Andrew Crowther

*Charles Sturt University
Leeds Parade, Orange, New South Wales 2800,
Australia*

Dr. Jacinta Kelly

*School of Nursing & Midwifery
24 Dolier St, Dublin 2
Ireland*

Dr. Jafar Alasad

*College of Nursing, King Saud bin Abdulaziz
University for Health Sciences
(MC 3105)
King Abdulaziz Medical City - National Guard
Health Affairs
P.O. Box 22490, Riyadh 11426,
Saudi Arabia*

Dr. Fintan Sheerin

*School of Nursing and Midwifery, Trinity College
Dublin,
24 D'Olier Street, Dublin 2.
Ireland*

Prof. Helen McCutcheon

*University of South Australia, School of Nursing &
Midwifery
GPO Box 2471, Adelaide, South Australia, 5001,
Australia*

Dr. Panagiotis Christopoulos MD,MSc,PhD,IFEPAG

*2nd Dept. Ob/Gyn
Medical School, University of Athens
1 Hariton street,
14564, N. Kifissia, Athens,
Greece.*

Dr. Arun Kumar

*Manipal College of Medical Sciences
Department of Biochemistry, Pokhara, Nepal
India*

Dr. Harunor Rashid

*Barts and the London Queen Mary's School of
Medicine and Dentistry, London
The Blizard Building, 4 Newark Street, London E1
2AT,
United Kingdom*

Editorial Board

Dr. Sawsan Majali

*Dar Al Hekma College
P.O.Box 34801, Jeddah 21478
Saudi Arabia*

Dr. Patricia L. Riley

*US Centers for Disease Control and Prevention (CDC)
1600 Clifton Road, NE
Mail Stop E-41
Corporate Square Bldg 1, Rm 2409
Atlanta, GA
30329-1902
USA*

Dr. Lucille van der Westhuizen

*University of Namibia
P/B 13301 Windhoek,
Namibia*

Dr. Imtiaz Wani

*S.M.H.S Hospital, Srinagar
Amira Kadal, Srinagar
India*

ARTICLE

- Uterine balloon tamponade for the management of uncontrolled postpartum hemorrhage
by midwives and family physicians** 109
Zaid Altawil, Emily de Redon, Hao Dinh, Daniela Suarez-Rebling and Thomas F. Burke,

Full Length Research Paper

Uterine balloon tamponade for the management of uncontrolled postpartum hemorrhage by midwives and family physicians

Zaid Altawil^{1,2*}, Emily de Redon², Hao Dinh², Daniela Suarez-Rebling² and Thomas F. Burke^{2,3,4}

¹Department of Emergency Medicine, Boston Medical Center, Boston, MA 02118, USA.

²Division of Global Health and Human Rights, Department of Emergency Medicine, Massachusetts General Hospital, Boston, MA 02114, USA.

³Harvard Medical School, Boston, MA 02115, USA.

⁴Harvard T.H. Chan School of Public Health, Boston, MA, 02115, USA.

Received 8 July, 2017; Accepted 14 August, 2017

Postpartum hemorrhage (PPH) is the leading cause of maternal morbidity and mortality world-wide. Although declining globally, maternal mortality is on the rise in the United States. Since uterine balloon tamponade (UBT) has been shown to be safe as well as highly effective in arresting PPH and halting shock progression, this study sought to clarify the practice and perceptions of UBT use among midwives and family physicians in the United States. A cross-sectional 12-item survey on the practice, experiences and perceptions of uterine balloon tamponade utilization for PPH management was conducted among midwives and family physicians that perform deliveries in the United States. The survey settings were one family medicine and two midwifery national conferences in the United States. One hundred and ninety-seven midwives (164; 83.2%) and family physicians (33; 16.8%) completed the surveys. The 197 providers had previously cared for a mean of 6.0 (SD= 18.8) uncontrolled PPH cases each, over the course of their career till date. Eighty (40.6%) of the 197 respondents had been involved with at least one hemorrhaging woman in which a UBT was placed, but only 13 (7.9%) of the 164 midwives and 6 (18.2%) of the 33 family physicians had ever placed a UBT device themselves. One hundred and thirty-one (73.6%) of 178 providers desired training on use of UBT. Midwives and family physicians in the United States care for women with uncontrolled PPH, however, infrequently place a UBT device. Most midwives and family physicians would like training on uterine balloon tamponade.

Key words: Uterine balloon tamponade, post-partum hemorrhage, maternal hemorrhage, emergency obstetric care.

INTRODUCTION

Postpartum hemorrhage (PPH) is the leading cause of maternal morbidity and mortality worldwide. It is estimated that PPH is responsible for 130,000 deaths and 2.6 million injured women across the globe annually (AbouZahr, 2003; World Health Organization, 2012; MacDorman et al., 2013). While many low and middle-

income countries have made considerable strides to reduce their maternal mortality ratios (MMR) over the past two decades, the MMR in the United States has worsened by 136% over the past 20 years (World Health Organization, 2014). The incidence of PPH in the United States has increased by over 30% since 2004 and today

it is estimated to occur in 2.9% of deliveries (Bateman et al., 2010). It is notable that one in ten women with PPH will fail first line therapies and are thus classified as having uncontrolled PPH (Ononge et al., 2016).

The most common cause of PPH is uterine atony. First-line treatment options for PPH from uterine atony include uterine massage, uterus and bladder emptying and uterotonic agents such as oxytocin, ergometrine, misoprostol and prostaglandin F2a (PGF2a) (World Health Organization, 2012). When first-line interventions are unsuccessful, PPH is called “uncontrolled”. Interventions that target uncontrolled PPH from atonic uterus include aortic compression, intravenous tranexamic acid (Collaborators, 2017), uterine balloon tamponade, uterine artery embolization, and fluid, blood and blood product resuscitation once hemorrhage control has been gained (World Health Organization, 2012). Surgical approaches to uncontrolled postpartum hemorrhage include internal iliac artery ligation, uterine compression sutures and emergency hysterectomy (El Senoun et al., 2011).

Over the past seven years, the authors have studied the impact of uterine balloon tamponade in resource-limited settings by studying efficacy, safety, cost-effectiveness, barriers and facilitators to optimal use, as well as models for scale (Natarajan et al., 2016; Burke et al., 2016). During this time, UBT use for uncontrolled PPH has been endorsed by the WHO, the International Federation of Obstetrics and Gynecology, the American College of Obstetrics and Gynecology, the Royal College of Obstetrics and Gynecology and the International Confederation of Midwives, among others, as the best evidence clinical intervention (World Health Organization, 2012; Doumouchtsis et al., 2007; Tindell et al., 2013; Lalonde et al., 2006; Mavrides et al., 2016).

The authors' work has shown that if applied in a timely fashion, uterine balloon tamponade arrests hemorrhage, halts shock progression (Burke et al., 2017), averts nearly all deaths and emergency hysterectomies (Pendleton et al., 2016; Burke et al., 2017), is “safe” and “highly cost-effective” (Mvundura et al., 2017), and can be placed successfully by any cadre, including illiterate front-line health workers (Natarajan et al., 2016). Deploying a UBT device involves placing a balloon within the uterine cavity and inflating it until hemorrhage ceases. A variety of uterine balloon devices are available, including the Sengstaken-Blakemore tube, Bakri balloon, Rusch balloon, Ebb (Belfort-Didley), and the condom catheter balloon (Pendleton et al., 2016; Dildy et al., 2014).

The authors believe that lessons learned in low- and middle-income countries may inform opportunities for improving health outcomes in highly resourced settings,

such as the United States and elsewhere (Tindell et al., 2013; Pendleton et al., 2016; Burke et al., 2016). The purpose of this study was to clarify the practice and perceptions of UBT use in the United States among midwives and family physicians.

MATERIALS AND METHODS

Ethical approval was obtained from the Partners Human Research Committee (Massachusetts General Hospital, Boston, MA, USA). A cross-sectional 12-item survey on the practice, experiences, and perceptions of uterine balloon tamponade utilization for PPH management was developed by the authors through an iterative process. Uncontrolled PPH was defined as hemorrhage unresponsive to a first dose of treatment uterotonics. Appropriate utilization of uterine balloon tamponade was defined as uterine placement of a UBT in each case of uncontrolled post-partum hemorrhage. Respondent credentials, location and years spent in practice, number of deliveries attended per month, number of uncontrolled PPH cases cared for and experience with uterine balloon tamponade were recorded.

Setting

The survey instrument was administered among midwives and family physicians in the United States at three national conferences: the American College of Nurse Midwives 60th Annual Meeting and Exhibition (Washington, DC, June 27, 2015); the American Academy of Family Physicians Family Medicine Experience (Denver, CO, September 29, 2015); and the Midwives Alliance of North America Annual Convention (Albuquerque, NM, October 15, 2015).

Study population

Conference regulations did not allow solicitation beyond the authors' exhibitor booth; therefore, the study participants were among those that voluntarily approached the booth. Providers were queried as to whether they provided obstetric delivery care. If the response was positive, the provider was invited to participate in the study and complete a survey. Oral consent was obtained and all surveys remained anonymous.

Inclusion and exclusion criteria

All providers that approached the authors' exhibitor booth were potential study participants. Exclusion criteria included health cadres other than midwives or family physicians, providers who did not practice obstetrics, students, providers who had less than one year of clinical experience and providers who primarily practiced outside United States.

Data analysis

Each of the 12 questions was analyzed individually. If no answer was provided to a question, the individual data point was removed

*Corresponding author. E-mail: zaidaltawil@gmail.com. Tel: +1-617-417-5406. Fax: +1-617-643-8772.

Author(s) agree that this article remain permanently open access under the terms of the [Creative Commons Attribution License 4.0 International License](https://creativecommons.org/licenses/by/4.0/)

Table 1. Interventions in the 198 described cases of uncontrolled PPH^a.

Intervention	Total (N=198) n(%)
B-lynch sutures	24 (12.1)
Uterine balloon tamponade	80 (40.4)
Uterine artery ligation	29 (14.6)
Hysterectomy	53 (26.8)
Dilation and curettage	9 (4.5)
Unspecified surgical intervention	3 (1.5)

^aRespondents could choose multiple answers; one, more than one, or none of the above interventions.

from both the numerator and the denominator. The registry was built using Microsoft Excel 2007 (Seattle, Washington, USA) and critically examined using standard descriptive analysis in Stata 13 (College station, Texas, USA).

RESULTS

Respondent characteristics

One hundred and ninety-seven providers met the inclusion criteria and completed the surveys. One hundred and sixty-four (83.3%) were midwives and 33 (16.7%), family physicians. The mean years of clinical practice of the 197 providers was 15.6 years (range: 1 to 45). One hundred and fifteen (70.1%) of the 164 midwives provided details of their training. Sixty (52.2%) of the 115 midwives were certified nurse midwives (CNM), 25 (21.7%) were certified professional midwives (CPM), and 18 (15.7%) were dual CPM and lay midwife (LM) certified. Among the 33 family physicians, five (15.2%) had pursued additional fellowship training in obstetric care and three (9.1%) were certified Advanced Life Support in Obstetrics (ALSO) providers. Of the 193 providers that described their practice location(s), 110 (57.0%) performed deliveries exclusively in hospitals, 63 (32.6%) in non-hospital settings (independent birthing centers and private homes) and 12 (6.2%) in both.

Experience with postpartum hemorrhage

One hundred and eighty-two (92.4%) of the 197 providers estimated the number of uncontrolled PPH cases they had cared for over the course of their career. The mean number of uncontrolled PPH cases per provider over the course of their career was 6.0 (SD= 18.8). In response to a request for a narrative on each of the provider's two most recent cases of uncontrolled PPH, 182 providers described a total of 198 women hemorrhaging uncontrollably (Table 1). Respondents provided information on one, two or no cases. A UBT device was placed in 80 (40.4%) of the 198 women with uncontrolled PPH. Seventy-six (38.4%) of the 198 cases of uncontrolled

PPH directly underwent surgical intervention without an attempt at a UBT device placement.

Nineteen (9.6%) of the 197 respondents had placed at least one UBT device at some point in their career. The 19 included 13 (7.9%) of the 164 midwives and 6 (18.2%) of the 33 family physicians. Two midwives reported having improvised a UBT device with a condom and Foley catheter.

Seventy-one of 110 (64.5%) respondents reported that the Bakri balloon was the most common uterine balloon at their facility. The next most common uterine balloon devices reported were the Foley catheter (8.2%) and the condom catheter (1.8%), followed by the Sengsten-Blakemore tube (0.9%), the Ebb balloon (0.9%) and the Rusch balloon (0.9%). One hundred and thirty-one (73.6%) of the 178 providers who responded that they had never placed a UBT device, wished to become trained on it in the future.

DISCUSSION

The 197 midwives and family physicians that participated in the survey collectively cared for almost 1,200 women with uncontrolled postpartum hemorrhage over the course of their careers. Although, UBT is widely endorsed by leading maternal health and obstetric physician organizations, only 9.6% of the surveyed providers had ever placed at least one. The surveyed providers reported that, in their experience, 38.4% of hemorrhaging women proceeded directly to emergency surgery with no UBT placement attempts. Most (73.5%) of the surveyed providers expressed a desire to become trained on UBT use.

Although, uterine balloon tamponade has been shown to be highly effective in the hands of all levels of healthcare providers (Tindell et al., 2013), very few had ever placed a UBT among this cohort of midwife and family physician obstetric providers despite considerable experience with women suffering from uncontrolled PPH. While this study was not designed to uncover barriers to appropriate use of uterine balloon tamponade, lack of knowledge and training were likely contributors to

suboptimal use. Since many of the interviewed providers performed deliveries outside the hospitals and since it is well known that early UBT placement leads to improved outcomes (Burke et al., 2017), it appears that there may be considerable untapped opportunity for strengthening the policies and clinical pathways for the emergency care of uncontrolled PPH.

During the course of the authors' research, it was inspiring to observe the near elimination of PPH deaths and emergency hysterectomies in major medical centers across several low-income countries, yet it is surprising and unsettling to learn about the significant number (38.4%) of hemorrhaging women reported by this cohort of providers, who were taken directly to emergency surgery without attempting uterine balloon placement. Previous research by the authors has clearly demonstrated that uterine balloon tamponade can both avert hysterectomies and other surgical interventions as well as arrest hemorrhage and shock progression (Pendleton et al., 2016; Burke et al., 2017). It appears there may have been numerous missed opportunities to avert emergency surgical intervention among those surveyed in this study.

The main limitations of this study were: being restricted to surveying only providers that were attracted to, and chose to visit the authors' exhibitor booth. Despite the sample size of 197 obstetric midwife and family practice providers, the surveyed providers collectively had considerable experience with uncontrolled PPH. While providers that were attracted to the booth related to the topic on PPH were more likely to be more knowledgeable on PPH, the findings demonstrated a striking lack of awareness and training on uterine balloon tamponade.

Conclusion

There may be considerable opportunity to improve appropriate use of uterine balloon tamponade across multiple levels of obstetric delivery providers in the United States.

CONFLICT OF INTERESTS


The authors have not declared any conflict of interests.

ACKNOWLEDGEMENT

This research was supported by the Ujenzi Charitable Trust.

REFERENCES

- AbouZahr C (2003). Global burden of maternal death and disability. *Br. Med. Bull.* 67(1):1-11.
- Bateman BT, Berman MF, Riley LE, Leffert LR (2010). The epidemiology of postpartum hemorrhage in a large, nationwide sample of deliveries. *Anesthesia Analg.* 110(5):1368-1373.
- Burke, TF, Ahn R, Nelson BD, Hines R, Kamara J, Oguttu M, Dulo L, Achieng E, B Achieng, Natarajan A, Maua J, Kargbo SAS, Altawil Z, Tester K, de Redon E, Niang M, Abdalla K, Eckardt MJ (2016). A postpartum haemorrhage package with condom uterine balloon tamponade: a prospective multi-centre case series in Kenya, Sierra Leone, Senegal, and Nepal. *BJOG* 123(9):1532-1540.
- Burke TF, Danso-Bamfo S, Guha M, Oguttu M, Tarimo V, Nelson BD (2017). Shock progression and survival after use of a condom uterine balloon tamponade package in women with uncontrolled postpartum hemorrhage. *Int. J. Gynecol. Obstet.* doi: 10.1002/ijgo.12251
- Collaborators WT (2017). Effect of early tranexamic acid administration on mortality, hysterectomy, and other morbidities in women with postpartum haemorrhage (WOMAN): an international, randomised, double-blind, placebo-controlled trial. *Lancet* 389(10084):2105-2116.
- Dildy GA, Belfort MA, Adair CD, Destefano K, Robinson D, Lam G, Strong TH, Polon C, Massaro R, Bukkapatnam J, Van Hook JW (2014). Initial experience with a dual-balloon catheter for the management of postpartum hemorrhage. *Am. J. Obstet. Gynecol.* 210(2):136-e1.
- Doumouchsis SK, Papageorghiou AT, Arulkumaran S (2007). Systematic review of conservative management of postpartum hemorrhage: what to do when medical treatment fails. *Obstet. Gynecol. Surv.* 62(8):540-547.
- El Senoun GA, Singh M, Mousa HA, Alfrevic Z (2011). Update on the new modalities on the prevention and management of postpartum haemorrhage. *Fetal Matern. Med. Rev.* 22(4):247-264.
- Lalonde A, Daviss BA, Acosta A, Herschderfer K (2006). Postpartum hemorrhage today: ICM/FIGO initiative 2004-2006. *Int. J. Gynecol. Obstet.* 94(3):243-253.
- Likis FE, Sathe NA, Morgans AK, Hartmann KE, Young JL, Carlson-Bremer D, Schorn M, Surawicz T, Andrews J (2015). Management of Postpartum Hemorrhage. Comparative Effectiveness Review No. 151. Agency for Healthcare Research and Quality. PMID: 26020092
- MacDorman MF, Declerq E, Matthews TJ (2013). Recent trends in out-of-hospital births in the United States. *J. Midwifery Womens Health* 58(5):494-501.
- Mavrides E, Allard S, Chandraharan E, Collins P, Green L, Hunt BJ, Riris S, Thomson AJ (2016). Prevention and management of postpartum haemorrhage. *BJOG* 124:e106-e149.
- Mvundura M, Kokonya D, Abu-Haydar E, Okoth E, Herrick T, Mukabi J, Carlson L, Oguttu M, Burke, T (2017). Cost-effectiveness of condom uterine balloon tamponade to control severe postpartum hemorrhage in Kenya. *Int. J. Gynecol. Obstet.* 137(2):185-191.
- Natarajan A, Alaska Pendleton A, Nelson BD, Ahn R, Oguttu M, Dulo L, Eckardt MJ, Burke TF (2016). Provider experiences with improvised uterine balloon tamponade for the management of uncontrolled postpartum hemorrhage in Kenya. *Int. J. Gynecol. Obstet.* 135(2):210-213.
- Ononge S, Mirembe F, Wandabwa J, Campbell OM (2016). Incidence and risk factors for postpartum hemorrhage in Uganda. *Reprod. Health* 13(1):38.
- Pendleton AA, Natarajan A, Ahn R, Nelson BD, Eckardt MJ, Burke TF (2016). A qualitative assessment of the impact of a uterine balloon tamponade package on decisions regarding the role of emergency hysterectomy in women with uncontrolled postpartum haemorrhage in Kenya and Senegal. *BMJ Open.* 6(1):p.e010083.
- Pendleton AA, Natarajan A, Ahn R, Nelson BD, Eckardt MJ, Burke TF (2016). Emergency hysterectomy for uncontrolled postpartum hemorrhage may be averted through uterine balloon tamponade in Kenya and Senegal. *Int. J. Gynecol. Obstet.* 133(1):124-124.
- Tindell K, Garfinkel R, Abu-Haydar E, Ahn R, Burke TF, Conn K, Eckardt M (2013). Uterine balloon tamponade for the treatment of postpartum hemorrhage in resource-poor settings: a systematic review. *BJOG* 120(1):5-14.
- World Health Organization (WHO) (2014). Trends in maternal mortality 1990 to 2013: estimates by WHO, UNICEF, UNFPA, The World Bank and the United Nations Population Division. ISBN 978 92 4 150722 6
- World Health Organization (WHO) (2012). WHO recommendations for the prevention and treatment of postpartum haemorrhage: evidence base. Geneva: World Health Organization. Available at: http://apps.who.int/iris/bitstream/10665/75411/1/9789241548502_eng.pdf



International Journal of Nursing and Midwifery

Related Journals Published by Academic Journals

- *International Journal of Medicine and Medical Sciences*
- *Journal of Medicinal Plant Research*
- *African Journal of Pharmacy and Pharmacology*
- *Journal of Dentistry and Oral Hygiene*
- *Medical Practice and Reviews*
- *Journal of Public Health and Epidemiology*

academicJournals