



International Journal of Medicine and Medical Sciences

Volume 9 Number 9 September 2017

ISSN 2006-9723



*Academic
Journals*

ABOUT IJMMS

The **International Journal of Medicine and Medical Sciences** is published monthly (one volume per year) by Academic Journals.

The **International Journal of Medicine and Medical Sciences (IJMMS)** provides rapid publication (monthly) of articles in all areas of Medicine and Medical Sciences such as:

Clinical Medicine: Internal Medicine, Surgery, Clinical Cancer Research, Clinical Pharmacology, Dermatology, Gynaecology, Paediatrics, Neurology, Psychiatry, Otorhinolaryngology, Ophthalmology, Dentistry, Tropical Medicine, Biomedical Engineering, Clinical Cardiovascular Research, Clinical Endocrinology, Clinical Pathophysiology, Clinical Immunology and Immunopathology, Clinical Nutritional Research, Geriatrics and Sport Medicine

Basic Medical Sciences: Biochemistry, Molecular Biology, Cellular Biology, Cytology, Genetics, Embryology, Developmental Biology, Radiobiology, Experimental Microbiology, Biophysics, Structural Research, Neurophysiology and Brain Research, Cardiovascular Research, Endocrinology, Physiology, Medical Microbiology

Experimental Medicine: Experimental Cancer Research, Pathophysiology, Immunology, Immunopathology, Nutritional Research, Vitaminology and Etiology

Preventive Medicine: Congenital Disorders, Mental Disorders, Psychosomatic Diseases, Addictive Diseases, Accidents, Cancer, Cardiovascular Diseases, Metabolic Disorders, Infectious Diseases, Diseases of Bones and Joints, Oral Preventive Medicine, Respiratory Diseases, Methods of Epidemiology and Other Preventive Medicine

Social Medicine: Group Medicine, Social Paediatrics, Medico-Social Problems of the Youth, Medico-Social Problems of the Elderly, Rehabilitation, Human Ecology, Environmental Toxicology, Dietetics, Occupational Medicine, Pharmacology, Ergonomy, Health Education, Public Health and Health Services and Medical Statistics The Journal welcomes the submission of manuscripts that meet the general criteria of significance and scientific excellence. Papers will be published approximately one month after acceptance. All articles published in IJMMS are peer-reviewed.

Contact

Editorial Office: ijmms@academicjournals.org

Desk: helpdesk@academicjournals.org

Website: <http://www.academicjournals.org/journal/IJMMS>

Submit manuscript online <http://ms.academicjournals.me/>

Editors

Dr. J. Ibekwe

*Acting Editor-in-chief,
International Journal of Medicine and Medical
Sciences Academic Journals
E-mail: ijmms.journals@gmail.com
<http://www.academicjournals.org/ijmms>*

Afrozul Haq

*Editor, Laboratory Medicine
Department of Laboratory Medicine
Sheikh Khalifa Medical City
P.O. Box 51900, ABU DHABI
United Arab Emirates*

Editorial Board

Chandrashekhar T. Sreeramareddy

*Department of Community Medicine,
P O Box No 155, Deep Heights
Manipal College of Medical Sciences,
Pokhara,
Nepal*

Sisira Hemananda Siribaddana

*259, Temple Road, Thalpathpitiya,
Nugegoda, 10250
Sri Lanka*

Dr. santi M. Mandal

*Internal Medicine
UTMB, Galveston, TX,
USA*

Konstantinos Tziomalos

*Department of Clinical Biochemistry
(Vascular Prevention Clinic),
Royal Free Hospital Campus,
University College Medical School, University College
London, London,
United Kingdom*

Cyril Chukwudi Dim

*Department of Obstetrics & Gynaecology
University of Nigeria Teaching Hospital (UNTH)
P.M.B. 01129, Enugu. 400001,
Nigeria*

Mojtaba Salouti

*School of Medical and Basic Sciences,
Islamic Azad University- Zanjan,
Iran*

Imtiaz Ahmed Wani

*Srinagar Kashmir, 190009,
India*

Professor Viroj Wiwanitkit

*Wiwanitkit House, Bangkhae,
Bangkok
Thailand 10160*

Dr. Srinivas Koduru

*Dept of Clinical Sciences
Collage of Health Sciences
University of Kentucky
Lexington USA*

Weiping Zhang

*Department of Oral Biology
Indiana University School of Dentistry
1121 West Michigan Street, DS 271
Indianapolis, IN 46202
USA*

Lisheng XU

*Ho Sin Hang Engineering Building
Department of Electronic Engineering
The Chinese University of Hong Kong
Shatin, N.T. Hong Kong,
China*

Dr. Mustafa Sahin

*Department of Endocrinology and Metabolism
Baskent University,
Ankara,
Turkey*

Dr. Harshdeep Joshi

*Maharishi Markandeshwar
Institute of Medical Sciences and Research
Ambala, (Haryana).
India.*

ARTICLE

Factors affecting prolonged intensive care unit stay in Nigist Eleni Mohammed Memorial Hospital, Hosanna, Southern Ethiopia

105

Mohammed Suleiman Obsa, Bedilu Girma Weji and Tahir Aman Worji

Full Length Research Paper

Factors affecting prolonged intensive care unit stay in Nigist Eleni Mohammed Memorial Hospital, Hosanna, Southern Ethiopia

Mohammed Suleiman Obsa^{1*}, Bedilu Girma Weji¹ and Tahir Aman Worji²

¹Department of Anesthesia, College of Health Science and Medicine, Wolaita Sodo University, P. O. Box 138, Wolaita Sodo, SNNR, Ethiopia.

²Department of Anesthesia, College of Health Science and Medicine, Arsi University, Asella, Ethiopia.

Received 17 April, 2017; Accepted 9 May, 2017

The reason for a long stay in the Intensive Care Unit (ICU) is unknown in most cases, but there are some risk factors that can prolong the patients' stay in the ICU. However, the threshold of prolonged length of stay is useful in analyzing the quality of health care and hospital costs. The main objective of this study was to assess the factors affecting prolonged intensive care unit stay in Nigist Eleni Mohammed Memorial Hospital from January, 2015 to January, 2016. Institutional based cross sectional study design was conducted. Census was used to include all study participants admitted to intensive care. Regular supervision and follow up was made. Data was entered into Epi info version 7 by investigators and was transported to statistical package for social sciences (SPSS) version 20 for analysis. Bivariate and multivariate analysis was used to identify factors associated with patient's outcome. A total of 280 patients were included in the study by census. The mean length of stay in the intensive care unit was 5.49 ± 4.36 . Majority of patients (61.4%) had poor nutrition status. In addition to their underlying diseases, majority of the patients in the intensive care units had one or more comorbidities. Poor nutritional status, male sex, presence of comorbidity and unplanned extubation was significantly associated with length of stays in intensive care unit at p-value less than 0.05. In this study, it was found that unconscious patients had about two times more likelihood to have prolonged length of stays in intensive care unit than conscious patients (AOR 1.724; 95% CI: 809 to 3.676). Higher proportions of patients stayed in the intensive care unit for more than three days. Therefore, the implication of this study was to recommend the decrease in length of intensive care unit stays in medically appropriate ways.

Key words: Intensive care unit, Ethiopia, length of stay, patient management.

INTRODUCTION

Length of stay (LOS) in the intensive care unit (ICU) is one of the most important factors that influence health management (Keenan et al., 1998; Halpern and Pastores,

2010). Prolonged stays in the intensive care unit are associated with high costs and burdens on patients, patients' families and in turn affect society at large. It

*Corresponding author. E-mail: msuleimaahn43@yahoo.com.

is associated with a complicated clinical course of the diseases with higher incidence of respiratory and renal insufficiency (Joskowiak et al., 2013). Therefore, it can decrease the chance of survival and reduce the quality of life (Lagercrantz et al., 2010), which may result in increased costs and reduced hospital beds available in the ICU (Arabi et al., 2002; Doering et al., 2000; Azarfarin et al., 2014).

A primary goal in intensive care is to decrease length of stay in a medically appropriate way which helps to improve the quality of medical care, reduce hospital cost and avoid excess use of resources (Jaber et al., 2006). The precise prediction of the length of ICU stay enables the physicians to obtain reliable information to provide a better care for patients, allocating funds to the health care systems and changing their routines (Azarfarin et al., 2014; Austin et al., 2002; De Cocker et al., 2011). Therefore, this study was conducted to identify potential factors of prolonged ICU stays that enable health care providers and managers to devise different strategies which in turn help to reduce length of stay (LOS) on one hand and improve quality of care for critically ill patients and reduce hospital cost on the other hand.

MATERIALS AND METHODS

Study setting

A cross-sectional study design was conducted from January, 2015 to January, 2016 at Nigist Eleni Mohammed Memorial Hospital (NEMMH) in Hosanna town. Hosanna town is an administrative city of Hadiya zone which is located at a distance of 230 km to the south of Addis Ababa, capital city of Ethiopia. The hospital has nine wards. There are adult and neonatal intensive care units in the hospital. All adult critical ill patients from different departments have been admitted to adult intensive care unit. This study excluded neonates as they were admitted to neonatal intensive care unit. The hospital has a total of 220 beds including eight and seven beds in neonatal and adult ICU, respectively.

Source population

This includes all charts of patients admitted to adult intensive care unit of Nigist Eleni Mohammed Memorial Hospital.

Study population

This includes selected charts of patients admitted to adult intensive care unit of Nigist Eleni Mohammed Memorial Hospital from January, 2015 to January, 2016.

Sample size determination and sampling technique

Census was used to include all eligible 280 patients admitted to adult intensive care unit of Nigist Eleni Mohammed Memorial Hospital from January, 2015 to January, 2016.

Data collection tools and procedure

Data was collected using structured questionnaires by two B. Sc.

anesthetists and supervised by one M. Sc. holder anesthetist. Patient's charts were reviewed. At the end of data collection, patient's charts were replaced into its original place properly.

Data quality assurance and data analyzing

The structured questionnaire prepared in English was used as a data collection tool. Data collectors and supervisors were trained on each items included in the study tools, objective, relevance of study, and right of respondents. During data collection, regular supervision and follow up were made. The data was entered into epi info version 7 and was exported to SPSS version 20 computer program for analysis. Bivariate and multivariate analyses were used to detect the effect of independent variable over length of ICU stays. A p-value of less than 0.05 was used as a cut of point for presence of association. Strength of association was measured by 95% confidence interval and/Odd ratio.

Definitions

Operational definitions

Prolonged ICU stays: Patients who stayed in the intensive care unit for more than three days.

Unplanned extubations: Premature removal of an endotracheal tube by a patient receiving mechanical ventilation support or unintentional removal of an endotracheal tube by staff during nursing and medical procedures.

Ethical consideration

Ethical clearance and approval was obtained from the Ethical Review Committee, Anesthesia Department, Wolaita Soddo University. Permission to conduct the work was obtained from the hospital. The obtained data was only used for the study purpose. Confidentiality and anonymity were ensured.

RESULTS

Socio-demographics and personal information

A total of 280 patients were enrolled in the study of which majority of them were male. Almost four fifth of the patients were found between age groups of 18 to 65 years. About 78.3% of the sampled respondents were married. The percentages of patients below 18 years and above 65 years were about 17.1 and 3.2%, respectively. Majority of them were from urban area. Patients in the intensive care unit can have single or multiple comorbidities. But, in this research, comorbidity diagnosis was divided into two categories, namely the presence and the absence of comorbidity. In addition to their underlying diseases, majority of the patients in the intensive care units developed comorbidities (Table 1).

Length of ICU stays

Mean length of stays in the intensive care unit was 5.49 ± 4.36 (minimum 1 and maximum 15) which included both

Table 1. Socio-demographics and personal information of patients admitted to adult intensive care unit of Nigist Eleni Mohammed Memorial Hospital, Hosanna, from January, 2015 to January, 2016.

Variable	Category	Frequency	Percentage
Sex	Male	132	47.1
	Female	83	29.6
Age group	<18	48	17.1
	18-65	223	79.6
	>65	9	3.2
Address	Urban	160	57.14
	Rural	110	42.86
Comorbidity	Present	263	88.0
	Abscent	17	5.7
Level of consciousness	Conscious	167	59.6
	Unconscious	113	40.4
Unplanned extubation	Yes	34	11.4
	No and/or not intubated	246	82.3

survivors and non-survivors. About 47.1% of the patients stayed in the intensive care unit for less than three days (Figure 1).

Nutrition status, age and gender among different categories of hospital stay

The height and weight of the patients used to calculate the body mass index (BMI), which is important to determine the nutritional status of the patients could not be obtained. Therefore, instead of using BMI to determine whether the patient is malnourished or not, the status of albumin level as indicators of nutritional status was used. Below the normal level of albumin (N<3.5 to 5.0 g/dl) was considered as poor nutrition status, while normal level was good nutrition status (N: 3.5 to 5.0 g/dl). According to the results of this study, majorities of patients (61.4%) had below normal albumin level which means that most of the patients in the intensive care units had poor nutrition status. Majority of the male patients stayed for more than three days and the reverse was true for female (Table 2).

Factors affecting length of stay in intensive care units

The result of multivariate analysis showed that sex, infection, nutritional, unplanned extubation had strong association with prolonged ICU stay at p-value of less

than 0.05. Unconscious patients had about two times more likely to have prolonged length of stays in intensive care unit than conscious patients (AOR 1.724; 95% CI: 0.809 to 3.676) (Table 3).

DISCUSSION

The reduction of length of stays in intensive care unit should be a major aim of medical staff as longer ICU stays increase the risk for infections and associated with a higher risk of hospital death (Vincent et al., 2009; Cavallazzi et al., 2012) and have immense costs (Wunsch et al., 2012). Some study revealed that social, psychological, and institutional factors have an impact on ICU length of stay. Lack of effective communication and the occurrence of conflicts between the physician and the patient's family can also lead to prolonged ICU stays (Higgins et al., 2003; Dimick et al., 2001). However, these cannot be identified in our study as psychological and social factors could not be recorded on patient's chart and it was considered as limitation of our study. Therefore, the study was conducted on other important risk factors available on patient's charts like age, sex, category of disease, nutritional status, presence of comorbidity, unplanned extubation and level of consciousness.

In these studies, the reason for a long stay in the ICU is unknown in most cases, but there are some risk factors that can prolong the patients' stay in the ICU. However, due to the differences among the populations and the




Figure 1. Length of ICU stays of patients admitted to adult intensive care unit of Nigist Eleni Mohammed Memorial Hospital, Hosanna, from January, 2015 to January, 2016.

Table 2. Relationship between sex and age of the patients and length of ICU stays at Nigist Eleni Mohammed Memorial Hospital, Hosanna, from January, 2015 to January, 2016.

Age in years		Length of ICU stays		Total
		≤3 days	>3 days	
Sex	Male	60	103	163
	Female	72	45	117
	Total	132	148	280
Age	<18	28	20	48
	18-65	70	93	223
	>65	34	25	9
	Total	132	148	280
Nutritional status	Good	49	47	96
	Poor	99	73	172
	Missing	4	8	12
	Total	152	128	280

countries, and the definition of prolonged ICU stay in different centers, the predictive models of a hospital may not be applicable in another hospital (Azarfarin et al., 2014; Curtis et al., 2002; Gardner et al., 2001).

In the present research, it was found that majority of the patients had prolonged ICU stays which is consistent with other similar study (Sugiarto and Darmawan, 2014). According to the result of present study poor nutritional status and presence of comorbidities had a significant relationship with length of stay which was consistent with other similar study. However, in this studies age did not show significant relationship with LOS. Another study also found that comorbidities influence not only the

outcome of patients but also it affects length of ICU stays (Ingeman et al., 2011; Luker et al., 2011). The study conducted on recent trends in the treatment of spontaneous intracerebral hemorrhage revealed that individual comorbidities could influence the outcome of patients and length of ICU stays (Ingeman et al., 2011). For example, some studies used hypertension or diabetes mellitus as outcome indicators, and other researchers found that respiratory failure or pneumonia influence the LOS and mortality rate of patients (Andaluz and Zuccarello, 2009; Szepesi et al., 2015). These patients utilize more hospital resources than those with a single comorbidity or no comorbidities, no matter whether

Table 3. Factors affecting length of stays in adult intensive care unit of Nigst Eleni Mohammed Memorial Hospital, Hosanna, from January 2015 to January 2016.

Variable	Df.	Sig.	AOR	95% C.I. for EXP(B)	
Male	1	0.002	0.323	0.159	0.653
Presence of comorbidity	1	0.021	0.223	0.062	0.796
Poor nutritional status	1	0.013	0.181	0.047	0.693
Unplanned extubation	1	0.000	81.836	11.541	580.319
Loss of consciousness	1	0.159	1.724	0.809	3.676

the comorbidities arise from previous disease history or complications. Although the previous disease history of patients cannot be changed, decreasing complications using quality control methods will lead to a better outcome and a shorter LOS (Ingeman et al., 2011).

The likely causes of prolonged ICU stays among patients who developed comorbidities were acute deterioration in vital sign of critically ill patients and the involvement of other vital organ in the disease process.

Limitations of this study were due to lack of availability of adequate risk factors on the patient's chart which might be important to thoroughly discuss risk factors of length of ICU stays and inadequate sample size.

Conclusions

The aim of intensive care medicine is to provide quality of care for patients while reducing length of stays in intensive care unit. In this study, poor nutritional status, male sex, presence of comorbidity and unplanned extubation had significant relation with LOS. There may also be social and psychological factors that influence patients' length of stays. Majorities of patients stayed in intensive care unit for a prolonged period. Therefore, it was recommended to reduce length of stay by preventing modifiable risk factors and improving the type and quality of communication. Further prospective studies on risk factors of length of stay should be conducted to come up with all potential risk factors of length of ICU stays.

CONFLICT OF INTERESTS

The authors have not declared any conflict of interest.

ACKNOWLEDGEMENTS

We would like to thank all the staff in the ICU and record keeper for their support during data collection.

Abbreviation

ICU, Intensive care unit; **LOS**, length of stays; **NEMMH**, Nigst Eleni Mohammed Memorial Hospital.

REFERENCES

- Andaluz N, Zuccarello M (2009). Recent trends in the treatment of spontaneous intracerebral hemorrhage: analysis of a nationwide inpatient database: Clinical article. *J. Neurosurg.* 110:403-410.
- Arabi Y, Venkatesh S, Haddad S, Alshimemeri A, Al malik S (2002). A prospective study of prolonged stay in the intensive care unit: predictors and impact on resource utilization. *Int. J. Qual Health Care* 14:403-410.
- Austin PC, Rothwell DM, TU JV (2002). A comparison of statistical modeling strategies for analyzing length of stay after CABG surgery. *Health Serv. Outcomes Res. Methodol.* 3:107-133.
- Azarfarin R, Ashouri N, Totonchi Z, Bakhshandeh H, Yaghoubi A (2014). Factors influencing prolonged ICU stay after open heart surgery. *Res. Cardiovasc. Med.* 3(4).
- Cavallazzi R, Saad M, Marik PE (2012). Delirium in the ICU: An overview. *Ann. Intensive Care* 2:49.
- Curtis JJ, Mckenny-Knox, CA, Wagner-mann CC (2002). Postcardiotomy centrifugal assist: a single surgeon's experience. *Artif Organs* 26:994-997.
- De cocker J, Messaoudi N, Stockman BA, Bossaert LL, Rodrigus IE (2011). Preoperative prediction of intensive care unit stays following cardiac surgery. *Eur. J. Cardio-Thoracic Surg.* 39:60-67.
- Dimick JB, Pronovost PJ, Heitmilller RF, Lipsitt PA (2001). Intensive care unit physician staffing is associated with decreased length of stay, hospital cost, and complications after esophageal resection. *Crit. Care Med.* 29:753-758.
- Doering LV, Esmailian F, Laks H (2000). Perioperative predictors of ICU and hospital costs in coronary artery bypass graft surgery. *Chest J.* 118:736-743.
- Gardner SC, Grunwald GK, Rumsfeld JS, Mackenzie T, Gao D, Perlin JB, McDonald G, Shroyer ALW (2001). Risk factors for intermediate-term survival after coronary artery bypass grafting. *Ann. thoracic Surg.* 72:2033-2037.
- Halpern NA, Pastores SM (2010). Critical care medicine in the United States 2000–2005: an analysis of bed numbers, occupancy rates, payer mix, and costs. *Crit. Care Med.* 38:65-71.
- Higgins TL, Mcgee WT, Steingrub JS, Rapoport J, Lemeshow S, Teres D (2003). Early indicators of prolonged intensive care unit stay: Impact of illness severity, physician staffing, and pre-intensive care unit length of stay. *Crit. Care Med.* 31:45-51.
- Ingeman A, Andersen G, Hundborg HH, Svendsen ML, Johnsen SP (2011). In-hospital medical complications, length of stay, and mortality among stroke unit patients. *Stroke* 110:610881.
- Jaber S, Amraoui J, Lefrant JY, Arich C, Cohendy R, Landreau L, Calvet Y, Capdevila X, Mahamat A, Eledjam JJ (2006). Clinical practice and risk factors for immediate complications of endotracheal intubation in the intensive care unit: a prospective, multiple-center study. *Crit. Care Med.* 34:2355-2361.
- Joskowiak D, Kappert U, Matschke K, Tugtekin S (2013). Prolonged intensive care unit stay of patients after cardiac surgery: initial clinical results and follow-up. *Thoracic Cardiovasc. Surg.* 61:701-707.
- Keenan SP, Massel D, Inman KJ, Sibbald WJ (1998). A systematic review of the cost-effectiveness of noncardiac transitional care units. *Chest* 113:172-177.
- Lagercrantz E, Lindblom D, Sartipy U (2010). Survival and quality of life in cardiac surgery patients with prolonged intensive care. *Ann.*

- Thoracic Surg. 89:490-495.
- Luker JA, Bernhardt J, Grimmer-somers KA (2011). Demographic and stroke-related factors as predictors of quality of acute stroke care provided by allied health professionals. *J. Multidiscipl. Healthcare* 4:247-259.
- Sugiarto N, Darmawan ES (2014). The Factors Affecting the Length of Stay in the Intensive Care Units of Pertamina Central Hospital in Indonesia Related to Healthcare Associated Infections. *J. US-China Med. Sci.* 11: 195-204.
- Szepesi R, Széll IK, Hortobágyi T, Kardos L, Nagy K, Láncki LI, Berényi E, Bereczki D, Csiba L (2015). New prognostic score for the prediction of 30-day outcome in spontaneous supratentorial cerebral haemorrhage. *BioMed Res. Int.* 2015:961085
- Vincent JL, Rello J, Marshall J, Silva E, Anzueto A, Martin CD, Moreno R, Lipman J, Gomersall C, Sakr Y (2009). International study of the prevalence and outcomes of infection in intensive care units. *Jama* 302(21):2323-2329.
- Wunsch H, Gershengorn H, Scales DC (2012). Economics of ICU organization and management. *Crit. Care Clin.* 28(1):25.



International Journal of Medicine and Medical Sciences

Related Journals Published by Academic Journals

- *Journal of Medicinal Plant Research*
- *African Journal of Pharmacy and Pharmacology*
- *Journal of Dentistry and Oral Hygiene*
- *International Journal of Nursing and Midwifery*
- *Journal of Parasitology and Vector Biology*
- *Journal of Pharmacognosy and Phytotherapy*
- *Journal of Toxicology and Environmental Health Sciences*

academicJournals