



International Journal of Science and Technology Education Research

Volume 8 Number 2 September 2017

ISSN 2141-6559



*Academic
Journals*

ABOUT IJSTER

The **International Journal of Science and Technology Education Research (IJSTER)** IJSTER is an open access journal. Abstracts and full texts of all articles published in the journals can be read online without any form of restriction

International Journal of Science and Technology Education Research (IJSTER) International Journal of Science and Technology Education Research (IJSTER) is a peer reviewed open access journal. The journal is published monthly and covers all areas of the subject.

Contact Us

Editorial Office: ijster@academicjournals.org

Help Desk: helpdesk@academicjournals.org

Website: <http://www.academicjournals.org/journal/IJSTER>

Submit manuscript online <http://ms.academicjournals.me/>

Editors

Prof. S. C. Garg

School of Sciences

Indira Gandhi National Open University

New Delhi,

India.

Assoc. Prof. Anita Seth

Institute of Engineering and Technology

DAVV University

Indore,

India.

Dr. Christine Redman

The University of Melbourne

Melbourne Graduate School of Education

Australia.

Dr. Laban P. Ayiro

Curriculum Research Services

Nairobi,

Kenya.

Paola Forcheri

National Research Council

Institute for Applied Mathematics and Information

Technologies

Italy.

Dr. Eduardo Galembeck

State University of Campinas

São Paulo,

Brazil.

ARTICLE

| | |
|--|-----------|
| Single basis function method for solving diffusion convection equations | 13 |
| Sunday Babuba | |

Full Length Research Paper

Single basis function method for solving diffusion convection equations

Sunday Babuba

Department of Mathematics, Federal University Dutse, Ibrahim Aliyu Bye-Pass, P. M. B. 7156, Dutse, Jigawa State, Nigeria.

Received 26 June, 2017; Accepted 24 August, 2017

In this study, we developed a new numerical finite difference method for solving various diffusion convection equations. The method involves reduction of the diffusion convection equations to a system of algebraic equations. By solving the system of algebraic equations we obtain the problem approximate solutions. The study of the numerical accuracy of the method has shown that the method provides similar results to the known explicit finite difference method for solving diffusion convection equations, but with fewer numbers of iterations.

Key words: Lines, multistep collocation, parabolic, Taylor's polynomial.

INTRODUCTION

In this study, we will deal with diffusion convection equations, where t and x are the time and space coordinates respectively, and the quantities h and k are the mesh sizes in the space and time directions.

We consider,

$$\frac{\partial U(x,t)}{\partial t} = V \frac{\partial^2 U(x,t)}{\partial x^2} + \bar{U} \frac{\partial U(x,t)}{\partial x}, \quad (1)$$

A special type of diffusion convection equation where $V, \bar{U} > 0$ are constants, defined for $a < x < b$ and positive time with appropriate initial and boundary conditions, and U is the temperature in the medium.

We are interested in the development of numerical techniques for solving diffusion convection equations. Recently, there is a growing interest concerning continuous numerical methods of solution for Ordinary Differential Equations (ODEs) (Adam and David, 2002; Awoyemi, 2002, 2003; Bao et al., 2003; Benner and Mena, 2004; Bensoussan et al., 2007; Motmans et al., 2005; Brown, 1979; Chawla and Katti, 1979; Cook, 1974; Crandall, 1955; Crane and Klopfenstein, 1965; Crank and Nicolson, 1947; Dahlquist and Bjorck, 1974; Dehghan, 2003; Dieci, 1992; Douglas, 1961; D' Yakonov and Ye, 1963; Eyaya, 2010; Fox, 1962; Penzl, 2000; Pierre, 2008; Richard and Albert, 1981; Richard et al., 2001; Saumaya et al., 2012; Yildiz and Subasi, 2001; Zheyin and Qiang, 2012). We are interested in the extension of a particular

E-mail: sundaydzupu@yahoo.com.

Authors agree that this article remain permanently open access under the terms of the [Creative Commons Attribution License 4.0 International License](https://creativecommons.org/licenses/by/4.0/)

continuous method to solve the diffusion convection equation. This is done based on the collocation and interpolation of the Partial Differential Equations (PDEs) directly over multi steps along lines but with reduction to a system of ODEs.

THE SOLUTION METHOD

We subdivide the interval $0 \leq x \leq b$ into N equal subintervals by the grid points $x_m = mh, m = 0, \dots, N$ where $Nh = b$. We want to obtain an approximation of the form

$$U(x, t) \approx U_K(x, t) = \sum_{r=0}^{k+1} a_r Q_r(x, t), \quad x_{i-1} \leq x \leq x_{i+1},$$

$$r = 0, 1, 2, \dots, k + 1 \tag{2}$$

Where $Q_r(x, t) = x^r t^r$ are canonical polynomials which are used as basic functions in the approximation, and a_r are parameters to be determined.

From the collocation equations

$$U_K(x_m, t) = \sum_{r=0}^k a_r Q_r(x_m, t) \tag{3}$$

where $m = i - 1, j, i + 1, j$

We can generate the following equations from Equation 3 as follows:

$$\left. \begin{aligned} U_k(x_{i-1}, t_j) &= a_0 + a_1 x_{i-1} t_j + \dots + a_k x_{i-1}^k t_j^k \approx U_{i-1, j} \\ U_k(x_i, t_j) &= a_0 + a_1 x_i t_j + \dots + a_k x_i^k t_j^k \approx U_{i, j} \\ U_k(x_{i+1}, t_j) &= a_0 + a_1 x_{i+1} t_j + \dots + a_k x_{i+1}^k t_j^k \approx U_{i+1, j} \end{aligned} \right\} \tag{4}$$

Writing Equation 4 as matrix in its augmented form, we have

$$\begin{bmatrix} 1 & x_{i-1} t_j & \dots & x_{i-1}^k t_j^k \\ 1 & x_i t_j & \dots & x_i^k t_j^k \\ 1 & x_{i+1} t_j & \dots & x_{i+1}^k t_j^k \end{bmatrix} \begin{bmatrix} a_0 \\ \dots \\ a_k \end{bmatrix} = \begin{bmatrix} U_{i-1, j} \\ U_{i, j} \\ U_{i+1, j} \end{bmatrix} \tag{5}$$

We solve Equation 5 for the value of a_k by Gaussian elimination method to obtain

$$a_k = \dots + \frac{U_{i+1, j} + U_{i-1, j} - U_{i, j}}{2ht^k_j},$$

From Equation (2) we can generate

$$U(x, y) = a_0 + a_1 xt + \dots + a_k x^k t^k \tag{6}$$

Putting the value of a_k in Equation 6, we obtain

$$U(x, t) = a_0 + a_1 xt + \dots + x^k t^k \left(\frac{U_{i+1, j} + U_{i-1, j} - 2U_{i, j}}{2ht^k_j} \right) \tag{7}$$

We take the first and second derivatives of Equation 7 with respect to x and obtain

$$U'(x, t) = a_1 t + \dots + k x^{k-1} t^k \left(\frac{U_{i+1, j} + U_{i-1, j} - 2U_{i, j}}{2ht^k_j} \right)$$

$$U''_i(x, t) = \dots + k(k-1) x^{k-2} t^k \left(\frac{U_{i+1, j} + U_{i-1, j} - 2U_{i, j}}{2ht^k_j} \right) \tag{8}$$

From the collocation Equation 3

$$U_K(x_n, t) = \sum_{r=0}^k a_r Q_r(x_n, t)$$

where $n = i - 1, j$ and $i + 1, j$

We generate the following equations as follows:

$$\left. \begin{aligned} U_k(x_{i-1}, t_j) &= a_0 + a_1 x_{i-1} t_j + \dots + a_k x_{i-1}^k t_j^k \approx U_{i-1, j} \\ U_k(x_{i+1}, t_j) &= a_0 + a_1 x_{i+1} t_j + \dots + a_k x_{i+1}^k t_j^k \approx U_{i+1, j} \end{aligned} \right\} \tag{9}$$

Writing Equation 9 as matrix in its augmented form, we have

$$\begin{bmatrix} 1 & x_{i-1} t_j & \dots & x_{i-1}^k t_j^k \\ 1 & x_{i+1} t_j & \dots & x_{i+1}^k t_j^k \end{bmatrix} \begin{bmatrix} a_0 \\ \dots \\ a_k \end{bmatrix} = \begin{bmatrix} U_{i-1, j} \\ U_{i+1, j} \end{bmatrix} \tag{10}$$

Solving Equation 10 for a_k we obtain

$$a_k = \frac{U_{i+1,j} - U_{i-1,j}}{2ht^k_j},$$

but from Equation 2 we obtain

$$U(x,t) = a_0 + a_1xt + \dots + a_kx^k t^k \tag{11}$$

Putting the value of a_k in Equation 11

$$U(x,t) = a_0 + a_1xt + \dots + x^k t^k \left(\frac{U_{i+1,j} - U_{i-1,j}}{2ht^k_j} \right) \tag{12}$$

Taking the first derivative of Equation 12 with respect to x , we obtain

$$U'(x,t) = k(x^{k-1}t^k) \left(\frac{U_{i+1,j} - U_{i-1,j}}{2ht^k_j} \right) \tag{13}$$

Similarly, by interchanging the roles of x and t we can obtain an approximation of the form

$$U(t,x) \approx U_k(t,x) = \sum_{r=0}^{k+1} a_r Q_r(t,x), \quad t_{j-1} \leq t \leq t_{j+1}, \tag{14}$$

$$r = 0, 1, 2, \dots, k+1$$

Again, from the collocation equation

$$U_K(x,t_g) = \sum_{r=0}^k a_r Q_r(x,t_g), \quad g = i, j \text{ and } i, j+1 \tag{15}$$

We have the following equations:

$$\left. \begin{aligned} U_k(t_j, x_i) &= a_0 + a_1 t_j x_i + \dots + a_k t_j^k x_i^k \approx U_{i,j} \\ U_k(t_{j+1}, x_i) &= a_0 + a_1 t_{j+1} x_i + \dots + a_k t_{j+1}^k x_i^k \approx U_{i,j+1} \end{aligned} \right\} \tag{16}$$

Writing Equation 16 as matrix in augmented form, we have

$$\begin{bmatrix} 1 & x_i t_j & \dots & x_i^k t_j^k \\ 1 & x_i t_{j+1} & \dots & x_i^k t_{j+1}^k \end{bmatrix} \begin{bmatrix} a_0 \\ \dots \\ \dots \\ a_k \end{bmatrix} = \begin{bmatrix} U_{i,j} \\ U_{i,j+1} \end{bmatrix} \tag{17}$$

We solve Equation 17 for a_k by using Gaussian elimination method

$$a_k = \dots + \frac{U_{i,j+1} - U_{i,j}}{2hx^k_j},$$

Again, from Equation 2 we obtain

$$U(t,x) = a_0 + a_1xt + \dots + a_kx^k t^k \tag{18}$$

Substituting the value of a_k in Equation 18 we have

$$U(t,x) = a_0 + a_1xt + \dots + x^k t^k \left(\frac{U_{i,j+1} - U_{i,j}}{2hx^k_i} \right) \tag{19}$$

Taking the first derivative of Equation 19 with respect to t , we obtain

$$U'_j(x,t) = \dots + kt^{k-1}x^k \left(\frac{U_{i,j+1} - U_{i,j}}{2hx^k_i} \right), \tag{20}$$

We collocate Equation (20) at $x = x_i$, Equations 8 and 13 at $t = t_j$; and substituting the resulting equations into Equation 1, we obtain a scheme that solves diffusion convection equations numerically.

ADVANTAGE OF THE NEW SCHEME OVER FINITE DIFFERENCE METHOD

The new scheme requires fewer numbers of iterations than the finite difference method to achieve the same results.

NUMERICAL EXAMPLES

Here, we will test the numerical accuracy of the new method by using the scheme to solve two test examples. We compute an approximate solution of Problems (1) and (2) at each time level (Tables 1 and 2). To achieve this, we truncate the polynomials after second degree.

SPECIFIC PROBLEM

Example 1

Here we use the scheme to approximate the solution to

Table 1. Result of action of the scheme on Problem 1.

| x | Exact solution $U(x,t)$ | Solution from New Method $U(x,t)$ | Solution from Finite Difference Method $U(x,t)$ | Error from Finite Difference Method | Error from New Method |
|------|-------------------------|-----------------------------------|---|-------------------------------------|-----------------------|
| 0 | 0 | 0 | 0 | 0 | 0 |
| 0.25 | 0.380814721 | 0.380862804 | 0.380862804 | $4.8 \times E-5$ | $4.8 \times E-5$ |
| 0.50 | 0.703723471 | 0.703742699 | 0.703742699 | $1.9 \times E-5$ | $1.9 \times E-5$ |
| 0.75 | 0.919471568 | 0.919484148 | 0.919484148 | $1.3 \times E-5$ | $1.3 \times E-5$ |
| 1.00 | 0.995167871 | 0.99524247 | 0.99524247 | $7.5 \times E-5$ | $7.5 \times E-5$ |
| 1.25 | 0.919471568 | 0.919484148 | 0.919484148 | $1.3 \times E-5$ | $1.3 \times E-5$ |
| 1.5 | 0.703723471 | 0.703742699 | 0.703742699 | $1.9 \times E-5$ | $1.9 \times E-5$ |
| 1.75 | 0.380814721 | 0.380862804 | 0.380862804 | $4.8 \times E-5$ | $4.8 \times E-5$ |
| 2.00 | 0 | 0 | 0 | 0 | 0 |

Table 2. Results of action of the scheme on Problem 2.

| x | Exact Solution $U(x,t)$ | Solution from Finite Difference method $U(x,t)$ | Solution from New Method $U(x,t)$ | Error from finite difference Method | Error from New Method |
|------|-------------------------|---|-----------------------------------|-------------------------------------|-----------------------|
| 0 | 0 | 0 | 0 | 0 | 0 |
| 0.10 | 0.30806537 | 0.30807172 | 0.30807172 | $6.3 \times E-6$ | $6.3 \times E-6$ |
| 0.20 | 0.585975167 | 0.58598723 | 0.58598723 | $1.20 \times E-5$ | $1.20 \times E-5$ |
| 0.30 | 0.806525626 | 0.80654224 | 0.80654224 | $1.7 \times E-5$ | $1.7 \times E-5$ |
| 0.40 | 0.948127737 | 0.948147264 | 0.948147264 | $2.0 \times E-5$ | $2.0 \times E-5$ |
| 0.50 | 0.9969205 | 0.996941032 | 0.996941032 | $2.1 \times E-5$ | $2.1 \times E-5$ |
| 0.60 | 0.948127737 | 0.948147264 | 0.948147264 | $2.0 \times E-5$ | $2.0 \times E-5$ |
| 0.70 | 0.806525626 | 0.80654224 | 0.80654224 | $1.7 \times E-5$ | $1.7 \times E-5$ |
| 0.80 | 0.585975167 | 0.58598723 | 0.58598723 | $1.20 \times E-5$ | $1.20 \times E-5$ |
| 0.90 | 0.30806537 | 0.30807172 | 0.30807172 | $6.3 \times E-6$ | $6.3 \times E-6$ |
| 1.00 | 0 | 0 | 0 | 0 | 0 |

the diffusion convection equation (Table 1)

$$\frac{\partial U}{\partial t} = \frac{\partial^2 U}{\partial x^2} + 2 \frac{\partial U}{\partial x} \quad 0 < x < 20, 0 < t < 20$$

$$U(x,0) = |10 - x|, 0 \leq x \leq 20$$

$$U(0,t) = U(20,t) = 0, \quad 0 \leq t \leq 20$$

Example 2

Here we use the scheme to approximate the solution to the diffusion convection equation (Table 2)

$$\frac{\partial U}{\partial t} = \frac{\partial^2 U}{\partial x^2} + 2 \frac{\partial U}{\partial x} \quad 0 < x < 20, 0 < t < 20$$

$$U(x,0) = \frac{x}{2}, 0 \leq x \leq 20$$

$$U(0,t) = 0, U(20,t) = 0, \quad 0 \leq t \leq 20$$

Conclusion

A numerical method which involves the discretization of the diffusion convection equations to a system of algebraic equations is formulated for solving diffusion convection problems. To check the accuracy of the numerical method, it is applied to solve two different test problems with known finite difference solutions. The overall numerical results obtained showed that the values obtained are the same with the results of the finite difference method. These confirmed the validity of the new numerical scheme and suggested that it is an interesting and viable numerical method.

CONFLICT OF INTERESTS

The author has not declared any conflict of interests.

REFERENCES

- Adam A, David R (2002). One dimensional heat equation. <http://www.ng/online.redwoods.cc.ca.us/instruct/darnold/deproj/sp02/.../paper.pdf>
- Awoyemi DO (2002). An Algorithmic collocation approach for direct solution of special fourth-order initial value problems of ordinary differential equations. *J. Nigerian Association of Math. Phys.* 6:271-284.
- Awoyemi DO (2003). A p-stable linear multistep method for solving general third order Ordinary differential equations. *Int. J. Comput. Math.* 80(8):987-993.
- Bao W, Jaksch P, Markowich PA (2003). Numerical solution of the Gross – Pitaevskii equation for Bose – Einstein condensation. *J. Compt. Phys.* 187(1):318-342.
- Benner P, Mena H (2004): BDF methods for large scale differential Riccati equations. *Proc. of Mathematical theory of network and systems. MTNS.* Edited by Moore, B. D., Motmans, B., Willems, J., Dooren, P.V. & Blondel, V.
- Bensoussan A, Da Prato G, Delfour M, Mitter S (2007). Representation and control of infinite dimensional systems. 2nd edition. Birkhauser: Boston, MA.
- Motmans B, Willems J, Dooren PV, Blondel V, Biazar J, Ebrahimi H (2005). An approximation to the solution of hyperbolic equation by a domain decomposition method and comparison with characteristics method. *Appl. Math. Comput.* 163:633-648.
- Brown PLT (1979). A transient heat conduction problem. *AICHE J.* 16:207-215.
- Chawla MM, Katti CP (1979). Finite difference methods for two-point boundary value problems involving high-order differential equations. *BIT.* 19:27-39.
- Cook RD (1974). *Concepts and Application of Finite Element Analysis:* NY: Wiley Eastern Limited.
- Crandall SH (1955). An optimum implicit recurrence formula for the heat conduction equation. *Q. Appl. Math.* 13(3):318-320.
- Crane RL, Klopfenstein RW (1965). A predictor – corrector algorithm with increased range of absolute stability. *JACM* 12:227-237.
- Crank J, Nicolson P (1947). A practical method for numerical evaluation of solutions of partial differential equations of heat conduction type. *Proc. Camb. Phil. Soc.* 6:32-50.
- Dahlquist G, Bjorck A (1974). *Numerical methods.* NY: Prentice Hall.
- Dehghan M (2003). Numerical solution of a parabolic equation with non-local boundary specification. *Appl. Math. Comput.* 145, 185-194.
- Dieci L (1992). *Numerical analysis.* SIAM J. 29(3):781-815.
- Douglas J (1961). A survey of numerical methods for parabolic differential equations in advances in computer II. Academic Press.
- D' Yakonov, Ye G (1963). On the application of disintegrating difference operators. *USSR Computational Mathematics and Mathematical Physics*, 3(2):511-515.
- Eyaya BE (2010). Computation of the matrix exponential with application to linear parabolic PDEs. http://www.dip.sun.ac.za/~eyaya/PGD-Essay-Template-2009_10.pdf
- Fox L (1962). *Numerical Solution of Ordinary and Partial Differential Equation.* New York: Pergamon.
- Penzl T (2000). A cyclic low-rank Smith method for large sparse Lyapunov equations. *SIAM J. Sci. Comput.* 21(4):1401-1418
- Pierre J (2008). Numerical solution of the dirichlet problem for elliptic parabolic equations. *SIAM J. Soc. Ind. Appl. Math.* 6(3):458-466.
- Richard LB, Albert C (1981). *Numerical analysis.* Berlin: Prindle, Weber and Schmidt, Inc.
- Richard L, Burden J, Douglas F (2001). *Numerical analysis.* Seventh ed., Berlin: Thomson Learning Academic Resource Center.
- Saumaya B, Neela N, Amiya YY (2012). Semi discrete Galerkin method for equations of Motion arising in Kelvin – Voigt model of viscoelastic fluid flow. *J. Pure Appl. Sci.* 3(2 & 3):321-343.
- Yildiz B, Subasi M (2001). On the optimal control problem for linear Schrodinger equation. *Appl. Math. Comput.* 121:373-381.
- Zheyin HR, Qiang X (2012). An approximation of incompressible miscible displacement in porous media by mixed finite elements and symmetric finite volume element method of characteristics. *Appl. Math. Computat.* Elsevier 143:654-672.

International Journal of Science and Technology Education Research

Related Journals Published by Academic Journals

- Journal of Media and Communication Studies
- International Journal of English and Literature
- Philosophical Papers and Reviews
- Educational Research and Reviews
- Journal of African Studies and Development
- Philosophical Papers and Reviews

academicJournals