



International Journal of Fisheries and Aquaculture

Volume 9 Number 9 September 2017

ISSN 2006-9839



*Academic
Journals*

ABOUT IJFA

The International Journal of Fisheries and Aquaculture (IJFA) (ISSN: 2006-9839) is an open access journal that provides rapid publication (monthly) of articles in all areas of the subject such as algaculture, Mariculture, fishery in terms of ecosystem health, Fisheries acoustics etc.

The Journal welcomes the submission of manuscripts that meet the general criteria of significance and scientific excellence. Papers will be published shortly after acceptance. All articles published in the IJFA are peer-reviewed.

Contact Us

Editorial Office: ijfa@academicjournals.org

Help Desk: helpdesk@academicjournals.org

Website: <http://www.academicjournals.org/journal/IJFA>

Submit manuscript online <http://ms.academicjournals.me/>

Editors

Dr. V.S. Chandrasekaran

*Central Institute of Brackishwater Aquaculture
Indian Council of Agricultural Research (ICAR)
Chennai,
India.*

Prof. Nihar Rajan Chattopadhyay

*Department of Aquaculture
Faculty of Fishery Sciences
West Bengal University of Animal & Fishery Sciences
West Bengal,
India.*

Dr. Lourdes Jimenez-Badillo

*Ecology and Fisheries Centre
Universidad Veracruzana
Veracruz,
México.*

Dr. Kostas Kapiris

*Institute of Marine Biological
Resources of H.C.M.R.
Athens,
Greece.*

Dr. Masoud Hedayatifard

*Department of Fisheries Sciences and Aquaculture
College of Agriculture and Natural Resources
Advanced Education Center
Islamic Azad University
Ghaemshahr,
Iran.*

Dr. Zhang Xiaoshuan

*China Agricultural University
Beijing,
China.*

Dr Joseph Selvin

*Marine Bioprospecting Lab
Dept of Microbiology
Bharathidasan University
Tiruchirappalli,
India.*

Editorial Board

Dr. Dada Adekunle Ayokanmi
*Department of Fisheries and Aquaculture
Federal University of Technology
Akure,
Nigeria.*

Dr. Ramasamy Harikrishnan
*Faculty of Marine Science
College of Ocean Sciences
Jeju National University
Jeju City,
South Korea.*

Dr. Kawser Ahmed
*Lab. of Ecology, Environment and Climate Change
Department of Fisheries
University of Dhaka
Dhaka,
Bangladesh.*

Dr. Maxwell Barson
*Biological Sciences Department
University of Zimbabwe
Harare,
Zimbabwe.*

Dr. Christopher Marlowe Arandela Caipang
*Faculty of Biosciences and Aquaculture
Bodø University College
Bodø,
Norway.*

Dr. William S. Davidson
*Department of Molecular Biology and Biochemistry
Simon Fraser University
Burnaby,
Canada.*

Dr. Babak Ghaednia
*Iran Shrimp Research Center (ISRC)
Bushehr,
Iran.*

Prof. Ratha Braja Kishore
*Department of Zoology
Biochemical Adaptation Laboratory
Banaras Hindu University
Varanasi,
India.*

Prof B. Sharma
*Department of Biochemistry
Centre for Biotechnology
University of Allahabad
Allahabad,
India.*

Dr. Sebastián Villasante
*Fisheries Economics and Natural Resources Research Unit
University of Santiago de Compostela
A Coruña,
Spain.*

Dr. Mohamed Hamed Yassien
*National Institute of Oceanography and Fisheries
Suez,
Egypt.*

Dr. Riaz Ahmad
*Department of Zoology
Aligarh Muslim University
Aligarh,
India.*

Dr. SM Nurul Amin
*Lab. Marine Science & Aquaculture
Institute of Bioscience
Universiti Putra Malaysia
Selangor,
Malaysia.*

Dr. Melcher Andreas
*University of Natural Resources and Applied Life Sciences
Vienna,
Austria.*

Dr. Ayotunde Ezekiel Olatunji
*Department of Fisheries and Aquatic Science
Cross River University of Technology
Calabar,
Nigeria.*

Dr. Ramachandra Bhatta
*Animal and Fisheries Sciences University
College of Fisheries
Kankanady,
India.*

Dr. Subha Ganguly
*Department of Fish Processing Technology
Faculty of Fishery Sciences
West Bengal University of Animal and Fishery Sciences
Chakgaria,
India.*

Dr. Yusuf Bozkurt
*Iskenderun Technical University
Faculty of Marine Sciences and Technology
Iskenderun Hatay,
Turkey.*

Prof. Rina Chakrabarti
*Aqua Research Lab
Department of Zoology
University of Delhi
Delhi,
India.*

Prof. Nihar Rajan Chattopadhyay
Faculty of Fishery Sciences
University of Animal and Fishery Sciences
Chakgaria,
India.

Dr. Sullip Kumar Majhi
Division of Fisheries
Indian Council of Agricultural Research Complex for Neh
Region
Meghalaya,
India.

Dr. Hooman Rahmati-Holasoo
Department of Aquatic Animal Health
Faculty of Veterinary Medicine
University of Tehran
Tehran,
Iran.

Dr. Liberty Napilan-Espectato
Institute of Fisheries Policy and Development Studies
College of Fisheries and Ocean Sciences
Iloilo,
Philippines.

Dr. Zeinab Abdel-Baki El-Greisy
National Institute of Oceanography and Fisheries (NIOF),
KayetBey,
Egypt.

Dr. Thanitha Thapanand-Chaidee
Department of Fishery Biology
Faculty of Fisheries
Kasetsart University
Chatujak,
Thailand.

Dr. Gabriel Ugwemorubong Ujagwung
Dept. of Fisheries and Aquatic Environment
Faculty of Agriculture
Rivers State University of Science and Technology
Port Harcourt,
Nigeria.

Dr. Margherita Gloria
School of Agriculture
Food Science and Veterinary Medicine
University College Dublin
Belfield,
Ireland.

Dr. Derya Guroy
Canakkale Onsekiz Mart University
Faculty of Fisheries
Aquaculture Department
Canakkale,
Turkey.

Dr. Daniela Giannetto
Department of Biology
Faculty of Sciences
Muğla Sıtkı Koçman University
Turkey.

Dr. Terngu Akaahan
Biological Sciences Department
Federal University of Agriculture Makurdi
Nigeria.

ARTICLE

Assessment of monthly physico-chemical properties and fish yields of two micro dams of Tigray Region, Northern Ethiopia	92
Haftom Zebib and Tsegay Teame	

Full Length Research Paper

Assessment of monthly physico-chemical properties and fish yields of two micro dams of Tigray Region, Northern Ethiopia

Haftom Zebib* and Tsegay Teame

Abergelle Agricultural Research Center, Tigray Agricultural Research Institute, Fishery and Other Aquatic Life Case Team, Abi-Adi, Ethiopia.

Received 25 November, 2016; Accepted 17 July, 2017

This study was conducted to assess some physico-chemical properties and yields of two micro dams (Korrir and Laelay Wukro) located in eastern zones of Tigray region (Northern Ethiopia). In each location, water samples and fish yield were examined from October 2013 to March 2014 once per month in each dams. The examined physico-chemical parameters were: Dissolved oxygen (DO), electrical conductivity (EC), total dissolved solids (TDS), water temperature, pH and transparency. The result showed that the monthly record values of DO, EC and TDS decreased while water temperature and transparency increased in studied months. Mean monthly pH value also fluctuated across months. The monthly mean value of DO, pH, temperature, EC, TDS and transparency were ranged: 3.20-7.70 mg/L, 8.23-9.14, 15.3-22.2°C, 284-353 μ S/cm, 1.47-2.11 mg/L and 19.00-39.00 cm in Korrir dam while in Laelay Wukro were: 3.44-5.70 mg/L, 7.93-8.61, 19.4-24.2°C, 209-462 mg/L, 110-270 μ S/cm and 18.00-40.20 cm, respectively. Fish yield was assessed by morphoedaphic index empirical models. There was no significant ($p < 0.05$) difference in fish yield and the average estimated productivity of Korrir and Laelay Wukro dams were 13.99 and 14.47 quintal/ha/year respectively. Therefore physico-chemical properties of these micro dams were good for fish production and fish farmers should practice utilization of the micro dams through good management of the water bodies.

Key words: Physico-chemical properties, fish yields, micro dams, morphoedaphic index.

INTRODUCTION

Water quality refers to all physical, chemical and biological characteristics of the water and plays an important role in the growth and survival of aquatic organisms. Relationship between water quality and aquatic productivity is a pre-requisite for obtaining optimum growth and production of aquatic organisms including fish. The water used for aquaculture and fish production would not give the desired production unless

the prevailing water quality parameters are optimum for the organism under culture (Bisht et al., 2013). Some of the physico-chemical parameters that are regularly measured within water bodies like pond include temperature, dissolved oxygen (DO), alkalinity, hardness, pH, electrical conductivity (EC), turbidity, total dissolved solids (TDS) and biological oxygen demand (USDA, 1996). Dissolved oxygen is required for respiration by

*Corresponding author. E-mail: hzhzebib6@gmail.com. Tel: +251344460741.

most aquatic animals. Apart from this, dissolved oxygen combined with other important elements such as Carbon, Sulphur, Nitrogen and Phosphorous that could have been toxicants in the absence of oxygen in the water bodies to form carbonate, sulphate, nitrate and phosphate respectively that constitute the required compounds for aquatic organisms for survival (Araoy, 2009). Each species of fish has a preferred or optimum temperature range where it grows best. At temperatures above or below optimum, fish growth is affected. Mortalities may occur at extreme temperatures (Piper et al., 1982). Turbidity determines light penetration in water. This in turn will have an effect on the temperature of the water and the amount of vegetation and algae that will grow in the pond, thus affecting the rate of photosynthesis and primary productivity (USDA, 1996; Environmental Review, 2008).

The annual fish production potential of Ethiopia based on empirical methods on individual lake surface area and mean depth of major water bodies was estimated to be 30,000 to 51,000 tons (FAO, 2003). The yield of fish, season change in fish density and the factors which influence them are important for the sustainable management of fisheries in the reservoirs (Tadesse et al., 2015). Currently, government of Ethiopia is engaged in harvesting rain water for the purpose of irrigation. Thus, in Tigray, more than 70 micro dams have been constructed for enhancing food production through integrated irrigation and fish farming (Tadesse et al., 2008). The governments have a strategy to make fishery and aquaculture sector as one on the method to reduce poverty in the country.

Considerable information exists on the limnology of the natural lakes of Ethiopia but little is known about the country's dams, rivers and reservoirs (Zenebe et al., 2012). Atakilt and Tadesse (2012) studied population dynamics and condition factor of *Oreochromis niloticus* L. in Korrir and Laelay Wukro dams. Tadesse et al. (2015) also reported condition factor of *O. niloticus* L. and yield in five tropical small and samples were collected in four months (December, 2012, March, April and May 2013). Araoy (2009) and Ovie et al. (2011) reported that monthly water physico-chemical variations have impact on fish growth and production potential. Therefore recent information on monitoring physico-chemical properties and estimating potential yields of these water bodies is useful to understand fish production potential, status, sustainability and select the most suitable way of area utilization. In this study the main physico-chemical water parameters (DO, pH, temperature, EC, TDS, transparency) and their potential fish yields of the *Korrir and Laelay Wukro* micro dams were examined.

MATERIALS AND METHODS

Description of study area

The study was conducted in Korrir and Laelay Wukro earthen micro

dams which are found in eastern zone of Tigray region constructed in 1995/1996 and 1997/1998, respectively. Korrir lies between Longitude 13°44'446" and Latitude 39°36'759" at altitude 2026 m while Laelay Wukro Longitude 13°48'331" and Latitude 39°36'735" at altitude of 2025 m (Figure 1).

Physico-chemical properties

The main physico-chemical parameters were measured from October 30, 2013 to March 30, 2014 once per month at three representative stations in the earthen micro dams and the average of reading record was used in each month. The pH was determined with Elmetron pH meter (Model: CP-411). The pH meter was calibrated using buffer solution of pH value of 7. Electrical conductivity and total dissolved solid was determined using Elmetron conductivity meter (Model: CC-411). Dissolved oxygen was measured using Elmetron oxygen meter (Model: CCO-401). Surface water temperature was measured by Elmetron oxygen meter installed with temperature electrode. The pH, conductivity, oxygen and temperature electrodes were immersed in water 6 cm below the water surface for about 4 to 5 min until stabilized reading is obtained. Transparency (black and white disk) was measured by lowering the disk by hand into the water to the depth at which it vanishes from sight. The difference of distance to vanishing and appeared point was recorded and then divided by two.

$$\text{Secchidisktransparency} = \left(\frac{A+B}{2} \right)$$

Where A = depth at which secchi disk disappear, and B = depth at which secchi disk reappears.

Fish yield estimation from morphoedaphic index (MEI)

The MEI has been used to estimate fish yield and production in reservoirs and lakes. Potential fish yield were estimated from three different empirical models developed by Henderson and Welcome (1974), Toews and Griffith (1979) and Marshall (1984). The area and depth were taken from literature and used in the empirical yield models. Where;

$$y = 14.3136 \text{MEI}^{0.4681} \text{ (Henderson and Welcome, 1974)}$$

$$\text{Log}(y) = 1.4071 + 0.3697 \text{Log}(\text{MEI}) - 0.00004565A \text{ (Toews and Griffith, 1979)}$$

$$\text{Ln}(Y) = 3.57 + 0.76 \ln(A) \text{ (Marshall, 1984)}$$

$$\text{Morphoedaphic index (MEI)} = \frac{\text{Conductivity} \left(\frac{\mu\text{mhos}}{\text{cm}} \right)}{\text{Mean depth}(m)}$$

Where, A = Surface area.

Statistical analysis

The one way ANOVA was used to test for significant variations between means of monthly physico-chemical parameters and fish yields using SAS institute and Cary, NC version 9.0. Duncan's multiple range tests was used to identify significant differences at




Figure 1. Map of the study areas.

significance level ($p < 0.05$).

RESULTS AND DISCUSSION

Dissolved oxygen

The mean DO value (4.77 and 4.85 mg/L) was observed in Korrir and Laelay Wukro dams respectively and monthly DO was decreased from October to March in the dams (Table 1). Tadesse et al. (2015) reported higher average DO value 7.03 mg/L from five tropical small dams (Korrir, Laelay Wukro, Mai Nigus, Mai Sessea and Mai Seye dams). Atakilt and Tadesse (2012) also reported 7.19 mg/L average DO value for Korrir and Laelay Wukro dams for one year data which were higher than the current study results. Mustapha (2008) observed low DO concentration value due to high temperature record and high rate of decomposition in the dry season which is similar with current findings. Boyd (1979) reported that DO concentrations of 3 to 12 mg/L will

promote the growth and survival of fish in reservoirs. The minimum DO of 5.0 mg/L was reported for tropical fishes by Saloom and Duncan (2005). Singh (2007) reported that the minimum concentration of DO, 4 mg/L should be maintained in fish ponds at all times. The water quality for any fish cultured in tropical region must be such that the DO concentration must not be less than 3 mg/L (Robert, 2007). Brian (2006) and Ita et al. (1995) noted that increased DO level is needed to support an increase in metabolic rates and reproduction. Dastagir et al. (2014) reported that DO requirements of any fish species varied with the age, temperature and concentration of minerals in water.

pH

The mean pH values recorded in the present study was 8.19 and 8.57 in Laelay Wukro and Korrir dams, and values fluctuated across months (Table 1). This mean values which was within the pH range of 6.5 to 9.0

Table 1. The value of monthly physico-chemical variations of the dams in the studied months.

Months	DO (mg/L)		pH		Temperature (°C)		EC (µS/cm)		TDS (mg/L)		Transparency (cm)	
	Korrir	Laelay Wukro	Korrir	Laelay Wukro	Korrir	Laelay Wukro	Korrir	Laelay Wukro	Korrir	Laelay Wukro	Korrir	Laelay Wukro
Oct	7.70 ^a	5.70 ^b	8.55	7.97	18.30	19.40	353	462	211	270	19.00 ^a	18.00 ^a
Nov	4.89 ^a	5.25 ^b	8.31	7.93	19.90	21.80	286	446	168	240	22.00 ^b	21.60 ^b
Dec	4.55 ^a	5.20 ^b	8.40	8.04	20.60	22.00	279	422	167	210	25.00 ^a	28.50 ^b
Jan	4.40 ^a	5.00 ^b	9.14	8.20	20.60	22.00	276	357	166	180	30.00 ^a	33.00 ^b
Feb	3.90 ^a	4.50 ^b	8.80	8.61	21.80	22.50	260	252	154	154	34.00	36.00
Mar	3.20 ^a	3.44 ^a	8.23	8.41	22.20	24.20	248	209	147	110	39.00	40.20
Min	3.20	3.44	8.23	7.93	18.30	19.40	248	209	147	110	19.00	18.00
Max	7.70	5.70	9.14	8.61	22.20	24.20	353	462	211	270	39.00	40.20
Mean	4.77±1.55 ^a	4.85±0.79 ^a	8.57±0.34 ^b	8.19±0.27 ^b	20.50±1.39 ^c	21.98±1.54 ^c	284±36.68 ^d	358±105.92 ^e	169±22.30 ^f	194±58.31 ^f	28.17±7.57 ^g	29.55±8.54 ^g

Mean values in column with the same letter are not significantly different ($p < 0.05$).

reported by Boyd and Tucker (1998) and Ali et al. (2000) is suitable for diverse fish production. Any variation beyond acceptable range could be fatal to many aquatic organisms (Iqbal et al., 2004). Depending on species, fish can generally be vulnerable or resistant to high or low pH (Ovie et al., 2011). Warm waters develop increased pH levels during daylight by photosynthesis and pH declines by the process of respiration at night (King, 1970).

Water temperature

The mean temperature record value (20.07 and 21.98°C) was increased across months in Korrir and Laelay Wukro dams respectively which were within range (20-30°C) reported by Boyd (1990). Atakilt and Tadesse (2012) found 18.88°C average temperature value for Korrir and Laelay Wukro dams for one year data. The above temperature range can still be considered as good for growth and body metabolism of the fishes in the water body. This is because intensity of metabolism of warm water fish is closely

associated with water temperature, that is, the higher the temperature, the higher the metabolic rate (Ovie et al., 2011). Variations in water temperature in the dry season can be attributed to intensified heat radiation and effect of harmattan (Olele and Ekelemu, 2008). Bishat et al. (2013) observed wide fluctuation in temperature in earthen pond, cemented ponds and open water body might be due to variation in volume of water.

EC

EC with mean value 284 and 358 µS/cm in Korrir and Laelay Wukro dams was consistent with the range 20 to 1500 µS/cm reported by Boyd and Tucker (1998) and Ali et al. (2000) for natural waters but higher than the value 201.56 µS/cm reported by Tadesse et al. (2015). The EC values observed decreased in the studied months. Roshinebegam and Selvakumar (2014) reported the higher EC value in some water bodies may be due to low flow of water, reduced volume and stagnation of water.

TDS

Mean TDS value (169 and 194 mg/L) was obtained in Korrir and Laelay Wukro dams respectively (Table 1). Tadesse et al. (2015) reported lower average TDS value of 101.28 mg/L from five tropical small dams (Korrir, Laelay Wukro, Mai Nigus, Mai Sessea and Mai Seye dams). A maximum value of 400 mg/L of TDS is permissible for a diverse fish population (Alikunhi, 1957). TDS concentration below 200 mg/L promoted even healthier spawning conditions (Deepak and Singh, 2014). Imoobe and Oboh (2003) and Olele and Ekelemu (2008) stated the higher TDS value in water bodies during dry season may be due to the evaporation of water and leaving a higher concentration of salt within a smaller volume of water.

Transparency

The mean transparency values were 28.17 and 29.55 cm in Korrir and Laelay Wukro dams

Table 2. Estimation of fish yields of the dams using different empirical models in the studied months.

Morphometric characteristics	Korrir dam	Laelay Wukro dam
Area (km ²)	0.14	0.01
Mean depth (m)	8.00	6.00
Conductivity (µm host/cm)	2.84	3.58
MEI	0.355	0.597
Models	Estimated productivity (Quintal/ha/year)	
Henderson and Welcome (1974)	8.81	11.24
Toews and Griffith (1979)	12.41	13.24
Marshal (1984)	20.76	18.92
Mean	13.99±6.13 ^a	14.47±3.98 ^a

Mean values in column with the same letter are not significantly different (p<0.05).

respectively which were lower than the value 39.75 cm reported by Tadesse et al. (2015) in five tropical small dams. Boyd (1982) reported the range (15-45 cm) for fish growth. Mustapha (2008) reported the range of Secchi disc visibility to be between 62 and to 162 cm which are higher compared with the current study results. These values reflect the depth of light penetration which is good for a shallow reservoir as plankton growth thus making food available to fish. Monthly transparency record value was increased from October to March in the dams. Mustapha (2008) and Sachinkumar Patil et al. (2013) reported that the increase transparency during dry season in water bodies may be due to gradual settlement of suspended particles at the bottom of the reservoir.

Table 2 show potential yield estimated of the dams. Results showed the lowest production yield (8.81 and 11.24 kg/ha/year) was obtained by Henderson and Welcome (1974) while highest production values (20.76 and 18.92 kg/ha/year) were according to Marshal (1984) models for Korrir and Laelay Wukro, respectively. The average of three empirical yield models were 13.99 and 14.47 quintal/ha/year which were higher than fish yield value (4.35 and 4.53 quintal/ha/year) according to Schlesinger and Regier (1982) model reported by Tadesse et al. (2015) in the Korrir and Laelay Wukro dams respectively.

Conclusion

The water physico-chemical properties of the dams were good for fish production and were within the range already documented though physico-chemical properties showed variations across months. Therefore fish farmers should practice utilization of the dams through good management of the water bodies.

CONFLICT OF INTERESTS

The authors have not declared any conflict of interests.

ACKNOWLEDGMENT

The authors greatly thank Tigray Agricultural Research Institute (TARI) for granting financial support and facilities to this study.

REFERENCES

- Ali M, Salam A, Azeem A, Shafique M, Khan BA (2000). Studies on the effect of seasonal variations on physical and chemical characteristics of mixed water from Rivers Ravi and Chenab at union site in Pakistan. *J. Res. BZ Univ. Multan.* 2:1-7.
- Alikunhi KH (1957). Fish culture in India. In *Bull. Ind. Council Agric. Res.* (20):144.
- Araoy PA (2009). The seasonal variation of pH and dissolved oxygen (DO₂) concentration in Asa lake Ilorin, Nigeria. *Int. J. Phy. Sci.* 4(5):271-274.
- Atakilt B, Tadesse D (2012). Population dynamics and condition factor of *Oreochromis niloticus* L. in two tropical small dams, Tigray (northern Ethiopia). *J. Agric. Sci. Technol.* 2:1062-1072.
- Bisht AS, Ali G, Rawat DS, Pandey NN (2013). Physico-chemical behaviors of three different water bodies of sub tropical Himalayan Region of India. *J. Ecol. Nat. Environ.* 5(12):387-395.
- Boyd CE (1990). Water quality in ponds for Aquaculture. Auburn University, Agric. Exptal. Sta., Auburn, Alabama. P 252.
- Boyd CE (1982). Water quality management of pond fish culture. Elsevier Sci. Pub. Co. Amsterdam-Oxford, New York, 318p.
- Boyd CE, Tucker CS (1998). Pond aquaculture and water quality management. Kluwer Academic Pub., London. pp. 44-48.
- Boyd CE (1979). Water quality in warm water fish ponds. Auburn Univ., Agriculture Experiment Station, Alabama. 359p.
- Brian O (2006). Dissolved Oxygen. *Environ. Qual.* 84:18766.
- Dastagir G, Narejo NT, Jalbani S (2014). Physico-chemical parameters and their variations in relation to fish production in Zhob River Balochistan Pakistan. *J. Anal. Environ. Chem.* 15:2.
- Deepak S, Singh NU (2014). The relationship between physico-chemical characteristics and fish production of Mod Sagar reservoir of Jhabua District, MP, India. *Res. J. Recent Sci.* 3:82-86.
- Environmental Review (ER) (2008). Coastal Zone Aquaculture Management. *Aquac. Strategic Plan* 5:1-14.
- FAO (2003). Fisheries statistics, <http://www.fao.org>. Accessed 13th Jan. 2006.
- Imoobe TOT, Obboh IP (2003). Physical and chemical hydrology of river Jamieson, Niger Delta, Nigeria. *Benin Sci. Digest* 1:105-119.
- Iqbal F, Ali M, Abdus Salam, Khan BA, Ahmad S, Qamar M, Kashif Umer (2004). Seasonal variations of physico-chemical characteristics of river soan water at Dhoak Pathan Bridge (Chakwal), Pakistan. *Intl. J. Agric. Biol.* 1:89-92.

- Ita EO, Balogun JK, Adimula AB (1995). A preliminary report of preimpoundment of fishes Survey of Goronyo Reservoir Sokoto. NIFFR Ann. Rep. pp. 35-38.
- King BDL (1970). The role of carbon in Eutrophication. J. Water Pollut. Control Federation 42:2035-2051.
- Mustapha MK (2008). Assessment of the water quality of Oyon reservoir, Offa, Nigeria, using selected physico-chemical parameters. Turk. J. Fish. Aquat. Sci. 8:309-319.
- Olele NF, Ekelemu JK (2008). Physico-chemical and periphyton/phytoplankton study of Onah lake, Asaba, Nigeria. Afr. J. Genet. Agric. 4(3):183-193.
- Ovie SI, Bwala RL, Ajayi O (2011). A preliminary study on limnological stock assessment, productivity and potential fish yield of Omi Dam, Nigeria. Afr. J. Environ. Sci. Technol. 5(11):956-963.
- Piper RG, McElwim IB, Prme LE, McCranen JP, Flower LG, Leonard JR (1982). Fish hatchery manage. J. Wildlife Manage. 2:6-10.
- Robert CS (2007). Water quality consideration for aquaculture. J. Anim. Ecol. 2:1-15.
- Roshinebegam K, Selvakumar S (2014). Seasonal changes in physico-chemical parameters of Mullai Periyar River, Tamil Nadu, India. Chem. Sci. Rev. Lett. 3(9):66-73.
- Sachinkumar Patil, Patil SS, Sathe TV (2013). Limnological status of Khanapur freshwater reservoir from Ajara Tahsil, Kolhapur district (ms), India. Int. J. Sci. Environ. Technol. 2:1163-1174.
- Saloom ME, Duncan RS (2005). Low dissolved oxygen levels reduce anti-predator behaviors of the freshwater Clam *Corbicula fluminea*. Fresh Water Biol. 50:1233-1238.
- Singh NP (2007). Guidelines for water quality management for fish culture in Tripura. Publication No. 29.
- Tadesse D, Tsehaye A, De Meester L, Afework M, Abreha G, Risch S. (2008). Limnological and ecological characteristics of tropical high land reservoirs in Tigray, northern Ethiopia. Hydrobiol. 610:193-200.
- Tadesse D, Atakilt B, Solomon K, Tsehaye A, Solomon A, Solomon T (2015). Yield and condition factor of *Oreochromis niloticus* L. in tropical small dams, Tigray, northern Ethiopia. World J. Fish Marine Sci. 7(1): 29-37.
- United States Department of Agriculture (USDA) (1996). Aquaculture outlook. The Rev. 4:26-28.
- Zenebe T, Abeba WG, Mulugeta J, Fekadu T Fasil D (2012). Effect of supplementary feeding of agro-industrial byproducts on the growth performance of Nile tilapia (*Oreochromis niloticus* L.) in concrete ponds Ethiopian. J. Biol. Sci. 11(1):29-41.



International Journal of Fisheries and Aquaculture

Related Journals Published by Academic Journals

- *Journal of Plant Breeding and Crop Science*
- *African Journal of Agricultural Research*
- *Journal of Horticulture and Forestry*
- *International Journal of Livestock Production*
- *International Journal of Fisheries and Aquaculture*
- *Journal of Cereals and Oilseeds*
- *Journal of Soil Science and Environmental Management*
- *Journal of Stored Products and Postharvest Research*

academicJournals