

Journal of Environmental Chemistry and Ecotoxicology

Volume 9 Number 5 November 2017

ISSN 2141-226X



*Academic
Journals*

ABOUT JECE

The **Journal of Environmental Chemistry and Ecotoxicology (JECE)** is published bi-monthly (one volume per year) by Academic Journals.

Journal of Environmental Chemistry and Ecotoxicology (JECE) is an open access journal that provides rapid publication (bi-monthly) of articles in all areas of the subject such as ocean acidification, pesticides use and regulation, arsenic induced genotoxicity by curcumin, heavy metals and its environmental effect etc. The Journal welcomes the submission of manuscripts that meet the general criteria of significance and scientific excellence. Papers will be published shortly after acceptance. All articles published in JECE are peer-reviewed.

Contact Us

Editorial Office:

jece@academicjournals.org

Help Desk:

helpdesk@academicjournals.org

Website:

<http://www.academicjournals.org/journal/JECE>

Submit manuscript online

<http://ms.academicjournals.me/>

Editors

Prof. Peter Massanyi

*Slovak University of Agriculture, Faculty of
Biotechnology and Food Sciences, Department of
Animal Physiology,
Management,
Tr. A. Hlinku 2, SK-949 76 Nitra, Slovak Republic,
Slovak Republic.*

Prof. Mostafa El-Sheekh

*Faculty of Science, Tanta University,
Tanta 31527,
Egypt.*

China

Prof. Minghua Zhou

*Nankai University,
No. 94, Road Weijin,*

*Nankai District,
Tianjin 300071,
China.*

Prof. Muhammad Abdul Rauf

*United Arab Emirates University,
United Arab Emirates.*

Prof. Shao Hongbo

*Qingdao University of Science Technology,
Zhengzhou Road 53, Qingdao266042, China ,
China.*

Prof. Ghasem D. Najafpour

*Oshirvani University of Technology
Babol, Iran
Iran.*

Prof. Toyin Ayodele Arowolo

*Department of Environmental Management &
Toxicology,
College of Environmental Resources*

*University of Agriculture,
P.M.B. 2240,
Abeokuta 110001,
Ogun State,
Nigeria.*

Dr. Xue Song Wang

*Department of Chemical Engineering,
Huaihai Institute of Technology,
CangWu Road 59#, Lianyungang, Jiangsu, PR.*

Dr. Mohamed Nageeb Rashed

*Aswan Faculty of Science, South Valley University,

Aswan,
Egypt.*

Prof. Hamayun Khan

*Department of Chemistry
Islamia College University
Peshawar-25120,
Pakistan.*

Editorial Board

Dr. Mohammad Al-Hwaiti

*Al-Hussein Bin Talal University
Environmental Engineering Department
O. Box (20) Ma'an-Jordan
Jordan.*

Prof. Ajai Kumar Srivastav

*DDU Gorakhpur University
Department of Zoology, DDU Gorakhpur University,
Gorakhpur,
INDIA.*

Nathaniel C. Añasco

*University of the Philippines Visayas
Miagao, Iloilo 5023 Philippines
Philippines.*

Prof. El-Refaie Kenawy

*King Saud University,
Faculty of Science,
Department of Chemistry,
Petrochemicals Research Chair,
B.O.Box 2455 Riyadh 11451 ,Saudi Arabia
Saudi Arabia.*

Dr. K. Senthil Kumar

*TÜV SÜD South Asia Pvt. Ltd.,
No: A-151, 2nd C Main, 2nd Stage, Peenya Industrial
Estate,
Bangalore 560058, Karnataka State, INDIA
India.*

Dr. Omotayo Sarafadeen Amuda

*Ladoke Akintola University of Technology
Ilorin Road, Ogbomoso,
Nigeria.*

Dr. Jitendra Pandey

*Banaras Hindu university
Environmental Science Division, Department of Botany,
Banaras Hindu university, Varanasi - 221005,
India.*

Dr. Soumya Chatterjey

*Defence Research Laboratory, Tezpur (DRDO)
Post Bag No. 2, Tezpur 784001, Assam,
India.*

Prof. Bechan Sharma

*University of Allahabad
Department of Biochemistry,
Allahabad-211002,
India.*

Dr. Ghousia Begum

*Indian Institute of Chemical Technology,
Hyderabad 500 007, A.P; India
Prof. Gang Yang
Key Laboratory of Forest Plant Ecology, Ministry of
Education,
Northeast Forestry University
26 Hexing Road, Harbin
China.*

Dr. Anindita Bhattacharyya

*Indian Statistical Institute
203, B. T. Road, Kolkata 700108, West Bengal,
India.*

Dr. Onome Davies

*Rivers State University of Science & Technology, Port
Harcourt, Rivers State
Dept. of Fisheries & Aquatic Environment, P.M.B. 5080,
Port Harcourt,
Nigeria.*

Dr. B. Stephen Inbaraj

*Fu Jen University
Department of Nutrition & food science,
Taipei 242, Taiwan
Environmental Chemistry & Analytical Chemistry
Taiwan.*

Dr. Hala A. Awney

*Institute of Graduate Studies and Research,
Alexandria University,
Egypt.*

Dr. Suhel Parvez

*Leibniz Institute for Neurobiology
Brennekestr. 6, Magdeburg 39118,
Germany.*

Dr. Mayalagu Rajkumar

*Institute of Oceanology, Chinese Academy of Sciences
7 Nanhai Road, Qingdao, 266071
China.*

Dr. Eldon Raj Rene

*University of La Coruna
Department of Chemical Engineering, Spain*

Vyacheslav Khavrus

*L. V. Pizarzhevskii Institute of physical chemistry of
NAS of the Ukraine (permanent position)
Ukraine.*

ARTICLE

- Seasonal variation of elemental, physicochemical and microbiological characteristics of the stream water around hospital waste dumpsite in Ilesa, South-western Nigeria 54**
O. T. Fakoya¹, E. A. Oluyemi, I. O. Olabanji and C. N. Fakorede

Full Length Research Paper

Seasonal variation of elemental, physicochemical and microbiological characteristics of the stream water around hospital waste dumpsite in Ilesa, South-western Nigeria

O. T. Fakoya¹, E. A. Oluyemi^{1*}, I. O. Olabanji¹ and C. N. Fakorede²¹Department of Chemistry, Obafemi Awolowo University, Ile-Ife, Osun State, Nigeria.²Department of Microbiology, Obafemi Awolowo University, Ile-Ife, Osun State, Nigeria.

Received 8 March, 2017; Accepted 18 August, 2017

This study assessed the seasonal variation of microbiological, physicochemical and heavy metal analysis of water samples from Ayao Stream around a hospital waste dumpsite. Four samples were collected downstream and analyzed for microbial count, coliforms, pH, electrical conductivity, BOD₅, alkalinity, total dissolved solids, water hardness, sulphate, nitrate, chloride and phosphate using standard method. Heavy metals (Fe, Mn, Cu, Zn, Cd and Pb) in the water were determined by Atomic Absorption Spectrophotometry. Results obtained showed that the stream was greatly polluted with microbial and coliform count of 2.1×10^4 to 3.6×10^5 CFU/ml, 1.8×10^2 to 8.0×10^2 CFU/ml, and 1.1×10^3 to $> 1.1 \times 10^3$ CFU/100 ml for bacteria, fungi and coliform count, respectively during both wet and dry season. The bacteria recovered from the samples included the antibiotic resistant *Pseudomonas* and *Klebsiella*. The mean values of pH (8.0 to 9.1) and BOD₅ (17.2 to 26.3) were obtained during dry and wet season, respectively and were found to be beyond the acceptable limit in drinking water. This result established a relationship between the BOD₅ and microbial load which indicate organic pollution through runoff input from the dumpsite into the stream. The concentration of Mn, Fe, Cu, Pb and Cd especially during rainy season was found to exceed the stipulated limit in drinking water. These metals could be bio accumulated to toxic level in aquatic organisms and end up in human through the food chain. This study shows that Ayao Stream is greatly polluted and unfit for domestic uses.

Key words: Dumpsite, heavy metal, microbial load, coliforms, hospital waste.

INTRODUCTION

Water is a universal solvent and one required by all living organisms for survival. In most communities in Nigeria, treated pipe-borne water is lacking, so groundwater and surface water from streams and rivers are the major

sources of domestic water. Water is contaminated when its acceptable quality has been altered through anthropogenic input by humans. As a result, the aesthetic value decreases having undesirable colour, taste, odour,

*Corresponding author. E-mail: tope4best2001@yahoo.com. Tel: +234-803-404-3291.

Author(s) agree that this article remain permanently open access under the terms of the [Creative Commons Attribution License 4.0 International License](https://creativecommons.org/licenses/by/4.0/)

turbidity, conductivity, organic matter content, harmful chemical contents (heavy metals, persistent organic pollutants) and infectious microorganisms (Abiona et al., 2012; Ogunfowokan et al., 2005). Major sources of surface water pollution are direct discharges from industrial effluents or sewage effluents or sewage treatment plant, run-off from hospital or municipal waste dumpsite, agricultural lands or other polluted land areas.

Hospital wastes are wastes generated within the hospital environment which include sharps, chemicals, pharmaceuticals, radioactive, laboratory, general and kitchen waste. This waste is majorly composed of liquid and solid materials which have effects on the element of the environment, altering its natural composition (Akter, 2000; WHO, 2005). The treatment and disposal method appropriate for hospital waste are incineration and sanitary landfills which is monitored against leachate seeping into the soil (Diaz et al., 2005). However, in Nigeria, more than 90% of the hospital waste generated is directly disposed on land because it is economical and there is lack of proper waste disposal management technique. There is also deficiency in the designation and identification of infectious waste, segregation, packaging, and final disposal of the waste (Mayet, 1993; Vivan et al., 2011). Hospital waste contains a great variety of chemicals, heavy metals, disinfectants and specific detergents resulting from diagnosis and research activities. Disposal of these wastes on open land area alters the natural composition of soil through the release of heavy metals, pesticides, and salts. Also, run-off from the surrounding land area causes extensive damage to the water quality characteristics, especially when microbial degradation of the anthropogenic input is unable to remove the pollutants fast enough to prevent environmental degradation. Release of heavy metals to the environment is of particular concern due to their persistence and toxicity to humans (Odewande and Abimbola, 2008; Khairiah et al., 2004). Manganese, nickel, selenium, zinc, copper, chromium, iron, lead and cadmium are typical examples of heavy metals. Among others, lead and cadmium are known to be toxic and carcinogenic at elevated levels. Uptake of heavy metals by plants or animals results in bioaccumulation which moves through the food chain and eventually ends up in man (Laurent et al., 2011). Hence, the adverse effect of surface water pollution is the creation of a harsh environment which adversely affects the activities of aquatic microflora and aquatic lives. Improper management of hospital waste may also contribute to oxygen demand and nutrient loading of the water and ultimately result in biological and environmental loss (Sharholly et al., 2008). The inter-relationship between the real public health effects and the range of hospital waste management processes regulated is an important cause of concern to the public at large today. There have been an increasing number of epidemiological research reports linking a range of ill-health outcomes to hospital waste

management installations. These studies include childhood cancers, birth abnormalities and liver cancers associated with incinerators and landfills facilities.

In this study, information has been provided on the physicochemical properties, heavy metal concentration and microbial activities of Ayao Stream located within the vicinity of the hospital with a view to establishing the pollution status of the stream as a result of anthropogenic input.

MATERIALS AND METHODS

Study area

The water samples were collected along the course of Ayao Stream situated at about 110 m from a hospital waste dumpsite (07° 36' 801" N and 004° 43' 984" E) with an elevation of 387 m in Ilesa East Local Government Area of Osun State (Figure 1).

Collection of water samples

Samples for microbial and physicochemical analysis were collected from five different locations (W1, W2, W3, W4, and control sample) of the Ayao Stream during rainy season between May and October, 2013 and dry season between November, 2013 and March, 2014. Samples W1, W2, W3 and W4 were taken at about 60 m from other downstream (after the dumpsite) along the Ayao Stream. The control sample was taken at about 100 m upstream before sample location W1. Samples for physicochemical and heavy metal analysis were collected in 1-L pre-cleaned plastic bottles, rinsed with the sample and filled. The samples were taken to the laboratory and refrigerated at about 4°C prior to analysis.

The physicochemical parameters analyzed are as follows: pH, conductivity, alkalinity, chloride, hardness, nitrate, sulphate and phosphate. Standard methods were followed in determining the aforementioned variables as described by APHA (1998).

The samples for microbial analysis were taken at about 10 cm below the water surface into 250-ml plastic bottles already disinfected with 75% ethanol and rinsed again with sterile distilled water. Each of the sampling containers was rinsed with water from sampling point twice before the collection of the water sample. The water samples were brought into the laboratory in plastic coolers with iced flakes and analysed within three hours of collection.

Acid digestion

Digestion of water was done according to standard method (APHA, 1998). 50 ml water sample was measured into an evaporating dish and digested by adding 5 ml concentrated HNO₃ and 1 ml HClO₄. The solution was heated on a steam bath until the volume was reduced to about 15 ml. The solution was allowed to cool, it was filtered and made up to 25-ml mark in a volumetric flask. Analysis for heavy metal concentration was carried out using the Perkin Elmer 400 atomic absorption spectrophotometer by flame atomization, using air-acetylene flame and single element hollow cathode lamp at the Central Science Laboratory of Obafemi Awolowo University, Ile-Ife, Nigeria.

Total microbial counts

Enumerations of total microbial (bacterial and fungal) counts were carried out using the serial dilution and pour plate method. Serial



Figure 1. Location of sampling water sites along Ayao Stream.

10-fold dilutions in sterile distilled water were carried out and 1 ml of each dilution was aseptically dispensed in sterile Petri-dishes in duplicates. A volume of 20 ml of molten nutrient agar (Lab M) cooled to 45°C for bacterial population or Malt Extract Agar (Lab M) for fungi was later added to each of the plates. The mixture was allowed to solidify and incubated inverted for 24 to 48 h at 35°C for bacteria. For fungi, the cultured plates were incubated un-invertedly and aerobically at 30°C for 5 to 7 days (until the plates showed no further increase in the number of fungal colonies).

The plates were observed for growth and selected for count after the expiration of the incubational periods. The cultured plates in which the number of colonies was between 30 and 300 were selected. The count obtained was multiplied by the dilution factor and expressed as colony forming unit (CFU) per ml of the original sample. This is the viable count.

Presumptive coliform count

The water samples were processed using the multiple fermentation tube method to determine the presumptive coliform count or most probable number (MPN) of coliforms based on standard methods (Senior, 2006). Suspensions from positive tubes were sub-cultured onto MacConkey agar and incubated at 37°C for 24 to 48 h, and the resulting colonies identified using standard procedures (Forbes et al., 2007).

Confirmatory test

The tubes showing gas and/or acid production (MPN tubes) were sub-cultured on Eosin Methylene blue (EMB) agar incubated at 35°C for 24 h and examined for typical colonies of *E. coli* (greenish metallic sheen). If typical colonies were seen, then the completed test was carried out.

Completed test (Eijkman test)

Eijkman's test, which is for the detection and enumeration of faecal coliform, involves the use of elevated temperature for the differentiation of faecal and non-faecal coliforms. Measured volume of water samples were inoculated into lactose broth in similar combinations as in the presumptive test for total coliforms and incubated at 44°C for 48 h. The number of positive tubes was recorded at the end of the incubation period and the MPN of faecal coliform was read from the MPN index table.

Isolation of pure cultures of microorganism

The distinct colonies from the mixed culture on each of the nutrient agar and the malt extract agar plates were picked and streaked onto fresh plates. Pure colonies of each bacterium and fungus were then picked for identification using standard biochemical tests which included Gram's stain, catalase, MRVP, citrate utilization, indole, oxidase, nitrate reduction, hydrogen sulphide production and sugar fermentation tests for bacteria.

RESULTS AND DISCUSSION

Physicochemical parameters in steam water

The mean value \pm standard deviation results of some physicochemical parameters analyzed in the stream water for the period of sampling are shown in Table 1.

The level of pH varied between 7.9 to 8.1 and 8.8 to 9.7 during dry and wet seasons, respectively. The mean pH values during wet season exceeded the Standard

Table 1. Physicochemical parameters of the stream water.

Location	pH		EC ($\mu\text{s/cm}$)		TDS (mg/L)		BOD (mg/L)		NO ₃ ⁻ (mg/L)	
	Dry	Wet	Dry	Wet	Dry	Wet	Dry	Wet	Dry	Wet
W1	8.0±0.7	9.0±1.1	88.2±6.4	153.3±8.3	95.5±4.6	54.9±9.3	10.1±2.3	15.5±5.1	3.2±0.7	4.7±2.3
W2	7.9±0.6	8.9±0.4	95.3±10.1	135.0±9.1	84.1±6.7	59.5±12.3	11.1±4.5	18.6±2.3	6.1±4.5	5.5±1.3
W3	8.1±0.6	8.8±0.5	88.1±14.3	131.0±10.2	81.8±10.2	55.0±7.9	20.2±1.7	40.1±1.1	5.5±9.2	2.1±0.8
W4	8.0±1.2	9.7±2.3	57.1±7.5	132.1±11.1	78.4±8.2	35.8±7.8	27.5±1.4	30.9±3.2	2.8±1.8	1.3±0.2
Ctrl Site	8.0±1.1	9.5±0.5	97.5±4.2	142.5±6.7	82.2±14.2	60.8±11.7	2.4±0.8	8.3±0.8	1.9±3.3	1.1±1.0
Mean	8.0	9.1	82.2	137.9	85.0	51.3	17.2	26.3	3.9	3.4
St. dev	0.08	0.41	17.0	10.43	7.41	10.6	8.22	11.3	1.64	2.02
WHO	6.5 - 8.5		400 $\mu\text{s/cm}$		1000 mg/L		3 - 6 mg/L		50.0 mg/L	
SON	6.5 - 8.5		1000 $\mu\text{s/cm}$		1000 mg/L		6.0 mg/L		10.0 mg/L	
Location	SO ₄ ²⁻ (mg/L)		PO ₄ ⁻ (mg/L)		Cl ⁻ (mg/L)		Water hardness (mg/L)		Alkalinity (mg/L)	
	Dry	Wet	Dry	Wet	Dry	Wet	Dry	Wet	Dry	Wet
W1	ND	ND	1.9±0.4	1.1±0.7	7.9±2.3	5.8±3.5	50.6±11.3	45.0±0.8	65±0.7	65±7.2
W2	ND	ND	2.6±1.1	1.3±0.1	7.7±0.6	6.2±1.9	48.4±15.8	41.3±2.1	60±1.4	65±2.3
W3	ND	ND	1.8±0.3	0.9±0.1	7.0±1.2	5.5±2.2	45.0±3.6	42.5±0.5	55±1.9	60±8.5
W4	ND	ND	2.3±1.1	1.2±1.1	5.7±0.8	6.0±1.6	48.4±1.4	43.3±3.4	50±4.4	65±2.1
Ctrl Site	ND	ND	2.1±1.7	0.7±2.3	4.5±2.3	6.5±7.8	49.5±8.9	43.3±2.6	45±2.8	75±7.6
Mean	-	-	2.15	1.13	7.1	5.90	48.1	43.0	57.5	63.8
St. dev	-	-	0.36	0.17	0.99	0.3	2.31	1.55	6.45	2.5
WHO	-		-		250 mg/L		500 mg/L		-	
SON	100.0 mg/L		5.0 mg/L		100 mg/L		150 mg/L		-	

N.D = Not detectable.

Organisation of Nigeria (SON, 2008) limit between 6.5 and 8.5 for drinking water. The high pH values suggest the presence of dissolved gases such as NH₃ and some alkali chemicals in the water resulting from runoff from the hospital dumpsite. Nevertheless, the high pH values in the study area suggest that the water may not influence the solubility of toxic metals in the water bodies. The electrical conductivity (EC) varied between 57.1 to 97.5 $\mu\text{s/cm}$ during dry season while higher EC values were obtained in the wet season (131 to 153.3 $\mu\text{s/cm}$). Higher values during wet season may be due to increased concentration of ionized substances in the water resulting from run-off of nutrients or chemicals from surrounding land area of the hospital waste dumpsite into the stream water. The mean EC values in both seasons were within the SON (2008) limit of 1000 $\mu\text{s/cm}$ in drinking water.

Total dissolved solids (TDS) varied between 78.4 to 95.5 mg/L and 35.8 to 60.8 mg/L during dry and wet seasons, respectively. Higher TDS value during dry season could be as a result of reduction in the volume of water, which results in high concentration of the mineral content. The results of TDS obtained in this water shows that the values are within the acceptable limit of 1000 mg/L (WHO, 2006) for domestic use regardless of other parameters. TDS level higher than 1000 mg/L in drinking

water increases turbidity of water, affecting the aesthetic value of water and making it unpalatable to taste.

BOD₅ is a measure of oxygen demand by microorganism while breaking down organic matter. The BOD₅ values range from 10.1 to 27.5 mg/L during dry season and 15.5 to 40.1 mg/L during wet season compared to 3.0 mg/L and 3.0 to 6.0 mg/L set by WHO (2006) and SON (2008), respectively. Higher BOD₅ values obtained during the wet season may result from leaching and runoff from land surrounding the dumpsite which increase the level of organic matter and activities of microorganism in the stream. As a result, the dissolved oxygen will decrease over time, promoting toxic algae blooms and leading to a destabilized aquatic ecosystem. Hence, BOD₅ result shows that the water is polluted and unfit for use for domestic purposes.

Chloride is a major ion in surface water. The mean concentrations of Cl⁻ in the stream water were minimal, well below the permissible limit of 100 mg/L (WHO, 2006) in water. Chloride occurred in the dry season in the range of 5.7 to 7.9 mg/L and 5.5 to 6.1 mg/L during wet season. The mean value of water hardness during dry season varied moderately between 45.0 to 50.6 mg/L and 41.3 to 45.0 mg/L during wet seasons. However, the values of water hardness fall into the category of soft water according to Radojevic and Bashkin (1999), which

suggest low solubility of toxic multivalent metals in water. The alkalinity values showed no observable variation in both seasons as it occurred in the range of 50 to 65 mg/L during dry season and 60 to 65 mg/L during wet season, although it is still within the limit of 100 mg/L allowed for potable water. Although, alkalinity has little public health significance (Radojevic and Bashkin, 1999), highly alkaline water is unpalatable and not good for domestic water supply.

The concentration of nitrate and phosphate varied between 2.8 to 6.1 mg/L and 1.8 to 2.6 mg/L, respectively during dry season and 1.3 to 5.5 mg/L and 0.9 to 1.3 mg/L during wet season. The levels of nitrate and phosphate in the samples did not exceed the SON (2008) safe limit of 10 and 5 mg/L, respectively. Low phosphate values were evidenced by the low chlorophyll in the stream water. Sulphate was below the detection limit of the instrument in all the samples.

Heavy metal concentration in stream water

The result obtained from the metal analysis of stream water samples in the study area using Atomic Absorption Spectroscopy are presented in Figure 2. Statistical T-test analysis for metal concentration during wet and dry seasons indicates that significant difference ($P > 0.05$) exist in Mn, Zn and Cd concentration in the stream water. The general trend observed for the metals are Fe>Mn>Zn>Cu>Cd>Pb for dry season; and Fe>Mn>Cu>Zn>Pb>Cd for wet season. The highest concentration value was recorded for Fe among other metals in both seasons. Iron (Fe) in the water samples ranged from 1 to 98 mg/L and 104 to 135 mg/L for dry and wet seasons, respectively, with the highest value occurring at sampling point W1 in both seasons. The mean concentration values for Fe in all samples for both seasons were higher than the WHO (2006) standard value of 0.3 mg/L for Fe in drinking water. High levels of iron found in the control sample during wet season suggest that there may be a direct discharge of pollutant substances into the water up stream. High Fe levels may result from runoff of land soil in complex with organic matter around the hospital waste dumpsite. The higher concentration of Fe obtained during wet season may be due to increased decomposition of organic matter in the hospital waste that is washed off into the stream water during rainfall. Also, iron could be leached from corroding metal which is abundant in the hospital dumpsite and eventually washed off into the stream. According to Canfield et al. (1984), Fe at high levels contributes to water coloration and reduces light penetration which is evident in the study area and may affect the biogeochemical cycling of organic matter, phosphorus and nitrogen. Photo-oxidation of organic matter is enhanced by Fe (Minor et al., 2006). Also, Fe can have direct and indirect toxic effects on aquatic biota.

Precipitation of Fe on fish gills can impair ion regulation and oxygen uptake (Peuranen et al., 1994). In humans, iron toxicity affects the red blood cells chemistry and delay normal and mental development in children (Ekiz, 2005).

Concentrations of Manganese (Mn) obtained in the water samples ranged between 2.79 to 10.48 mg/L and 4.14 to 6.28 mg/L for dry and wet seasons, respectively. These values exceed the WHO (2006) standard value of 0.5 mg/L for Mn in drinking water. Anthropogenic sources of manganese include mining and smelting, engineering, traffic and agriculture. It is also used in the manufacture of steel, glass, dry batteries and chemicals. Manganese can be released to the stream water by leaching and runoff of these materials from the dumpsite. Excessive amount of Mn causes seizure activity, poor bone formation, impair fertility, birth defect in humans and irreversible brain diseases with prominent psychological and neurological disturbances (Agency for Toxic Substances and Disease Registry (ATSDR), 2000; United States Environmental Protection Agency (USEPA), 1984).

Concentrations of Copper in the water range between 0.32 to 3.19 mg/L in the dry season while it was found in the range of 3.68 to 3.74 mg/L in the wet season. Higher concentration values of copper during the wet season exceeded the WHO (2006) stipulated safe limit of 2.0 mg/L. The levels of zinc in water were within the WHO limit of 5 mg/L in drinking water.

Lead, a toxic heavy metal was below detection limit in the samples collected during the dry season. However, during wet season it appeared at high levels above the WHO (2006) stipulated limit of 0.05 mg/L in the water, as it ranged from 0.58 to 0.85 mg/L. High levels of exposure to Pb may result in toxic biochemical effects in humans which in turn cause problems in the synthesis of haemoglobin, affects the kidneys, gastrointestinal tract, joints and reproductive system, and could cause acute or chronic damage to the nervous system (Goyer, 1986).

Cadmium in water occurred at high levels ranging from 0.08 to 0.23 mg/L during dry season, but was below the detection limit in the samples during wet season. The values for dry season were above the safe limit of 0.003 mg/L Cd stipulated by WHO (2006). The absence of Cd during the wet season might be due to the effect of rainfall which increased the volume of water and water flow rate, eventually lowering the concentration of Cd. Excessive amount of Cd is known to cause damage to body cells by increasing their permeability and enhancing the accumulation of other heavy metals. It is also known to cause kidney damage and bone pains in man (Jarup, 2003). This result shows a higher concentration of the metals in the stream during wet season which indicate there was more discharge through surface runoff during rain which is enhanced by the sloppy topography of the land area. With exception of zinc, higher levels of Fe, Mn, Cu, Cd and Pb in the stream water suggest




Figure 2. Mean seasonal distribution of selected heavy metal concentrations in the stream water.

anthropogenic input through run-off from the hospital waste dumpsite (Figures 3 and 4). Also, anoxic condition in the water with respect to the BOD₅ level may enhance the solubility of some of these metals which increase the leaching of metallic objects in the hospital dumpsite.

Other anthropogenic sources that may be responsible for the presence of high values of these metals in the study samples include atmospheric deposition and the use of agrochemicals in nearby farmlands as shown in Figure 2. This result indicates that the aquatic biota may



Figure 3. Hospital waste dumpsite.



(a) the dumpsite



(b). A representative of the sampling points W2 - W5. The locations are covered by vegetation in Figure 1

Figure 4. Hospital Dumpsite and Stream water.

bioaccumulate these metals to toxic levels and enters the food chain to human. Hence, the assessment of metal concentration in Ayao Stream shows that the water is polluted and therefore unfit for domestic and recreational purposes.

Dominant genera of microorganisms isolated from sampling points of Ayao Stream

A total of 23 strains of bacteria representing 9 genera and 7 species of fungi were isolated from the water samples

Table 2. Biochemical characterization of the bacterial isolates obtained from the water samples.

Isolate	Cell shape	Gram reaction	Oxidase	Catalase	TSI reaction	Sim reaction	Citrate utilization	Mr	Vp	Sugar fermentation					O-f or h & l	Nitrate reduction	Possible organism (s)
										Glucose	Maltose	Manitol	Sucrose	Lactose			
A	SR	-	-	+	YG YMC	---	+	+	-	YG	YG	YG	YG	YG	F	+	<i>Escherichia</i> sp.
B	MLR	-	-	+	YGNC NC	---	+	+	+	YG	YG	YG	YG	YG	F	+	<i>Klebsiella</i> sp.
C	MLR	-	-	++	Y NC NC	---	+	+	-	Y	NC	NC	NC	NC	F	+	<i>Klebsiella</i> sp.
D	MLR	-	-	-	Y NC N	+++	+	+	+	Y	Y	Y	Y	Y	F	+	<i>Klebsiella</i> sp.
E	SR	-	+	++	NC NCNC	-+-	+	-	-	NC	NC	NC	NC	NC	OX	+	<i>P. aeruginosa</i>
F	MLR	-	-	++	YG NCNC	---	-	+	+	YG	YG	YG	YG	YG	F	+	<i>Klebsiellasp.</i>
G	SR	-	-	+	YG YNC	+++	-	+	-	YG	YG	YG	YG	YG	F	+	<i>Escherichia</i> sp.
H	MLR	-	-	+	Y NC NC	---	-	+	-	Y	NC	NC	NC	NC	F	+	<i>Klebsiellasp.</i>
I	SR	-	+	+	NC NCNC	-+-	+	-	-	NC	NC	NC	NC	NC	OX	+	<i>P. aeruginosa</i>
J	SR	-	-	+	YG YNC	-+-	+	-	+	YG	YG	YG	YG	YG	F	+	<i>Enterobacter</i> sp.
K	MLR	-	-	+++	Y NC NC	---	+	+	-	Y	NC	NC	Y	NC	F	+	<i>Klebsiella</i> sp.
L	Cocci	+	-	+++	NC Y NC	---	-	+	+	Y	Y	Y	Y	Y	OX	-	<i>Aerococcus</i> sp.
M	SR	-	-	++	YG Y NC	++-	-	+	-	YG	YG	YG	YG	YG	F	+	<i>Escherichia</i> sp.
N	MLR	-	-	+++	YG NCNC	---	-	-	+	YG	YG	YG	YG	YG	F	+	<i>Klebsiella</i> sp.
O	MLR	-	-	+++	YG NCNC	---	-	+	+	YG	YG	YG	YG	YG	F	+	<i>Klebsiella</i> sp.
P	LR	-	-	-	Y NC NC	---	-	-	-	Y	NC	NC	NC	NC	None	-	<i>Acinetobacter</i> sp.
Q	SR	-	-	+	YGYNC	++-	-	+	-	YG	YG	YG	YG	YG	F	+	<i>Escherichia</i> sp.
R	MLR	-	-	++	Y NC NC	---	+	+	-	Y	NC	NC	NC	NC	F	+	<i>Klebsiella</i> sp.
S	SR	-	-	+	NC NCNC	---	-	-	-	NC	NC	NC	NC	NC	OX	+	<i>P. aeruginosa</i>
T	SR	-	-	+	Y NCNC	++-	-	+	-	Y	Y	Y	Y	Y	F	+	<i>Shigella</i> sp.
U	MLR	-	-	-	Y NCNC	++-	-	+	+	Y	NC	NC	NC	NC	F	+	<i>Proteus</i> sp.
V	SR	-	+	++	NC NCNC	-+-	+	-	-	NC	NC	NC	NC	NC	OX	+	<i>P. aeruginosa</i>
W	Filamentous	+	-	++	NCNCNC	---	-	-	-	NC	NC	NC	NC	NC	None	+	<i>Streptomyces</i> sp.

of Ayao Stream as shown in Tables 2 and 3. The dominant bacteria genera in the upstream samples were *Pseudomonas*, *Enterobacter*, *Aerococcus* and *Acinetobacter* while *Klebsiella*, *Escherichia*, *Shigella* and *Proteus* were the major genera isolated in downstream water samples.

Six fungal species which included *Rhizopus japonicus*, *Pullularia pollulans*, *Trichoderma* sp., *Absidia* sp., *Aspergillus glaucus* and *Aspergillus fumigatus* were isolated from both the upstream and downstream water samples while additional *Mucor mucedo* was isolated from the control site

sample.

Microbiological parameters of stream water

The microbiological parameters of the Ayao Stream (upstream and downstream) sampled and analysed with bacterial, coliform bacterial, faecal coliform and fungal counts are presented in Table 4. The downstream samples (W3 and W4) contained higher bacterial (4.7×10^4 - 3.6×10^5 CFU/ml) and fungal (4.0×10^2 - 8.0×10^2 CFU/ml)

counts in both dry and wet seasons. A higher bacterial and fungal load was observed during the wet season in all the samples, which is most likely due to the surface runoff of decaying organic matter which suitably introduced more organisms and increased microbial activity in the stream. Similar findings were reported by Adewoye (2010) and Fakorede et al. (2013). However, the control samples (upstream) contained lesser loads of bacteria count (2.3×10^4 and 2.1×10^4 CFU/ml) and fungal count (3.6×10^2 and 1.0×10^2 CFU/ml) when compared with the downstream samples for

Table 3. Microorganisms obtained from sampling points of Ayao Stream.

Microorganism	W1	W2	W3	W4	Control point
Bacteria					
<i>Escherichia</i> sp.	-	-	+	+	-
<i>Klebsiella</i> sp.	-	-	+	+	-
<i>P. aeruginosa</i>	+	+	+	+	+
<i>Enterobacter</i> sp.	+	+	+	-	+
<i>Aerococcus</i> sp.	+	+	+	-	+
<i>Acinetobacter</i> sp.	+	+	-	+	-
<i>Shigella</i> sp.	-	-	+	+	-
<i>Proteus</i> sp.	-	-	+	+	-
<i>Streptomyces</i> sp.	-	-	-	-	+
Fungi					
<i>Rhizopus japonicus</i>	+	+	-	-	+
<i>Pullularia pollulans</i>	+	+	+	+	-
<i>Trichoderma</i> sp.	+	+	+	+	+
<i>Absidia</i> sp.	+	+	-	-	-
<i>Aspergillusglaucus</i>	-	+	+	+	+
<i>A. fumigates</i>	-	+	+	+	+
<i>Mucormucedo</i>	-	-	-	-	+

Table 4. Microbiological parameters of the upstream and downstream water sampled from Ayao Stream.

Location	Total bacterial count (CFU/ml) dry season		Coliform count (CFU/100 ml)		Faecal coliform count (CFU/100 ml)		Total fungal count (CFU/ml)	
	Wet	Dry	Wet	Dry	Wet	Dry	Wet	Dry
W1	5.0×10^4	2.1×10^4	$>1.11 \times 10^4$	1.1×10^3	0	0	4.6×10^2	2.1×10^2
W2	5.2×10^4	2.6×10^4	$>1.1 \times 10^3$	1.1×10^3	0	0	5.2×10^2	1.8×10^2
W3	3.6×10^5	5.0×10^4	$>1.1 \times 10^3$	$>1.1 \times 10^3$	0	0	8.0×10^2	4.2×10^2
W4	3.3×10^5	4.7×10^4	$>1.1 \times 10^3$	$>1.1 \times 10^3$	0	0	7.6×10^2	4.0×10^2
Control site (Upstream)	2.3×10^4	2.1×10^4	1.1×10^3	1.1×10^3	0	0	3.6×10^2	1.0×10^2

wet and dry seasons respectively. The low microbial counts in the control samples suggest that the runoff from the dumpsite is a point source of pollution. Also, the microbial load was observed to increase with increasing distance from the dumpsite (downstream). This is in agreement with the BOD₅ value which could result from lower water volume and flow rate downstream. Hence, the microbial load is more concentrated.

The coliform bacterial counts ranged from 1.1×10^3 to $>1.1 \times 10^3$ CFU/100 ml. The result of the coliform count in the water samples is suggestive of the fact that the dumpsite contributed to the presence of coliform bacteria in the water bodies. Therefore, none of the water bodies is safe for drinking since their bacterial and coliform bacterial densities greatly exceeded the World Health Organization (WHO, 2004) standard of 10 CFU/ml in drinking water.

Faecal coliforms were however, not isolated from all the samples collected indicating that faecal wastes were

not being introduced into water body from the dumpsite. The bacterial strains isolated from the samples belong to the genera *Escherichia*, *Klebsiella*, *Pseudomonas*, *Acinetobacter*, *Streptomyces*, *Proteus*, *Shigella*, *Enterobacter* and *Aerococcus*. Some of these bacteria are implicated in infections of wounds, burns, ulcer and urinary tract. The presence of these microbes is of public health importance because of their potentials as pathogens and the fact that they can be multi resistant to antibiotics which may make the choice of antibiotics for treatment of infections arising from them a herculean task. The fungi isolated from these samples especially *Trichoderma* sp. are implicated in human skin diseases.

Conclusion

From the results obtained in this study, the values of physicochemical parameters in stream water were within

the standard limit stipulated by WHO (2006) and SON (2008). However, significant pollution was indicated by the high values of pH and BOD especially during wet season above the standard limit in water. High BOD value in stream water indicates organic pollution due to run-off input from the hospital dumpsite and agricultural land which increase microbial decomposition of organic matter in water. On the other hand, the total metal concentration of Fe, Mn, Cu, Pb and Cd in stream water were higher than the standard set by WHO (2006). This suggests that there could be danger of bioaccumulation of toxic metals for aquatic life which eventually leads to toxicity in the biota over time.


The microbiological results revealed that the Ayao Stream was greatly polluted by the run-off from the dumpsite into the stream as the samples contained very high microbial loads throughout the sampling period. The coliform count indicate that the water bodies is unsafe for drinking since their bacterial and coliform bacterial densities greatly exceeded the acceptable limit by World Health Organization (WHO) in drinking water. Certain bacteria and fungi strain isolated from the stream water have the potential to become pathogens and can be multi resistant to antibiotics. This study is particularly of public health significance because this stream flowed outside the hospital premises and it is being used by a good number of people including farmers for irrigation, washing, bathing and other domestic purposes. This suggests that hospital waste in the environment is a major health and environmental threat, which call for a proper regulatory system of its disposal.

CONFLICT OF INTERESTS

The authors have not declared any conflict of interests.

REFERENCES

- Abiona OO, Adebajo JA, Anifowose AJ, Abdulla JO, Bamigbelu OR (2012). Impact Assessment of pollution from Metal Concentration in water and Fish. *Nat. Sci.* 8:10.
- Adewoye SO (2010). Effects of detergent effluent discharges on the aspect of water quality of Asa River, Ilorin, Nigeria. *Agric. Biol. J. N. Am.* 1(4):731-736.
- Akter N (2000). Medical waste management: A review. Pathumthani, Thailand. Asian Institute of Technology. Retrieved from: <http://www.engconsult.com/BEN/papers/Paperanasima>.
- APHA (1998). Standard Methods for the Examination of Water and Wastewater, 18th Edition. Washington. 1220:45-60.
- ATSDR (2000). Toxicological profile for manganese (update). Draft for public comment. Atlanta, GA, US Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry.
- Canfield DE, Linda SB, Hodgson LM (1984). Relations between color and some limnological characteristics of Florida lakes. *Water Res. Bull.* 20:323-329.
- Diaz LF, Savage, GM, Eggerth LL (2005). Alternative for the treatment and disposal of healthcare waste in developing countries. *Waste Manage.* 25:626-637.
- Ekiz C (2005). The effect of iron deficiency anaemia on the function of the immune system. *Hematol. J.* 5:579
- Fakorede CN, Igbenughu OA, Odeyemi O (2013). The Effect of a Nigerian Brewery Effluent on Two Receiving Streams. *Nat. Sci.* 11(5):95-102.
- Forbes BA, Sahm BF, Weissfeld AS (2007). *Bailey & Scott's diagnostic microbiology*. 12th ed. St. Louis, MO, USA, Mosby Elsevier.
- Goyer RA (1986). Toxic effects of metals. In: Cassarett and Doull's Toxicology. The basic science of poisons. 3rd ed. McGraw-Hill, New York.
- Jarup L (2003). Hazards of Heavy Metal Contamination. *Br. Med Bull.* 68:167-182.
- Khairiah T, Zalifah MK, Yin YH, Aminath A (2004). The Uptake of Heavy Metals by Fruit type Vegetables, Grown in Selected Agricultural Areas. *Pak. J. Biol. Sci.* 7(2), 1438-1442.
- Laurent M, Pascal RO, Jean PT (2011). Heavy metals in soil on spoil heap of an abandoned lead ore treatment plant, SE Congo-Brazzaville. *Afr. J. Environ. Sci. Tech.* 5(2):89-97.
- Mayet MAG (1993). Domestic Waste Generation in the Urban Core of the Durban Functional Region. Masters Thesis, Department Of Civil Engineering, University Of Kwazulu.
- Minor EC, Pothen J, Dalzell BJ, Abdulla H, Mopper K (2006). Effects of salinity changes on the photodegradation and ultraviolet-visible absorbance of terrestrial dissolved organic matter. *Limnol. Oceanogr.* 51:2181-2186.
- Odewande AA, Abimbola AF (2008). Contamination indices and heavy metal concentrations in urban soil of Ibadan metropolis, Southwestern Nigeria. *Environ. Geochem Health* (30):243-254.
- Ogunfowokan AO, Okoh EK, Adenuga AA, Asubiojo OI (2005). An assessment of the impact of point source pollution from University sewage treatment Oxidation pond on a receiving stream – a preliminary study. *J. Appl. Sci.* 5(1):36-43.
- Peuranen S, Vuorinen PJ, Vuorinen M, Hollender A (1994). The effects of iron, humic acids and low pH on the gills and physiology of brown trout (*Salmo Trutta*). *Ann. Zool. Fenn.* 31:389-396.
- Radojevic M, Bashkin VN (1999). *Practical Environmental Analysis*. The Royal Society of Chemistry, Cambridge, UK, P 466.
- Senior BW (2006). Examination of water, milk, food and air. In: Colle J.G., Duguid J.P, FASTER A.G., Marmion, B.P, Simmons, A. (Eds). Mackie and McCartney Practical Medical Microbiology, pp. 883-892.
- Sharholly M, Ahmad MK, Mahmood G, Trivedi RC (2008). Municipal Solid Waste Management in Indian Cities – A review. *Waste Manage.* 28(2):459-467.
- Standard Organisation of Nigerian (SON) (2008). Nigerian Standard for Drinking Water Quality. ICS 13.060.20 Standard Organisation of Nigerian, SON.
- Vivan EL, Blamah NV, Ezemokwe I, Okafor CI (2011). An Assessment of Hospital Waste Management in Some Selected Hospitals in Zonkwa District of Zangon-Kataf Local Government Area of Kaduna State. *Int. J. Environ. Stud.* 1(2):23-28.
- World Health Organization (WHO) (2004). *World Health Organization. Guidelines for Drinking water Quality. Third Edition, Vol. 1. Recommendation WHO: Geneva, Switzerland 2004.* P 515.
- World Health Organization (WHO) (2005). *Safe Healthcare Waste Management Policy by WHO.* *Waste Manage.* 25:568-569.
- World Health Organization (WHO) (2006). *Guidelines for Drinking water Quality: Incorporation First Addendum. Vol 1, Recommendations, Geneva, Switzerland- 3rd Ed.*
- US Environmental Protection Agency (USEPA) (1984). *Health assessment document for manganese*. Final draft. Cincinnati, OH, US Environmental Protection Agency, Office of Research and Development. (Report No. EPA/600/8-83/013F).

A hand is shown holding a glass flask containing a blue liquid, positioned over a patch of soil. The background is slightly blurred, showing green plants and brown earth. The text is overlaid on a semi-transparent dark grey rectangle.

Journal of Environmental Chemistry and Ecotoxicology

Related Journals Published by Academic Journals

- *African Journal of Pure and Applied Chemistry*
- *International Journal of Physical Sciences*
- *Journal of Geology and Mining Research*
- *African Journal of Mathematics and Computer Science Research*
- *Journal of Internet and Information Systems*
- *Journal of Oceanography and Marine Science*
- *Journal of Petroleum Technology and Alternative Fuels*

academicJournals