



African Journal of Pharmacy and Pharmacology

Volume 11 Number 46, 15 December, 2017
ISSN 1996-0816



*Academic
Journals*

ABOUT AJPP

The **African Journal of Pharmacy and Pharmacology (AJPP)** is published weekly (one volume per year) by Academic Journals.

African Journal of Pharmacy and Pharmacology (AJPP) is an open access journal that provides rapid publication (weekly) of articles in all areas of Pharmaceutical Science such as Pharmaceutical Microbiology, Pharmaceutical Raw Material Science, Formulations, Molecular modeling, Health sector Reforms, Drug Delivery, Pharmacokinetics and Pharmacodynamics, Pharmacognosy, Social and Administrative Pharmacy, Pharmaceutics and Pharmaceutical Microbiology, Herbal Medicines research, Pharmaceutical Raw Materials development/utilization, Novel drug delivery systems, Polymer/Cosmetic Science, Food/Drug Interaction, Herbal drugs evaluation, Physical Pharmaceutics, Medication management, Cosmetic Science, pharmaceuticals, pharmacology, pharmaceutical research etc. The Journal welcomes the submission of manuscripts that meet the general criteria of significance and scientific excellence. Papers will be published shortly after acceptance. All articles published in AJPP are peer-reviewed.

Contact Us

Editorial Office: ajpp@academicjournals.org

Help Desk: helpdesk@academicjournals.org

Website: <http://www.academicjournals.org/journal/AJPP>

Submit manuscript online <http://ms.academicjournals.me/>

Editors

Himanshu Gupta

*Department of Pharmacy Practice
University of Toledo
Toledo, OH
USA.*

Prof. Zhe-Sheng Chen

*College of Pharmacy and Health Sciences
St. John's University
New York,
USA.*

Dr. Huma Ikram

*Neurochemistry and Biochemical
Neuropharmacology Research Unit,
Department of Biochemistry,
University of Karachi
Karachi-75270
Pakistan*

Dr. Shreesh Kumar Ojha

*Molecular Cardiovascular Research Program
College of Medicine
Arizona Health Sciences Center
University of Arizona
Arizona,
USA.*

Dr. Vitor Engracia Valenti

*Departamento de Fonoaudiologia
Faculdade de Filosofia e Ciências,
UNESP
Brazil.*

Dr. Caroline Wagner

*Universidade Federal do Pampa
Avenida Pedro Anunciação
Brazil.*

Dr. Ravi Shankar Shukla

*Macromolecule and Vaccine Stabilization Center
Department of Pharmaceutical Chemistry
University of Kansas
USA.*

Associate Editors

Dr. B. Ravishankar

*SDM Centre for Ayurveda and Allied Sciences,
SDM College of Ayurveda Campus,
Karnataka
India.*

Dr. Natchimuthu Karmegam

*Department of Botany,
Government Arts College,
Tamil Nadu,
India.*

Dr. Manal Moustafa Zaki

*Department of Veterinary Hygiene and
Management
Faculty of Veterinary Medicine,
Cairo University
Giza,
Egypt.*

Prof. George G. Nomikos

*Takeda Global Research & Development Center
USA.*

Prof. Mahmoud Mohamed El-Mas

*Department of Pharmacology,
Faculty of Pharmacy
University of Alexandria,
Alexandria,
Egypt.*

Dr. Kiran K. Akula

*Electrophysiology & Neuropharmacology Research
Unit
Department of Biology & Biochemistry
University of Houston
Houston, TX
USA.*

Editorial Board

Prof. Fen Jicai

School of life science, Xinjiang University, China.

Dr. Ana Laura Nicoletti Carvalho

Av. Dr. Arnaldo, 455, São Paulo, SP. Brazil.

Dr. Ming-hui Zhao

*Professor of Medicine
Director of Renal Division, Department of Medicine
Peking University First Hospital
Beijing 100034
PR. China.*

Prof. Ji Junjun

Guangdong Cardiovascular Institute, Guangdong General Hospital, Guangdong Academy of Medical Sciences, China.

Prof. Yan Zhang

*Faculty of Engineering and Applied Science,
Memorial University of Newfoundland,
Canada.*

Dr. Naoufel Madani

*Medical Intensive Care Unit
University hospital Ibn Sina, Univesity Mohamed V
Souissi, Rabat,
Morocco.*

Dr. Dong Hui

Department of Gynaecology and Obstetrics, the 1st hospital, NanFang University, China.

Prof. Ma Hui

School of Medicine, Lanzhou University, China.

Prof. Gu HuiJun

School of Medicine, Taizhou university, China.

Dr. Chan Kim Wei

*Research Officer
Laboratory of Molecular Biomedicine,
Institute of Bioscience, Universiti Putra,
Malaysia.*

Dr. Fen Cun

Professor, Department of Pharmacology, Xinjiang University, China.

Dr. Sirajunnisa Razack

Department of Chemical Engineering, Annamalai University, Annamalai Nagar, Tamilnadu, India.

Prof. Ehab S. EL Desoky

Professor of pharmacology, Faculty of Medicine Assiut University, Assiut, Egypt.

Dr. Yakisich, J. Sebastian

*Assistant Professor, Department of Clinical Neuroscience R54
Karolinska University Hospital, Huddinge
141 86 Stockholm ,
Sweden.*

Prof. Dr. Andrei N. Tchernitchin

*Head, Laboratory of Experimental Endocrinology and Environmental Pathology LEEPA
University of Chile Medical School,
Chile.*

Dr. Sirajunnisa Razack

*Department of Chemical Engineering,
Annamalai University, Annamalai Nagar, Tamilnadu, India.*

Dr. Yasar Tatar

*Marmara University,
Turkey.*

Dr Nafisa Hassan Ali

*Assistant Professor, Dow institute of medical technology
Dow University of Health Sciences, Chand bbi Road, Karachi,
Pakistan.*

Dr. Krishnan Namboori P. K.

*Computational Chemistry Group, Computational Engineering and Networking,
Amrita Vishwa Vidyapeetham, Amritanagar, Coimbatore-641 112
India.*

Prof. Osman Ghani

*University of Sargodha,
Pakistan.*

Dr. Liu Xiaoji

*School of Medicine, Shihezi University,
China.*

ARTICLE

GC-MS analysis of leaf, stem-bark and root extracts of <i>Alstonia boonei</i> Babatunde, O.	577
---	------------

Full Length Research Paper

GC-MS analysis of leaf, stem-bark and root extracts of *Alstonia boonei*

Babatunde, O.

Department of Chemical Sciences, Ajayi Crowther University, Oyo, Nigeria.

Received 28 October, 2017; Accepted 4 December, 2017

Alstonia boonei De Wild is a medicinal plant commonly found in West African and is popularly known as God's tree. The plant parts have been traditionally used as a painkiller, antimicrobial, antimalarial and antidiabetic which have also been proved scientifically. Previous studies revealed little information on phytochemical components of *A. boonei*. The present work aims at investigating and comparing the chemical components of the leaf, the stem-bark and the root of the plant in order to provide sufficient baseline information for isolation work on medicinal components of the plant. Leaf, stem-bark and root extracts of *A. boonei* were prepared by maceration using 1:1 EtOAc/MeOH. The crude extract was successively macerated with hexane, dichloromethane (DCM) and methanol. Thin layer chromatography (TLC) by 2,2-diphenyl-1-picrylhydrazyl (DPPH) (TLC-DPPH) analysis was used to screening out DCM fraction for further analysis. Gas chromatography and mass spectroscopy studies were performed to profile phytochemical constituent of the plant. The GC-MS analysis of DCM extract of the leaf revealed ten chemical components with Eugenol as major component (54.58%); DCM extract of the stem-bark showed forty one components with alpha-amyrin (32.25%) while DCM extract of the root revealed twenty components with 1,2-benzenedicarboxylic acid (49.2%) as major component. This study shows that the *A. boonei* extracts of the leaf, stem-bark and root consist of different types of compounds with few components common to two of the parts. Quantitatively, common phytochemicals decrease from leave to root. The most prominent compounds identified by GC/MS were Eugenol, benzenedicarboxylic acid and alpha-amyrin.

Key words: Phytochemicals, *A. boonei*, gas chromatography/mass spectroscopic studies, Eugenol

INTRODUCTION

Alstonia boonei De Wild belongs to the family Apocynaceae. It is a herbal medicinal plant of West African origin, popularly known as God's tree, cheese wood and known locally among Yoruba in Nigeria as Ahun. All the parts of the plant are very useful but the

thick bark cut from the matured tree is the part that is most commonly used for therapeutic purposes. Therapeutically, beta-amyrin and alpha-amyrin acetate isolated from the stem bark of *A. boonei* have been found to possess anti-inflammatory properties (Okoye et al.,

E-mail: o.babatunde@acu.edu.ng

Author(s) agree that this article remain permanently open access under the terms of the [Creative Commons Attribution License 4.0 International License](https://creativecommons.org/licenses/by/4.0/)




Figure 1. SL = Hex LC = MeOH, EL = DCM

2014) Antimicrobial activities of various fractions of stem bark of *A. boonei* were also tested against some microbes (Bello et al., 2009; Amole and Ilori, 2010; Misra et al., 2011). A wide array of chemical compounds have been isolated from *A. boonei*. These include alkaloids, tannins, iridoids, and triterpenoids which have been demonstrated to possess some pharmacological activities (Akinmoladun et al., 2007). The alkaloids isolated from the plant include echitamine and other alkaloids, and the triterpenes b-amyrin, lupenol, and ursolic acid have all been isolated from leaves and stem bark (Adotey et al., 2012). Echitamine possess a battery of pharmacological and autonomic activities (Ojewole, 1983, 1984) including anticancer activities (Adotey et al., 2012; Ashok et al., 2015). Moronkola and Kunle (2012) reported essential oil compositions of the leaf, stem bark, and root of *A. boonei*. (Z)-9-Octadecenoic acid was found to be the most abundant volatile oil in the leaf and stem bark while methyl (7 E)- 7-octadecenoate was the most abundant in the root. The aim of this study was to investigate the distribution trend of *A. boonei* chemical composition

MATERIALS AND METHODS

Sample preparation

A. boonei leaf, stem-bark and root were collected at a farmland in Akinmarin village, Oyo and were identified at the Department of Botany, University of Ibadan, Ibadan. The air-dried plant materials: (204.7 g) leaves, (282.5 g) stem-bark, and (102.9 g) root of, *A. boonei*, were exhaustively extracted separately with 1:1 EtOAc:MeOH for 3 days. The extracts were filtered separately, concentrated on rotary evaporator to about 50 ml and evaporated to dryness under vacuum. Each extract was successively macerated with hexane, dichloromethane and methanol. Obtained fractions were evaporated to dryness and stored in a refrigerator at +4°C until use.

Antioxidant screening

The method of Mensor et al. (2001) with Kotze and Eloff (2002) methods were adopted to screen fractions. Fractions obtained from leaves were developed on aluminium-backed thin layer chromatography (TLC) using solvent ratio: chloroform/ethyl acetate/formic acid (5:4:1) (intermediate polarity/acidic). Chromatogram was sprayed with 0.2%, 2,2, diphenyl-2-picryl-hydrazyl to detect antioxidant properties of various fractions.

Gas chromatography (GC)-mass spectrometer (MS) analysis

GC-MS: Hewlett-Packard 5890 gas chromatograph, combined with a Jeol JMS-HX 110 mass spectrometer with source at 270°C at 70 eV. Injector was set at 270°C with splitting ratio 1:30. The analysis was performed on the aforementioned programme on equivalent column HP-5 (25 m × 0.22 mm and 0.25 μm). A mass spectral survey was performed using the NIST mass spectral search program.

RESULTS AND DISCUSSION

Qualitative analysis of antioxidant activity was used to screen out less potent fraction. The dichloromethane fraction displayed highest antioxidant activity after spraying chromatogram with DPPH because of the intensity of yellow colour produce. Dichloromethane (DCM) was then selected for phyto-constituent analysis. Thin layer chromatogram of hexane, dichloromethane and methanol fractions obtained from the leaves is shown in Figure 1. GC-MS analyses of DCM fraction of the leaf, the stem-bark and the root column of *A. boonei* are presented in Table 1. Constituents were listed in order of elution from HP-5 capillary column (Table 1). Gas chromatography-mass spectrometry analysis of the leaf extract resulted in the identification of 10 compounds, 41 compounds in stem-bark extract and 20 compounds in root extract.

Table 1. Chemical composition of leaf, stem-bark and root of *Alstonia boonei*.

S/N	Constituent	% composition		
		Leaf	Stem-bark	Root
1	Eugenol	54.58(9.730)	0.14(9.722)	-
2	4-Tetradecene	-	-	1.39(10.185)
3	Caryophyllene	6.99(10.631)	-	-
4	Phenol	2.01(12.148)	-	0.56(11.970)
5	1H-2-Benzopyran-1-one	-	0.54(12.434)	-
6	Cyclododecane	-	0.28(12.943)	-
7	Dichloroacetic acid	-	-	4.83(12.943)
8	Hexadecane	-	0.16(13.033)	1.32(13.029)
9	2H-Pyran-2-one	-	1.54(13.161)	-
10	2-Ethyl-3-methoxypyrazine	-	0.34(14.334)	-
11	1-Octadecene	-	0.37(15.324)	6.59(15.323)
12	Octadecane	-	0.15(15.398)	1.35(15.392)
13	Cyclononasiloxane	-	0.20(15.713)	-
14	Bicyclo(3.1.1)heptane	2.61(15.833)	-	-
15	1,2-Benzenedicarboxylic acid	-	-	49.20(16.210)
16	Phthalic acid	-	0.15(16.216)	-
17	7,9-Di-tert-butyl-1-oxaspiro(4,5) deca-6,9-diene-2,8-dione	-	-	1.13(16.777)
18	Hexadecanoic acid	2.13(16.765)	0.67(16.777)	-
19	n-Hexadecanoic acid	5.40(17.183)	1.40(17.303)	-
20	Dibutyl phthalate	-	-	2.61(17.200)
21	6H-Furo[2',3':4,5] oxazolo [3,2-a]pyrimidin-6-one	-	0.24(17.338)	-
22	Heptafluorobutyric acid	-	0.25(17.458)	-
23	n-Nonadecanol-1	-	0.28(18.391)	5.90(17.452)
24	Eicosane	-	0.16(20.365)	1.34(17.509)
25	Ethanone	-	-	1.30(17.767)
26	1-Heneicosanol	2.88(17.950)	-	-
27	Behenic alcohol	-	-	3.21(18.373)
28	9,12-Octadecadienoic acid	-	0.34(18.488)	-
29	11-Octadecenoic acid	-	0.26(18.545)	-
30	Phytol	1.63(18.665)	-	-
31	3-Isopropoxy-1,1,1,7,7,7-hexamethyl-3,5,5-tris(trimethylsiloxy)tetrasiloxane	-	0.33(18.820)	-
32	Oleic Acid	-	0.46(18.986)	-
33	1-Octadecene	-	0.19(19.192)	-
34	1-Docosene	13.28(22.585)	-	3.65(19.392)
35	2-Eicosanol	-	0.14(19.398)	-
36	Docosane	-	-	0.66(19.438)
37	6-Acetamido-2-methylbenzothiazole	-	0.78(20.090)	-
38	Cyclotetracosane	-	-	2.56(21.337)
39	Hexasiloxane	-	0.40(21.686)	-
40	2(3H)-Furanone	-	-	0.82(22.430)
41	1-Heptadecene	-	-	0.92(22.659)
42	Tetratetracontane	-	0.95(22.693)	-
43	Oxirane	-	0.27(23.186)	-
44	Olean-12-ene	-	0.63(24.948)	-
45	1S,6R,9S)-5,5,9,10-Tetramethyltricyclo [7.3.0.0(1,6)] dodec-10(11)-en	-	1.29(25.245)	-
46	12-Oleanen-3-yl acetate	-	0.66(25.366)	-
47	2H-Cyclopropa[a]naphthalen-2-one	-	1.10(25.474)	-
48	Heptadecane	-	-	1.53(26.304)
49	Stigmasterol	-	0.22(29.182)	-
50	2,6,10,14,18,22-Tetracosahexaene	8.50(29.829)	-	-

Table 1. Cont'd

51	5(1H)-Azulenone	-	1.53(30.401)	-
52	Urs-12-en-24-oic acid	-	0.50(30.447)	-
53	Urs-12-en-3-ol	-	7.76(30.922)	-
54	2-Isopropenyl-4a,8-dimethyl-1,2,3, 4,4a,5,6,8a-octahydronaphthalene	-	2.44(32.009)	-
55	Nonacosane	-	0.68(32.524)	-
56	Azuleno[6,5-b]furan-2,5-dione	-	-	4.15(33.439)
57	Bicyclo[2.2.1]heptane	-	3.89(33.582)	-
58	alpha.-Amyrin	-	32.25(35.276)	-
59	Pyrrolo[2,3-b]indole	-	4.05(36.049)	-
60	2(1H)Naphthalenone	-	10.32(36.094)	-
61	9,19-Cycloergost-24(28)-en-3-ol	-	9.13(37.771)	-

Note: Values in bracket are retention time

Table 2. Comparative analysis of chemical profiles of the whole *Alstonia boonei* plant.

S/N	Constituent	% composition		
		Leaf	Stem-bark	Root
1	Eugenol	54.58(9.730)	0.14(9.722)	-
2	Phenol	2.01(12.148)	-	0.56(11.970)
3	Hexadecane	-	0.16(13.033)	1.32(13.029)
4	1-Octadecene	-	0.37(15.324)	6.59(15.323)
5	Octadecane	-	0.15(15.398)	1.35(15.392)
6	Hexadecanoic acid	2.13(16.765)	0.67(16.777)	-
7	n-Hexadecanoic acid	5.40(17.183)	1.40(17.303)	-
8	n-Nonadecanol-1	-	0.28(18.391)	5.90(17.452)
9	Eicosane	-	0.16(20.365)	1.34(17.509)
10	1-Docosene	13.28(22.585)	-	3.65(19.392)

Note: Values in bracket are retention time

Chemical compounds that were common to the three parts of the plant were presented in Table 2. A comparative analysis of chemical profiles of the leaf and stem-bark showed that hexadecanoic acid and n-hexadecanoic acid were present in the two parts but in higher proportion in the leaf while phenol and 1-Docosene were present in the leaf and the root. Hexadecane, 1-Octadecene, Octadecane, n-Nonadecanol-1 and Eicosane were present in both stem-bark and root. There is decrease in quantity of common phytochemicals from leaf to root and root to stem. The results revealed that Eugenol (54.58), 1-Docosene (13.28), 2,6,10,14,18,22-Tetracosahexaene (8.50), Caryophyllene (6.99) and n-Hexadecanoic acid (5.40) were found as the five major components in the leaf extract, the minor compound was Phytol. The major constituents of stem-bark extract were alpha-amyrin (32.25), 2(1H) Naphthalenone (10.32), 9,19-Cycloergost-24(28)-en-3-ol (9.13) and Urs-12-en-3-ol (7.76) while the major component found in root were 1,2-Benzenedicarboxylic acid (49.20), 1-Octadecene(6.59)

And n-Nonadecanol-1(5.90).

Jaganathan and Supriyanto (2012) reported that eugenol possess anticancer activity against various types of cancers. Bioactivity experiments revealed positive anti-cancer activity of 1, 2-Benzenedicarboxylic acid on PC3, MCF, HCT-116, A549, and MIAPACA cell lines (Save et al., 2015). Beta-Amyrin and alpha-amyrin acetate isolated from the stem bark of *A. boonei* display profound anti-inflammatory activity (Nkeoma et al., 2014).

Conclusion

The present study represents the comprehensive analysis of phyto-constituents of *A. boonei* leaf, stem-bark, and root DCM extracts. Percentage composition of common phytochemicals decreases from leaf to root. Previous study revealed that the three main components of *A. boonei* identified by GC-MS exhibit anticancer activity. Therefore, the results of this study will form the basis for selection of plant part for further investigation in

the potential drug discovery.

CONFLICT OF INTEREST

The author has not declared any conflict of interest.

REFERENCES

- Adotey JPK, Adukpo GE, Boahen YO, Armah FA (2012). A review of the ethnobotany and pharmacological importance of *Alstonia boonei* De Wild (Apocynaceae). *Int. Scholar Res. Net* 1:1-9.
- Akinmoladun AC, Ibukun EO, Afor E (2007). Chemical constituents and antioxidant activity of *Alstonia boonei*. *Afri J Biotech* 6(10):1197-201.
- Amole OO and Ilori OO (2010). Antimicrobial Activity Of The Aqueous And Ethanolic Extracts Of The Stem Bark Of *Alstonia boonei*. *Int. J. Phytopharmacol.* 1(2):119-123.
- Ashok R1, Lokesh SJ, Hari Kiran (2015) Studies on the anticataract activity of echitamine, an alkaloid from the stem bark of *Alstonia scholaris*. *Der Pharmacia Lettre* 7(5):197-199
- Bello IS, Oduola T, Adeosun OG, Omisore NOA, Raheem GO, Ademosun AA (2009) Evaluation of Antimalarial Activity of Various Fractions of *Morinda lucida* Leaf Extract and *Alstonia boonei* Stem Bark. *Glob. J. Pharmacol.* 3(3):163-165.
- Jaganathan SK, Supriyanto E (2012) Antiproliferative and molecular mechanism of eugenol-induced apoptosis in cancer cells. *Molecules* 17(6):6290-6304.
- Kotze M, Eloff JN (2002). Extraction of antibacterial compounds from *Combretum microphyllum* (Combretaceae). *South Afr. J. Bot.* 68(1):62-67.
- Mensor LL, Fabio SM, Gildor GL, Alexander SR, Tereza CD, Cintia SC, Suzane GL (2001). Screening of Brazilian plant extracts for antioxidant activity by the use of DPPH free radical methods. *Phytother. Res.* 15(2):127-130.
- Misra CS, Pratyush K, Sagadevan, LDM, James J, Veettil AKT, Thankamani V (2011). A comparative study on phytochemical screening and antibacterial activity of roots of *Alstonia scholaris* with the roots, leaves and stem bark. *Int. J. Res. Phytochem. Pharmacol.* 1(2):77-82.
- Moronkola DO, Kunle OF (2012). Essential oil compositions of leaf, stem bark and root of *Alstonia boonei* de Wild (Apocynaceae). *Int. J. Biol. Pharm. Res.* 3(1):51-60.
- Nkeoma NO, Daniel LA, Henry NO, Emmanuel EI, Chukwuemeka SN, Festus BCO (2014) beta-Amyrin and alpha-amyrin acetate isolated from the stem bark of *Alstonia boonei* display profound anti-inflammatory activity. *Pharm. Biol.* 52(11):1478-1486
- Ojewole JAO (1983). Autonomic pharmacology of echitamine, an alkaloid from *Alstonia boonei* De Wild. *Fitoterapia* 54(3):99-113.
- Ojewole JAO (1984) Studies on the pharmacology of echitamine, an alkaloid from the stem bark of *Alstonia boonei* L. (Apocynaceae). *Int. J. Crude Drug Res.* 22(3):121-143.
- Save SA, Lokhande RS, Chowdhary AS (2015) Determination of 1, 2-Benzenedicarboxylic acid, bis (2-ethylhexyl) ester from the twigs of *Thevetia peruviana* as a Colwell Biomarker. *J. Innov. Pharm. Biol. Sci.* 2(3):349-362.



African Journal of Pharmacy and Pharmacology

Related Journals Published by Academic Journals

- *Journal of Medicinal Plant Research*
- *African Journal of Pharmacy and Pharmacology*
- *Journal of Dentistry and Oral Hygiene*
- *International Journal of Nursing and Midwifery*
- *Journal of Parasitology and Vector Biology*
- *Journal of Pharmacognosy and Phytotherapy*
- *Journal of Toxicology and Environmental Health Sciences*

academicJournals