



African Journal of Pure and Applied Chemistry

Volume 11 Number 6 December 2017

ISSN 1996-0840



*Academic
Journals*

ABOUT AJPAC

The **African Journal of Pure and Applied Chemistry (AJPAC)** is an open access journal that publishes research analysis and inquiry into issues of importance to the science community. Articles in AJPAC examine emerging trends and concerns in the areas of theoretical chemistry (quantum chemistry), supramolecular and macromolecular chemistry, relationships between chemistry and environment, and chemicals and medicine, organometallic compounds and complexes, chemical synthesis and properties, chemicals and biological matters, polymer synthesis and properties, nanomaterials and nanosystems, electrochemistry and biosensors, chemistry and industry, chemistry and biomaterials, advances in chemical analysis, instrumentation, speciation, bioavailability. The goal of AJPAC is to broaden the knowledge of scientists and academicians by promoting free access and provide valuable insight to chemistry-related information, research and ideas. AJPAC is a bimonthly publication and all articles are peer-reviewed.

African Journal of Pure and Applied Chemistry (AJPAC) is published twice a month (one volume per year) by Academic Journals.

Contact Us

Editorial Office: ajpac@academicjournals.org

Help Desk: helpdesk@academicjournals.org

Website: <http://www.academicjournals.org/journal/AJPAC>

Submit manuscript online <http://ms.academicjournals.me/>.

Editors

Prof. Tebello Nyokong

*Acting Editor
Chemistry Department
Rhodes University
Grahamstown 6140,
South Africa.*

Prof. F. Tafesse

*Associate Editor
Associate professor
Inorganic chemistry
University of South Africa
South Africa.*

Editorial Board

Dr. Fatima Ahmed Al-Qadri

*Asst. Professor
Chemistry Department
Sana'a University
Republic of Yemen.*

Dr. Aida El-Azzouny

*National Research Center
(NRC, Pharmaceutical and
Drug Industries Research Division)
Dokki-Cairo, 12622-Egypt.*

Dr. Santosh Bahadur Singh

*Department of Chemistry
University of Allahabad
Allahabad, India.*

Dr. Gökhan Gece

*Department of Chemistry
Bursa Technical University
Bursa, Turkey.*

Dr. Francisco Torrens

*Institute for Molecular Science
University of Valencia
Paterna Building Institutes
P. O. Box 22085
E-46071 Valencia
Spain.*

Dr. Erum Shoeb

*Asst. Professor
Department of Genetics
University of Karachi
Karachi-75270
Pakistan.*

Dr. Ishaat Mohammad Khan

*Physical Research Laboratory
Department of Chemistry
Aligarh Muslim University
Aligarh 202002, India.*

Prof. Jean-Claude Bunzli

*Department of Chemistry
Swiss Federal Institute of Technology Lausanne
(EPFL)
Institute of Chemical Sciences and Engineering
BCH 1402
CH-1015 Lausanne (Switzerland).*

Mrinmoy Chakrabarti

*Department of Chemistry,
Texas A&M University
415 Nagle Street, College Station, TX 77840
USA.*

Dr. Geoffrey Akien

*430 Eisenhower Drive, Apartment B-2,
Lawrence, Kansas 66049,
United States.*

Prof. Anil Srivastava

*Jubilant Chemsys Ltd.,
B-34, Sector-58,
Noida 201301 (UP),
India.*

ARTICLE

- Determination of mercury and cadmium levels in omega-3 food supplements available on the Ghanaian market** 50
Adolf Oti-Boakye, Akwasi Acheampong, Ohene Gyang Nathan,
Akorfa Akosua Agbosu and Amoah Charles Agyei

Full Length Research Paper

Determination of mercury and cadmium levels in omega-3 food supplements available on the Ghanaian market

Adolf Oti-Boakye^{1*}, Akwasi Acheampong¹, Ohene Gyang Nathan², Akorfa Akosua Agbosu² and Amoah Charles Agyei³

¹Department of Chemistry, Kwame Nkrumah University of Science and Technology, Kumasi, Ghana.

²Department of Science, Akrokeri College of Education, Ghana.

³Department of Science, Ola College of Education, Ghana.

Received 4 February, 2016; Accepted 17 March, 2017

The numerous reputed health benefits of the omega-3s (eicosapentaenoic acid [EPA] and docosahexaenoic acid [DHA]), particularly, their cardio-protective effects have led to the manufacture of omega-3 supplements by various pharmaceutical companies resulting in their flooding of the Ghanaian market. Coldwater fishes which are the primary sources of the omega-3 fatty acids are known to have high levels of mercury and cadmium in them. There is therefore the potential of mercury and cadmium poisoning in the course of people taking the omega-3 food supplements. Mercury and cadmium levels in ten products of omega-3 food supplements have been determined in order to ascertain their safety for human consumption. All the levels of mercury and cadmium determined were within the acceptable limits stipulated by Food and Agriculture Organization and World Health Organization, and therefore do not pose any health threat to consumers.

Key words: Metals, omega-3, cardio-protective, pharmaceutical.

INTRODUCTION

Omega-3 fatty acids belong to a class of fatty acids that are called essential fatty acids (EFAs). They are so called because the body cannot produce them and thus must be obtained from the diet (Weber et al., 2006). Three major nutritionally important omega-3 fatty acids that are ingested in foods and used by the body are alpha-linolenic acid (ALA), eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA) with ALA being the primary omega-3 fatty acid (Weber et al., 1986). EPA and DHA

have been receiving a lot of attention because of their cardio-protective and other so called "pleiotropic" effects (Weber et al., 1986).

The primary source of omega-3 has been the oils of cold-water fish, such as tuna, salmon, trout, herring, sardines, bass, swordfish and mackerel, etc., which in turn derive the omega-3s from feeding on the algae, the ultimate biological source (Weber et al., 2006). Studies have revealed that since these coldwater fishes are

*Corresponding author. E-mail: oticasty@yahoo.com. Tel: +233202520284.

Table 1. Recovery test for metal analysis.

| Sample weight | Percentage recovery | |
|--------------------------|---------------------|---------|
| | Mercury | Cadmium |
| 25 ng | 88.00 | 90.00 |
| 50 ng | 90.00 | 94.00 |
| 100 ng | 90.00 | 92.00 |
| Mean percentage recovery | 89.33 | 92.00 |

known to have high levels of mercury and cadmium in them, there is the likelihood of mercury and cadmium poisoning in the course of people taking the omega-3s from these sources (Mahaffey, 2004).

There is a strong link between micronutrient uptake by fish and their products, and the impact of contaminations on humans (De Leonardis et al., 2000; Yuzbasi et al., 2003). Mercury and cadmium belong to the class of toxic heavy metals in our environment and have no known biological function in the body (Zhang and Wong, 2007; WHO, 2008).

Humans are most commonly exposed to these metals primarily by eating fish (and fish products) and marine mammals (e.g., whales, seals) that may contain some of these metals in their tissues (International Programme on Chemical Safety [IPCS], 1990; WHO, 2008; Sallsten et al., 1996).

Thus, the benefits of the omega-3 food supplements may be completely reversed if it is found to have high levels of these toxic metals in them (Stillwell and Wassall, 2003; United States Environmental Protection Agency, 2004). Levels of mercury and cadmium in the omega-3 supplements consumed should therefore be of importance and concern. There is lack of data on the levels of mercury and cadmium in omega-3 supplement products imported into Ghana. It is in the light of this that this work was done to measure their levels in various omega-3 supplement products available on the Ghanaian market and ensure their safety for human consumption.

MATERIALS AND METHODS

Sampling and sample preparation

A total of 100 samples of omega-3 food supplement soft gels comprising ten different brands were purchased from retail outlets on the Ghanaian Market between November, 2009 and April, 2010. Samples were transported to the laboratory, coded for easy identification and then stored in the fridge until time for analysis.

Digestion procedure for the determination of metals

The omega-3 samples were digested for the total metal determination by an open flask procedure (Akagi and Nishimura, 1991). One soft gel sample was put in a 50 ml volumetric digestion flask. 1 ml distilled water, 2 ml HNO₃:HClO₄ (1:1) and 5 ml H₂SO₄ were then added in turn and the mixture was heated within the range of 195 to 205°C for 30 min. The sample solution was then cooled and diluted to 50 ml with double distilled water.

Determination of the metals

Determination of mercury in all the digests was carried out by cold vapor atomic absorption spectrophotometer using an automatic Mercury Analyzer Model HG-5000 (Sanso Seisakusho Co., Ltd, Japan). The procedure is as described by Adimado and Baah (2002).

Determination of cadmium in all the digests was done using SOLAAR (S Series 711239 v1.23) Flame Atomic Absorption Spectrometer.

Recovery of the metals was determined by adding 25, 50 and 100 ng to two different samples of omega-3. The samples were taken through the digestion procedure. The resulting solutions were analyzed for mercury and cadmium concentration using the appropriate methods. Percentage recovery for the spiked solutions was then calculated.

RESULTS AND DISCUSSION

The mean percentage recoveries (Table 1) for mercury and cadmium were 89.33 and 92.00%, respectively signifying the reliability of the method.

The results showing the mean levels of mercury and cadmium in the omega-3 samples are shown in Table 2. The mean daily intakes of the metals are also shown in Table 3. The calculations for the daily metal intakes were based on the recommended daily dosages stated on the products.

Heavy metal content in fish is a good indicator for human exposure to metal bioaccumulation and has been confirmed in many publications (Love et al., 2003; Goyer, 1997; Adimado and Baah, 2002). This means that the consumption of fish products could be a source of human exposure to heavy metals. The main source of the omega-3 oils are the deep sea fishes. The level of heavy metal accumulation in fishes therefore has a direct correlation to the level of heavy metal in the omega-3 product. The mean metal concentrations (µg/g) recorded in this study are comparable to the concentration (µg/g) obtained by Vanaja et al. (2007) and Kotb et al. (1991) who determined the concentrations of trace metals in some brands of fish oil supplements and the levels of some heavy metals in omega-3 fatty acids in popular species of Arabian Gulf fish, respectively (Vanaja et al., 2007; Kotb et al., 1991).

The different levels of the metals in the omega-3 products may be due to the fact that the oils were from different fishes from different sources at perhaps different

Table 2. Average concentration of metals ($\mu\text{g/g}$) in the omega-3 samples.

| Omega-3 samples | Mercury | Cadmium |
|-----------------|-------------------|-------------------|
| A | 0.017 \pm 0.003 | 0.387 \pm 0.062 |
| B | 0.093 \pm 0.002 | 0.729 \pm 0.014 |
| C | 0.021 \pm 0.003 | 0.995 \pm 0.047 |
| D | 0.273 \pm 0.005 | 1.226 \pm 0.110 |
| E | 0.123 \pm 0.004 | 0.785 \pm 0.103 |
| F | 0.658 \pm 0.001 | 0.788 \pm 0.014 |
| G | 0.018 \pm 0.005 | 0.762 \pm 0.030 |
| H | 0.027 \pm 0.008 | 0.531 \pm 0.025 |
| I | 0.428 \pm 0.002 | 0.762 \pm 0.035 |
| J | 0.029 \pm 0.004 | 0.611 \pm 0.063 |

Table 3. Mean daily intake of metals in the omega-3 samples ($\mu\text{g/g}$).

| Omega-3 sample | Stated daily dosage on product (Capsule) | Mercury | Cadmium |
|----------------|--|---------|---------|
| A | 2 | 0.033 | 0.774 |
| B | 1 | 0.093 | 0.729 |
| C | 1 | 0.021 | 0.995 |
| D | 3 | 0.818 | 3.678 |
| E | 6 | 0.738 | 4.712 |
| F | 1 | 0.658 | 0.789 |
| G | 3 | 0.053 | 2.287 |
| H | 1 | 0.027 | 0.531 |
| I | 1 | 0.428 | 0.762 |
| J | 1 | 0.029 | 0.611 |

trophic levels (Love et al., 2003), at different factors such as organic matter content, pH, seasonal changes, microbial activities, regional variations and hydrologic conditions as well as the rate of atmospheric deposition (Lindquist, 1991). Different methods of extraction or refining and thermal treatment could also account for the difference in the metal concentrations (Seidler, 1987).

Conclusion

The calculated daily intakes determined for the metals were below the daily intakes recommended by the Joint FAO/WHO expert committee on food supplements and additives showing that the omega-3 food supplements analyzed in this study are not likely to pose any health risk to the public from taking the omega-3 products as far as mercury and cadmium are concerned.

CONFLICT OF INTERESTS

The authors have not declared any conflict of interests.

REFERENCES

- Adimado AA, Baah DA (2002). Mercury in human blood, urine hair, nail and fish from Ankobra and Tano river Basin in South Western Ghana. *Bull. Environ. Contam. Toxicol.* 68:339-346.
- Akagi H, Nishimura H (1991). Speciation of Mercury in the environment. In: T. Suzuki T, Imura N, Clarkson TW (Eds), *Advances in mercury Toxicology*, Plenum press, USA. pp. 53-76.
- De Leonardis A, Macciola V, De Felice M (2000). Copper and iron determination in edible vegetable oils by graphite furnace atomic absorption spectrometry after extraction with diluted nitric acid. *IJFST* 35:371-375.
- Goyer AR (1997). Toxic metals and essential metal interactions. *Annu Rev Nutr.* 17:37-50.
- IPCS (International Programme on Chemical Safety), (1990). Methylmercury, World Health Organization. Geneva. EHC, 101.
- Kotb AR, Hadeed A, Al-Baker AA (1991). Omega-3 PUFAs and Heavy metal content in some popular species of Arabian Gulf fishes. *Food Chem.* 40:185-190.
- Lindquist O (1991) Mercury in Swedish environment. *Water Air Soil Publs.* 55(1-2):1-261.
- Love JL, Rush GM, McGrath H (2003). Total mercury and methylmercury levels in some New Zealand commercial marine fish species. *FAC* 120:37-43.
- Mahaffey KR (2004). Fish and shellfish as dietary sources of methylmercury and the omega-3 fatty acids, eicosahexaenoic acid and docosahexaenoic acid: risks and benefits. *Environ. Res.* 95(3):414-428.

- Sallsten G, Thoren J, Barregard L, Schutz A, Skarping G (1996). Long-term use nicotine chewing gum and mercury exposure from dental amalgam fillings. *J. Dent. Res.* 75:594-598.
- Seidler T (1987). Effect of additives and thermal treatment on the content of Nitrogen compounds and nutritive value of hake meat. *Die Nahrung* 31(10):959-70.
- Stillwell W, Wassall SR (2003). Docosahexaenoic acids membrane properties of unique fatty acids. *J. Am. Diet. Assoc.* 107:1599-1611.
- US EPA (United States Environmental Protection Agency), (2004). What you need to know about mercury in fish and shellfish. EPA-823-F-04-009 2.
- Vanaja S, Driscoll B, Obenarf R (2007). Trace elements in fish and fish oil supplements. *The Application Notebook, Atomic Spectroscopy, Spex Certiprep.* 123:13-16.
- Weber PC, Fischer S, von Schacky C (1986). The conversion of dietary eicosapentaenoic acid to prostaglandins and leukotrienes in man. *Prog. Lipid Res.* 25:273-276.
- Weber HS, Selini D, Huber G (2006). Prevention of cardiovascular diseases and highly concentrated n-3 polyunsaturated fatty acids (PUFAs). *Herz.* 31(3):24-30.
- WHO (World Health Organization) (2008). Elemental mercury and inorganic mercury compounds: Human health aspects. *CICAD* 61:12-65.
- Yuzbasi N, Sezgin E, Yildirim M, Yildirim Z (2003). Survey of lead, cadmium, iron, copper and zinc in Kasar cheese. *FAC* 20:464-469.
- Zhang I, Wong MH (2007). Environmental mercury contamination in China: sources and impacts. *Environ. Int.* 33:108-121.



African Journal of Pure and Applied Chemistry

Related Journals Published by Academic Journals

- *African Journal of Mathematics and Computer Science Research*
- *International Journal of the Physical Sciences*
- *Journal of Geology and Mining Research Technology*
- *Journal of Environmental Chemistry and Ecotoxicology*
- *Journal of Internet and Information Systems*
- *Journal of Oceanography and Marine Science*
- *Journal of Petroleum Technology and Alternative Fuels*

academicJournals