

A close-up photograph of a fish's head, likely a species of sea bream, characterized by its vibrant red color and numerous white spots. The fish is shown in profile, facing left, against a solid blue background. The image is framed with rounded corners.

# International Journal of Fisheries and Aquaculture

Volume 9 Number 10 November 2017

ISSN 2006-9839



*Academic  
Journals*

## ABOUT IJFA

**The International Journal of Fisheries and Aquaculture (IJFA)** (ISSN: 2006-9839) is an open access journal that provides rapid publication (monthly) of articles in all areas of the subject such as algaculture, Mariculture, fishery in terms of ecosystem health, Fisheries acoustics etc.

The Journal welcomes the submission of manuscripts that meet the general criteria of significance and scientific excellence. Papers will be published shortly after acceptance. All articles published in the IJFA are peer-reviewed.

### Contact Us

**Editorial Office:** [ijfa@academicjournals.org](mailto:ijfa@academicjournals.org)

**Help Desk:** [helpdesk@academicjournals.org](mailto:helpdesk@academicjournals.org)

**Website:** <http://www.academicjournals.org/journal/IJFA>

**Submit manuscript online** <http://ms.academicjournals.me/>

## Editors

**Dr. V.S. Chandrasekaran**

*Central Institute of Brackishwater Aquaculture  
Indian Council of Agricultural Research (ICAR)  
Chennai,  
India.*

**Prof. Nihar Rajan Chattopadhyay**

*Department of Aquaculture  
Faculty of Fishery Sciences  
West Bengal University of Animal & Fishery Sciences  
West Bengal,  
India.*

**Dr. Lourdes Jimenez-Badillo**

*Ecology and Fisheries Centre  
Universidad Veracruzana  
Veracruz,  
México.*

**Dr. Kostas Kapiris**

*Institute of Marine Biological  
Resources of H.C.M.R.  
Athens,  
Greece.*

**Dr. Masoud Hedayatifard**

*Department of Fisheries Sciences and Aquaculture  
College of Agriculture and Natural Resources  
Advanced Education Center  
Islamic Azad University  
Ghaemshahr,  
Iran.*

**Dr. Zhang Xiaoshuan**

*China Agricultural University  
Beijing,  
China.*

**Dr Joseph Selvin**

*Marine Bioprospecting Lab  
Dept of Microbiology  
Bharathidasan University  
Tiruchirappalli,  
India.*

## Editorial Board

**Dr. Dada Adekunle Ayokanmi**  
*Department of Fisheries and Aquaculture  
Federal University of Technology  
Akure,  
Nigeria.*

**Dr. Ramasamy Harikrishnan**  
*Faculty of Marine Science  
College of Ocean Sciences  
Jeju National University  
Jeju City,  
South Korea.*

**Dr. Kawser Ahmed**  
*Lab. of Ecology, Environment and Climate Change  
Department of Fisheries  
University of Dhaka  
Dhaka,  
Bangladesh.*

**Dr. Maxwell Barson**  
*Biological Sciences Department  
University of Zimbabwe  
Harare,  
Zimbabwe.*

**Dr. Christopher Marlowe Arandela Caipang**  
*Faculty of Biosciences and Aquaculture  
Bodø University College  
Bodø,  
Norway.*

**Dr. William S. Davidson**  
*Department of Molecular Biology and Biochemistry  
Simon Fraser University  
Burnaby,  
Canada.*

**Dr. Babak Ghaednia**  
*Iran Shrimp Research Center (ISRC)  
Bushehr,  
Iran.*

**Prof. Ratha Braja Kishore**  
*Department of Zoology  
Biochemical Adaptation Laboratory  
Banaras Hindu University  
Varanasi,  
India.*

**Prof B. Sharma**  
*Department of Biochemistry  
Centre for Biotechnology  
University of Allahabad  
Allahabad,  
India.*

**Dr. Sebastián Villasante**  
*Fisheries Economics and Natural Resources Research Unit  
University of Santiago de Compostela  
A Coruña,  
Spain.*

**Dr. Mohamed Hamed Yassien**  
*National Institute of Oceanography and Fisheries  
Suez,  
Egypt.*

**Dr. Riaz Ahmad**  
*Department of Zoology  
Aligarh Muslim University  
Aligarh,  
India.*

**Dr. SM Nurul Amin**  
*Lab. Marine Science & Aquaculture  
Institute of Bioscience  
Universiti Putra Malaysia  
Selangor,  
Malaysia.*

**Dr. Melcher Andreas**  
*University of Natural Resources and Applied Life Sciences  
Vienna,  
Austria.*

**Dr. Ayotunde Ezekiel Olatunji**  
*Department of Fisheries and Aquatic Science  
Cross River University of Technology  
Calabar,  
Nigeria.*

**Dr. Ramachandra Bhatta**  
*Animal and Fisheries Sciences University  
College of Fisheries  
Kankanady,  
India.*

**Dr. Subha Ganguly**  
*Department of Fish Processing Technology  
Faculty of Fishery Sciences  
West Bengal University of Animal and Fishery Sciences  
Chakgaria,  
India.*

**Dr. Yusuf Bozkurt**  
*Iskenderun Technical University  
Faculty of Marine Sciences and Technology  
Iskenderun Hatay,  
Turkey.*

**Prof. Rina Chakrabarti**  
*Aqua Research Lab  
Department of Zoology  
University of Delhi  
Delhi,  
India.*



**Prof. Nihar Rajan Chattopadhyay**  
*Faculty of Fishery Sciences*  
*University of Animal and Fishery Sciences*  
*Chakgaria,*  
*India.*

**Dr. Sullip Kumar Majhi**  
*Division of Fisheries*  
*Indian Council of Agricultural Research Complex for Neh*  
*Region*  
*Meghalaya,*  
*India.*

**Dr. Hooman Rahmati-Holasoo**  
*Department of Aquatic Animal Health*  
*Faculty of Veterinary Medicine*  
*University of Tehran*  
*Tehran,*  
*Iran.*

**Dr. Liberty Napilan-Espectato**  
*Institute of Fisheries Policy and Development Studies*  
*College of Fisheries and Ocean Sciences*  
*Iloilo,*  
*Philippines.*

**Dr. Zeinab Abdel-Baki El-Greisy**  
*National Institute of Oceanography and Fisheries (NIOF),*  
*KayetBey,*  
*Egypt.*

**Dr. Thanitha Thapanand-Chaidee**  
*Department of Fishery Biology*  
*Faculty of Fisheries*  
*Kasetsart University*  
*Chatujak,*  
*Thailand.*

**Dr. Gabriel Ugwemorubong Ujagwung**  
*Dept. of Fisheries and Aquatic Environment*  
*Faculty of Agriculture*  
*Rivers State University of Science and Technology*  
*Port Harcourt,*  
*Nigeria.*

**Dr. Margherita Gloria**  
*School of Agriculture*  
*Food Science and Veterinary Medicine*  
*University College Dublin*  
*Belfield,*  
*Ireland.*

**Dr. Derya Guroy**  
*Canakkale Onsekiz Mart University*  
*Faculty of Fisheries*  
*Aquaculture Department*  
*Canakkale,*  
*Turkey.*

**Dr. Daniela Giannetto**  
*Department of Biology*  
*Faculty of Sciences*  
*Muğla Sıtkı Koçman University*  
*Turkey.*

**Dr. Terngu Akaahan**  
*Biological Sciences Department*  
*Federal University of Agriculture Makurdi*  
*Nigeria.*

**ARTICLE**

**Physical losses of fish along the value chain in Zambia: A case study of Barotse Floodplain**

**98**

Kefi A. S., Cole S. M., Kaminski A. M., Ward A. and Mkandawire N. L.

*Full Length Research Paper*

## Physical losses of fish along the value chain in Zambia: A case study of Barotse Floodplain

Kefi A. S.<sup>1\*</sup>, Cole S. M.<sup>2</sup>, Kaminski A. M.<sup>2</sup>, Ward A.<sup>3</sup> and Mkandawire N. L.<sup>4</sup>

<sup>1</sup>Department of Fisheries, P. O. Box 350100, Chilanga, Zambia.

<sup>2</sup>World Fish, Unit D, Suite No.2, Counting House Square, Thabo Mbeki Road, Lusaka, Zambia.

<sup>3</sup>Independent Consultant, 1 Coes Cottages, Beaney's Lane, Shottenden, Kent, CT4 8JA, United Kingdom.

<sup>4</sup>Department of Food Science and Technology University of Zambia, Lusaka, Zambia.

Received 2 July, 2017; Accepted 29 August, 2017

The study was conducted in three districts of Western Province of Zambia namely; Mongu, Nalolo and Senanga that lie along the Zambezi River in the Barotse Floodplain. The study employed two main data tools aimed at understanding the extent of post-harvest losses in the fishing, processing and trading nodes in the fish value chain study sites. The tools were the Exploratory Fish Loss Assessment Method (EFLAM) study and a Quantitative Loss Assessment Method (QLAM) survey. The study found that physical fish losses occur at three nodes in the value chain and differ significantly ( $P < 0.05$ ) between the nodes. On average, the processors lose the largest volume of fish (7.42%) followed by the fish traders (2.9%). The fishers experience the least physical losses at 2% although, this is not significantly different ( $P > 0.05$ ) from the fish lost at trading node. The major cause of physical loss was found to be breakages at processing and trading nodes. There is need to introduce improved processing technologies that can reduce breakages. Furthermore, economic and nutrient losses should also be profiled to fully understand the total losses that occur within the Barotse Floodplain fish value chain.

**Key words:** Fish, physical losses, fishing, processing, trading.

### INTRODUCTION

The fisheries subsector that includes both aquaculture and capture fisheries is important in the economy of Zambia as it supports around one million people both directly and indirectly (Central Statistics Office, 2014). Furthermore, fish contributes more than 53.4% of animal protein in the diets of Zambians (FAO, 2012) and is recognised for its nutritional quality and health benefits that have been documented elsewhere (Kawarazuka, 2010; Beveridge et al., 2013; Béné et al., 2015). Although

global fish consumption per capita has grown from 9.9 kg in the 1960s to a record high of 20 kg in 2014 (FAO, 2016), the scenario in Zambia has remained subdued at 9.1 kg lower than in 1970s (Department of Fisheries, 2017). However, the fish supply per capita statistics for Zambia are calculated from the catch data from capture fisheries and production data from aquaculture, as well as including the imports. Post-harvest losses (PHL) occur along the fish value chain (FVC), yet are not accounted

\*Corresponding author. E-mail: [askefi@yahoo.com](mailto:askefi@yahoo.com). Tel: +260 969426244; +2609797255620.

for in these statistics. Fish is a highly perishable commodity and is, therefore, likely that fish supply per capita is even lower than what is recorded. Fish losses affect negatively on the amount of fish consumed by the people and income generated by the actors along the fish value chain. The widening gap between supply and demand due to annual human population growth of around 3% in Zambia means that there is every need to reduce the losses that occur during harvest through trading.

According to de Lucia and Assennato (1994), PHL refers to measurable quantitative and qualitative food loss in the post-harvest food system. Although fish losses along the value chain have not been determined in Zambia, they result in annual economic losses of \$2 to \$5 billion in sub-Saharan Africa (Béné, 2011). In Africa it is estimated that about 10% of the total weight of the world's fish catch is lost with small scale fish processing losing as much as 40% (FAO, 1984; Moes, 1980). It is reported, however, that around 70% of the total loss is due to quality losses and that in sub-Saharan Africa only around 5% of the total small-scale production is totally wasted/discarded (FAO, 1996; Akande and Diei-Ouadi, 2010).

Fish losses are categorised into two main types: the quality and physical losses. Quality loss results in the fish value chain actors sell their fish for a low price due to spoilage or damage. These fish can also be sold for a lower price due to the forces of the market (if the demand becomes lower than supply) (Diei-Ouadi and Mgawe, 2011). On the other hand physical losses are those that result in fish becoming unsaleable due to severe insect infestation and breakages or poor post-harvest handling. In this case, no income is gained and no fish is consumed since the fish is physically lost.

The Barotse floodplain in the Western Province of Zambia is the second largest floodplain in Zambia, after the Bangweulu complex. It helps meet important resource needs for its human inhabitants including their cultural and socio-economic needs ([https://en.wikipedia.org/wiki/Barotse\\_Floodplain](https://en.wikipedia.org/wiki/Barotse_Floodplain)). The floodplain starts from the Zambezi River's confluence with the Kabompo and Lungwebungu Rivers in the north, to a point about 230 km south, above the Ngonye falls and south of Senanga. The Zambezi River that runs through the floodplain is the largest River in the country and is important in the communication and transportation of goods and services in the floodplain. In fact the country's name Zambia is derived from the Zambezi River. The flood plain is an important fishery harbouring 80 fish species and employing an estimated 4,350 fishers (Department of Fisheries, Zambia, 2012). According to the Department of Fisheries (DOF) (2016), a total of 7, 714 metric tons of fish were caught from the upper Zambezi fishery.

Although, the catch per unit effort has declined in the last three decades from 3.1 to 1.2 tons/fisher due to the

increased number of fishers (Kefi and Mofya, 2015), the fisheries is still the main economic stay and source of livelihood for many dwellers along the floodplain. Subsistence and commercial fishing, and angling tourism, are major activities throughout the system, which includes several large lakes and numerous, highly productive floodplains (Tweddle, 2010; Tweddle et al., 2015). Fish provides an important source of income and protein. Consumption of fish by local residents in Western Province is estimated at about five times the national average (Baidu-Forson et al., 2014). The quantities of the fish lost between harvesting and consumption are thought to occur, but exact details remain unknown and post-harvest losses are not recognised in government databases or in national policy. This weakens the role that fish plays in the food and nutrition security of people in the region and country on the whole (Longley and Kruijssen, 2014).

The scope of this paper is to describe a study that quantitatively assessed the amounts of the physical fish post – harvest losses incurred by the actors (fishers, processors and traders) along the fish value chain (FVC), in order to shed light on this important component that should be addressed in fisheries management and policy debates. In the context of this paper FVC is defined as a full range of activities that the fish undergoes through harvest to consumption with the major nodes being fishing, processing and trading. In addition, an attempt has been made to determine the causes of the physical losses at each of the FVC nodes (excluding consumption) in the Barotse floodplain of Western Province, Zambia.

## MATERIALS AND METHODS

### Study areas


The study was conducted in three districts of Western Province; Mongu, Nalolo and Senanga that lie along the Zambezi River in the Barotse Floodplain (Figure 1). Six fishing camps were selected purposively in a multistage sampling technique. A fishing camp is a seasonal, temporary site where women and men conduct fishing, processing and trading activities as the water levels recede between May and October every year.

In Mongu District, Liyoyeli, Nebubela and Mukakani were the fishing camps selected while Marana was the only site in Nalolo district. Tangatanga and Matula were the sites chosen in Senanga District. The sites were selected based on the following criteria: number of fishers / processors / traders distance to the market and the availability of fish based on the catch assessment surveys conducted by Department of Fisheries (DOF). However, the main markets within the three districts were also sites for fish traders.

### Data tools and sampling

This study employed two main data tools aimed at understanding the fishing, fish processing and fish trading in the study districts. These were Exploratory Fish Loss Assessment Method (EFLAM) and Quantitative Loss Assessment Method (QLAM) tools. Prior to





**Figure 1.** Map of Western Province showing the districts where the study was conducted.

Source: [https://en.wikipedia.org/wiki/Western\\_Province,\\_Zambia](https://en.wikipedia.org/wiki/Western_Province,_Zambia).

data collection, ethical clearance was sought from the University of Zambia, Humanities and Social Sciences Research Ethics Board (HSSREC) and consent was sought from respondents by trained enumerators. These enumerators (Department of Fisheries extension officers) were trained in data collection that included pre-testing the survey in Mongu District.

In order to calculate the size of the sample for the fishers, processors and traders, the methods employed by Cochran (1963) were used:

$$n = 1.96^2 \cdot (0.5) \cdot (1 - 0.5) / 0.05^2 = 385 \text{ fishers, processors and traders.}$$

Where  $n = Z^2 pq / e^2$  the sample size,  $z$  is the abscissa of the normal curve that cuts off an area at  $\alpha$  at the tails ( $1 - \alpha$  equals the desired confidence level e.g. 95%, therefore the precision is  $\pm 0.05\%$ ),  $e$  is the desired level of precision,  $p$  is the estimated proportion and  $q$  is  $1-p$ . We assumed that there were a lot of fishers, processors and traders and we took a highest variability (50%) (Israel, 2013).

However, a total of 207 fishers, 110 processors and 110 traders (total 427) were sampled with 32 people classified as being both fishers and processors. Interviews and FGDs from the EFLAM were spread over the period of three months (April to June 2015) before the QLAM across the three Districts was administered.

#### **Exploratory fish loss assessment method and quantitative loss assessment method tools**

In order to estimate the biomass losses, understand fish losses and people's experiences and attitudes around fish losses, an Exploratory Fish Loss Assessment Method (EFLAM) study was undertaken. EFLAM is a qualitative research tool that includes

semi-structured interviews with individuals and Focus Group Discussions (FGDs) with groups of people.

Formally known as Informal Fish Loss Assessment Method (IFLAM), it helps to develop a qualitative understanding of losses and provides indicative quantitative data on the post-harvest fish losses (Diei-Ouadi and Mgawe, 2011). A key activity was a stakeholder meeting in Mongu (mainly involving the fishers, processors, traders, government workers and traditional leaders) which was held in February 2015. The fishers, processors and traders attending were drawn from all the selected fishing camps. The EFLAM was used to generate qualitative and indicative quantitative PHL data, that was used to develop the quantitative tool the Quantitative Loss Assessment Method (QLAM) (Annex), based on the methods described by Diei-Ouadi and Mgawe (2011). The QLAM was administered to fishers, processors and traders.

The QLAM is a formal questionnaire survey approach to quantitatively estimate the quantity of losses, type of losses incurred, reasons for losses and the variables that create loss, such as the type of fishing gear used or fish processing methods, etc. According to Akande and Diei-Ouadi (2010), QLAM is used to provide quantitative data on a wide range of issues that enables validation of data over a wide geographical area. In order to capture the effect of seasonality, the QLAM was administered for the entire fishing season from July to November 2015. Prior to data collection, the enumerators were trained in data collection and the questionnaires were pre tested in Mongu district to check any ambiguities that contained in the tools.

#### **Statistical analysis**

The physical loss of fish was calculated from the QLAM by subtracting the total amount of fish that was discarded before consignment to move onto the next node. Prior to all the measuring

**Table 1.** Percent of physical losses (Mean±SD) of fish at fishing node (N = 207).

District	Fish loss (%)	Minimum (%)	Maximum (%)
Mongu	0.81±4.69 <sup>a</sup>	0	33.3
Nalolo	1.93±3.82 <sup>ab</sup>	0	12.5
Senanga	3.67±10.88 <sup>bc</sup>	0	50
Average	2.03±7.70	0	50

Values with superscripts in a column are significantly different ( $P < 0.05$ ).

**Table 2.** Percent of physical losses (Mean±SD) of fish at processing node (N = 110).

District	Fish loss (%)	Minimum (%)	Maximum (%)
Mongu	13.81±19.45 <sup>b</sup>	0	50.00
Nalolo	12.16±10.31 <sup>b</sup>	0	33.33
Senanga	5.11±10.68 <sup>a</sup>	0	50.00
Average	7.42±12.73	0	50.00

Values with superscripts in a column are significantly different ( $P < 0.05$ ).

units, fish lost were standardised by transforming all the local measuring units into metric units. This was done by weighing the local units in triplicates and taking the average weight for each unit.

One way Analysis of Variance (ANOVA) was used to test the percent that physical fish was loss at every node. The Duncan Multiple Range Test (DMRT) was used to separate the means if significant ( $P < 0.05$ ).

## RESULTS

### Physical losses of fish at fishing node along the value chain

There were significant differences ( $P < 0.05$ ) in the physical fish losses incurred at the fishing node among the surveyed districts. Senanga fishers recorded the highest amount of fish loss with Mongu recording the lowest physical losses. On average, losses were 2.03±7.70% at the fishing node with a lot of variation as evidenced by the large standard deviation. Some fishers lose up to half of their catch (50%) as observed in Senanga District (Table 1).

### Causes of fish physical losses at fishing node along the value chain

Due to a small number of fishers (N = 23) who reported to have lost fish, no statistical analysis was conducted on the causes of the losses. However, the reasons for the losses mentioned were spoilage, squashing due to poor handling and pest infestation.

### Physical losses of fish at processing node along the value chain


The total physical fish losses that were experienced at the processing level were averaged 7.42±12.73%. Senanga district processors experience significantly ( $P < 0.05$ ) lower physical losses than the processors in the other two districts under the study. Although Mongu district processors had the highest percentage of physical losses of fish, they did not differ ( $P > 0.05$ ) with the losses experienced by Nalolo district (Table 2).

### Causes of fish physical losses at processing node along the value chain

Three major causes of physical fish losses were identified by the processors. The study revealed that breakages contributed significantly ( $P < 0.05$ ) to the fish losses. Of the sample, 72.2% of the respondents indicated breakages as the main reason for loss whilst 19.4% attributed the losses to spoilage, although this has no significant different ( $P > 0.05$ ) from pest infestation (8.3%)(Figure 2).

### Physical losses of fish at trading node along the value chain

On average, the physical fish losses at the trading node amount to 2.87±8.26%. There were significant differences ( $P < 0.05$ ) in the physical fish losses at the



**Figure 2.** Causes of fish losses at processing along the value chain (values with superscripts at the top of the bars are significantly different ( $P < 0.05$ )).

**Table 3.** Percent of physical losses (Mean $\pm$ SD) of fish at trading node (N = 110).

District	Fish loss (%)	Minimum (%)	Maximum (%)
Mongu	4.13 $\pm$ 9.19 <sup>b</sup>	0	33.33
Nalolo	8.33 $\pm$ 8.33 <sup>b</sup>	0	16.67
Senanga	1.55 $\pm$ 7.16 <sup>a</sup>	0	40.00
Average	2.87 $\pm$ 8.26	0	40.00

Values with superscripts in a column are significant different ( $P < 0.05$ ).

trading node among the three districts. Nalolo experienced the highest fish losses followed by Mongu. The smallest losses were found in Senanga estimated at 1.55 $\pm$ 7.16% with a high variation in responses (Table 3).

#### **Causes of fish physical losses at trading node along the value chain**

The survey revealed that there were only two causes of fish losses during trading and these were breakages and spoilage. Although there were no significant differences ( $P > 0.05$ ) between the two causes of fish loss, the former (58.2%) contributed more to losses than the latter (41.8%).

#### **DISCUSSION**

The study of fish PHL along the value chain is the first of its kind in the Zambian fisheries subsector. And the results of this study are compared with previous fish loss

assessments conducted in other parts of the world. There is a limited literature, on fish PHL globally and studies often combine with the physical and quality losses. This study revealed that fish physical losses occur throughout the value chain in the Barotse Floodplain, since every node recorded a positive loss. The estimated cumulative PHL of fish was 12.32%, with processors incurring the highest percentage loss of 7.42 $\pm$ 12.73%.

The total losses translated to around six million United States dollars based on the fish catches was reported by Department of Fisheries (DOF) (2017). Fishers' consignments of fish experienced the lowest percentage of losses on average, and thus have the least liability in the value chain. Percentages of fish losses reported in other countries vary. According to Béné et al. (2015) the total estimated PHL of fish is around 20% in small-scale fisheries, to FAO (1996), fish losses in sub-Saharan Africa may be around 5 percent of the total artisanal production, while for the West African Region others indicate losses which occur between 10 and 20% (McConnery, 1994). Akande and Diei-Ouadi (2010), who studied PHL in five countries in sub-Saharan Africa,

estimated PHL of fish at 5%, although physical losses in some countries were as high as 40%.

Therefore, the cumulative percentage loss found in the districts along the floodplain fishery is lower than the global average but within what has been reported in the region. It should be noted that discarding fish in artisanal fisheries is not common, where post-harvest methods typically utilize a wide range of species, including those that are of low commercial value (Akande and Diei-Ouadi, 2010). Nonetheless, such physical losses result in lost incomes for value chain actors and ultimately lost fish for consumers within and outside the province, including lost opportunities acquire nutritional benefits from these fish.

The major causes of fish losses in this setting were found to be node dependent. Breakages were found to be the most significant reason why physical losses occur at the processing and trading nodes. This can be attributed to the poor methods employed during processing and likely poor storage and handling in route or during trading. There is need to introduce improved processing, storage, and handling methods and technologies that can reduce breakages during processing, storage, transportation and retailing. The introduction of the cold chain within the region which provides primary preservation is key to reduce the losses incurred by the value chain actors especially those that deal in fresh fish.

Furthermore, economic and nutrient losses should be profiled to understand the total losses that occur within the value chain in the Barotse Floodplain. There is need for further studies of the fish losses according to fish forms or products in the fish value chain.

## CONFLICTS OF INTERESTS

The authors have not declared any conflict of interests.

## ACKNOWLEDGMENTS

We are indebted to the International Development Research Center (IDRC) and the Australian Centre for International Agriculture Research (ACIAR) for funding the project “Improving Livelihood Security and Gender Relations in Rural Zambia and Malawi through Post-Harvest Fish Value Chain Innovations and Social Change Interventions”. It is through the project that this study was conducted. Many thanks go to fishers, processors and traders who accepted to be interviewed.

## REFERENCES

Akande GR, Diei-Ouadi Y (2010). Post-harvest losses in small-scale fisheries. Case studies in five sub-Saharan African countries. FAO Fisheries and Aquaculture Technical Paper No. 550. Rome, Italy.

Baidu-Forson JJ, Phiri N, Ngu'ni D, Mulele S, Simainga S, Situmo J, Ndiyoi M, Wahl C, Gambone F, Mulanda A, Syatwinda G (2014).

Assessment of agrobiodiversity resources in the Borotse flood plain, Zambia. CGIAR Research Program on Aquatic Agricultural Systems. Penang, Malaysia. Working Paper: AAS-2014-12.

Béné C (2011). CAADP and Fisheries Policy in Africa: are we aiming for the right reform? Policy Brief 40, Future Agriculture. Available at: <http://www.future-agricultures.org/>

Béné C, Barange M, Subasinghe R, Pinstrup-Andersen P, Merino G, Hemre GI, Williams M (2015). Feeding 9 billion by 2050 – Putting fish back on the menu. *Food Secur.* 7(2):261-274.

Beveridge MC, Thilsted SH, Phillips MJ, Metian M, Troell M, Hall SJ (2013). Meeting the food and nutrition needs of the poor: The role of fish and the opportunities and challenges emerging from the rise of aquaculture. *J. Fish Biol.* 83:1067-1084.

Central Statistics Office (2014). Census of Population and Housing report, CSO, Lusaka, Zambia.

Cochran WG (1963). *Sampling Techniques*, 2nd Ed., New York: John Wiley and Sons, Inc.

De Lucia M, Assennato D (1994). Agricultural engineering in development: post-harvest operations and management of foodgrains. L'après-recolte des grains: organisation et techniques. La ingegneria agraria en el desarrollo: manejo y tratamiento de granos poscosecha; organización y técnicas (No. FAO ASB-93). FAO, Roma (Italia).

Diei-Ouadi Y, Mgawe YI (2011). Post-harvest fish loss assessment in small-scale fisheries. Food and Agriculture Organization of The United Nations. FAO Fisheries and aquaculture technical Paper, Rome, Italy.

Department of Fisheries (2016). 2014 Annual Report for Department of Fisheries. Department of Fisheries, Chilanga, Zambia.

Department of Fisheries (2017). 2016 Annual Report for Department of Fisheries. Department of Fisheries, Chilanga, Zambia.

Department of Fisheries (2012). 2011 Annual Report for Department of Fisheries. Department of Fisheries, Chilanga, Zambia.

Food and Agriculture Organization (FAO) (1984). Prévention des pertes de poisson traité. FAO Doc. Techn. Pêches 219:84.

Food and Agriculture Organization (FAO) (1996). Rome Declaration on World Food Security and the World Food Summit Plan of Action. (Available at: <http://www.fao.org/docrep/003/w3613e/w3613e00.HTM>).

Food and Agriculture Organization (FAO) (2012). The State of World Fisheries and Aquaculture, Rome, Italy.

Food and Agriculture Organization (FAO) (2016). The State of World Fisheries and Aquaculture, Rome, Italy. [https://en.wikipedia.org/wiki/Barotse\\_Floodplain](https://en.wikipedia.org/wiki/Barotse_Floodplain). Retrieved on 18<sup>th</sup> August 2017 [https://en.wikipedia.org/wiki/Western\\_Province](https://en.wikipedia.org/wiki/Western_Province), Zambia. Retrieved on 18<sup>th</sup> August 2017

Israel GD (2013). Determining Sample Size. Univ Florida IFAS Ext. 2013, 1-5. Available at: <http://edis.ifas.ufl.edu>.

Longley C, Kruijssen F (2014). Fish Value Chains in the Barotse, Western Province Zambia. Program Technical Report, WorldFish, Penang.

Kawarazuka N (2010). The contribution of fish intake, aquaculture, and small-scale fisheries to improving nutrition: A literature review. The WorldFish Center Working Paper No.2106. The WorldFish Center, Penang, Malaysia.

Kefi AS, Mofya RM (2015). The fisheries sector in Zambia: status, management, and challenges. Technical Paper No. 3. IAPRI, Lusaka, Zambia.

McConnery KS (1994). Tackling the problem of post-harvest losses in the artisanal fishing. *The Courier* 147:95-98.

Moes JE (1980). Reduction of spoilage of fish caught in Lake Chad. Project report NIR/74/001/F/01/12.

Tweddle D (2010). Overview of the Zambezi River System: Its history, fish fauna, fisheries, and conservation. *Aquat Ecosyst Health Manag.* 13:224-240.

Tweddle D, Cowx IG, Peel RA, Weyl OLF (2015). Challenges in fisheries management in the Zambezi, one of the great rivers of Africa. *Fish. Manag. Ecol.* 22:99-111.

**Annex: QLAM Questionnaire**

1. Date of interview...../...../.....  
 Enumerator's Name.....
2. CultiAF site: [ ]  
 Mukakani=1 Nebubela=2 Liyoyelo=3 Marana=4 Matula=5 Tangatanga=6
3. Name of fishing camp :.....
4. Name of respondent :..... 5. Gender: [ ] Female = 0 Male = 1
6. Contact # :.....(optional)
7. Marital status : [ ] Single=1 Married=2 Polygynous=3  
 Cohabiting=4 Divorced=5 Widowed=6
8. Education – years of schooling: .....years or grade
9. Where is your permanent place of residence? .....
10. Did you fish, process, and/or trade fish in last 7 days? YES / NO  
 If yes continue; if no, finish the interview and thank the respondent.
11. What fish-related activities did you engage in during the last 7 days? (underline or circle)

- |   |   |  |
|---|---|--|
| <p><b>1. Fishing</b></p> <p>a. Multifilament gill net</p> <p>b. Monofilament gill net</p> <p>c. Trap</p> <p>d. Hook</p> <p>e. Spear &amp; light</p> <p>f. Seine net</p> <p>g. Other:.....</p> | <p><b>2. Processing</b></p> <p>a. smoking</p> <p>b. sun drying</p> <p>c. frying</p> <p>d. salting</p> <p>e. Other:.....</p> | <p><b>3. Trading (buying and selling fish)</b></p> <p>a. fresh fish – no ice</p> <p>b. fresh fish – with ice</p> <p>c. sun dried fish</p> <p>d. smoked fish</p> <p>e. fried fish</p> <p>f. salted fish</p> <p>g. Other:.....</p> |
|---|---|--|

**Instructions for the enumerator**

*If the respondent has done fishing in the past 7 days, please complete Section B.  
 If the respondent has done processing in the past 7 days, please complete Section C.  
 If the respondent has done processing and trading, please complete Section C.  
 If the respondent has done trading (but not processing), please complete Section D.*

*All questions in Sections B, C and D refer to the most recent catch or 'batch' or 'consignment' of fish that was caught, processed, or bought and then sold. In addition, all responses within each section should refer to a single species of fish (any species that is most predominant among the different species that might have been caught, processed or bought). All responses within each section should also refer to a single product type (e.g. fresh, smoked, sun-dried). It is necessary to be very clear with the respondent about which species and product type is being referred to from the start of each section to avoid confusion.*



**SECTION B: FISHING.**

***This section should only be completed by those who have done fishing in the past 7 days.***

B.1 What was the most predominant species of fish caught the **last time** you went fishing?

Species: [ ] *Lipapati=1 Nembele=2 Ndombe =3 Other=4 (specify).....*

B.2. Approximately how much fish of this species did you catch the **last time**?

Unit:..... Number of units:..... [*Unit might be pieces, bafa , plate, etc*]

B.3. Was any of the fish of this species lost from the time of catching to the time of selling (if sold fresh) or processing (if processed)? YES / NO [*If Yes, proceed to Qu. B.4. If no, go to Qu. B.6.*]

B.4. How much fish of this species did you lose **last time** you went fishing?

Unit:.....Number of units:.....

B.5. How did you lose this fish (physical loss)?

Main reason:.....

Other reasons\*:.....

B.6. Was the fish of this species that you caught the last time processed? YES / NO  
 [*If yes, please go to Section C. If no, please continue with the remaining questions in Section B.*]

B.7. If this fish was sold fresh, where (place) did you sell this fish **last time**? .....

B.8. What was the highest price you received for this fish **last time**? ZK.....Unit.....

B.9. How much fish of this species did you sell for a high price last time?

Unit..... Number of units.....

B.10. What was the lowest price you received for this fish **last time**? ZK.....Unit.....

B.11. How much fish of this species did you sell for a low price **last time**?

Unit.....Number of units.....

B.12. Why did you sell fish for a low price **last time**?

Main reason:.....

Other reasons\*:.....

[*Include issues related to transport type, time, packaging etc.*]

\* *Remember to probe for non technical and gender related causes as well*

**SECTION C. PROCESSING. *This section should only be completed by those who have done processing in the past 7 days, or those who have done processing and trading (see Qu. 11).***

C.1 What was the most predominant species of fish processed the **last time** you were processing?

Species: [ ] *Lipapati=1 Nembele=2 Ndombe =3 Other=4 (specify).....*

C.2. What was the product that you processed of this species: [ ]  
*Fresh=1 Smoked=2 Sundried=3 Salted=4 Other=5 (specify).....*

C.3. Approximately how much fish of this species did you process in this way the **last time** you processed and sold processed fish?

Unit:..... Number of units:..... [Unit might be pieces, bafa , plate, etc]

C.4 Was any of the fish of this species lost the last time you were processing, from the time of processing up to the time of sale? YES / NO. If Yes, proceed to Qu. C.5. If no, go to Question C.7.

C.5. How much fish of this species did you lose the **last time** you processed and sold fish?

Unit:..... Number of units:.....

C.6. How did you lose this fish (physical loss)?

Main reason:.....

Other reasons\*:.....

C.7. Where (place) did you sell the processed fish of this species **last time**? .....

C.8. What was the highest price you received for this fish **last time**? ZK.....Unit.....

C.9. How much fish of this species did you sell for a high price last time?

Unit..... Number of units.....

C.10. What was the lowest price you received for this fish **last time**? ZK.....Unit.....

C.11. How much fish of this species did you sell for a low price **last time**?

Unit..... Number of units.....

C.12. Why did you sell fish for a low price **last time**?

Main reason:.....

Other reasons\*:.....

[Include issues related to transport type, time, packaging etc.]

\* Remember to probe for non technical and gender related causes as well

**SECTION D. TRADING. This section should only be completed by those who have traded but not processed fish in the past 7 days (see Qu. 11).**

D.1 What was the most predominant species of fish purchased the **last time** you bought fish?

Species: [ ] Lipapati=1 Nembele=2 Ndombe =3 Other=4 (specify).....

D.2. In what form was the fish when you bought it?

Form: [ ] Fresh=1 Smoked=2 Sundried=3 Salted=4 Other=5 (specify).....

D.3. Approximately how much fish of this species and form did you purchase the **last** time you bought fish for trading purposes?

Unit..... No. units..... NB: Unit might be pieces, bafa , meda, plate, etc.

D.4 Was any fish of this species lost from the time of purchase up to the time of sale? YES / NO.

[If yes, proceed to Question D.5. If no, go to Question D.7.].

D.5. How much fish of this species did you lose **last time**?

Unit:..... Number of units:.....

D.6. How did you lose fish (physical loss)?

Main reason:.....

Other reasons\*:.....

D.7. Where (place) did you sell the fish of this species **last time**? .....

D.8. What was the highest price you received for this fish **last time**? ZK.....Unit.....

D.9. How much fish of this species did you sell for a high price last time?

Unit..... Number of units.....

D.10. What was the lowest price you received for this fish **last time**? ZK.....Unit.....

D.11. How much fish of this species did you sell for a low price **last time**?

Unit..... Number of units.....

D.12. Why did you sell fish for a low price **last time**?


Main reason:.....

Other reasons\*:.....

[Include issues related to transport type, time, packaging etc.]

\* Remember to probe for non technical and gender related causes as well

**THANK THE RESPONDENT FOR THEIR TIME**



# International Journal of Fisheries and Aquaculture

## Related Journals Published by Academic Journals

- *Journal of Plant Breeding and Crop Science*
- *African Journal of Agricultural Research*
- *Journal of Horticulture and Forestry*
- *International Journal of Livestock Production*
- *International Journal of Fisheries and Aquaculture*
- *Journal of Cereals and Oilseeds*
- *Journal of Soil Science and Environmental Management*
- *Journal of Stored Products and Postharvest Research*

**academicJournals**