

Journal of Plant Breeding and Crop Science

Volume 9 Number 11 November 2017

ISSN 2006-9758



*Academic
Journals*

ABOUT JPBCS

The **Journal of Plant Breeding and Crop Science (JPBCS)** is published monthly (one volume per year) by Academic Journals.

The **Journal of Plant Breeding and Crop Science (JPBCS) (ISSN: 2006-9758)** is an open access journal that provides rapid publication (monthly) of articles in all areas of the subject such as Sustainable use of plant protection products, Agronomic and molecular evaluation of recombinant inbred lines (RILs) of lentil, Pollen behaviour and fertilization impairment in plants, Development of a fast and reliable ozone screening method in rice etc.

The Journal welcomes the submission of manuscripts that meet the general criteria of significance and scientific excellence. Papers will be published shortly after acceptance. All articles published in JPBCS are peer-reviewed.

Contact Us

Editorial Office: jpbcs@academicjournals.org

Help Desk: helpdesk@academicjournals.org

Website: <http://www.academicjournals.org/journal/JPBCS>

Submit manuscript online <http://ms.academicjournals.me/>

Editors

Dr. Munir Aziz Noah Turk *Crop Production*
Department, Faculty of Agriculture
Jordan University of Science & Technology
Irbid, Jordan
E-mail: jpbcs@acadjourn.org
<http://www.academicjournals.org/jpbcs>

Dr. B.Sasikumar
ITEC Expert (Spices Technology) National
Agril.Res.Inst.,
Mon Repos,ECD,Guyana" India

Dr. Abdul Jaleel Cheruth
Stress Physiology Lab, Department of
Botany, Annamalai University, Annamalaiagar -
608
002, Tamilnadu,
PO Box No- 15711, AL-AIN, UAE, India

Dr. S. Paulsamy
Kongunadu Arts and Science College, Coimbatore
- 641 029,
India

Dr. Ivana Maksimovic
Department of Field and Vegetable Crops
Faculty of Agriculture, University of Novi sad,
Serbia

Dr. Aboul-Ata E Aboul-Ata
Plant Virus and Mycoplasma Res. Sec.,
Plant Path. Res. Inst., ARC, PO Box 12619, Giza,
Egypt

Dr. Lusike A. Wasilwa
Kenya Agricultural Research Institute P. O. Box
57811-00200, Nairobi, Kenya

Dr. Neeraj Verma University of California Riverside, CA
92521, USA

Dr. Yongsheng Liu
Research Center for Bio-resource and Eco-
environment
College of Life Science,
Sichuan University, Chengdu 610064, P. R. China

Editorial Board

Dr. Hadia Ahmed Mohamed Moustafa Heikal
Genetic Engineering & Biotechnology Research, Institute
(GEBRI),
Sadat City, Menoufiya University
Egypt

Dr. Nambangia Justin Okolle
Research Entomologist,
African Research Center on Bananas and Plantains
(CARBAP)
Njombe,
Cameroon

Dr. Nihaluddin Mari
Rice Research Institute Dokri, District Larkana, Sindh, Pakistan

Dr. Veronica Sanda Chedea
Department of Chemistry and Biochemistry,
University of Agricultural Sciences and Veterinary Medicine
(USAMV),
Cluj-Napoca, str. Manastur 3-5, 400372 Cluj-Napoca
Romania

Dr. Marku Elda
Tirana University,
Faculty of Natural Sciences, Chemistry Department, Tirana
Albania

Dr. Mershad Zeinalabedini
ABRII Agricultural Biotechnology Research,
Institute of Iran
Iran

Dr. Md. Mainul Hasan
Visiting Fellow (Plant Cell Biotechnology Lab.): 2008-
Present: MU
Department of Agricultural Botany, Faculty of Agriculture,
Patuakhali Science and Technology University (PSTU),
Bangladesh
Thailand

Dr. Amr Farouk Abdelkhalik Moustafa
Rice Research and Training Center, 33717. Sakha. Kafr
El-Shiekh, Egypt

Prof P.B. Kirti
Department of Plant Sciences, University of Hyderabad,
Hyderabad - 500 046,
India

Dr. Abdel Gabar Eltayeb
University of Sudan,
College of Agricultural Studies, Crop Science Department,
P.O. Box 71 Shambat, Khartoum North
Sudan

ARTICLES

Assessment of the inheritance pattern of the novel "female only flower" trait in *Jatropha curcas* L. as potential for hybrid seed production 208
Gongora-Canul, C. C., Aguilera-Cauch, E. A. and Martinez-Sebastián, G.

Estimates of combining ability for resistance to pod shattering in soybean (*Glycine max* (L.) Merrill) genotypes 217
F. A. Umar, M. S. Mohammed, M. Oyekunle, I. S. Usman, M. N. Ishaq and S. N. Dachi

Full Length Research Paper

Assessment of the inheritance pattern of the novel "female only flower" trait in *Jatropha curcas* L. as potential for hybrid seed production

Gongora-Canul, C. C.^{1*}, Aguilera-Cauchich, E. A.² and Martinez-Sebastián, G.²

¹Universidad Anáhuac Mayab, División de Ingeniería y Ciencias Exactas. Carretera Mérida-Progreso Km 15.5, interior Km 2 Carretera a Chablekal, Oncá, Cordemex, CP 97310, Mérida, Yucatán, México.

²Agroindustria Alternativa del Sureste - Grupo Lodemo. Calle 60 diagonal, Num. 490, CP: 97300 Parque Industrial Yucatán, México.

Received 8 October, 2017; Accepted 17 November, 2017

Jatropha curcas L. (JLC) is a perennial shrub, originating in Central and South America and widely distributed in the tropical regions of the world. It is important in the bioenergy industries due to the characteristics of its oil which can be transformed into high quality biofuel for the substitution of diesel and jet fuel. The native gene, female only flower (FOF), is a desirable trait in the production of hybrid seed of *J. curcas* L. In this study, inbreeding was induced in eight accessions from wild, S₁, S₂ to S₃ endogamy level, and two wild accessions naturally expressing the FOF trait were used as females to cross with male accessions (FM) in order to estimate segregation. The Chi-Square test was used to determine if the FOF trait follows a Mendelian inheritance. Results strongly support the assumption that FOF trait follows a recessive monogenic Mendelian inheritance pattern since, in the inbreeding process of the trait, it segregated a phenotypic frequency of 3:1 (FM:FOF) and in crosses, a frequency of 1:1 (FM:FOF) was segregated. This study potentially contributes to furthering the commercial production of JLC hybrid seed, given that the plants with FOF trait can be used as female parents to be naturally pollinated in isolated plots, thereby reducing labor and cost.

Key words: Floral biology, heredity, inbreeding, *Jatropha curcas*, native gene.

INTRODUCTION

The physic nut *Jatropha curcas* L. (JCL) has gained attention in recent years as a feedstock source for biodiesel and biojet fuel, as its seed contains 30 to 40% oil (King et al., 2009). However, the crop is still being developed especially the production of commercial hybrids. Commercial hybrid seed production results from

crossing two parental inbreds, homozygous or heterozygous (Johnson, 1966; McRobert et al., 2014). This raises the importance of understanding the flowering biology of JLC. The mating system of JLC is complex; however, it is considered to be a monoecious species, also presenting atypical inflorescences (gynoecious,

*Corresponding author. E-mail: carlos.gongora@agroindustria.com.mx.

androecious, andromonoecious, androgynomonocious) (Adriano-Anaya et al., 2016), producing plants with female and male flowers: FM, plants with female only flowers: FOF and occasionally plants with hermaphrodite flowers: HF (Heller, 1996), or even producing seeds asexually by apomixes (Bressan et al., 2013; Ahton and Quenum, 2012). The dioecious type (gynoecious plants) with the female only flower trait (FOF) has been naturally observed in JLC without any breeding process (Heller, 1996; Pecina-Quintero et al., 2011; Adriano-Anaya et al., 2016; Ovando-Medina et al., 2013) and can potentially be used as female parent for hybrid seed production. The production of hybrid seed is one of the most expensive and labor-intensive processes in the seed industry (McRobert et al., 2014). The pollination for a cross, especially when homozygous inbreds are used, needs to be controlled as this guarantees the desirable combination of inbred traits and avoids pollen contamination on the female inbred parent by self-pollination or with pollen other than that of the desired male contributed by insects (Vaknin et al., 2012), this results in a uniform F1 population offspring, and expresses heterosis for yield (Birchler et al., 2010; Perez-Prat and van Lookeren, 2002). Pollination should be efficient and economical and depends on factors such as the synchronization of the flowering between the female and male parents in an interplanting row plot (Haque et al., 2012) and contamination of the female parent with foreign pollen (Masuda et al., 2010). Some strategies to reduce female pollen contamination include the emasculation by detaselling of the female parent, as in corn (Stevens et al., 2004), the usage of the cytoplasmic male sterility trait (CMS) (Yamagishi and Bhat, 2014) or the transgenic genes based on nuclear-encoded male sterility as the seed production technology (SPT) (Wu et al., 2016).

Although there are studies on JLC flowering biology (Heller, 1996; Pecina-Quintero et al., 2011; Adriano-Anaya et al., 2016; Costa et al., 2016; Tang et al., 2016; Li et al., 2017), including a published patent WO2011084867 A2 (Rotter, 2010), little is known about the type of inheritance patterns of the FOF trait. The objective of this work was to elucidate the heredity of the FOF native trait through the inbreeding and crossing process of JCL accessions for further potential usage as female parent for commercial hybrid seed production.

MATERIALS AND METHODS

Location of the experiment

The study was carried out in the Jatronergy Research Center, property of Lodemo Company, located in Espita, Yucatan, Mexico. The surrounding vegetation in the experimental areas is mainly tropical forest. The plantations are part of a breeding program aimed at improving JLC for biodiesel production. The soil types in the experimental areas are young soils with shallow depth, mostly stony soil with calcareous origin, with pH between 6.3 to 7.3 and poor organic matter content between 6.7 to 8.9% (Bautista et al.,

2005; Bautista et al., 2015).

Genetic materials

A total of 23 accessions of JCL, all property of Jatronergy germplasm bank with origins from the Americas and Mexico, were included in the evaluation method. Table 1 shows the morphological description carried out from 2012 to 2014. All accessions were maintained *in vivo*, by planting cuttings in the field.

Determination of female only flower heredity by inbreeding

Inbreeding wild accessions

A total of eight wild accessions were included in the process of inbreeding to determine the segregation of FOF trait in different inbred populations (Table 2). In parallel, the FOF trait segregation was estimated in a single wild accession, ID 4, by self-pollination up to S₃ endogamy level by using only FM segregating plants (Table 3). In order to represent the endogamy process and differentiate between populations, a hypothetical pedigree was used; the first self-pollination was represented by the letter "X", the number position references a high level of inbreeding for example S₁: (4)X; S₂: (4)X6, and so forth. The number also represents the selected plant among all individuals in the populations.

Experimental design

The eight wild accessions were planted in the germplasm bank area of the company in 2012. Each individual accession was planted in rows of 15 plants per row with 3 m among rows and 2.5 m between plants. All replicated plants per accessions were cuttings. Subsequently, following Jatronergy protocols, inbred seeds were obtained by self-pollination from each wild accession with endogamy levels of S₁, S₂ or S₃. The inbreeding process S₁ was carried out and seeds were harvested in 2012 and S₂ and S₃ in 2013. All inbred seeds were planted in a separate plot in April to May 2014. An inbred population was composed of 10 or 30 plants, each planted at 3.0 m among rows and 2.5 m between plants. Some rows could have more than one population. The plants were drip irrigated, fertilized, and weeded according to the protocols of the company.

During the inbreeding process, the expected inheritance ratio was calculated assuming a Mendelian monogenic inheritance, where the FOF trait represents a homozygous recessive gene (*ff*) and the FM trait, both homozygous and heterozygous dominant gene (*FF* or *Ff*).

Heredity by crossing FOF female parental with FM male parental (F1 inheritance)

Genetic material

Two wild genotypes of JCL accessions, ID 20 and 24 (Table 1), were selected in the germplasm bank with previous knowledge of naturally expressing the FOF trait as observed by other authors (Adriano-Anaya, 2016; Ovando-Medina et al., 2013); both materials were used as female parents and crossed with 21 and 20 wild accessions, respectively (Table 1) with FM trait used as males to produce F1 hybrids (Tables 4 and 5).

Experimental design

The crosses were carried out and harvested according to Jatronergy protocols in 2012. All hybrid harvested seeds were

Table 1. Morphological description of 23 accessions of *Jatropha curcas* L. included in the evaluation of heredity Female Only Flower (FOF) trait.

Accession ID	TF	PH (m)	SGF ¹	LR ²	LL ²	SS ³	SR ²	SW (g)	ER ⁴
1	FM	3.68	2	0.876	5.00	2	1.730	0.914	7
2	FM	3.31	1	0.916	5.00	2	1.678	0.800	7
3	FM	3.15	2	0.834	5.00	2	1.687	0.931	7
4	FM	2.42	3	0.855	2.50	2	1.664	0.752	5
5	FM	3.40	2	0.984	5.00	1	1.572	0.818	5
6	FM	3.05	2	0.857	5.00	1	1.574	0.826	7
7	FM	2.77	2	0.839	4.75	2	1.655	0.837	7
8	FM	2.10	3	0.899	5.00	2	1.760	0.740	5
9	FM	3.55	1	0.926	4.75	2	1.691	0.837	7
10	FM	2.35	2	0.830	4.75	1	1.627	0.904	7
11	FM	1.96	3	0.822	5.00	2	1.753	0.671	5
12	FM	1.64	3	0.866	5.00	2	1.783	0.712	5
13	FM	1.65	2	0.893	5.00	3	1.736	0.648	7
14	FM	2.59	2	0.934	5.25	2	1.825	0.805	7
15	FM	2.78	2	0.878	4.00	2	1.596	0.858	2
16	FM	2.16	3	1.014	5.00	2	1.757	0.748	5
17	FM	2.92	2	0.877	4.75	2	1.616	0.788	5
18	FM	2.03	2	0.870	5.00	2	1.734	0.802	7
19	FM	1.78	2	0.901	4.75	2	1.773	0.766	7
20	FOF	2.99	2	0.869	5.00	2	1.578	0.846	7
22	FM	2.92	2	0.904	5.00	2	1.726	0.887	7
23	FM	2.45	2	0.933	4.00	2	1.605	0.774	7
24	FOF	2.93	2	0.877	4.00	1	1.517	0.981	7
Average	-	2.63	-	0.889	4.75	-	1.680	0.811	-
SD ±	-	0.60	-	0.05	0.59	-	0.08	0.080	-

TF, Type of flowering; FM, Female and male flowers; FOF, Female Only Flowers; PH, Plant height; SGF, Stem growth form; LR, leaf relationship (length/width); LL, leaf lobes; SS, seed shape; SR, seed relationship; SW, seed weight; ER, endocarp rugosity; ¹qualitative, 1, Erect; 2, Semierect; 3, Open; ²Quantitative numerical; ³Semiquantitative; 1, Oblong; 2, elliptical; 3, obovate; ⁴ Qualitative; 1, Absent; 5, medium; 7, High. Accession ID 21 was not included in the morphological description analysis.

Table 2. Segregation of FOF plants of several populations per accessions of *Jatropha curcas* L. under process of inbreeding planted in Yucatan, Mexico, during 2014 and 2015.

Accession ID	Inbreeding level by population Endogamy level	N	Number of FM plants	Number of FOF plants	FOF trait (%)	E (FM/FOF)	χ^2	P<F
4	S ₁	35	25	10	29	18.7/6.25	3.0	0.083
4	S ₂	148	127	21	14	111/37.0	9.2	0.002
4	S ₃	199	152	47	24	149.2/49.7	0.20	0.653
6	S ₂	4	3	1	25	3.0/1.0	>0.0	0.99
7	S ₂	125	101	24	19	93.7/31.2	2.24	0.134
9	S ₂	16	15	1	6	12.0/4	3.0	0.083
13	S ₂	13	11	2	15	9.75/3.2	0.64	0.423
16	S ₂	21	20	1	5	15.75/5.25	4.87	0.032
19	S ₂	16	14	2	25	12.0/4.0	1.33	0.248
22	S ₂	24	22	2	8	18.0/6.0	3.55	0.059

N, Total plants in the population; FM, plants with female and male flowers; FOF, plants with female only flowers.

Table 3. Segregation of FOF plants by tracking one accession of *Jatropha curcas* L. under process of inbreeding planted in Yucatan, Mexico, during 2014 and 2015.

¹ Endogamy Population	Endogamy level	N	Number of FM plants	Number of FOF plants	FOF trait (%)	E (FM/FOF)	χ^2	<i>P</i> < <i>F</i>
² (4)	S ₀	5	5	0	0	-	-	-
(4)X	S ₁	9	-	-	-	-	-	-
<u>(4)X1</u>	S ₂	10	9	1	10	7.5/2.5	1.200	0.273
(4)X11	S ₃	5	4	1	20	3.7/1.2	0.067	0.796
<u>(4)X2</u>	S ₂	44	35	9	20	33/11	0.485	0.486
(4)X21	S ₃	4	4	0	0	3.0/1.0	1.330	0.248
(4)X22	S ₃	4	4	0	0	3.0/1.0	1.330	0.248
(4)X23	S ₃	30	18	12	40	22.5/7.5	3.600	0.058
(4)X24	S ₃	5	5	0	0	3.75/1.25	1.660	0.197
(4)X25	S ₃	5	5	0	0	3.75/1.25	1.660	0.197
<u>(4)X3</u>	S ₂	34	27	7	21	25.5/8.5	0.353	0.552
(4)X31	S ₃	27	27	0	0	20.25/6.75	9.000	0.003
(4)X32	S ₃	5	5	0	0	3.75/1.25	1.660	0.197
(4)X33	S ₃	56	37	19	34	42/14	2.381	0.123
(4)X34	S ₃	4	4	0	0	3.0/1.0	1.330	0.248

¹Underlined represents founder population at a certain endogamy level; ²Wild plant; N, Total plants in the population; E, expected value; FM, plants with female and male flowers; FOF, plants with female only flowers.

planted in a separated plot in August 2013. A hybrid population composed of 10 or 30 plants; each one planted at 3.0 m among rows and 2.5 m between plants. Some rows could have more than one population. The plants were drip irrigated, fertilized, and weeded according to the protocols of the company.

Data collection

Traits recorded were those relating to flowering biology at a single plant level; type of inflorescence per plant (FM or FOF plants). The data were recorded from August to October in 2014 and 2015 for methodologies of inbreeding and crossings.

Genetic diversity

For assessment of the genetic differences between accessions, eight morphological traits (Table 1) were included in a cluster analysis using Euclidean distance and UPGMA method, with Primer v6 (Clarke and Goyler, 2006). This was to demonstrate evidence of genetic diversity for the crosses to produce F1 and parents used to produce F1 were no siblings.

Data analysis for segregation of FOF trait

For the two methodologies described above, in order to estimate proportions of segregation of FOF plants, the χ^2 (Chi-Square) test was used to validate the existence of Mendelian inheritance of the FOF trait; this is a simple method of quantifying the various deviations expected from observed frequencies, or how well the observed data fits the predictions values if a hypothesis is true. A statistical significance of $p=0.05$ was used.

RESULTS

FOF heredity by inbreeding of several accessions

All ten populations from eight different *J. curcas* L.

accessions in S₁, S₂ and S₃ levels of endogamy showed segregation of FOF plants ranging from 5 to 29% in proportion to all their number of plants. The values of χ^2 ranged from 0.0 to 9.2 and only two populations out of ten in S₂ level of endogamy had values less than $P < 0.05$, indicating that mating is random and that segregation and independent assortment resulted in a substantial statistical deviation between observed and expected values of FOF traits plants. In the rest of the populations there was not rejection of the null hypothesis, indicating that observed values were the same as the expected values (no substantial statistical deviations). The hypothesized phenotypic ratio of FM:FOF traits was 3:1 (75%:25%), with eight out of ten populations of JCL accession confirming a monogenic Mendelian inheritance of the recessive homozygous allele *ff*. Given that nine populations segregated the FOF trait in less than 29%, this indicates that the alleles of the plants which were self-crossed were in a heterozygous state *Ff*, and none of them had homozygous dominant alleles *FF* for FM trait (Table 2).

FOF heredity by tracking the inbreeding process of one accession

The wild JCL accession, subjected to an inbreeding process from wild to S₃ endogamy level, showed segregation of plants with FM and FOF traits in their offspring. The founder S₁ population (4)X is a group of bulk plants subjected to an inbreeding process; however, in S₂ and S₃ endogamy levels it was possible to track the segregation (Table 3).

Table 4. Hybrids F1 from crossing a wild accession ID 20 of *Jatropha curcas* L. with naturally occurring FOF trait with other 21 accessions with FM trait.

Hybrid (F1) (Acc. ID)	N	Number of FM plants	Number of FOF plants	% FOF plants	E FM plants	E FOF plants	χ^2	$P < F$
A1 (20x1)	10	0.00	10	100.00	5.00	5.00	10.0	0.002
A2 (20x2)	9	3.00	6	66.67	4.50	4.50	1.0	0.317
A3 (20x3)	10	5.00	5	50.00	5.00	5.00	0.0	1.000
A4 (20x4)	10	6.00	4	40.00	5.00	5.00	0.4	0.527
A5 (20x5)	10	7.00	3	30.00	5.00	5.00	1.6	0.206
A6 (20x6)	10	7.00	3	30.00	5.00	5.00	1.6	0.206
A7 (20x7)	10	7.00	8	20.00	5.00	5.00	3.6	0.057
A8 (20x8)	10	7.00	3	30.00	5.00	5.00	1.6	0.206
A9 (20x9)	10	6.00	4	40.00	5.00	5.00	0.4	0.527
A10 (20x10)	10	9.00	1	10.00	5.00	5.00	6.4	0.011
A11 (20x11)	10	10.00	0	0.00	5.00	5.00	10.0	0.002
A12 (20x12)	10	10.00	0	0.00	5.00	5.00	10.0	0.002
A13 (20x13)	10	10.00	0	0.00	5.00	5.00	10.0	0.002
A14 (20x14)	10	10.00	0	0.00	5.00	5.00	10.0	0.002
A15 (20x15)	10	10.00	0	0.00	5.00	5.00	10.0	0.002
A16 (20x16)	10	10.00	0	0.00	5.00	5.00	10.0	0.002
A17 (20x17)	10	10.00	0	0.00	5.00	5.00	10.0	0.002
A18 (20x18)	9	9.00	0	0.00	4.50	4.50	9.0	0.003
A19 (20x19)	10	10.00	0	0.00	5.00	5.00	10.0	0.002
A20 (20x22)	10	6.00	4	40.00	5.00	5.00	0.4	0.527
A21 (20x23)	10	10.00	0	0.00	5.00	5.00	10.0	0.002

N, Total plants; FM, plants with female and male flowers; FOF, plants with female only flowers.

From all 15 populations with different endogamy levels S_1 , S_2 and S_3 , only six populations segregated FOF plants. The percentage of FOF segregation ranged from 0 to 40% and the χ^2 values ranged from 0.067 to 3.60. According to the statistical significance for χ^2 test only two populations showed values of $P < 0.05$ indicating that that observed values differ from expected values (substantial statistical deviations).

The S_2 population (4)X1 and the next endogamy level S_3 population offspring (4)X11 segregated FOF plants. The S_2 population (4)X2 segregated FOF plants, however, only one population out of 5 of its S_3 endogamy level offspring segregated FOF plants. The S_2 population (4)X3 segregated FOF plants, however, only one population out of 4 of its S_3 endogamy level offspring segregated FOF plants.

The hypothesized phenotypic ratio of FM:FOF plants of 3:1 (75%:25%) was achieved in 13 out of 14 populations of JCL, according to the χ^2 values and $P < 0.05$, confirming a Mendelian inheritance of a recessive homozygous allele *ff*.

The populations in S_3 level of endogamy without segregation of FOF plants may have homozygous dominant allele *FF* or perhaps the small sample size or number of plants produced was too low, even though the

self-pollinated plant had heterozygous alleles *Ff* (Table 3).

Crossing accessions naturally expressing FOF trait accession with FM accessions

The cluster analysis showed phenotypic and genotypic differences among the accessions by using all traits; twenty-two accessions were grouped in two clusters ("a" and "b"), and two accessions separated ("4" and "15") from all evaluated accessions, with a range of distances of 8, 7, 13 and 20, respectively (Figure 1). Cluster "b" grouped the maximum number of accessions (total of 15) and the accessions with the FOF trait. The dendrogram showed a genetic contrast between all evaluated accessions, even among accessions with FOF trait within the same group (Figure 1). The results indicated high genetic variability.

The F1 hybrids of JCL, product of crossing a female accession ID 20 with FOF trait x 21 males, showed FOF segregations ranging from 0 to 100%. However, only eleven out of twenty-one F1 hybrids showed segregation of FOF plants (Table 4).

Three F1 hybrids had values $\geq 50\%$ of FOF segregation

Table 5. Hybrids F1 from crossing a wild accession ID 24 of *Jatropha curcas* L. with naturally occurring FOF trait with 20 other accessions with the FM trait.

Hybrid (F1) (Acc. ID)	N	Number of FM plants	Number of FOF plants	% FOF plants	E plants	FM plants	E plants	FOF plants	χ^2	<i>P</i> < <i>F</i>
B1 (24x1)	1	1	0	0.00	0.50	0.50	1.00	0.317		
B2 (24x2)	8	7	1	12.5	4.00	4.00	4.5	0.034		
B3 (24x3)	10	2	8	80.00	5.00	5.00	3.60	0.058		
B4 (24x4)	4	2	2	50.00	2.00	2.00	0.00	1.000		
B5 (24x5)	3	0	3	100.00	1.50	1.50	3.00	0.083		
B6 (24x6)	10	7	3	30.00	5.00	5.00	1.60	0.206		
B7 (24x7)	10	8	2	20.00	5.00	5.00	3.60	0.058		
B8 (24x8)	7	5	2	28.57	3.50	3.50	1.29	0.257		
B9 (24x9)	6	5	1	16.67	3.00	3.00	2.66	0.120		
B10 (24x10)	3	3	0	0.00	1.50	1.50	3.00	0.083		
B11(24x11)	9	9	0	0.00	4.50	4.50	9.00	0.003		
B12 (24x12)	10	10	0	0.00	5.00	5.00	10.0	0.002		
B13 (24x13)	10	10	0	0.00	5.00	5.00	10.00	0.002		
B14 (24x14)	10	10	0	10.00	5.00	5.00	10.00	0.002		
B15 (24x15)	10	8	2	20.00	5.00	5.00	3.60	0.058		
B16 (24x16)	10	10	0	0.00	5.00	5.00	10.00	0.002		
B17 (24x17)	10	10	0	0.00	5.00	5.00	10.00	0.002		
B18 (24x18)	10	10	0	0.00	5.00	5.00	10.00	0.002		
B19 (24x19)	9	9	0	0.00	4.50	4.50	9.00	0.003		
B20(24x21)	10	10	0	0.00	5.00	5.00	10.00	0.002		

N, Total plants; FM, plants with female and male flowers; FOF, plants with female only flowers.




Figure 1. Dendrogram UPGMA showing the morphological diversity between the 23 accessions of *Jatropha curcas* L. evaluated. FM, Female and male flowers; FOF, female only flowers.

plants. The values of χ^2 including all F1 hybrids ranged from 0.0 to 10.0 and the $P < 0.002$ to 1.0. Of eleven F1 hybrids that segregated FOF plants, only two (A1 and A10) had a $P < 0.05$, indicating that observed values differ from expected values (substantial statistical deviations). The nine remaining hybrids with $P < 0.05$ presented a FOF segregation less than 50% (Table 4).

The F1 hybrids that segregated FOF plants reached values from 10 up to 100% which confirmed the hypostatized phenotypic ratio of FM:FOF plants of 1:1 (50%:50%) respectively and also confirmed the Mendelian inheritance of a recessive homozygous allele *ff* in the female parent when crossed with a male with heterozygous alleles *Ff*. However, in the case of F1 hybrids without segregation of FOF plants, this could be due to the small sample size or perhaps the number of plants produced was too low, even though the male parent may have heterozygous alleles *Ff* or the male parent had homozygous dominant allele *FF* (Table 4).

The F1 hybrids of JCL, product of crossing the accession ID 24 with FOF trait with 20 males, showed FOF segregations ranging from 0 to 100%. However, only nine out of 20 F1 hybrids showed segregation of FOF plants (Table 5). Three F1 hybrids had values $\geq 50\%$ of FOF segregation plants. The values of χ^2 including all F1 hybrids ranged from 0.0 to 10.0 and the P from 0.002 to 1.0. From nine F1 hybrids that segregated FOF plants only one cross, B12, had a $P < 0.05$, indicating that observed values differ from expected values (substantial statistical deviations). The remaining eight hybrids with $P < 0.05$ had FOF segregation less than 50%.

The crosses that segregated FOF plants reached values from 10 up to 100%, confirming the hypothesized phenotypic ratio of FM:FOF plants of 1:1 (50%:50%), respectively, and thereby confirming the Mendelian inheritance of a recessive homozygous allele *ff* in the female parent when crossed with a male with heterozygous alleles *Ff*. However, in the case of crosses without segregation of FOF plants, this could be due to the sample size or the low number of plants produced, even though the male parent may have heterozygous alleles *Ff* or the male parent had homozygous dominant allele *FF* (Table 5).

DISCUSSION

Inbreeding several populations

Segregation of FOF trait naturally occurred in almost all wild populations of JCL when induced to an inbreeding process. Segregation of less than 30% of FOF plants in S_1 , S_2 and S_3 levels of endogamy, confirmed that this trait is inherited at a phenotypic ratio of 75% of FM and 25% FOF, according to Mendelian genetics, since monogenic inheritance implies segregation at a single locus (Elston et al., 2012). Similarly, in tobacco, the native gene *fw*

expressing the double-flowering follows an inheritance of a monogenic recessive trait when it is homozygous (Zainol and Stimart, 2001). However, for the FOF segregation, more than 25% in S_1 could be due to the law of maximum segregation according to Mendel's laws. For all S_2 populations FOF segregation was $\leq 25\%$. The FOF trait variability ranged between 5 to 29%, which may also be due to factors relating to sampling size, since the seeds for planting, which were selected randomly from one single plant or seed number and produced by self-pollination, were not enough. It is likely therefore that the protandry and protogyny reported in JLC may play a role in the amount of seed production, as reported by Negussie et al. (2014). Plants expressing FOF trait for two consecutive years give support to the belief that this trait is not environment dependent, as in the case of cytoplasmic male sterility (CMS) (Fan and Stefansson, 1986).

Inbreeding of one population

Segregation of FOF traits occurred in one wild population of JCL, when induced by the inbreeding process; it also occurred in S_1 , S_2 , and S_3 endogamy levels and the segregation of less than 40% of FOF plants in all endogamy levels confirmed that this trait is inherited at a phenotypic ratio of 75% of FM and 25% of FOF plants according to Mendelian laws. However, not all plants of a population segregated FOF plants; this could be due to a selection of plants with FM trait to continue the inbreeding process that have heterozygous alleles *Ff* for the FOF trait or even homozygous alleles *FF* for FM trait that may not segregate in the further endogamy dose and relating to sampling size. Occurrence of FOF trait has been observed in wild accessions of JLC (Adrano-Anaya et al., 2016; Ovando-Medina et al., 2013); this may occur due to the type of JLC reproduction which can produce seed by self-pollination (geitonogamy) or even by xenogamy when crossing half or full sibling plants (Ahoton and Quenum, 2012). Inbreeding depression in JCL did not affect survival and fertility although the possible presence of recessive deleterious mutations has been reported by Charlesworth and Willis (2009).

Crossing accession naturally expressing FOF trait with FM accessions

Wild accessions of JCL (ID 20 and 24) naturally express the FOF trait and, when crossed with wild accessions with the FM trait, can segregate FOF plants. However, the percentage of that segregation can range from 0 up to 100%. Segregation of $>50\%$ of FOF trait indicates a phenotypic ratio of 50% of this trait according to Mendel's laws. The range of the variable may be due to factors such as sampling size and zygosity of the alleles, either

Ff or *FF*, in the male parents (Fairbanks and Rytting, 2001).

Although JCL is a monoecious plant (Noor Camellia et al., 2012), the origin of those two JCL accession natural expressing FOF trait in this study, may be occurred by natural inbreeding by geitonogamy or by xenogamy (sibling cross or cross with two genetically different plants) (Fresnedo-Ramirez, 2013; Rincón-Rabanales et al., 2016). Sibling crosses may occur when an indehiscent fruit produces three sibling plants which germinate and develop in close proximity and cross with each other. Ahoton and Quenum (2012) also reported that JCL can produce seed by self-pollination, inducing natural /spontaneous autogamy (geitonogamy). If two wild plants have the FOF trait in a heterozygous allele they can also produce FOF trait plants. Kaur et al. (2011) reported up to 72.2% of natural self-pollination and 36.3% of apomixes which is associated with a high level of endogamy in JCL.

In nature, it is likely that plants do not produce the FOF trait due to the fact that at the moment of crossing, one parent has homozygous dominant FM trait alleles (*FF*) and the other heterozygous FOF trait alleles (*Ff*), thus, the offspring will never phenotypically express the FOF trait. There is also evidence of seed produced by apomixes; in this case, the plant produced under this mechanism could affect the expression of FOF trait (Ahoton and Quenum, 2012).

The inheritance of this trait, therefore, is nuclear and not cytoplasmic, as with the CMS in other crops, which is an environment-dependent trait. Some reports mention that sexual ratio in *Jatropha* flowering is influenced by temperature (Kaur et al., 2011) or by the mode of reproduction (Nietsche et al., 2014) however, there is no strong data available to support this statement. Nietsche et al. (2014) reported that extreme high temperatures produced flower abortions. The FOF trait, therefore, has the potential to produce JCL hybrid seed without the risk of compromising sterility in the female parent or causing contamination, while also reducing the cost by avoiding detasseling.

Fresnedo-Ramirez and Orozco-Ramirez (2013) has reported that flower sex ratio in JCL is genetically controlled and can be altered by photoperiod, relative humidity, temperature, soil nutrient availability, exogenously applied growth regulators and by an epigenetic control. Plant growth regulators such as benzyladenine and gibberellic acid could affect number of flowers and branching but would not modify the genetics (Costa et al., 2016). Also, some genes play a role in regulating flowering; the gene *jcLFY* in JCL regulates flower identity, flower organ patterns and fruit shape (Tang et al., 2016), *JcTFL1* genes play redundant roles in repressing flowering in *Jatropha* (Li et al., 2017). Flower sex ratio, therefore, is not a conservative trait. The inbreeding process of JCL could be a method to derive FOF trait plants used as female plants in a hybrid

production.

The genetic diversity observed in this study formed two clusters that could be divergent for a hybridization program, similar to that reported by Gohil and Pandya (2008) in India, although they evaluated nine genotypes for a breeding program, which were grouped into five clusters. They proposed that only clusters III and IV were the most divergent. The results in this study may be attributed to a wide diversity among groups, since all accessions proceed from America, center of origin for JCL (Heller, 1996), where the possibility to naturally express a homozygous recessive allele for FOF trait is high, as reported (Adriano-Anaya et al., 2016; Ovando-Medina et al., 2013). This genetic diversity is a prerequisite to achieve a successful breeding program (Chikara et al., 2013).

A morphological study of seed traits, based on non-hierarchical Euclidean cluster analysis, in 24 accessions from India, revealed 6 clusters; the maximum inter-cluster distance was 5.1 and the minimum was 2.4 (Kaushik et al., 2007); the American and Mexican accessions evaluated in this study revealed greater genetic diversity, as was also reported by Fresnedo-Ramírez and Orozco-Ramirez (2013). Based on our results, FOF is a monogenic homozygous recessive trait with potential for use in the production of commercial volumes of JCL hybrid seed.

CONFLICT OF INTERESTS

The authors declare that they have no conflict of interest.

REFERENCES

- Adriano-Anaya MdL, Pérez-Castillo E, Salvador-Figueroa M, Ruiz-González S, Vázquez-Ovando A, Grajales-Conesa J, Ovando-Medina I (2016). Sex expression and floral diversity in *Jatropha curcas*: a population study in its center of origin. PeerJ 4: e2071.
- Ahoton LE, Quenum F (2012). Floral Biology and Hybridization Potential of Nine Accessions of Physic Nut (*Jatropha curcas*) L. Originating from three continents. Tropicult. 30(4):193-198.
- Bautista F, Frausto O, Ihl T, Aguilar Y (2015). Actualización del mapa de suelos del Estado de Yucatán, México: Enfoque Geomorfopedológico y WRB. Ecosist. Rec. Agropec. 2(6):303-315.
- Bautista F, Palma-López D, Huchin-Malta W (2005). Actualización de la clasificación de los suelos del estado de Yucatán. In: Bautista F, Palacio G (eds.) Caracterización y Manejo de los Suelos de la Península de Yucatán: Implicaciones Agropecuarias, Forestales y Ambientales. Universidad Autónoma de Campeche, Universidad Autónoma de Yucatán, 2005.
- Birchler JA, Yao H, Chudalayandi S, Vaiman D, Veitia RA (2010). Heterosis. Plant Cell 22:2105-2112.
- Bressan EA, Sebbenn AM, Ferreira RR et al (2013). *Jatropha curcas* L. (Euphorbiaceae) exhibits a mixed mating system, high correlated mating and apomixes. Tree Genet. Genome 9(4):1089-1097.
- Costa AP, Vendrame W, Nietsche S, Crane J, Moore K, Schaffer B (2016). Bunching, flowering and fruiting of *Jatropha curcas* treated with ethephon or benzyladenine and gibberellins. Ann. Braz. Acad. Sci. 88(2):989-998.
- Charlesworth D, Willis JH (2009). The genetics of inbreeding depression. Nature Rev. Genet. 10(11):783-796.
- Chikara J, Prakash AR, Mastan SG, Ghosh A (2013). Genetic

- improvement in *Jatropha curcas* through selection and breeding. In: Bahadur B, Sujatha M, Carels N. (eds) *Jatropha*, Challenges for a New Energy Crop. Springer, New York, NY.
- Clarke KR, Gorley RN (2006). PRIMER v6: User Manual/Tutorial. PRIMER-E, Plymouth.
- Elston RC, Satagopan JM, Sun S (2012). Genetic Terminology. *Methods in Mol. Biol.* 850:1-9.
- Fan Z, Stefansson BR (1986). Influence of temperature on sterility of two cytoplasmic male-sterility system in rape (*Brassica napus* L.). *Can. J. Plant Sci.* 66:221-227.
- Fairbanks D, Rytting B (2001). Mendelian controversies: a botanical and historical review. *Am. J. Bot.* 88(5):737-752.
- Fresnedo-Ramírez J (2013). The floral biology of *Jatropha curcas* L. – a review. *Trop. Plant Biol.* 6:1-15.
- Fresnedo-Ramírez J, Orozco-Ramírez Q (2013). Diversity and distribution of genus *Jatropha* in Mexico. *Genet. Resour. Crop Evol.* 60(3):1087-1104.
- Gohil RH, Pandya JB (2008). Genetic diversity assessment in physic nut (*Jatropha curcas* L.). *Int. J. Plant Prod.* 2(4):321-326.
- Haque MA, Monir uz Z, Rahman MS, Alam QM (2012). Profitability of hybrid maize (*Zea mays* L.) seed production under contract farming in Bangladesh: A farm level study. *Bang. J. Agric. Res.* 37(2):327-334.
- Heller J (1996). Physic nut. *Jatropha curcas* L. Promoting the conservation and use of underutilized and neglected crops. 1. Institute of Plant Genetics and Crop Plant Research, Gatersleben/International Plant Genetic Resources Institute, Rome.
- Johnson AG (1966). Inbreeding and the production of commercial F1 hybrid seed of brussels sprouts. *Euphytica* 15(1):68-79.
- Kaur K, Dhillon GPS, Gill RIS (2011). Floral biology and breeding system of *Jatropha curcas* in North-Western India. *J. Trop. For. Sci.* 23(1):4-9.
- Kaushik N, Kumar K, Kumar S, Kaushik N, Roy S (2007). Genetic variability and divergence studies in seed trait and oil content of *Jatropha* (*Jatropha curcas* L.) accessions. *Biomass Bioener.* 31(7):497-502.
- King AJ, He W, Cuevas JA, Freudenberger M, Ramiaramananana D, Graham IA (2009). Potential of *Jatropha curcas* as a source of renewable oil and animal feed. *J. Exp. Bot.* 60(10):2897-2905.
- Li C, Fu Q, Niu L, Luo L, Chen J, Xu ZF (2017). Three TFL1 homologues regulate floral initiation in the biofuel plant *Jatropha curcas*. *Sci. Rep.* 7:43090.
- Masuda M, Ojiewo CO, Nagai M, Murakami K, Masinde PW, Yu W (2010). Simulating hybrid-seed contamination risk with selfed seeds from residual fertility in a male-sterile T-4 mutant tomato, *Solanum lycopersicum* L. *J. Jap. Soc. Hort. Sci.* 79(1):34-39.
- McRobert JF, Setimela PS, Gethi J, Worku M (2014). Maize hybrid seed Production Manual, CIMMYT, México.
- Negussie A, Achten WMJ, Verboven HAF, Hermy M, Muys B (2014). Floral display and effects of natural and artificial pollination on fruiting and seed yield of the tropical biofuel crop *Jatropha curcas* L. *GCB Bioenerg.* 6(3):210-218.
- Nietsche S, Vendrame WA, Crane J H, Pereira MCT (2014). Assessment of reproductive characteristics of *Jatropha curcas* L. in south Florida. *GCB Bioenerg.* 6(4):351-359.
- Noor Camellia NA, Thohirah LA, Abdullah NAP (2012). Floral biology, flowering behaviour and fruit set development of *Jatropha curcas* L. in Malaysia. *Pertanika J. Trop. Agric. Sci.* 35(4):725-736.
- Ovando-Medina I, Adriano-Anaya L, Vázquez-Ovando A, Ruiz-González S, Rincón-Rabanales M, Salvador-Figueroa M (2013). Genetic Diversity of *Jatropha curcas* in Southern Mexico. In: Bahadur B., Sujatha M., Carels N. (eds) *Jatropha*, Challenges for a New Energy Crop. Springer, New York, NY.
- Pecina-Quintero V, Anaya-López JL, Zamarripa A, Montes N, Núñez C, Solís J, Aguilar-Rangel M, Gil H, Méjia D (2011). Molecular characterization of *Jatropha curcas* L. genetic resources from Chiapas, México through AFLP markers. *Biomass Bioenerg.* 35(5):1897-1905.
- Perez-Prat E, van Lookeren MM (2002). Hybrid seed production and the challenge of propagating male-sterile plants. *Trends Plant. Sci.* 7(5):199-203.
- Rincón-Rabanales M, Vargas-López LI, Adriano-Anaya L, Vázquez-Ovando A, Salvador-Figueroa M, Ovando-Medina I (2016). Reproductive biology of the biofuel plant *Jatropha curcas* in its center of origin. *Peer J* 4:1-12.
- Rotter R (2010). *Jatropha* hybrids through female only trait. U.S. Patent No. US2010062525.
- Stevens WE, Berberich SA, Sheckell PA, Wiltse CC, Halsey ME, Horak MJ, Dunn DJ (2004). Optimizing Pollen Confinement in Maize Grown for Regulated Products. *Hortic. Sci.* 44(6):2146-2153.
- Tang M, Tao YB, Fu Q, Song Y, Niu L, Xu ZF (2016). An ortholog of LEAFY in *Jatropha curcas* regulates flowering time and floral organ development. *Sci. Rep.* 6:1-15. DOI:10.1038/srep37306
- Vaknin Y (2012). The significance of pollination services for biodiesel feedstocks, with special reference to *Jatropha curcas* L.: A review. *Bioenerg. Res.* 5(1):32-40.
- Wu Y, Fox TW, Trimnell MR, Wang L, Xu RJ, Cigan AM, Huffman GA, Garnaat CW, Hershey H, Albertsen MC (2016). Development of a novel recessive genetic male sterility system for hybrid seed production in maize and other cross-pollinating crops. *Plant Biotechnol. J.* 14(3):1046-1054.
- Yamagishi H, Bhat SR (2014). Cytoplasmic male sterility in Brassicaceae crops. *Breed. Sci.* 64(1):38-47.
- Zainol R, Stimart DP (2001). A monogenic recessive gene, *fw*, conditions flower doubling in *Nicotiana glauca*. *Hortsci.* 36(1):128-130.

Full Length Research Paper

Estimates of combining ability for resistance to pod shattering in soybean (*Glycine max* (L.) Merrill) genotypes

F. A. Umar^{1*}, M. S. Mohammed², M. Oyekunle², I. S. Usman², M. N. Ishaq¹ and S. N. Dachi¹

¹National Cereals Research Institute, P. M. B. 8, Badeggi, Niger State, Nigeria.

²Department of Plant Science, Ahmadu Bello University, Zaria, Kaduna State, Nigeria.

Received 16 August, 2017; Accepted 4 October, 2017

Combining ability estimates were studied for pod shattering and other agronomic traits in eight parental soybean genotypes of different pod shattering group namely: susceptible to pod shattering, moderately resistant to pod shattering were crossed in a 4 x 4 North Carolina mating design II to generate 16 crosses, due to missing stands encountered in some crosses, 9 successful F_{1s} hybrids were obtained in the present study and evaluated along with the parents to estimate the mode of gene action controlling pod shattering in soybean and the combining ability for pod shattering and other agronomic traits in soybean. The mean square from the analysis of variance for the ten traits measured showed highly significant differences ($p < 0.01$) among the genotypes. This has demonstrated the existence of genetic variability among soybean genotypes with pod shattering under additive gene action. TGX1955-10E, NG/AD/11/08/023 and NG/SA/07/100 were good general combiners for resistance to pod shattering, plant height, days to 50% flowering and days to maturity while NG/MR/11/11/060, NGB00/08, TGX1740-1F and NG/SA/07/055 are good general combiner for number of seeds/ pod, pod length, hundred seed weight and grain yield. NG/SA/07/100 x TGX1740-1F, NG/MR/11/11/060 x NGB00129 and NG/MR/11/11/060 x NG/AD/11/08/023 had positive specific combining ability effects for number of branches/ plant, number of pods/plant, pod length, number of seeds/pod, hundred seed weight and grain yield. NG/MR/11/11/060 X NG/SA/07/055 had negative SCA effects for plant height. This implied that these hybrids performed better than the parents GCA effects. This suggests that the cross combination can be advanced for selection in later generation.

Key words: Combining ability effects, soybean genotypes, pod shattering, Samaru.

INTRODUCTION

Soybean ($2n = 40$), an important leguminous and a miracle crop of the world, has its origin in North-Eastern China (Indu, 2014). It is among the major industrial and

food crops grown in every continent. It is an economically important crop with an average of 40% protein and 20% oil content (Context Network and Sahel Capital, 2016).

*Corresponding author. E-mail: fatima4destiny19@yahoo.com.

Like other important economic crops, one of the major problems associated with soybean production in the tropical and sub-tropical ecology is pod shattering, an important physiological constrain in soybean production and its productivity.

Pod shattering is the opening of mature pod along the dorsal or ventral sutures and followed by seed dispersal when the crop reaches maturity and during harvesting (Bara et al., 2013). The extent of yield loss that could result due to pod shattering in soybean may range from 34 to 100% (Krisnawati and Adie, 2016). Resistance to pod shattering is the most important factor for the improvement of soybean, especially in tropics. The Nigerian climate is characterized by abundant sunshine, which provides an ideal environment for soybeans production, hence will increase in pod shattering and resulting in significant yield losses in soybean.

Understanding the mode of gene action controlling resistance to pod shattering is crucial for designing appropriate breeding strategy for the development of pod shattering resistance. Earlier studies have generated different results on the genetics of pod shattering. Caviness (1969) reported that pod shattering is a qualitative heritable trait with multiple genes governing the trait. Haruna (2010) reported that inheritance of resistance to pod shattering was under the influence of either duplicate recessive or dominant and recessive epistasis depending on the parental genotypes used in the cross.

Exploiting genetic variability in soybean through effective crosses between adapted varieties bearing pod shattering trait and cultivars with resistance to pod shattering will provide information for the development of high yielding and resistant varieties that will increase the productivity and utilization of soybean. This study was therefore undertaken to determine the mode of gene action controlling pod shattering, to estimate the general combining ability (GCA) effects of the parents and specific combining ability (SCA) effects of the hybrids.

MATERIALS AND METHODS

Eight soybean genotypes which consisted of two susceptible, two moderately resistant to pod shattering and four resistant to pod shattering were crossed in a 4 × 4 North Carolina mating design II (NC11) to generate 16 F₁s hybrids during the dry season of 2016, but due to missing stands, 9 successful F₁s were generated for this study and evaluated along with the 8 parents. This was planted in the screen house of Institute for Agriculture Research (IAR), Samaru Ahmadu Bello University, Zaria (Latitude 11°11'N and 7°38' E, 600 m above sea level) which is located in the Northern Guinea Savannah Zone of Nigeria, with mean annual rainfall is about 1045 mm during the rainy season and were laid-out in a randomized complete block design (RCBD) with three replications. Each experimental unit consisted of a pot of 24.5 × 25.5 cm², diameter and height, respectively with each pot containing 10 kg of sandy-loam soil. Four seeds were sown per pot. All recommended agronomic practices like weeding, fertilizer application of 30 kg P₂O₅/ha in form of single super phosphate (SSP-18%) and a single

spray of insecticide Cypermethrin + Dimethoate at the rate of 100 ml in 1.5 L of water for any infestation from pre-flowering through post-flowering phases were carried out to maintain healthy plants. Data were collected on number of days to flowering, number of branches per plant, number of pods per plant, plant height at maturity, number of seeds per pod, 100 seeds weight, days to maturity, pod length, and grain yield.

The level of resistance to pod shattering was evaluated using the field-screening method (Helms, 1994) on all the genotypes tested. This was scored on pods of matured genotypes (95% of pods turn tan or grey). Pod shattering score was taken at one, two and three weeks after maturity. The percentage of pod shattering was then determined on a scale of 1 to 5 used by Asian Vegetable Research and Development Centre (AVRDC, 1979) in which, 1 = 0% shattering, 2 = 1 to 10%, 3 = 11 to 25%, 4 = 26 to 50%, and 5 ≥ 50%. The shattering score was classified as follows: 1 = highly resistant; 2 = resistant; 3 = moderately resistant; 4 = susceptible; and 5 = highly susceptible (AVRDC, 1979). Percent pod shattering was estimated using the following formula:

$$\frac{\text{Number of shattered pods per plant}}{\text{Total number of pods per plant}} \times 100$$

The combining ability analysis and the estimates of GCA and SCA effects were done using NCD II for Model I method based on the procedure described by Comstock and Robinson (1948) using Statistical Analysis System (SAS) Package (2002). The significant differences among GCA and SCA effects were tested using the formula of Singh and Chaunghary (1977).

RESULTS AND DISCUSSION

Response of F₁ population and parental genotypes to pod shattering

The results in Table 1 show the mean square for pod shattering and other agronomic traits in soybean genotypes. All traits were highly significant (p<0.01) indicating the existence of genetic diversity among parental genotypes and their progeny. This implied that selection for desirable soybean genotypes could be made using these traits in a soybean breeding program. Similar result was reported by Nassar (2013).

Combining ability analysis

Mean square for both GCA and SCA were significant for all studied characters (Table 2), except for number of pods per plant which showed no significant difference by female parents. The significant difference observed for SCA and GCA for all the traits studied in the present study indicates the importance of additive and non-additive gene actions in the inheritance of the studied traits, though each component's contribution may vary according to each traits. The high GCA/SCA ratio obtained from the mean square due to GCA for pod shattering indicates that additive gene action was more important in the control of this trait. This is similar to the

Table 1. Mean square of all traits for NC11 crosses evaluated at Samaru, Zaria in 2016.

Source of variation	Df	Days to 50% flowering	Plant height (cm)	Number of branches/plant	Number of pods/plant	Number of seeds/pod	Pod length (cm)	Hundred seed weight (g)	Days to maturity	Grain yield (kg)	Shattering score
Genotype	15	2191.75**	3986.40**	53.62**	4368.84**	3.77**	9.61**	73.22**	7146.75**	2242228.27**	9.73**
REP	2	14.06	120.89	6.81	5.77	0.02	0.11	0.06	3.77	139.65	0.02
Error	30	11.06	164.07	3.23	515.39	0.11	0.08	0.06	4.84	98.87	0.02

**Significant at 1% level of probability, *significant at 5% level of probability, ns= not significant.

Table 2. Mean square for combining ability and GCA/SCA ratio for resistance to pod shattering and other agronomic traits in soybean genotypes evaluated at Samaru 2016.

Source of variation	Df	Days to 50% flowering	Plant height (cm)	No. of braches/plant	No. of pods/plant	No. of seeds/pod	Pod length (cm)	Days to maturity	Hundred seeds weight (g)	Grain yield (kg)	Pod shattering score
Rep	2	14.06	120.90	6.81	5.77	0.021	0.11	3.77	0.05	139.65	0.02
Female	3	1646.74**	3791.47**	12.61*	1136.74	3.92**	8.59**	6034.69**	36.31**	1141836.44**	19.83**
Male	3	1754.52**	2656.24**	73.06**	4886.19**	3.47**	10.53**	5829.81**	105.46**	2365948.25**	5.33**
M*F	9	2519.17**	4494.76**	60.81**	5273.76**	3.82**	9.64**	7956.42**	76.56**	2567785.55**	7.83**
Error	30	11.06	164.07	3.23	515.39	0.110	0.08	4.84	0.05	98.87	0.02
GCA/SCA		1.35	1.43	1.41	1.14	1.93	1.98	1.49	1.85	1.37	3.21

*Significant difference observed among the mean ($P < 0.05$); **highly significant difference observed among mean ($P < 0.01$). GCA: General combining ability; SCA: specific combining ability.

findings of Tiwari and Bhatnagar (1991) who reported that pod shattering is controlled by additive gene action. The preponderance of additive effects observed for days to 50% flowering, days to maturity, number of braches/plant, number of pods/plant, 100 seeds weight is in agreement with the findings of Thakare et al. (2017), Shiv et al. (2011), and Nassar (2013). The significance of additive gene effects for number of seed/pod indicates that additive gene action was more important in the control of this trait. This result is in agreement with the report of Nassar (2013). These results revealed that the traits studied can be exploited effectively by selection in breeding program.

Estimation of GCA effects

Female parents TGX1955-10E and NG/SA/07/100 showed considerable highly significant negative GCA effects for shattering score, plant height, days to 50% flowering and days to maturity, hence they are good combiners. TGX1955-10E and NG/SA/07/100 showed undesirable significant negative GCA effects for number of seeds/pod, pod length and hundred seed weight, hence they are poor combiner. NG/MR/11/11/060 and NGB00/08 showed significant positive GCA effects for number of seeds/pod, pod length, hundred seed weight and grain yield. This suggests that these parents could be utilized in

breeding for improved yield in soybean (Table 3).

Male parents TGX1740-1F and NG/SA/07/055 showed desirable significant positive GCA effects for number of seeds/pod, pod length, hundred seed weight and grain yield and undesirable significant positive GCA effects for shattering score, days to 50% flowering and days to maturity; hence, they are poor combiner for these traits. Male parent NG/AD/11/08/023 had significant negative GCA effects for shattering score, plant height, days to 50% flowering, days to maturity; hence it is a good combiner. Male parent NGB00129 showed highly significant negative GCA effects for number of branches, pod length, hundred seed weight and grain yield. The

Table 3. General combining ability effects for male and female parents for resistance to pod shattering and other agronomic traits in soybean at Samaru in 2016.

Parent	Days to 50% flowering	Plant height (cm)	No. of branches/plant	Number of pod/plant	Number of seeds/pod	Pod length (cm)	Days to maturity	Hundred seed weight (g)	Grain yield (kg)	Pod shattering score
Female										
TGX1955-10E(P1)	-4.73**	-13.15**	-0.67	-17.31**	-0.21*	-0.17*	-7.79**	-0.38**	180.06**	-1.17**
NG/MR/11/11/060(P2)	12.69**	15.85**	0.83	15.85*	0.38**	0.51**	19.46**	1.05**	381.53**	0.50**
NGBOO/08(P3)	5.94**	14.77**	0.92**	0.69	0.54**	0.78**	16.54**	1.62**	108.04**	1.58**
NG/SA/07/100(P4)	-13.90**	-17.48**	-1.08*	0.77	-0.71**	-1.12**	-28.21**	-2.28**	-309.52	-0.92**
SE(gi)	0.96	3.70	0.52	6.55	0.10	0.08	0.63	0.07	2.90	0.04
SE(gi-gj)	1.36	5.23	0.73	9.27	0.13	0.11	0.85	0.09	4.06	0.06
Male										
TGX1740-1F(P5)	10.10**	8.85*	3.3**	28.52**	0.38**	0.756**	17.13**	2.62**	549.65**	0.33**
NG/SA/07/055(P6)	8.69**	-5.81	0.5	-6.56	0.46**	0.731**	16.46**	2.27**	34.62**	0.33**
NG/AD/11/08/023(P7)	-15.90**	-18.06**	-2.3**	-20.48*	-0.71**	-1.20**	-29.63**	-3.55**	534.50**	-1.00**
NGBOO129(P8)	-2.90**	15.02**	-1.4**	-1.48	-0.13	-0.29**	-3.96**	-1.33**	-49.77**	0.33**
SE(gi)	0.96	3.70	0.52	6.55	0.10	0.08	0.63	0.07	2.90	0.04
SE(gi-gj)	1.36	5.23	0.73	9.27	0.13	0.11	0.85	0.09	4.06	0.06

*Significant difference observed among the mean ($P < 0.05$); **Highly significant difference observed among mean ($p < 0.01$).

desirable significant negative GCA effect for pod shattering recorded by female parents, viz., TGX1955-10E and NG/SA/07/100 and a male parent (NG/AD/11/08/023) implied that these genotypes are good general combiners. Significant negative GCA effects recorded for plant height, days to 50% flowering and days to maturity by two female parents and male parent suggests that these genotypes are good general combiners for earliness in hybridization. The preponderance of additive gene effects observed for the studied traits implied that crossing between two good general combiners for pod shattering and other agronomic traits studied will produce hybrids with good specific combining ability. This is according to Daniel et al. (2006) who reported that two good general combiners that are governed by additive \times additive gene actions will

produce hybrids with good specific combining ability.

Estimation of SCA effects

The estimation of SCA effects provides opportunities to isolate crosses where all traits are in the most desirable combinations (Ercan and Mehmet 2005). The specific combining ability effects in Table 4 revealed that crosses NG/SA/07/100 \times TGX1740-1F, NG/MR/11/11/060 \times NGBOO129 and NG/MR/11/11/060 \times NG/AD/11/08/023 showed desirable significant positive SCA effects for number of branches/plant, number of pods/plant, pod length, number of seeds/pod, hundred seeds weight, and grain yield. NGBOO/08 \times NG/SA/07/055, TGX1955-10E \times

NG/SA/07/055, NG/MR/11/11/060 \times NG/AD/11/08/023, NG/MR/11/11/060 \times NGBOO129, NG/SA/07/100 \times TGX1740-1F and TGX1955-10E \times TGX1740-1F showed significant positive SCA effects for days to 50% flowering and days to maturity. NG/MR/11/11/060 \times NG/SA/07/055 showed a desirable significant negative SCA effects for plant height.

The high significant positive SCA effects obtained by cross NG/MR/11/11/060 \times NGBOO129 for grain yield and NG/SA/07/100 \times TGX1740-1F and NG/MR/11/11/060 \times NGBOO129 for number of pods/plant, number of seeds/pod, pod length and hundred seed weight in which NG/MR/11/11/060 and TGX1740-1F were good general combiner and agrees with Cruz et al. (2004) who reported that selection for good estimates of SCA should focus on cross combinations that involve at least

Table 4. Specific combining ability effects for crosses for resistance to pod shattering and other agronomic traits in soybean at Samaru in 2016.

Crosses	Days to 50% flowering	Plant height (cm)	Number of branches/plant	Number of pod/plant	Number of seeds/pod	Pod length (cm)	Days to maturity	Hundred seed weight (g)	Grain yield (kg)	Shattering Score
NGBOO/08 × TG 1740-1F (P3 × P5)	1.65	-8.44	1.25	-5.94	-0.13	-0.07	4.88*	-3.80**	-99.3**	0.42**
TGX1955-10E × TG X1740-1F (P1 × P5)	14.31**	26.15*	0.83	13.73	0.96**	1.58**	26.54**	1.08**	881.2**	0.17
NG/MR/11/11/060 × NG/SA/07/055 (P2 × P6)	3.31*	-16.85*	0.75	6.94	0.29*	0.49**	2.96*	1.60**	142.6**	0.50**
NGBOO/08 × NG/SA/07/055 (P3 × P6)	4.73**	14.90*	0.33	11.44	0.46	0.59**	7.54**	-0.17	594.4**	0.42**
TGX1955-10E × NG/SA/07/055 (P1 × P6)	16.06**	12.81*	2.58	14.60	0.21	0.47**	31.21**	3.83**	-198.1**	-1.50**
NG/MR/11/11/060 × NG/AD/11/08/023 (P2 × P7)	27.56**	32.40*	4.92**	19.35	1.13**	1.69**	52.04**	4.15**	456.2**	1.50**
NG/MR/11/11/060 × NGBOO129 (P2 × P8)	21.23**	43.31**	2.67**	47.02**	0.54**	1.01**	35.04**	3.20**	1146.2**	0.50**
NGBOO/08 × NGBOO129 (P3 × P8)	12.98**	24.40*	1.25	1.19	0.71**	1.00**	27.96**	2.03**	-107.8**	1.42**
NG/SA/07/100 × TG X1740-1F (P4 × P5)	36.15**	41.15**	6.25**	67.98**	1.13**	1.69**	58.63**	6.38**	963.0**	1.92**
SE(Sij)	1.92	7.40	1.04	13.11	0.19	0.16	1.27	0.13	5.74	0.08
SE(Sij-Ski)	2.71	10.46	1.47	18.54	0.26	0.23	1.80	0.19	8.12	0.11

*,**Significant difference at 0.05 and 0.01 levels of probabilities.

one parent which has shown good effect of GCA. Kadams et al. (1999) also reported that hybrid with high SCA effects involved one or both of the good general combiners as parents. The absence of negative SCA effects for shattering score, days to 50% flowering and days to maturity indicates that the hybrids obtained for the present study do not fit in for these traits.

Mean performance values for parents and their F_{1s}

The mean performance of the parental genotypes and their crosses for the studied traits are shown in Table 5. The result revealed that pod shattering score among parents and hybrids ranged from 1 (highly resistant) for NG/AD/11/08/023, NG/SA/07/055, NGBOO129, TGX1955-10E, TGX1955-10E × NG/SA/07/055 and TGX1955-10E × TGX1740-1F to 5 (highly susceptible) for NGBOO/08 and NGBOO/08 × NGBOO129. Grain yield varied from 314.3 for NG/SA/07/100 to

2291.7 kg for hybrid NG/MR/11/11/060 × NGBOO129 with a mean of 1283.04. Days to 50% flowering ranged from 43.3 for NG/SA/07/055 to 61.7 for NG/SA/07/100 × TGX1740-1F with a mean of 50.92. Number of branches varied from 4 for NG/SA/07/100 to 13 for hybrid NG/SA/07/100 × TGX1740-1F with a mean of 6.94. Plant height ranged from 27 cm for hybrid NG/MR/11/11/060 × NG/SA/07/055 to 108 cm for NG/MR/11/11/060 × NGBOO129 with a mean of 62.61. Number of pods/plant ranged from 17 for NG/SA/07/100 to 128 for hybrid NG/SA/07/100 × TGX1740-1F with a mean of 47.25. Hundred seed weight g ranged from 5 g for NG/MR/11/11/060 to 12 g for NG/SA/07/100 × TGX1740-1F. Days to maturity ranged from 87 for NGBOO129 to 104 for NG/MR/11/11/060 × NGBOO129 with a mean of 94.3. Pod length varied from 3 to 4 cm with a mean of 3.7 cm. The highest mean performance recorded by some crosses for grain yield, number of branches, number of pods/plant, and 100 seed weight which was higher than all parental genotypes was similar to the findings of Saul et al.

(2017). This may be as a result of recombination of additive alleles or interaction between two alleles of two different genes due to a wide variation between genotypes of their parents (Marama et al., 2009).

Conclusion

The study has revealed that both additive and non-additive gene effects were important in the inheritance of pod shattering and other agronomic traits. However, additive gene effect was more important implying that selection could be effective. The good general combiners for resistance to pod shattering, plant height, days to 50% flowering and days to maturity were TGX1955-10E, NG/AD/11/08/023 and NG/SA/07/100, while NG/MR/11/11/060, NGBOO/08, TGX1740-1F and NG/SA/07/055 were good general combiner for number of seeds/pod, pod length, hundred seed weight and grain yield. Therefore, these parents are

Table 5. Mean performances of parents and the F₁s evaluated for Resistant to Pod Shattering and other agronomic traits using field screening methods.

Genotype	Days to 50% flowering	No. of branch/plant	Plant height (cm)	No. of pods/plant	No. of seeds/pod	Pod length (cm)	Days to maturity	Hundred seeds weight (g)	Grain yield (kg)	Pod shattering score
NG/AD/11/08/023	46.7	6	59.7	59.0	3.0	4.1	103.3	7.2	1822.4	1
NGBOO129	48.7	6.3	79.0	33.3	2.7	3.7	87.0	9.2	841.9	1
TGX1955-10E	45.3	8.0	52.3	30.0	3.0	4.4	90.3	10.2	921.1	1
NG/SA/07/055	43.3	5.7	72.7	43.0	2.7	3.7	90.7	10.4	902.6	1
NG/SA/07/100	60.7	4.3	54.0	17.0	2.7	3.2	94.0	12.0	314.3	3
NGBOO/08	44.3	6.0	76.7	31.7	2.7	4.2	93.3	7.8	1723.9	5
TGX1740-1F	48.3	7.0	61.7	29.0	2.7	3.8	98.3	11.0	1442.3	2.3
NG/MR/11/11/060	59.3	6.7	62.0	18.7	3.0	4.4	90.3	5.3	823.3	4.3
NG/MR/11/11/060 × NG/AD/11/08/023	53.7	7.7	64.3	43.7	2.0	2.9	95.3	6.9	1117.0	2.7
NG/MR/11/11/060 × NG/SA/07/055	54.0	6.3	27.3	55.0	2.3	3.7	92.3	10.2	1372.5	3
NG/MR/11/11/060 × NGBOO129	60.3	6.3	108.3	90.3	2.0	3.2	104.0	8.2	2291.7	3
NG/SA/07/100 × TGX1740-1F	61.7	12.7	66.7	128.7	2.0	3.3	101.0	12.0	2016.9	3
NGBOO/08 × NG/SA/07/055	48.7	6.0	58.0	44.3	2.7	4.0	94.0	9.0	1550.8	4
NGBOO/08 × NGBOO129	45.3	5.0	88.3	29.3	2.3	3.4	94.0	7.6	764.2	5
NGBOO/08 × TGX1740-1F	47.0	9.7	49.3	54.7	2.0	3.4	92.0	11.0	1372.1	4
TGX1955-10E × NG/SA/07/055	49.3	6.7	28.0	39.3	1.7	3.0	93.3	11.0	470.2	1
TGX1955-10E × TGX1740-1F	49.0	7.7	56.0	56.3	2.3	4.1	89.3	8.6	2064.5	1
CV%	6.84	30.55	23.28	59.96	27.14	14.40	2.79	2.79	0.74	12.38
Mean	50.92	6.94	62.61	47.25	2.45	3.67	94.27	9.27	1283.04	2.67
LSD	5.79	3.52	24.24	47.13	1.11	0.88	4.37	0.43	15.80	0.55

recommended as source of pod shattering resistance and high yielding for soybean breeding program, while the good crosses for SCA effects for yield and yield components are NG/SA/07/100 × TGX1740-1F, NG/MR/11/11/060 × NGBOO129 and NG/MR/11/11/060 × NG/AD/11/08/023. These crosses should be advanced for selection in later generations.

REFERENCES

- Asian Vegetable Research Development Centre (AVRDC) (1979). International Co-operator's Guide. Suggested cultural practices for soybean. Asian Vegetable Research Development Centre, Taiwan. Robinson, H.F. No.79-112.
- Bara N, Khare D, Srivastava AN (2013). Studies on the factors affecting pod shattering in soybean. *Ind. J. Genetics*. 73(3):270-277.
- Context Network and Sahel Capital (2016). Early Generation Seed System, Study feed the future: Building Capacity for African Agricultural Transformation (Africa Lead II). pp. 51-52.
- Caviness CE (1969). Heritability of pod dehiscence and its association with some agronomic characters in soybeans. *J. Crop Sci.* 9:207-209.
- Cruz CD, Regazzi AD, Carnero PCS (2004). Modelos biometricos aplicados ao melhoramento genetico. Vicosa: Universidade Federal de Vicosa. P 480.
- Daniel OI, Oloyede HT, Adeniji OT, Ojo DK, Adegbite AE (2006). Genetic analysis of earliness and yield in elite parental lines and hybrids of tropical maize (*Zea mays* L.). *J. Genet. Breed.* 60:289-296
- Ercan C, Mehmet AA (2005). Combining Ability and Heterosis for Grain Yield and Some Yield Components in Pea (*Pisum sativum* L.). *Pak. J. Biol. Sci.* 8(10):1447-1452.
- Haruna M (2010). Genetic analysis of resistance to pod shattering in soybean (*Glycine max.* (L.) Merrill). An MSc. Thesis Submitted to the Department of Crop and Soil Sciences, Faculty of Agriculture, Kwame Nkrumah University of Science and Technology, Kumasi. pp. 50-55.
- Helms TC (1994). Greenhouse and field evaluation of pod dehiscence in soybean. *Can. J. Plant Sci.* 74(4):699-701.
- Indu B (2014). Gene Action for Yield and Related Traits in Soybean and Development of Interspecific Hybrids Involving Wild Species. A PhD. Thesis Submitted to the Department

- of Plant Breeding and Genetics, Chaudhary Sarwan Kumar University, India. pp. 82-83.
- Kadams AM, Ugherughe PO, Abakura J.B (1999). Genetic Correlation and Path Coefficient International Analysis of Yield and its Components in Wheat. Nig. J. Trop. Agric. 1:107-114.
- Krisnawati A, Adie MM (2016). Pod shattering resistance in different soybean genotypes. Proceedings of ILETRI National Seminar. ILETRI, IAARD, Malang. [Indonesian].
- Maramé F, Desalegne L, Fininsa C, Sagvald R (2009). Genetic analysis for some plant and fruit traits, and its implication for a breeding progeny of hot ppepper. Hereditas 146:131-140.
- Nassar MAA (2013). Heterosis and combining ability for yield and its components in some crosses of soybean. Aust. J. Basic Appl. Sci. 7(1):566-572.
- Saul EM, Mildred OS, Kassim S, Boris A, Patrick R, Paul G, Wales S, Richard E (2017). Estimates of combining ability and heritability in cowpea genotypes under drought stress and non-stress conditions. J. Plant Breed. Crop Sci. 9(2):10-18.
- Shiv D, Sajeela N, Bhadana VP, Sharma PR (2011). Gene action for yield and its components in soybean (*Glycine max* (L.) Merrill). Int. J. Plant Res. 24(1):89-92.
- Singh RK, Chaudhary BD (1977). Biometrical Methods in Quantitative Genetic Analysis. Kalyani Pub., Ludhiana
- Thakare DS, Chimote MP, Deshmukh MP, Bhailume MS, Adsul AT (2017). Inheritance of yield and yield components in Soybean (*Glycine max* (L.) Merrill.) J. Plant Breed. 8(1):176-181.

The background of the entire page is a photograph of a garden. In the foreground, a pair of bright blue rubber boots sits on a reddish-brown brick path. To the right, a woven basket is filled with several onions. The background is a soft-focus view of green plants and a wooden fence.

Journal of Plant Breeding and Crop Science

Related Journals Published by Academic Journals

- *African Journal of Agricultural Research*
- *Journal of Horticulture and Forestry*
- *Journal of Cereals and Oilseeds*
- *International Journal of Livestock Production*
- *International Journal of Fisheries and Aquaculture*
- *Journal of Development and Agricultural Economics*

academicJournals