

A syringe is shown injecting a clear liquid into the stem of a large, ripe red strawberry. The strawberry is placed on a wooden surface. A yellow sign with the word 'DISURB' is pinned to the surface behind the strawberry. The background is a wooden table.

International Journal of Plant Physiology and Biochemistry

Volume 9 Number 1, April 2017

ISSN 2141-2162



*Academic
Journals*

ABOUT IJPPB

The **International Journal of Plant Physiology and Biochemistry (IJPPB)** (ISSN 2141-2162) is published Monthly (one volume per year) by AcademicJournals.

International Journal of Plant Physiology and Biochemistry (IJPPB) provides rapid publication (monthly) of articles in all areas of the subject such as plant hormones, seed biology, plant DNA repair, Concepts of target cells in plant differentiation etc.

The Journal welcomes the submission of manuscripts that meet the general criteria of significance and scientific excellence. Papers will be published shortly after acceptance. All articles published in IJPPB are peer-reviewed.

Contact Us

Editorial Office: ijppb@academicjournals.org

Help Desk: helpdesk@academicjournals.org

Website: <http://academicjournals.org/IJPPB>

Submit manuscript online <http://ms.academicjournals.me/>

Editors

Prof Dr. Ishrak Khafagi

*Faculty of Science,
Suez Canal University,
Ismailia,
Egypt*

Prof. Mohamed Mahgoub Azooz

*Biology Department
Faculty of Science,
King Faisal University,
Saudi Arabia.*

Dr. Bansal Parveen

*National Institute of Ayurvedic Pharmaceutical
Research
Moti Bagh Road,
Patiala-(Punjab) India.*

Prof. Bechan Sharma

*Department of Biochemistry,
University of Allahabad,
Faculty of Science,
Allahabad-211002,
India.*

Editor Board

Prof. Weimin Zhang

*Guangdong Institute of Microbiology 100 Central
Xian lie Road, Guangzhou, Guangdong 510070,
China.*

Dr. Xu Suxia

*Fujian Institute of Subtropical Botany, 780-800,
Jiahe Road,
Xiamen,
China361006,
China.*

Dr. Adaku Vivien Iwueke

*Department of Biochemistry,
Federal University of Technology,
Owerri
Nigeria.*

Ass. Prof. Turgay CELIK

*Gulhane Military Medical Academy,
School of Medicine,
Department of Cardiology,
Turkey.*

Dr.Topik Hidayat

*Department of Biology Education
Indonesia University of Education (UPI)
Jalan Dr. Setiabudhi 229 Bandung 40154 Indonesia
Indonesia.*

Dr.Tariq Mahmood

*Quaid-i-Azam University,
Department of Plant Sciences,
Quaid-i-Azam University,
Islamabad,
Pakistan.*

Dr.Neveen B. Talaat

*Department of Plant Physiology,
Faculty of Agriculture,
Cairo University,
Egypt.*

Dr. Sudhamoy Mandal

*Fulbright Visiting Fellow
Department of Plant Pathology
University of Nebraska Lincoln USA*

Asso. Prof. Chankova Stehka Georgieva

*Central Laboratory of General Ecology,
Bulg Acad
Sci 1113 Sofia, 2 Gagarin str,
Bulgaria.*

Shijie Han

*Center of Forestry,
Institute of Applied Ecology,
Chinese Academy of Sciences,
72 Wenhua Road, Shenyang City,
Liaoning Province 110016, PR China.*

Szu-Chuan Shen

*Department of Medical Nutrition,
I-Shou University Yanchao Township,
Kaohsiung County 824, Taiwan.*

Seddik Khennouf

*Dept of Biology,
Faculty of Science
University Ferhat Abbas,
SETIF, 19000, ALGERIA.*

Saranyu Khammuang

*Department of Chemistry,
Faculty of Science,
Mahasarakham University,
Thailand.*

Samir Ranjan Sikdar

*Bose Institute
P-1/12, C.I.T. Scheme VII M,
Kolkata 700 054,
India.*

Dr. M. Abdul Salam

*Department of Agronomy,
College of Agriculture,
Kerala Agricultural University,
Vellayani 695 522, Trivandrum, Kerala.*

Dr.Saeed Aminzadeh

*National Institute of Genetic Engineering and
Biotechnology (NIGEB)
Shahrak-e-Pajooheh Km 15, Tehran-Karaj Highway,
Tehran, I.R.
Iran.*

Dr.Ruzica Stricevic

*Faculty of
Agriculture,
University of
Belgrade
Nemanjina 6, Zemun, 11080,
Serbia.*

Rumbos Christos

*University of Thessaly
Fytokoy Str, 384 46
Volos, Greece.*

Dr. Özge Zencir

*Kemah Vocational Training School,
Erzincan University,
Kemah, Erzincan,
Turkey.*

Riyazali Zafarali Sayyed

*SI P Arts, GBP Science & STSKVS Comm. College
SHAHADA Dist
Nandurbar, Maharashtra,
India.*

Raul Rodriguez-Herrera

*Universidad Autonoma de
Coahuila School of Chemistry
Blvd. V. Carranza y González Lobo s/n Col
República Saltillo Coahuila México.*

Dr. A. H. M. Mahbubur

Rahman *Rajshahi University,
Bangladesh. Department of
Botany, University of Rajshahi,
Bangladesh.*

Paul S. Marley

*Department of Crop Protection,
IAR/FOA Ahmadu Bello University,
P.M.B. 1044, Samaru, Zaria,
Nigeria.*

Patrick Addo-Fordjour

*Department of Theoretical and Applied Biology,
Kwame Nkrumah University of Science
And Technology (Knust),
Kumasi,
Ghana.*

Battu.Prasanna

Reddy Nosch Labs
*Pvt Ltd Hyderabad,
India.*

Noureddine Benkeblia

*UWI - Department of Life Sciences
Mona Campus,
Kingston 7,
Jamaica.*

Keutgen, Norbert

*Uniwersytet Technologiczno-Przyrodniczy
im. Jana i Jędrzeja Śniadeckich w Bydgoszczy
Kadra Katedry Fizjologii Roslin
(Institute of Plant Physiology)
ul. Bernardynska 6/8, 85-029 Bydgoszcz,
Poland.*

Nicholas E. Korres

*University College Cork,
Environmental Research Institute.
Lee Road, Cork,
Ireland.*

Dr Naveen Kumar

*University of Florida
2685 SR 29 N SWFREC/IFAS/UFL,
Immokalee, FL34142,
USA.*

Dr Modala Venkateswarlu

*Seribiotech research Laboratory,
Kodathi Carmelaram post,
Bangalore.*

Mirza Hasanuzzaman

*Department of Agronomy,
Faculty of Agriculture,
Sher-e-Bangla Agricultural University,
Dhaka-1207, Bangladesh.*

Maybelle Gaballah

*National Research Centre,
El Behoos street, Dokki, Cairo.*

Mauro Guida Santos

*Universidade Federal de Pernambuco Street
Moraes Rego – CDU – CCB – Botany Department,
s/n. 50670-901. Pernambuco State, Brazil.*

Marcelo Rodrigo Pace *University of Sao Paulo Rua do
Matão, 277, Cidade Universitária São Paulo, Brazil.*

Marcelo Francisco Pompelli

*Federal University of Pernambuco
Department of Botany, Profª Moraes Rego Av.,
Recife – PE – Brazil, 50670-901.*

Luca Catalina Mariana

*University of Bucharest, Faculty of Biology, Dept
of Biochemistry and Molecular Biology
Spl. Independentei, no.91-95, Bucharest 5,
Romania.*

Lin Wang

*Institute of Biostatistics, Fudan University
220 Handan Road, Shanghai 200433,
China genetics, microbiology
China.*

Li Qiang

*Institute of karst geology, MLR
50 Qixing Road,
China.*

Dr. Ayanakumar Kumar

*C.Abdul Hakeem College of Engg. & Tech.,
Melvisharam-632 509, Vellore Dist, Tamil Nadu,
INDIA.*

P. Krishnamoorthy

*P.G. AND RESEARCH DEPARTMENT OF ZOOLOGY
RAJAH SERFOJI GOVT. COLLEGE.
India.*

Hare Krishna

*Central Institute of Temperate Horticulture-
Regional Station,
Mukteshwar-263 138, District-
Nainital, Uttarakhand, India.*

K.G. Mandal

*Directorate of Water Management (formerly
Water Technology Centre for Eastern Region)
Indian Council of Agricultural Research
C.S. Pur, Bhubaneswar-751023, ORISSA, INDIA.*

Dr. Jukta Adhikari

*Presidency College
86/1, College Street, Kolkata – 700 073, India.*

Jorge Teixeira

*Botany Department, Faculty of Sciences,
University of Porto,
Edifício FC4, Rua do Campo Alegre, S/N, 4169-
007
Porto, Portugal.*

Johnson Toyin Fasinmirin

*Federal University of Technology, Akure,
Nigeria Department of Agricultural
Engineering, FUT, P.M.B. 704, Akure, Ondo
State, Nigeria..*

Joel K. Ransom

*North Dakota State University
166 Loftsgard Hall, Department of Plant
Sciences, NDSU Dept. 7670, PO Box 6050,
Fargo, ND 58108-
6050.*

João Claudio Damasceno de Sá

*UENF
Av. José Carlos Pereira Pinto, 39. Pq. Vicente
Dias. Campos RJ. Brazil.*

Jalal Jalali Sendi

*University of Guilan
Department of Plant Protection, university of
Guilan, Rasht, Iran.*

Iúri Drumond Louro

*Universidade Federal do Espírito Santo
Rua Horácio Andrade de Carvalho, 210, Victoria,
ES, 29052-620, Brazil.*

Hong Bo Guo

*Northwest A and F University
22 Xinong, Yangling 712100, Shaanxi, PR China.*

Harsukh P. Gajera

*Junagadh Agricultural University Department of
Biochemistry, College of Agriculture, JAU,
Junagadh- 362 001, Gujarat, India.*

Hanan Abdel Fattah El-Sadawy

*National Research Center
El-Buhoth St., Dokki, Giza, Egypt.*

Assist. Prof .Azime KÜÇÜKGÜL GÜLEÇ *Tunceli
University Fisheries Faculty 62000, Tunceli/TURKEY.*

Greg T. Hannig

*DuPont
1090 Elkton Road Newark, DE 19711.*

Gilberto Santos Andrade

*Instituto de Biotecnologia Aplicada a Agropecuária
(BIOAGRO), Departamento de Biologia Animal,
Universidade Federal de Viçosa, Viçosa, MG
36571- 000, Brazil.*

Dr. T. Muthukumar

*Department of Botany, Bharathiar
University Coimbatore -641 046, Tamilnadu,
India.*

Kunjupillai Vijayan

*Institute of Plant and Microbial Biology
Academia Sinica, Taipei, Taiwan-115, ROC.
Taiwan.*

Badre Alam

*National Research Centre For Agroforestry
Gwalior Road, Jhansi-284003, U.P., India.*

Abeer Essam El-Din Mahmoud

*Biochemistry Department
Genetic Engineering & Biotechnology Division
National Research Center El Tahrir St., El
Dokki 12622, Cairo, Egypt.*

Qazi Fariduddin

*Aligarh Muslim University
Department of Botany, Aligarh 202 002, India.*

Darmawan Darma

*Faculty of Agriculture, Andalas University
Kampus Limau Manis Padang-25163, Indonesia.*

Barbara Chaves

Institute for Agricultural and Fisheries Research.

Sudhamoy Mandal

*Central Horticultural Experiment Station
(ICAR) Aiginia, Bhubaneswar,
PIN-751019.*

Cavit Bircan

*Adnan Menderes University
Faculty of
Agriculture
Department of Food
Engineering
09100/Aydin/Turkey*

Carlos Alberto Ortega-Ojeda

*Central University of Ecuador. Faculty of Agriculture Sciences. Quito, Ecuador
Calle 12 # 29 B - 78, Apto. 102 F, Unidad Residencial
Colseguros, Cali, Colombia.*

Brian Wade Jamandre

*National Taiwan University
Rm. 622, life science bldg., NTU, no.1,
sec.4, Roosevelt rd. Taipei 10617, Taiwan (ROC).*

Bitá Naseri

*Agricultural Research Institute
Department of Plant Protection, Agricultural Research Institute, PO Box 45195474, Zanjan, Iran..*

Behzad Kaviani

*Adeyemi Oluyomi Stephen Bells
University of Technology
Chemical Sciences Department, Km 8 Ididroko Road, Ota, Ogun
State, Nigeria.*

Ajayi Adedayo Olajide

*Adekunle Ajasin University
Dept. of Microbiology, P.M.B 01, Akungba-Akoko, Ondo State, Nigeria.*

Alexandre Igor Azevedo Pereira (Pereira, A.I.A.)

Universidade Federal de Viçosa, Departamento de Biologia Animal, Programa de Pós-Graduação em Entomologia. 36570-000, Viçosa, Minas Gerais State, Brazil.

Gilberto Santos Andrade

Instituto de Biotecnologia Aplicada a Agropecuária (BIOAGRO), Departamento de Biologia Animal, Universidade Federal de Viçosa, Viçosa, MG 36571-000, Brazil.

Pradeep. A.R., Ph.D

*Seribiotech Research Laboratory
Carmelaram.P.O; Bangalore, INDIA.*

Azamal Husen

*University of Gondar
Department of Biology, Faculty of Natural Sciences, University of Gondar
P.O. Box #196, Gondar, Ethiopia.*

Muhammad Aslam

*University College of Agriculture, Bahauddin Zakariya University
Multan 60800, Pakistan.*

Autumn J. Smith

Sam Houston State University, Texas.

La Sara

*Haluoleo University
Kampus Baru Tridharma, Kendari, Southeast Sulawesi, Indonesia.*

Aliyu Mohammed

Department of Human Physiology, ABU, Zaria. Nigeria.

Prof. EL-Said Ahmed AL-Sayed Ragab *National Research Institute of Astronomy and Geophysics, 11421, Helwan, Egypt.*

Shnoudy Anwar Bakhoun

National Institute of Oceanography & Fisheries (NIOF), Egypt.

Antonio Americo Barbosa Viana

*Embrapa Recursos Geneticos e Biotecnologia PBI-LPP1
PqEB Final W/5 Norte, Brasilia, DF – Brazil*

Dr.Shirish Rajmalwar

National Research Laboratory for Conservation, Shirish Rajmalwar, LIG Plot No. 43, Mhada colony, Wardha – 442001, (MS) India.

Dr. Amaresh Chandra

*Universidade Federal de Viçosa, Departamento de Biologia Animal, Programa de Pós-Graduação em Entomologia.
36570-000, Viçosa,
Minas Gerais State,
Brazil.*

Dr. Atul Kumar

*GB PANT University of Agriculture & Technology
Department of Basic Science, College of Forestry &
Hill Agriculture, HILL CAMPUS, PO Ranichauri, Tehri
Garhwal, Uttarakhand State, India.*

Prof. Levenko Boris

*Natl. Botanical Gardens, NAS of Ukraine
01014 Kiev, 1 Timiryasevska st. Ukraine.*

Dr. Dionisio G. Alvindia

*Bureau of Postharvest Research and Extension
CLSU Compound, Science City of Munoz,
Nueva Ecija 3120, Philippines.*

Dr. Bhoopander Giri

*University of Delhi
Department of Botany, SSNC (University of
Delhi) Alipur, Delhi 110036, India.*

Dr. Anjali Sood

*University of Delhi
Department of Botany, University of Delhi, Delhi-
110 007, INDIA.*

Dr. A. K. Verma

*G.B. Pant University of Agriculture & Technology,
Pantnagar, Department of Biochemistry, College
of
Basic Sciences,
India.*

Dr. Anjana Jajoo

*School of Life Science, Devi Ahilya
University, Indore, DAVV
Khandwa Road campus, Indore 452 017,
M.P., India.*

Dr. Deepak Ganjewala

*Vellore Institute of Technology University
55 Thennaraam Street, Vellore-632 014 (T.N.),
India.*

Dr. Geetha Govind

*Max-Planck-Institute for Chemical Ecology
Hans-Knöll Straße 8, 07745 Jena, Germany.*

Dr. Hossam El-Din Saad El-Beltagi

*Biochemistry Department, Faculty of Agriculture,
Cairo University, Giza, Egypt, P.O.Box 12613 Egypt..*

Prof. Dr. Md. Shahidul Haque

*Dept. of Biochemistry and Molecular Biology
University of Rajshahi, Rajshahi-6205,
Bangladesh Bangladesh.*

DR. P.K.NAGAR

*Retired Senior Scientist, IHBT, Palampur, (H.P.),
B.21/115-10A Batuk Dham Colony, Kamachha,
Varanasi 221 010, INDIA.*

Dr. Satyawati Sharma

*Indian Institute of Technology
Centre for Rural Development & Technology, IIT
Delhi-110016
Biomass Production on waste land, India.*

Dr. Uğur Çakılcıoğlu

*Firat University Elazığ/TURKEY
Cumhuriyet M. Malatya C.
No:50/A.*

Prof. Abdelrhani Elachqar

*Faculty of Sciences Dhar El Mahraz, Fez,
Morocco BP 1796, Fès-Atlas, Fès, Maroc,
Morocco.*

Ass. Prof. Jianfeng Xu

*Arkansas State University
PO Box 639, State University, AR 72467 USA.*

Ass.Prof. Jin Xu

*Center for Agricultural Resources Research,
Institute
of Genetics and Developmental Biology,
Chinese Academy of Sciences
Huaizhong RD 286, Shijiazhuang, HeBei, China.*

José Carlos Rebuglio Velloso Ph.D

*PARANÁ STATE UNIVERSITY OF PONTA GROSSA
(Universidade Estadual de Ponta Grossa – UEPG)
General Carlos Cavalcanti Avenue, 4748,
Uvaranas, Ponta Grossa/PR – PO box 84030-900*

Dr. Krouma Abdelmajid

*Centre of Biotechnology, Borj Cedria Ecopark
BP 901, Hammam-Lif 2050, Tunisia
College of Science and Arts, Qassim University, BP
53, Al-Rass 3330353, Qassim, Saudi Arabia
Saudi Arabia*

Dr. Majid Rostami

*Malayer University
Department of Agriculture and Natural
Resources, Postal code: 65719- 95863,
University of Malayer Malayer, Iran.*

Dr. Mohammad Nasir Khan

*Aligarh Muslim University, Aligarh, INDIA Plant
Physiology Section, Department of
Botany, Aligarh Muslim University, Aligarh-
202 002, U.P., India.*

Prof. N.K.Matta

*Kurukshetra University
Department of Botany, Kurukshetra
University, Kurukshetra 136119, INDIA.*

Dr. Naceur Djebali

*Centre of Biotechnology Borj-Cedria
(CBBC) BP 901, Hammam-Lif 2050
Tunisia.*

Dr. Nader Chaparzadeh

*Azarbaijan University of Tarbiat Moallem,
Tabriz, Iran.*

Nautiyal Prakash Chandra

*Directorate Of Groundnut Research (Icar)
Post box, No. 5, Ivnagar Road, Junagadh-362001,
Gujarat, India.*

Prof. Hussein Fawzy Hussein Abouzienna

*National Research Center
Botany Department, National Research
Center, Elbhoss Street, Dokki, Cairo, Egypt.*

Dr. D.E. Chandrashekar Rao

*National Research Council Canada / Plant
Biotechnology Institute (NRC-PBI)
110 Gymnasium Place / Saskatoon, Saskatchewan
S7N 0W9
Canada.*

Dr. S.R Madhan Shankar

*PSG College of Arts & Science
Civil Aerodrome Post, Coimbatore-641 014, India.*

Prof. Dr. Safdar Hussain Shah

*Institute of Biotechnology and Genetic Engineering
NWFP, Agricultural University Peshawar, Pakistan.*

Prof. Dr. Md. Shahidul Haque

*Dept. of Biochemistry and Molecular Biology
University of Rajshahi, Rajshahi-6205, Bangladesh.*

Dr. Sivakumar Swaminathan

*Iowa State University (ISU)
G-319, Agronomy Department, ISU, Ames, Iowa -
50011, USA*

Dr. Subrahmanyam Desiraju

*Directorate of Rice Research (ICAR)
Plant Physiology Division, Rajendranagar,
Hyderabad-500030, A.P. India.*

Dr. Tariq Aziz Dr. Deepak Ganjewala

*University of Agriculture, Faisalabad, Sub-Campus
Depalpur, Dist. Okara, Pakistan.*

Dr. Thangavel

Palaniswamy SUN
YAT-SEN UNIVERSITY
GUANGZHOU, PR CHINA.

Yi-Ping Chen Ph.D

*Institute of Earth Environment,
Chinese Academy of Science Fenghui S.R, 10,
Xi'an Hi-Tech Zone, Xi'an, Chnia.*

Saha Prasenjit

*The Samuel Roberts Noble Foundation
2510 Sam Noble Parkway, Ardmore, Ok USA.*

Abdul Khaliq Ph.D

*Department of Agronomy
University of Agriculture
Faisalabad 38040, Pakistan.*

Dr. Arafat Abdel Hamed abdel Latef

*Assistant Professor of Plant
physiology Botany Department
Faculty of Science at Qena
South Valley
University Egypt.*

Dr. Ahmad Bybordi

*Research Center of Agriculture and Natural
Resources of East Azarbaijan
Member of Scientific Board of Research Center of
Agriculture and Natural Resources of East
Azarbaijan,
Tabriz. Iran.*

Dr. Arijit Sinhababu

Bankura Christian College (under –The University of Burdwan)

Department of Botany, Bankura Christian College, P.O. + Dist. Bankura, Pin.-722101, West, Bengal, India.

Dr. Maria Alejandra Equiza

*University of Alberta,
4-51 Earth Sciences Building, Dept. Renewable Resources, University of Alberta, Edmonton, AB T6G 2E3, Canada*

Dr. Suphla Bajpai Gupta

Indian Institute of integrative Medicine –CSIR, Scientist, Plant biotechnology division, Canal Road, Jammu, Jammu & Kashmir, India-180001. India.

Dr. Linga R Gutha

*Washington State University,
2410 N Bunn Road, Prosser, WA 99350, USA.*

Dr. Medhat Mekhail Tawfik

*National Research Center,
El Bohooth Str. Dokki, Giza. Egypt, PO Box 12311, Egypt.*

Dr. Rafiq Islam

The Ohio State University South Centers, 1864 Shyville Road, Piketon, OH 45661.

Dr. Rakesh Kumar

*V.S.P. Govt. P.G. College, Kairana, Muzaffarnagar (Uttar Pradesh),
Department of Botany, V.S.P. Govt. P.G. College, Kairana, Muzaffarnagar (Uttar Pradesh), India-247774.*

Dr. Ivan Sestari

*University of São Paulo,
Av. Pádua Dias, 11: CP 9. CEP 13418-900.*

D.Sc. Rachel Fatima Gagliardi

*State University of Rio de Janeiro,
Rua São Francisco Xavier, 524 – PHLC sala 602.*

Dr. Ullas Pedmale

*Salk Institute for Biological Studies,
10010 N Torrey Pines RD, La Jolla, CA 92037.*

Dr. Allah Bakhsh Dr. Deepak Ganjewala

*Department of Field Crops, Faculty of Agriculture, University of Ankara,
Apartment No. 12/10, Sanatorym Caddesi, Kalaba, Kecioren, Ankara,
Turkey.*

Dr. Atilgan Atilgan

*Suleyman Demirel University, Agriculture Faculty,
Department of Agricultural Structures and Irrigation, Isparta,
Turkey.*

Mr. Andrej Pilipovic

*University of Novi Sad – Institute of Lowland Forestry and Environment,
Antona Cehova 13, 21000 Novi Sad, Serbi.*

Dr. Zulfiqar Ahmad Saqib

*Institute of Soil and Environmental Sciences,
University of Agriculture, Faisalabad,
Civil Line Road, Faisalabad,
Pakistan.*

MS. C. Mehrnoush Eskandari Torbaghan

*North Khorasan Agricultural & Natural Resource Research Center (NKANRRC)
P.O. Box: 94155-1416, No. 52, Hassan Kallate Alley, Tarbiyat St., Mother Sq. Bojnourd, Iran.*

Dr. Vinod Kumar

*Department of Zoology & Environmental Science, Gurukula Kangri University,
Haridwar-249404 (UK), India.*

Dr . Panda Tribhubana

*Kalahandi Institute for Tribiology and Ethnobiology(KITE),
At-Jilingdar, PO-Deydar, Dist-Kalahandi,Odisha, India,766014, India*

Dr. Sabarinath Sundaram

*Institute of Developmental and Molecular Biology,
Texas A&M University,
Biological Sciences
Building West Suite
403.*

Dr. Diogo Pineda Rivelli

*University of São Paulo,
Av. Prof. Lineu Prestes 580, São Paulo, SP, 05508-000.*

Dr. Qiang Wang

*Virginia Tech, 427
Latham Hall.*

Dr. Foteini Hassiotou

*University of Western Australia, 35
Stirling Highway, Crawley, WA
6009, Australia.*

Dr. Nivedita Sahu

*Indian Institute of Chemical
Technology, Chemical Biology
Laboratory
(NaturalProductChemistry), Uppal
Road, Hyderabad-500607.*

Dr. Mohammad Anwar Hossain *Bangladesh
Agricultural University, Assistant Professor,
Dept. of Genetics and Plant Breeding,
Bangladesh Agricultural University,
Mymensingh-2202, Bangladesh.*

Dr. Ahmad Ali

*National Institute of Pharmaceutical Education
& Research,
Dept of Biotechnology, NIPER, Jandaha Road,
Hajipur, Bihar, India, Pin – 844102, India.*

Mr. Karthikkumar V

*Annamalai University,
Department of Biochemistry & Biotechnology.*

Dr.K.Rajendiran

*Dept of plant science, Tagore Govt. college,
9, 4th cross, Tagore Nagar, Pondicherry – 605
008, India.*

Dr. V. Balakrishnan *K.S.Rangasamy College of
Technology, Department of Biotechnology, KSR Kalvi
nagar, Tiruchengode- 637215, Tamilnadu, India.*

Dr. NourAli Sajedi

*Department of Agronomy and plant Breeding,
Islamic Azad University, Arak Branch, Arak,
Iran.*

(Dr) Ms. Rachel Predeepa

*Not Applicable ,
2/387 Gokul Nagar, Kannanenthal Madurai.*

Dr. Rajendra Gyawali

*Department of Pharmacy and Biology, Kathmandu
University,
Dhulikhel,
Nepal.*

Ms. Rocheli de Souza

*UFRGS,
Porto Alegre,
Brazil.*

Dr. Om Prakash Verma

*Sam Higginbottom Institute of Agriculture,
Technology & Sciences (Formerly Allahabad
Agricultural Institute), Allahabad, U.P.,
Department of Molecular & Cellular Engineering,
Jacob School of Biotechnology & Bioengineering,
India.*

Dr. Ashwani Kumar

*JMIT, Radaur,
Department of Biotechnology, JMIT, Radaur-
135133, Haryana, India.*

Dr. Sarfaraz F. A. Al-Bamarny

*University of Duhok, College of Agriculture, Dept.
of Horticulture,
Duhok, Iraqi Kurdistan Region,
Iraq.*

Prof. Wafaa Mohamed Shukry Abdel Meamem

*Dammam University - Saudi Arabia,
Faculty of Science for Girl.
Biology Department,
P.O.Box: 838 Dammam 31113,
Saudi Arabia.*

Dr. Stephka G. Chankova

*Institute of Biodiversity and ecosystem Research,
BAS,
2 Gagarin str, 1113
Sofia, Bulgaria.*

Dr. Nana Ewusi-Mensah

*Kwame Nkrumah University of Science and Technology,
Dept. of Crop and Soil Sciences, Faculty of Agriculture, KNUST, Kumasi.*

Dr. Mukesh Lokanath Chavan

*K.r.c. College of horticulture, arabhavi 591 310, karnataka,
University of horticultura sciences, bagakot, India.*

Dr. Maiti Parthapratim

Dept. of Botany Midnapore College, Midnapore-721101, Paschim Medinipur, West Bengal, India.

Mr. Mohammad Anwar Hossain

*Kagawa University (Present), Bangladesh Agricultural University (Permanent)
Lab. of Plant Stress Responses, Faculty of Agriculture, Kagawa University, Miki-cho, Kitagun, Kagawa 761 0795, Japan.*

Dr. Antonia Tathiana Batista Dutra

*Universidade Federal do Ceará,
Av. Humberto Monte s/n – Pici Bloco 907, laboratório 1080.*

Dr. Kuntal Das

*St. John's Pharmacy College,
#6, II Main, 9th Cross, Vijayanagar, Bangalore-104. India.*

Dr. Amitava Rakshit

*Banaras Hindu University,
Department of Soil Science & Agril Chemistry.*

Dr. Kranthi Kiran Mandadi

*Texas A&M University,
2132 TAMU, Peterson-Rm408, College Station, Texas-77840, USA.*

Dr. Monica Butnariu

*Banat's University of Agricultural Sciences and Veterinary Medicine from Timisoara,
Chemistry and Vegetal Biochemistry Department, Calea Aradului no.119, 300645 Timisoara, Romania.*

Dr. Ahmad Bybordi

*East Azarbaijan Research Center for Agriculture and Natural Resources,
Tabriz, Iran.*

Dr. Haiwei Gu

903 Fifth St., West Lafayette, IN 47906.

Dr. Hu Yanbo

*Northeast Forestry University,
26# Hexing Road, Xiangfang District, Harbin city, 150040, P.R., China.*

Dr. Arash Kianianmomeni

Institute of Biology / Humboldt-University Berlin, Invalidenstr. 42.

Dr. Zvonko Pacanoski

Faculty for Agriculture Sciences and Food, Boul. Aleksandar Makedonski bb, 1000 Skopje, R.of Macedonia..

Dr. Lingjuan Zheng

Department of Organismic Biology, University of Salzburg, Hellbrunnenstraße 34, 5020, Salzburg, Austria.

Dr. Md. Mokter Hossain

Department of Horticulture, Bangladesh Agricultural University, Mymensingh-2202, Bangladesh.

Dr. Forouzandeh Soltani

Department of Horticultural Sciences, College of Agriculture and Natural Resources, University, of Tehran, Daneshkadeh Street, Karaj 31587-11167, Iran.

Dr. M.C.Harish

*Bharathiar University,
Department of Biotechnology, Coimbatore, India.*

Dr. Zong-shen Zhang

*School of Biological Engineering, Dalian Polytechnic University,
Qinggongyuan, Ganjingzi District, Dalian, China, postcode 116034.*

Prof. T. V. Ramana Rao

*B R Doshi School of Biosciences,
Sardar Patel University, Vallabh Vidyanagar,
Gujarat, India.*

Dr. Sanjeev Chandel

*Baba Isher Singh Institute of Sciences &
Technology, Gagra (Moga), Punjab, India.*

Dr. Kuladip Jana

*Bose Institute Centenary Campus, P 1/12, C.I.T.
Scheme VIIM, Kolkata-700 054, India.*

Prof. Ljubinko Jovanovic

*University Educons,
Faculty for Ecological Agriculture, Sremska
Kamenica, Vojvode Putnika 87, Serbia.*

Dr. Luis F. Goulao

*Instituto de Investigacao Cientifica Tropical
[Tropical Research Institute] Eco-Bio / IICT, Av. da
Republica - Quinta do Marques, 2784-505 Oeiras,
Portugal.*

Dr. Lucky K. Attri

*College of Punjabi University Patiala, E-41,
Sector-14, Panjab University, Chandigarh.*

Prof. Bassam Taha Yasseen

*Flat 307 Point Red,
146 Midland Road, Luton, LU2
OBL, UK.*

Dr. Massimo Piccotto

*Tecna S.r.l.,
Area Science Park, Loc. Padriciano, 99, I-34149
Trieste, Italy.*

ARTICLE

Biochemical and physiological constituents and their correlation in wheat (*Triticum aestivum* L.) genotypes under high temperature at different development stages

1

H. R. Ramani, M. K. Mandavia, R. A. Dave, R. P. Bambharolia, H. Silungwe and N. H. Garaniya

Full Length Research Paper

Biochemical and physiological constituents and their correlation in wheat (*Triticum aestivum* L.) genotypes under high temperature at different development stages

H. R. Ramani^{1*}, M. K. Mandavia¹, R. A. Dave¹, R. P. Bambharolia², H. Silungwe³ and N. H. Garaniya¹

¹Department of Biochemistry, Junagadh Agricultural University, Junagadh-362001, Gujarat, India.

²College of Agriculture, Navsari Agricultural University, Waghai-394730, Gujarat, India.

³Department of Food science and Technology, University of Venda, Limpopo province, P/bag X5050, Thohoyandou 0950, South Africa

Received 4 November, 2015; Accepted 5 July, 2016

Wheat is the most popular and staple food for millions of people. It is severely affected by heat stress in many countries. Vegetative growth and reproductive phases in wheat differ in their sensitivity to temperature. Heat tolerant (GW-190) and Heat susceptible (J-2010-11) genotypes grown up to tillering and grain filling stages and Heat treatments (40°C and 45°C for 2h and 4h) were given using Heating House. After heat treatment samples were collected, the biochemical and physiological analysis such as Protein, Proline, Glycine betaine, Membrane stability, Relative water content, Germination percent, Seed vigour and Heat tolerant index were performed. Protein, proline and glycine betaine were found significantly highest 34.06 mg/g Fr. Wt., 13.70 mg% and 902.24 µg/F. wt respectively in heat tolerant genotype GW-190 at 45°C for 4 h at tillering stage. Membrane stability and relative water content were found significantly highest 56.83% and 86.15 % respectively in heat tolerant genotype GW-190 at tillering stage. Germination percent, Seed vigor and Heat tolerance index were found higher in control group of Heat tolerant GW-190 genotypes. Where, all the biochemical and physiological contents were found lower in heat susceptible J-2010-11 genotype. From the above results it was concluded that GW-190 was heat tolerant genotypes which is suitable for grown in area of high temperature and J-2010-11 was found heat susceptible.

Key words: Relative water content, membrane stability, seed index, heat tolerant index, correlation

INTRODUCTION

Wheat is one of the world's most popular crops. It is the staple food for millions of people and forms an important

part of many people's daily diet (Curtis et al., 2002). The yield and quality of cereals are severely affected by heat

*Corresponding author. E-mail: hiteshramani2007@gmail.com.

stress in many countries. Wheat genotypes express a differential response to chronic heat as well as a heat shock (Kamal et al., 2010). Temperature requirements and temperature extremes varies widely for different cultivars of the same species, and among species. The reproductive phase of many crop species is relatively more sensitive to heat stress than the vegetative phase (Hall, 1992; Martiniello et al., 2011).

Vegetative growth and the reproductive phase in wheat, differ in their sensitivity to temperature (Almeselmani et al., 2011; Chakrabarti et al., 2011; Karmanenko et al., 2011; Khakwani et al., 2011; Al-Karaki, 2012; Hakim et al., 2012; Hossain et al., 2012; Hossain et al., 2012; Noorka and Teixeira, 2012; Noorka et al., 2013). The total area of wheat production affected by some or other form of heat stress is estimated to be 65 to 70 million ha. Of these, 7 million ha are grown under continual heat stress (Reynolds, et al., 2001). High temperature stress is a major growth restraining factor for most crop plants. Long term or even a temporary exposure to high temperature can change different metabolic functions, thereby affecting various plant parts including leaves, flower buds and roots (Tsukaguchi et al., 2003; Iwaya-Inoue et al., 2004). Most of the changes are noticeable in the cellular membranes. The metabolic changes include changes in the activities or structures of enzymes and biochemical contents. Heat stress affects the enzymes from metabolic compartments such as mitochondria (Nash et al., 1982), cytoplasm (Laurie et al., 1990) or chloroplasts (Dionisio et al., 1999). Thus, high temperature is believed to produce an array of changes in plants. Under heat stress, biochemical and physiological constitutions changed in heat tolerant and heat susceptible genotypes.

MATERIALS AND METHODS

Heat tolerant (GW-190) and heat susceptible (J-2010-11) wheat genotypes were selected from fourteen genotypes for experiment (Ramani et al., 2015). One set was grown up to tillering stage (35 days) and divided into two groups, control and heat treatments. Other set was grown up to Grain filling stage (around 100 days) and also divided into two sets, control and heat treatments. Heat treatments (40°C and 45°C for 2 h and 4 h) were given using Heating House at tillering and grain filling stages and samples were collected under ice condition. After heat treatments the leaf samples were taken out and leaves were weighed and then transferred immediately to the respective extracting medium for biochemical and physiological analysis.

Estimation of total protein

True protein was estimated by the standard method of Lowry et al., 1951. The samples (0.1 ml) and standard BSA (0.5 - 3 ml from 0.1 mg/ml BSA stock) were taken in a series of test tubes and the volume was made up to 3.0 ml with distilled water. Then 5 ml solution-C was added, mixed well and incubated at room temperature for 10 mins. Then 0.5 ml Folin - Ciocalteu reagent was also added mixed immediately and incubated in dark for 30

mins. A reagent blank was prepared by taking only reagents and volume made up with distilled water. Light blue color was observed after incubation which measured at 660 nm in a spectrophotometer. The protein content was calculated by taking Bovine serum albumin as standard.

Estimation of free proline

Proline was estimated by using acid ninhydrin method of Bates et al., 1973. The samples (0.3 ml) and standard proline (0.1 to 0.6 ml from 0.05 mg/ml proline stock) were taken in a series of test tubes and the volume was made up to 1.0 ml with distilled water. Then 2 ml glacial acetic acid and 2 ml acid ninhydrin reagent were added. Then tubes were kept in boiling water bath for 1 hour. The tubes were cooled in running water at room temperature. After that 4 ml toluene was added. The absorbance was recorded from toluene phase at 520 nm in spectrophotometer. The free proline was calculated by taking Proline as standard.

Estimation of glycine betaine

Glycine betaine was estimated by Grieve and Grantta, 1983, 0.5g leaf powder was shaken in 20 ml distilled water for 24h at 25 °C. The samples were filtered and supernatant was used for analysis. 0.5 ml of supernatant and equal volume of 2N H₂SO₄ were kept in ice water for 1 h. 0.2 ml of cold KI-I₂ was added and reactants were gently vortex. The tubes were kept at 0 to 4°C for 16 h and centrifuge at 10,000 rpm for 15 minute at 0°C. Supernatant were discarded. A periodical crystal were dissolved in 9 ml of 1, 2-dichloroethane. After 2h, the absorbance measured at 365nm using glycine betaine as standard.

Estimation of membrane stability

0.2 g of leaves of uniform size from control and treated plants were taken in tubes containing 30 ml of distilled water. The treatment tubes were then covered with plastic wrap and incubated in a thermostatically controlled water bath for 15 min at 53°C. The control tubes were maintained at 25 °C for the same period. After the elevated temperature treatment, the treatment tubes were quickly cooled down to 25°C. Both tubes were then stored in an incubator overnight for 18 h at 10°C to allow the diffusion of electrolytes from the discs. The tubes were then brought to 25 °C, inverted several times to mix the contents and initial conductance was measured using an electrical conductivity meter. After completing the measurement, tubes were covered and kept at 95 °C for 10 min to completely kill the leaf tissue. Then tubes were cooled down to 25 °C, mixed thoroughly, and the final conductance was measured. Membrane stability was calculated as described in Martineau et al., 1979 method. Membrane Stability Index= $(1 - T1/T2) \times 100$, Where, T1 = Initial conductance of sample kept at 53°C, T2 = Final conductance of sample kept at 95°C.

Estimation of relative water content

0.2 g fresh leaves were weighed to record fresh weight (FW), followed by dipping half of their portion in Petri dish containing 30 ml distilled water for 12 h. The leaves were blotted to wipe off excess water, weighed to record fully turgid weight (TW), and Subjected to oven drying at 70°C for 8 h to record the dry weight (DW). Relative water content (RWC) was calculated as per formula and expressed as: $RWC = [FW - DW] \times 100 / [TW - DW]$ of standard method of Turner, 1986.

Table 1. Biochemical and physiological constituents of heat tolerant and heat susceptible wheat genotypes at tillering and grain filling stages.

Sr. No	Genotypes	Heat treatment (°C)	Protein(mg/g.f.wt)				Proline (mg %)				Glycine betaine (µg/f.wt)				Membrane stability (%)				Relative water content (%)			
			Tillering		Grain filling		Tillering		Grain filling		Tillering		Grain filling		Tillering		Grain filling		Tillering		Grain filling	
			Time (h)		Time (h)		Time (h)		Time (h)		Time (h)		Time (h)		Time (h)		Time (h)		Time (h)		Time (h)	
			2	4	2	4	2	4	2	4	2	4	2	4	2	4	2	4	2	4	2	4
1	GW-190	Control	19.49	19.57	11.86	11.66	1.88	1.93	1.59	1.61	185.55	185.93	156.35	156.12	56.83	55.65	53.49	53.23	86.15	86.07	56.53	56.42
		40	24.91	27.07	13.74	18.44	3.58	6.17	3.07	5.22	335.49	644.93	267.09	449.49	52.68	48.84	43.29	37.26	80.01	74.79	46.82	44.61
		45	29.01	34.06	19.65	20.84	11.07	13.70	8.32	11.00	439.72	902.24	342.01	771.96	43.55	31.58	35.80	33.04	67.37	58.41	37.51	29.29
2	J-2010-11	Control	17.88	18.06	11.23	11.66	1.66	1.73	1.30	1.38	150.22	149.83	117.32	117.26	51.23	51.37	45.29	46.72	79.19	79.37	51.73	51.67
		40	20.77	25.68	12.60	13.31	2.86	4.92	2.55	4.52	211.72	556.98	188.92	361.55	42.53	41.22	35.26	31.43	66.79	62.91	45.00	31.24
		45	28.03	29.05	15.14	16.92	8.59	10.19	7.24	9.29	315.95	765.44	227.23	635.16	37.92	37.32	36.96	35.33	53.45	46.14	28.21	23.45
CD@ 5%			0.78		N.S.		N.S.		0.16		N.S.		N.S.		N.S.		N.S.		N.S.		3.31	
CV %			1.91		1.90		0.98		0.21		5.15		6.45		3.95		4.6		2.21		4.69	

*table show the mean values of three replication, N.S. = none significant.

Estimation of germination percent

Seed were surface sterilized with 0.1% HgCl solution for 2 min and then washed with distilled water. Twenty five seeds each were sown on Whatman filter paper No. 1 bed kept in petri dishes and watered with distilled water. These petri dishes were kept for germination at 22°C (Control), 40°C and 45°C in seed germinator for 7 days to germinate. The filter paper beds were moistened whenever necessary. The germination was recorded after 48 hours and after 7 days. The germination percentage and rate of germination was calculated as per formula given by Almaghrabi, (2012).

Germination Percentage (GP) = $\frac{Ng}{Nt} \times 100$,

Where, Ng = Total number of germinated seeds, Nt = Total number of seeds evaluated.

Estimation of seed vigor

Seeds were allowed to germinate for 7 days after 7 days, the seedling length (shoot and root length) was recorded. The seedling vigor was calculated as per the formula described by Buriro et al., 2011.

Seedling Vigor = Germination percentage × Seedling length at 7 days.

Estimation of heat tolerant index

Heat tolerance index (HTI) was calculated as per the formula given by Fernandez (1992). Heat tolerance index (HTI) = $\frac{(G_{si} \times G_{pi})}{Gp^2}$, Where, G_{si} = germination of each genotype under stress condition, G_{pi} = Germination of each genotype under control condition, Gp^2 = Average of germination in all genotypes under control condition.

Correlation of biochemical and physiological constituents

Mean values of biochemical and physiological constituents were used for correlation analysis using SPSS 16.0 version statistical software.

RESULTS AND DISCUSSION

Results shown in Table 1 to 3 revealed different

levels of the biochemical and physiological constituents and their correlation between heat tolerant and heat susceptible genotypes under high temperature stress at different development stages.

Protein content

Protein was found significantly highest 34.06 mg/g in heat tolerant genotype GW-190 at 45°C for 4 h at tillering stage. The heat susceptible genotype J-2010-11 recorded the lowest (11.23 mg/g. wt.) at grain filling stage. When plants were kept for 2 h, protein content was found lower in both genotypes at both stages as compared to 4 h of heat treatment. All the control plants of heat tolerant GW-190 and susceptible genotypes J-2010-11 showed lower protein as compared to treated plants at both stages. Compared to grain filling stage the protein content was found higher at tillering stage in heat tolerant and heat susceptible

genotypes. As the temperature and duration of temperature treatment increased, the protein was increased in both genotypes at tillering and grain filling stages (Table 1). Plants cope with heat stress in a complex manner, where heat shock proteins (HSPs) which might play a central role in the complex cellular network (Baniwal et al., 2004).

Differential expression of carbohydrate biosynthesis and metabolic pathway enzymes was also inferred in plants that need high energy to cope with heat stress. The responses of rice seedlings to different high-temperature stresses at 35, 40 and 45°C for 48 h. At 35°C, some protective mechanisms were activated to maintain the photosynthetic capability. At 40°C, antioxidative pathways were also active so, the overall protein content increased with temperature (Han et al., 2009). When seven day old rice seedlings encountered high-temperature stress at 45°C, in addition to those induced at 35°C and 40°C, heat shock proteins were effectively induced so the protein content also increased as temperature increased. The appearance of new proteins in wheat shoots subjected to the high temperature stress are heat shock proteins of molecular weights 111, 90, 70, 45, 32, 24 and 8 kDa and they were accumulated under high temperature (Khalid and Devaraj, 2011).

Proline content

Proline was found significantly highest 13.70 mg% in heat tolerant genotype GW-190 at 45°C for 4 h at tillering stage. The heat susceptible genotype J-2010-11 recorded the lowest (1.30 mg%) at grain filling stage. When plants were kept for 2 h, proline content was found lower in both genotypes at both stages as compared to 4 h of heat treatment. All the control plants of heat tolerant GW-190 and susceptible genotypes J-2010-11 showed lower proline as compared to treated plants at both stages. Compared to grain filling stage, the proline content was found higher at tillering stage in heat tolerant and heat susceptible genotypes. As the temperature and duration of temperature treatment increased the proline content was also increased in both genotypes at tillering and grain filling stages (Table 1). Accumulation of certain low molecular mass organic compounds, generally called compatible solutes or osmoprotectants, is a key adaptive mechanism in many plants grown under abiotic stress including extreme temperature (Hare, et al., 1998). A variety of osmolytes may accumulate in different plant species under stress. These include sugars and sugar alcohols (polyols), proline, tertiary and quaternary ammonium compounds and tertiary sulphonium compounds (Sairam and Tyagi, 2004). At high temperature (35°C) the heat tolerant genotypes produced more than double (> 200%) proline than that of 25°C but the heat

susceptible genotypes produced less quantity of proline at 35°C when compared to that in heat susceptible genotypes at 25°C (Ahmed and Hasan, 2011). Proline content was remarkably declined at different stages of growth with lowest accumulation at seed hardening stage (Kumar et al., 2012). Compared to heat sensitive genotypes, proline content were higher in tolerant genotypes under late sown conditions (Dhyani et al., 2013).

Glycine betaine

Glycine betaine was found significantly highest 902.24 µg/F. wt in heat tolerant genotype GW-190 at 45°C for 4 h at tillering stage. The heat susceptible genotype J-2010-11 recorded the lowest (117.32 µg/F. wt) at grain filling stage. When plants were kept for 2 h glycine betaine was found lower in both genotypes at both stages as compared to 4 h of heat treatment. All the control plants of heat tolerant GW-190 and susceptible genotypes J-2010-11 showed lower glycine betaine as compared to treated plants at both stages. Compared to grain filling stage the glycine betaine content was found higher at tillering stage in heat tolerant and heat susceptible genotypes. As the temperature and duration of temperature treatment increased the glycine betaine content was also increased in both genotypes at tillering and grain filling stages (Table 1). Glycine betaine is an amphoteric quaternary amine that plays an important role as osmoprotectant in wheat plants and accumulated under a range of abiotic stresses including high temperature. Stress, caused a significant increase in Glycine betaine content of wheat genotypes at tillering and anthesis stages. The responses of wheat genotypes to drought, heat stress condition and analyzed mechanism of glycine betaine were involved in an improvement of wheat tolerance (Wang et al., 2010).

Membrane stability

Membrane stability was found significantly highest 56.83% in heat tolerant genotype GW-190 at tillering stage. The heat susceptible genotype J-2010-11 recorded the lowest (31.43%) at grain filling stage. When plants were kept for 2 h, membrane stability was found higher in both genotypes at both stages as compared to 4 h of heat treatment. All the control plants of heat tolerant GW-190 and susceptible genotypes J-2010-11 showed higher membrane stability when compared to treated plants at both stage. Compared to grain filling stage the membrane stability was found higher at tillering stage in heat tolerant and heat susceptible genotypes. As the temperature and duration of temperature treatment increased the membrane stability was also decreased in both genotypes

Table 2. Germination per cent, Seed vigor and heat tolerant index of heat tolerant and heat susceptible wheat genotypes under high temperature.

Sr. No	Genotypes	Heat treatment (°C)	Germination (%)	Seed vigor	Heat tolerant index
1	GW-190	Control	100	1945	1.00
		40	92.67	1190	0.97
		45	90.67	980	0.89
2	J-2010-11	Control	100	1871	1.00
		40	70.33	820	0.87
		45	62.67	658	0.67

at tillering and grain filling stages (Table 1). Based on membrane thermostability (MT) test, varieties took maximum heat killing time and were classified as heat tolerant, three varieties are moderately tolerant and the rest took the shortest heat killing time and are considered as heat sensitive (Skider, et al., 2001). Thermo tolerant wheat genotypes showed higher Membrane stability index when compared to thermo sensitive wheat genotypes at different development stages (Mohammadi, et al., 2007). The Membrane thermostability was significantly affected by different growth stages. Membrane stability of wheat genotypes decreased during the later developmental stages. Membrane stability of flag leaf at the early milk stage was significantly correlated with grain yield (Yildirim et al., 2009).

Relative water content

Relative water content was found significantly highest 86.15% in heat tolerant genotype GW-190 at tillering stage. The heat susceptible genotype J-2010-11 records the lowest (23.45%) at grain filling stage. When plants were kept for 2 h, relative water content was found higher in both genotypes at both stages as compared to 4 h of heat treatment. All the control plants of heat tolerant GW-190 and susceptible genotypes J-2010-11 showed higher relative water content compared to treated plants at both stage. Compared to grain filling stage the relative water content was found higher at tillering stage in heat tolerant and heat susceptible genotypes. As the temperature and duration of temperature treatment increased the relative water content was also decreased in both genotypes at tillering and grain filling stages (Table 1). The rise in temperature during late sowing significantly decreased leaf relative content at 8 and 23 days after anthesis in wheat (Sairam et al., 2000). This is in accordance with present results. Wheat genotypes had lower relative water content values at early sowing and late sowing date compared to normal sowing date (Bhesaniya, 2005). Relative water content in root was not affected by higher temperature at all stages of wheat seedling. While

coleoptiles relative water content decreased at all stages of seedling development (Savicka and Skute, 2012).

Germination

The mean germination percentage recorded significantly higher 100% germination in control of GW-190 followed by (92.67%) and (90.67%) at 40°C and 45°C respectively while significantly lowest value was observed with genotype J-2010-11 (62.67%) at 45°C (Table 2). As the temperature increased the germination of seeds decreased in both genotypes but the ratio of decreased in heat susceptible genotype J-2010-11 was more. Germination characters increased with increasing temperature from 15 to 35°C in all wheat genotypes but heat tolerant ability was not based on speed of germination (Hasan et al., 2004). Germination characteristics in heat tolerant cultivars increased with increasing temperature as compared to heat sensitive cultivars (Sikder and Paul, 2010). However, high temperature treatment at 30 to 40°C, germination was reduced in rice (Prasanth et al., 2012) and at 27°C and 30°C temperature treatments, germination was reduced in barley (Ghazi, et al., 2007).

Seed vigor

The mean seed vigor was recorded significantly higher (1945) in control of GW-190 followed by (1190) and (980) at 40°C and 45°C respectively while significantly lowest value was observed with genotype J-2010-11 (658) at 45°C. As the temperature increased the seed vigor decreased in both genotypes but the ratio of decreased in heat susceptible genotype J-2010-11 was more (Table 2). Seedling vigor index was increased with increasing temperature (10 to 30°C) and maximum was recorded at 20°C and 30°C (Buriro et al., 2011). The seedling vigor is adversely affected by heat stress (36°C) which resulted in reduced shoot and root length (Grass and Burris, 1995).

Generally shoot and root growth of heat susceptible

Table 3. Pearson correlations between biochemical and physiological constituents in heat tolerant and susceptible wheat genotypes under high temperature stress.

	Protein	Proline	Glycine betaine	Membrane stability	Relative water content	Germination percent	Seed vigor	Heat tolerance index
Protein	1.000							
Proline	0.960**	1.000						
Glycine betaine	0.994**	0.950**	1.000					
Membrane stability	-0.750*	-0.808*	-0.803*	1.000				
Relative water content	-0.721	-0.818*	-0.768*	0.962**	1.000			
Germination percent	-0.368	-0.452	-0.454	0.801*	0.861*	1.000		
Seed vigor	-0.735	-0.732*	-0.804*	0.944**	0.920**	0.868*	1.000	
Heat tolerance index	-0.531	-0.661	-0.580	0.811*	0.934**	0.913**	0.817*	1.000

*correlation is significant at 0.05, ** correlation is significant at 0.01 (1-tailed).

genotype in relation to length is affected more than those of heat tolerant genotype at higher temperature-35°C (Hasan et al., 2004).

Heat tolerant index

The mean Heat tolerant index was recorded significantly higher (1.00) in control of GW-190 followed by (0.97) and (0.89) at 40°C and 45°C respectively while significantly lowest value was observed with genotype J-2010-11 (0.67) at 45°C. As the temperature increased the seed vigor decreased in both genotypes but the ratio of decreased in heat susceptible genotype J-2010-11 was more (Table 2). Stress tolerance index (STI) and stress susceptible index (SSI) were more accurate criteria for the selection of heat tolerant and high yielding genotypes by growing under normal and stress condition (Khodarahmpouret et al., 2011).

Correlation analysis

One heat tolerant and one heat susceptible

genotypes GW-190 and J-2010-11 respectively were used for correlation analysis (Pearson correlation test) including biochemical and physiological parameters as listed earlier (Table 3). Results of correlation analysis under heat stress condition showed that, protein had highly significant and positive correlation with proline and glycine betaine content whereas negatively and significant correlation with membrane stability. Also protein content was negatively and non-significantly correlated with relative water content, germination percentage, seed vigor and heat tolerant index. Proline content had high significant positive correlation with glycine betaine and membrane stability, while relative water content and seed vigor were negatively and significantly correlated whereas, germination percent and heat tolerant index were negatively and non-significantly correlated. Glycine betaine had significant negative correlation with membrane stability, relative water content and seed vigor, Membrane stability had highly significant positive correlation with relative water content and seed vigor, and positive significant correlation with germination and heat tolerant index. Relative

water content was positively and significantly correlated with germination percent, seed vigor and heat tolerant index. Seed vigor was positively correlated with heat tolerant index (Table 3). Result clearly indicated that relative water content and membrane stability was positively correlated and more effective indicator for screening of heat tolerant genotypes. Protein content, proline and glycine betaine content were positively correlated, they were higher in heat tolerant genotypes. Membrane stability index was good indicators of stress tolerance (Sairam and Srivastava, 2001).

Conclusion

Some abiotic constraints such as high temperature, cold, drought, high concentrations of toxic minerals and salinity results in, severe loss of crop yield and quality. At both tillering and grain filling stages, Protein, proline, glycine betaine content were higher in tolerant genotype as compared to susceptible genotypes. Also these content were increased with increase in temperature in both genotypes. But the amount of increase was higher

in tolerant genotype when compared to susceptible genotypes. Membrane stability, relative water content, heat tolerant index, seed vigor and germination were higher in tolerant genotypes when compared to susceptible genotypes but as temperature increased, they were decreased in both the genotypes at both stages. Also biochemical and physiological content was significantly co-related.

Conflict of interest

The authors have not declared any conflict of interest

REFERENCES

- Ahmed JU, Hasan MA (2011). Evaluation of seedling proline content of wheat genotypes in relation to heat tolerance. *Bangladesh J. Bot.* 40(1):17-22.
- Al-Karaki GN (2012). Phenological development-yield relationships in durum wheat cultivars under terminal high temperature stress in semiarid conditions. *Agron.* pp 7.
- Almeselmani M, Teixeira da Silva JA, Deshmukh P (2011). Stability of different physiological characters, yield and yield components under high temperature stress in tolerant and susceptible wheat genotypes. *Fruit Veg. Cereal Sci. Biotechnol.* 5:86-92.
- Baniwal SK, Bharti K, Chan KY, Fauth M, Ganguli A, Kotak S, Mishra SK, Nover L, Port M, Scharf KD, Tripp J, Weber C, Zielinski D, Von Koskull-Doring P (2004). Heat stress response in plants: a complex game with chaperones and more than twenty heat stress transcription factors. *J. Biol. Sci.* 29:471-487.
- Bates LS, Waldren RP, Teare ID (1973). Rapid determination of free proline for water stress studies. *Plant Soil* 39:205-207.
- Bhesaniya SV (2005). Effect of high temperature on biochemical changes in different genotypes of wheat (*Triticum aestivum* L.). M. Sc. Thesis (Unpublished). Junagadh Agricultural University, Junagadh, Gujarat, India.
- Buriro M, Oad FC, Kerio MI, Tunio S, Gandahi AW, Hassan SWU, Oad SM (2011). Wheat seed germination under the influence of temperature regimes. *Sarhad J. Agric.* 27(4):539-543.
- Chakrabarti B, Singh SD, Nagarajan S, Aggarwal PK (2011). Impact of temperature on phenology and pollen sterility of wheat varieties. *Aust. J. Crop Sci.* 5(8):1039-1043.
- Curtis BC, Rajaram S, Macpherson HG (2002). Bread wheat: FAO. *Plant Production and Protection Series* 30:12-15.
- Dhyani K Ansari MW, Rao YR, Verma RS, Shukla A, Tuteja N (2013). Comparative physiological response of wheat genotypes under terminal heat stress. *Plant Signal. Behav.* 8(6):1-6.
- Dionisio ML, Shono M, Tobita S (1999). Effect of proline and betain on heat inactivation of ribulose 1,5- bisphosphate carboxylase/oxygenase in crude extract of rice seedling. *Photosynthetica* 36(4):557-563.
- Fernandez GCJ (1992). Effective selection criteria for assessing stress tolerance. In: Kuo, C. G. eds. *Proceedings of the International Symposium on Adaptation of Vegetables and Other Food Crops in Temperature and Water Stress.* Publication, Tainan, Taiwan.
- Ghazi NA, Asadulla A, Yahia O (2007). Seed germination and early growth of three barley cultivars as affected by temperature and water stress. *Am. Euro. J. Agric. Environ. Sci.* 2(2):112-117.
- Grass L, Burriss JS (1995). Effect of heat stress during seed development and maturation on wheat (*Triticum durum* L.) seed quality. I. Seed germination and seedling vigor. *Can. J. Plant Sci.* 75:821-829.
- Grieve CM, Grattan SR (1983). Rapid assay for determination of water soluble quartnary ammonium compounds. *Plant Soil* 70:303-307.
- Hakim MA, Hossain A, Teixeira da Silva JA, Zvolinsky VP, Khan MM (2012). Yield, protein and starch content of 20 wheat (*Triticum aestivum* L.) genotypes exposed to high temperature under late sowing conditions. *J. Sci. Res.* 4(2):477-489.
- Hall AE (1992). Breeding for heat tolerance. *Plant Breed. Rev.* 10:129-168.
- Han F, Chen H, Li XJ, Yang MF, Liu GS, Shen SH (2009). A comparative proteomic analysis of rice seedling under various high temperature stresses. *Biochem. Biophys. Acta* 1794:1625-1634.
- Hare PD, Cress WA, Staden JV (1998). Dissecting the roles of osmolyte accumulation during stress. *Plant Cell Environ.* 21:535-553.
- Hasan MA, Ahmed JU, Hossain T, Hossain MM, Ullah MA (2004). Germination characteristics and seed reserve mobilization during germination of different wheat genotypes under variable temperature regimes. *J. Nation Sci. Foundation Sri Lanka* 32(3, 4):97-107.
- Hossain A, Teixeira da Silva JA, Lozovskaya MV, Zvolinsky VP, Mukhortov I (2012). High temperature combined with drought affect rainfed spring wheat and barley in south-eastern Russia: Yield, relative performance and heat susceptibility index. *J. Plant Breed. Crop Sci.* 4(11):184-196.
- Iwaya-Inoue M, Matsui R, Fukuyama M (2004). Cold- or heat-tolerance of leaves and roots in perennial ryegrass determined by ¹H-NMR. *Plant Prod. Sci.* 7(2):118-128.
- Kamal AHM, Kim KH, Shin KH, Choi JS, Baik BK, Tsujimoto H, Hwa Young Heo HY, Park CS, Woo SH (2010). Abiotic stress responsive proteins of wheat grain determined using proteomics technique. *Aust. J. Crop Sci.* 4(3):196-208.
- Karmanenko NM, Osipova LV, Nilovskaya NT (2011). Acid, cold and drought tolerance in cereal. *Russ. Acad. Sci.* 37(5):354-357.
- Khakwani AA, Dennett MD, Munir M (2011). Drought tolerance screening of wheat varieties by inducing water stress conditions. *J. Sci. Technol.* 33(2):135-142.
- Khalid MN, Devaraj VR (2011). Antioxidant and other biochemical defense responses of *Macrotyloma uniflorum* (Lam.) Verdc. (Horse gram) induced by high temperature and salt stress. *Braz. J. Plant Physiol.* 23(3):187-195.
- Khodarahmpouret Z, Choukan R, Bihanta MR, Hervan EM (2011). Determination of the best heat stress tolerance indices in maize (*Zea mays* L.) inbred lines and hybrids under khuzestan province conditions. *J. Agric. Sci. Technol.* 13:111-121.
- Kumar RR, Goswami S, Sharma SK, Singh K, Gadpayle KA, Kumar N, Rai GK, Singh M, Rai RD (2012). Protection against heat stress in wheat involves change in cell membrane stability, antioxidant enzymes, osmolyte, H₂O₂ and transcript of heat shock protein. *Int. J. Plant Physiol. Biochem.* 4(4):83-91.
- Laurie S, Stewart GR (1990). The Effects of Compatible Solutes on the Heat Stability of Glutamine Synthetase from Chickpeas Grown under Different Nitrogen and Temperature Regimes. *J. Exp. Bot.* 41(11):1415-1422.
- Lowry OH, Rose rough AL, Farr AL, Randall RJ (1951). Protein measurement with the Folin Phenol reagent. *J. Biol. Chem.* 193(1):265-275.
- Martineau PM, Williams JH, Speght JE (1979). Temperature tolerance in soybeans-II. Evaluation of segregating population for membrane thermo-stability. *Crop Sci.* 19:79-81.
- Martiniello P, Teixeira da Silva JA (2011). Physiological and bio-agronomical aspects involved in growth and yield components of cultivated forage species in Mediterranean environments: A review. *Eur. J. Plant Sci. Biotechnol.* 5(2):64-98.
- Mohammadi V, Bihanta MR, Zali AA (2007). Evaluation of screening techniques for Heat tolerance in wheat. *Pak. J. Biol. Sci.* 10(6):887-892.
- Nash D, Paleg LG, Wiskich JJ (1982). Effect of Proline, Betaine and Some Other Solutes on the Heat Stability of Mitochondrial Enzymes. *Aust. J. Plant Physiol.* 9(1): 47-57.
- Noorka IR, Teixeira da Silva JA (2012). Mechanistic insight of water stress-induced aggregation in wheat (*Triticum aestivum* L.) quality. *Note Sci. Biol.* 4(4):32-38.
- Noorka IR, Batool A, Rauf S, Teixeira da Silva JA, Ashraf E (2013). Estimation of heterosis in wheat (*Triticum aestivum* L.) under contrasting water regimes. *Int. J. Plant Breed.* 7(1):55-60.
- Prasanth VV, Chakravarthi DVNT, Kiran V, Rao YV, Panigrahy M,

- Mangrauthia SK, Viraktamath BC, Subrahmanyam D, Voleti SR, Sarla N (2012). Evaluation of rice germplasm and introgression lines for heat tolerance. *Annals Biol. Res.* 3(11):5060-5068.
- Ramani HR, Mandavia MK, Khunt AG, Golakiya BA (2015). Evaluation of wheat genotypes (*Triticum aestivum* L.) against heat stress based on their biochemical and physiological responses. *India J. Agric. Biochem.* 27(2):133-137.
- Reynolds MP, Nagarajan S, Razzaque MA, Ajeeb OOA (2001). Heat tolerance. In: M.P. Reynolds, J. I. Ortiz-Monasterio and A. McNab, Eds, Application of physiology in wheat breeding, CIMMYT, Mexico, pp. 124-135.
- Sairam RK, Srivastava GC (2001). Water stress tolerance of wheat (*Triticum aestivum* L.): variations in hydrogen peroxide accumulation and antioxidant activity in tolerant and susceptible genotypes. *J. Agron. Crop Sci.* 186:63-70.
- Sairam RK, Tyagi A (2004). Physiology and molecular biology of salinity stress tolerance in plants. *Curr. Sci.* 86(3):407-421.
- Sairam RK, Srivastava GC, Saxena DC (2000). Increased antioxidant activity under elevated temperature: a mechanism of heat stress tolerance in wheat genotypes. *Biologia-Plantarum (Czech Republic.)* 43(2):245-251.
- Savicka M, Skute N (2012). Some morphological, physiological and biochemical characteristics of wheat seedling *Triticum aestivum* L. organs after high-temperature treatment. *Ekologija* 58(1):9-21.
- Sikder S, Paul NK (2010). Study of influence of temperature regimes on germination characteristics and seed reserves mobilization in wheat. *Afr. J. Plant Sci.* 4(10):401-408.
- Skider S, Ahmed JU, Hossain T (2001). Heat Tolerance and Relative Yield Performance Of Wheat Varieties Under Late Seeded Conditions. *Indian J. Agric. Res.* 35(3):141-148.
- Tsukaguchi TY, Kawamitsu H, Takeda K, Suzuki K, Egawa Y (2003). Water status of flower buds and leaves as affected by high temperature in heat-tolerant and heat-sensitive cultivars of snap bean (*Phaseolus vulgaris* L.). *Plant Prod. Sci.* 6:24-27.
- Turner NC (1986). Crop water deficit: a decade of progress. *Advance Agronomy*, 39, Almaghrabi, O. A. (2012). Impact of drought stress on germination and seedling growth parameters of some wheat cultivars. *Life Sci. J.* 9(1):590-598.
- Wang GP, Zhang XY, Li F, Luo Y, Wang W (2010). Over accumulation of glycine betaine enhances tolerance to drought and heat stress in wheat leaves in the protection at photosynthesis. *Photosynthetica* 48:117-126.
- Yildirim M, Bahar B, Mujde KOC, Barutcular C (2009). Membrane Thermal Stability at Different Developmental Stages of Spring Wheat Genotypes and Their Diallel Cross Populations. *J. Agric. Sci.* 15(4):293-300.

A photograph of a ripe red strawberry on a wooden cutting board. A green marker is being used to pierce the strawberry. The background is a wooden surface. The text is overlaid on a semi-transparent dark brown band.

International Journal of Plant Physiology and Biochemistry

Related Journals Published by Academic Journals

- *African Journal Of Plant Science*
- *Journal Of Microbiology And Antimicrobials*
- *International Journal Of Biotechnology And Molecular Biology Research*
- *International Journal Of Biodiversity And Conservation*

academicJournals