

OPEN ACCESS



Journal of
**Pharmacognosy and
Phytotherapy**

June 2018
ISSN 2141-2502
DOI: 10.5897/JPP
www.academicjournals.org



**ACADEMIC
JOURNALS**
expand your knowledge

ABOUT JPP

The **Journal of Pharmacognosy and Phytotherapy (JPP)** is published monthly (one volume per year) by Academic Journals.

The **Journal of Pharmacognosy and Phytotherapy (JPP)** is an open access journal that provides rapid publication (monthly) of articles in all areas of the subject such as ethnobotany, phytochemistry, ethnopharmacology, zoopharmacognosy, medical anthropology etc.

The Journal welcomes the submission of manuscripts that meet the general criteria of significance and scientific excellence. Papers will be published shortly after acceptance. All articles published in JPP are peer-reviewed.

Contact Us

Editorial Office: jpp@academicjournals.org

Help Desk: helpdesk@academicjournals.org

Website: <http://www.academicjournals.org/journal/JPP>

Submit manuscript online <http://ms.academicjournals.me/>

Editors

Dr. (Mrs) Banasri Hazra

Research Scientist (U.G.C.)
Department of Pharmaceutical Technology
Jadavpur University
Calcutta - 700032
India

Dr. Yuanxiong Deng

Dept of Pharmaceutical Science
School of Medicine
Hunan Normal University
Tongzipo Road 371, Changsha 410013,
Hunan China

Prof. Maha Aboul Ela

Beirut Arab University, Faculty of Pharmacy, Beirut
Campus

Dr. S. RAJESWARA REDDY

Assistant Professor, Division of Animal Biotechnology
Department of Biotechnology, School of Herbal
Studies and Naturo Sciences,
Dravidian University, Kuppam – 517 425, A.P.
India

Dr. Mekhfi Hassane

University Mohammed the First, Faculty of Sciences,
Department of biology, Oujda, Morocco
Morocco

Dr. Arun Kumar Tripathi

Central Institute of Medicinal and Aromatic Plants
P.O. CIMAP, LUCKNOW-226015,
India

Dr. Wesley Lyeverton Correia Ribeiro

Universidade Estadual do Ceará, Faculdade
de Veterinária/Laboratório de Doenças
Parasitárias Av. Paranjana, 1700
Itaperi - Fortaleza
60740-903, CE - Brazil

Dr. Maryam Sarwat

C/O A.M. Khan, House No. 195

Dr. Yong-Jiang Xu

Saw Swee Hock School of Public Health,
National University of Singapore, Singapore.

Prof. Dr. Adeolu Alex Adedapo

Department of Veterinary Physiology,
Biochemistry and Pharmacology
University of Ibadan, Nigeria

Dr. Joana S. Amaral

Campus de Sta Apolónia,
Ap. 1134, 5301-857 Bragança,
Portugal

Dr. Asad Ullah Khan

Interdisciplinary Biotechnology UNIT
Aligarh Muslim University,
India

Dr. Sunday Ene-ojo Atawodi

Biochemistry Department
Ahmadu Bello University
Zaria, Nigeria

Prof. Fukai Bao

Department of Microbiology and Immunology,
Kunming Medical College
China

Dr. Bhaskar C Behera

Agharkar Research Institute
Dept. of Science & Technology,
Plant Science Division
India

Prof. R. Balakrishna Bhat

Walter Sisulu University
Department of Botany
Mthatha, South Africa

Dr. Mohammad Nazrul Islam Bhuiyan

BCSIR Laboratories;
Chittagong cantonment;
Chittagong-4220;
Bangladesh

Dr. Baojun Bruce Xu

Beijing Normal University-Hong Kong Baptist
University United International College Zhuhai,
Guangdong Province,
China

Dr. Hamad H. Issa

Department of Physical Sciences,
School of natural Sciences,
The University of Dodoma,
Tanzania

Dr. Gagan Deep

Department of Pharmaceutical Sciences
School of Pharmacy,
University of Colorado Denver,
Colorado,
USA

Dr. Fengguo Xu

Dept of Epidemiology and Public Health,
Yong Loo Lin School of Medicine,
National University of Singapore,
Singapore

Dr. Haitao Lv

Medicine and Endocrinology,
Albert Einstein College of Medicine,
Yeshiva University,
USA

Hassane MEKHF

University Mohammed the First,
Faculty of Sciences,
Department of biology,
Laboratory of Physiology and Ethnopharmacology,
Morocco

Dr. Subhash C. Mandal

Division of Pharmacognosy
Pharmacognosy and Phytotherapy Research
Laboratory,
Department of Pharmaceutical Technology,
Jadavpur University,
India.

Dr. Adibe Maxwell Ogochukwu

Clinical Pharmacy and Pharmacy Management,
Faculty of Pharmaceutical Sciences,
University of Nigeria, Nsukka
Enugu state,
Nigeria.

Dr. Odukoya, Olukemi Abiodun

Department of Pharmacognosy,
Faculty of Pharmacy
University of Lagos.
Nigeria.

Dr. Qinxue Richard Ding

Medical Center at Stanford University,
Palo Alto,
USA

Dr. Sulejman Redžić

Faculty of Science of the University of Sarajevo
33-35 Zmaja od Bosne St.,
Sarajevo,
Bosnia and Herzegovina

Dr. Michal Tomczyk

Medical University of Bialystok,
Faculty of Pharmacy,
Department of Pharmacognosy,
Poland

Dr. Ugur Çakilcioglu

Firat University,
Faculty of Science and Arts,
Department of Biology,
Elazig
Turkey

Prof. Samson Sibanda

National University of Science and Technology
Cnr Gwanda Road/Cecil Avenue,
Ascot, Bulawayo,
Zimbabwe

Journal of Pharmacognosy and Phytotherapy

Table of Contents: Volume 10 Number 6 June 2018

ARTICLE

Fungal metagenomes in polyherbal medicines used for the treatment of tuberculosis

102

Elizabeth Bosede Famewo, Anna Maria Clarke and Anthony Jide Afolayan

Full Length Research Paper

Fungal metagenomes in polyherbal medicines used for the treatment of tuberculosis

Elizabeth Bosede Famewo¹, Anna Maria Clarke¹ and Anthony Jide Afolayan^{2*}

¹Microbial Pathogenicity and Molecular Epidemiology Research Group (MPMERG), Faculty of Science and Agriculture, University of Fort Hare, Alice 5700, South Africa.

²Medicinal Plants and Economic Development (MPED) Research Niche Area, Faculty of Science and Agriculture, University of Fort Hare, Alice 5700, South Africa.

Received 24 January, 2018; Accepted 22 March, 2018

The traditional systems of medicine have significantly become more accepted in the developed and developing countries due to their curative property, less toxicity and minimal side effects. However, several studies have shown that they are associated with microbial contaminants. This study aimed at identifying fungi in nine polyherbal medicines used for the treatment of tuberculosis in Eastern Cape Province, South Africa. Sequences of fungi DNA that encodes internal transcribed spacer (ITS) region were retrieved from the remedies. The ITS region of the fungal rRNA operon was amplified using ITS1 and ITS4 primers. The amplicons were visualized on agarose gel electrophoresis, followed by end repair and adaptor ligation. They were further purified and quantified using Library Preparation kit NEBNext® UltraT DNA Library Prep Kit for illumina and run on illumina's MiSeq platform. The study revealed that the polyherbal medicines are contaminated with fungi species. The predominant mycoflora obtained belongs to different genera or species of fungi. They include *Aspergillus*, *Penicillium*, *Alternaria*, *Candida*, *Ramularia*, *Cladosporium* and *Malassezia* among others. Some of these organisms are capable of causing infections in immunocompromised patients. Thus, the study identified various fungal contaminants in polyherbal remedies sold to tuberculosis patients in five communities in the Eastern Cape Province of South Africa.

Key words: Fungi, internal transcribed spacer (ITS) region, polyherbal medicines, public health, tuberculosis.

INTRODUCTION

The use of polyherbal formulations for therapeutic purposes has significantly increased in the developed and developing countries because of their curative property, less toxicity and minimal side effects (Binu, 2008; Devi et al., 2010). These benefits have made the

usage of herbal medicines to be intertwined with that of modern medicine, thus, increasing the global market by 7% annually (Dubey et al., 2008). Traditional healers, especially those in low-income countries make use of various herbal preparations for the treatment and

*Corresponding author. E-mail: aafolayan@ufh.ac.za. Tel: +27 82 202 2167. Fax: +27 866 282 295.

management of ailments such as wound infection, skin diseases, diabetes, diarrhoea, urinary tract infections, stomach illnesses and tuberculosis (Louw et al., 2002; Buwa and Afolayan, 2009).

Tuberculosis (TB), caused by a bacterium called *Mycobacterium tuberculosis* is the major killer among the infectious diseases and it is the ninth-leading cause of death worldwide. An estimate of 10.4 million new TB cases were reported in 2016, of this population, 1.7 million deaths including human immunodeficiency virus-TB co-infected individual were recorded (WHO, 2017). Seven countries have been implicated to have the highest burden of TB cases, thus responsible for 64% of the world TB burden. These countries include India, Indonesia, China, Philippines, Pakistan, Nigeria and South Africa (WHO, 2017). According to the Statistics for South Africa, TB is a significant public health challenge accounting for 7.2% of all death in 2016 followed by diabetes mellitus (SSA, 2017). The report of WHO (2016) gave an estimated incidence of 454,000 cases of active TB in South Africa; this implies that, about 0.8% of the 54 million South Africa populations develop active TB diseases (WHO, 2016).

In South Africa, about three million people make use of herbal remedies for their health care purposes especially for the treatment of infection disease such as TB (Louw et al., 2002; Elujoba et al., 2005). Despite the increasing use of herbal preparations and the global expansion of the market, safety is of great concern. Some studies have revealed that due to unscientific mode of harvesting, drying, transportation, cleaning and handling of these herbal preparations, the raw plants are prone to infestations and exposed them to different kinds of microbial contaminants (Stevic et al., 2012). The dominating contaminants are the bacterial endospores and fungal spores while the remaining are heavy metals and viruses originating from the soil (Adeleye et al., 2005; Kaume et al., 2012; Ting et al., 2013; Noor et al., 2014).

A few surveillance studies (Czech et al., 2001; Tassaneeyakul et al., 2004; Okunlola et al., 2007; Kulshrestha et al., 2008) have been conducted and have shown the presence of microbial contaminants in herbal preparations. Walther et al. (2016) investigated 109 traditional liquid herbal medicinal products in Mwanza city; the findings revealed that 81.7% of the samples were contaminated with fecal coliforms. The microbial quality of some oral liquid herbal medicines marketed in Ile-Ife conducted showed that 90% of the samples carried microbial loads beyond officially permissible limits (Igbeneghu and Lamikanra, 2016). Also, the quality control of hypoglycemic herbal preparations investigated in Nairobi investigated has shown that the preparations are contaminated with both bacterial and fungal contaminants (Chege et al., 2015).

Siakrwar et al. (2014) isolated and identified a wide spectrum of fungi including *Aspergillus*, *Penicillium*, *Alternaria*, *Rhizopus* and *Syncephalastrum* species in 15

medicinal plants. Toma and Abdulla (2013) found that most of the fungal species detected in different types of spices and medicinal plants were *Aspergillus* spp. and *Penicillium* spp. while *Stachybotrys* spp., *Syncephalastrum racemocum*, *Uocladium botrytis*, *Alternaria alternata*, *Cladosporium lignicolum* and *Gliocladium catenulatum* were less frequently detected. Quality assessment of aqueous herbal/medicinal products has shown that the most abundant fungi were from *Cladosporium herbarum*. This was then followed by *Aspergillus* spp., *Saccharomyces kluyverii*, *Rhodotorulla minuta*, *Candida membranifasciens* and *Sporobolomyces salmonicolor* (Osei-Adjei et al., 2013).

The presence of numerous fungal species in herbal preparations can be harmful to consumers. Thus, in order to safe guard the health of the consumers, this study aimed at identifying the presence of different fungi in polyherbal medicines used for the treatment of tuberculosis in Amathole District Municipality, Eastern Cape Province, South Africa.

MATERIALS AND METHODS

Sample collection

This study is a secondary data analysis of the first authors' research project "Ethno-medicinal documentation of polyherbal medicines used for the treatment of tuberculosis in Amathole District Municipality of the Eastern Cape Province, South Africa" where information about the herbs used for the preparation of this remedies are revealed (Famewo et al., 2017). A total of nine different polyherbal medicines were purchased from the traditional herbal healers in five different communities, namely; East London (EL), King Williams Town (KWT), Hogsback (HB), Alice (AL) and Fort Beaufort (FB) as shown in Figure 1. These remedies were liquid preparations and each of them was already homogenized and packaged in a 2 L container by the herbal healers. Each remedy was labelled and coded according to the place of collection. The number of remedies obtained in this study was due to the fact that only a few traditional healers treat and sell the remedies for the treatment of tuberculosis. They claim to have acquired the knowledge from their ancestors and this knowledge is being transferred from one generation to another. The samples were transported to Medicinal Plants and Economic Development (MPED) Research Centre Microbiology Laboratory for analysis.

DNA extraction

A modified method of Dei-Tutuwa et al. (2014) was used for the fungal DNA extraction. One milliliter of each sample was pipetted into Eppendorf tubes and centrifuged at 12500 rpm for 10 min, the supernatant was discarded and the cell pellets was collected. The total fungal DNA was extracted using ZR Fungal/Bacterial DNA MiniPrep™ Kit (Zymo Research, USA) according to the manufacturer instructions.

Amplification of fungal DNA using polymerase chain reaction (PCR)

The assay was conducted using the internal transcribed spacer (ITS) region of the fungal genome which is highly variable among




Figure 1. Map of Amathole District Municipality.
Source: Famewo et al. (2016).

species or even populations of the same species (Schoch et al., 2012). This region lies between the 18S small subunit (SSU) and 28S large subunit (LSU) ribosomal RNA (rRNA) genes, which also contains two non-coding spacer regions (ITS-A and ITS-B) separated by the 5.8S rRNA gene. The total genomic fungal DNA was amplified using forward ITS1 (5'-TCCGTAGGTGAACCTGCGG-3') and reverse primers ITS4 (5'-TCCTCCGCTTATTGATATGC-3') (White et al., 1990). PCR reaction was carried out in a final volume of 25 μ L consisting 5 μ L template DNA, 12.5 μ L of Q5[®] Hot start High-Fidelity 2X PCR Master Mix (New England Biolabs, USA), 1 μ L of 10 pmol each of the oligonucleotide primers (Inqaba Biotech, SA) and 5.5 μ L of nuclease free water). Reactions were performed in the thermocycler (Bio-Rad Mycycler, USA) under the following conditions: initial denaturation at 95°C for 2 min, followed by 30 cycles at 95°C for 20 s, 55°C for 30 s, and 72°C for 30 s, and a final elongation at 72°C for 5 min (Kuo et al., 2005). In order to confirm the PCR products size, 5 μ L of the amplicons was analyzed by gel electrophoresis in 1% agarose (Merck, SA) stained with 3 μ L ethidium bromide (Sigma-Aldrich, USA). A 100 bp DNA ladder (Thermo Scientific, EU) Lithuania) was included for band size estimation purposes. All gels were run in 0.5X TBE buffer at 95 V for 1 h and visualized by UV trans-illumination (Alliance 4.7, France).

Purification of amplicons and sequencing

The amplicons were purified using the Agencourt[®] Ampure[®] XP bead protocol (Beckman Coulter, USA). The amplicon libraries were purified using the Agencourt[®] Ampure[®] XP bead protocol (Beckman Coulter, USA). Library concentration was measured using Nebnext

Library quant kit (New England Biolabs, USA) and quality validated using Agilent 2100 Bioanalyser (Agilent Technologies, USA). The samples were pooled in equimolar concentrations and diluted to 4 nM based on library concentrations and calculated amplicon sizes. The library pool was sequenced on a MiSeq[™] (Illumina, USA) using a MiSeq[™] Reagent kit V3 600 cycles PE (Illumina, USA). The final pooled library was at 10 pM with 20% PhiX as control amplicon sequencing protocol. Each sample was sequenced in the sense and antisense directions using ITS1 and ITS4 primers (White et al., 1990; Kozich et al., 2013).

Data analysis

The relative frequency of different fungal phylum in each of the remedy, percentage occurrence of each fungal family and the abundance of fungal families was calculated according to Girridher and Ready (1997).

$\% \text{ of frequency} = \frac{\text{Number of observation in which a species appeared}}{\text{Total number of observation}} \times 100$

RESULTS

Distribution of different fungal phylum in each of the polyherbal remedies

The mycological analysis of the nine polyherbal remedies




Figure 2. Relative frequencies of contaminating organisms in each of the polyherbal remedy.

revealed that all the herbal preparations are contaminated with different fungal phylum and species. The predominant fungal phylum identified in all the herbal remedies was *Ascomycota* followed by *Basidiomycota*. The presence of 5% *Glomeromycota* was identified in KWTb and HBss remedies, while 5% *Zygomycota* was found in KWTc remedy (Figure 2).

Distribution of different fungal families in all the polyherbal remedies

A total of 43 different fungal families were identified in all the nine polyherbal remedies. Members of the families *Davidiellaceae*, *Mycosphaerellaceae*, *Trichocomaceae*, *Pleosporaceae* and *Saccharomycetaceae* were identified in all the remedies (Figure 3). This was followed by *Debaryomycetaceae*, *Malasseziaceae*, *Dothioraceae*, *Herpotrichiellaceae* and *Tremellaceae*.

The abundance of each fungal family in all the remedies is as shown in Figure 4. Members of the family *Trichocomaceae* were the most abundant followed by *Mycosphaerellaceae*, *Pleosporaceae*, *Saccharomycetaceae*, *Tremellaceae*, *Davidiellaceae*, *Malasseziaceae* and *Sporidiobolaceae*.

Distribution of different fungal genera and species in each of the polyherbal remedies

By analyzing each of the herbal preparation (supplementary material), KWTa remedy was mainly contaminated with species of *Debaryomyces*, *Penicillium* and *Ramularia*. However, *Rhodotorula*, *Cladosporium*, *Ramularia*, *Candida* and *Malassezia* species were found in KWTb remedy. The remedy from KWTc was contaminated with *Ramularia*, *Candida*, *Cryptococcus*, *Rhodotorula*, *Cystofilobasidium* and *Mucor* species.

While AL remedy was mostly dominated with *Ramularia*, *Aspergillus*, *Penicillium*, *Candida*, *Rhinochadiella* and *Parmotrema* species; *Cladosporium*, *Ramularia*, *Alternaria*, *Candida* and *Malassezia* were identified in EL remedy. The presence of *Cladosporium*, *Ramularia*, *Aspergillus*, *Penicillium*, *Alternaria* and *Candida* were detected in FB remedy.

While species of *Cladosporium*, *Ramularia*, *Aspergillus* and *Penicillium* dominated the mycofloral of HBfs remedy; HBss remedy was contaminated with *Cladosporium*, *Ramularia* and *Aspergillus*. However, *Cladosporium*, *Ramularia*, *Aspergillus*, *Penicillium*, *Alternaria* and *Wallemia* species were detected in HBts remedy.




Figure 3. Percentage occurrence of each fungal family identified in the polyherbal remedy.




Figure 4. The abundance of fungal families in all the polyherbal remedies.

Across all the herbal remedies, the most predominant mycoflora obtained was distributed in four different genera which comprised of *Candida*, *Cladosporium*, *Ramularia* and *Alternaria*. This was followed by *Aspergillus* and *Penicillium* which were found in seven and eight remedies, respectively (supplementary

material).

DISCUSSION

The use of polyherbal medicines for the treatment and

management of various illnesses is part of the health-care culture in South Africa. These remedies are prepared locally by traditional healers by combining two or more parts of medicinal plants such as the root, leaf, stem, flower and seed. The results obtained in the present study revealed the presence of different fungi contaminants particularly moulds and yeast in the polyherbal remedies, which are used for the treatment of TB. The presences of these organisms may pose potential health risks to tuberculosis patients considering their immunocompromised status. The high fungal population in each of the remedies is an indication of low environmental sanitation and unhygienic standard of processing these herbal medicines. All the remedies were stored at room temperature in the shops and there were no expiry date written on them, thus, the storage condition could have encouraged the growth of these fungal species.

Many of the fungal species identified are naturally inhabitant of the soil and some are plant pathogens. According to Sharma (2001), fungal contamination of herbal preparations mainly occurs during a slow drying process. Inadequate drying or postharvest storage of the herbs under a high relative humidity and favourable temperature promotes the growth of these microbes (Sharma, 2001). Also, the unscientific methods of collection, unsuitable transportation and prolonged storage of the plants and inadequate hygiene of the handler could trigger the growth of organisms in herbal medicines (Sago et al., 2009; Stević et al., 2012).

The presence of mould such as *Aspergillus* and *Penicillium* spp. in seven and eight polyherbal remedies, respectively, could be attributed to the growth of these organisms in the herbs before the medicinal plants were completely dried (Stevic et al., 2012). Both species of *Aspergillus* and *Penicillium* have been associated with food poisoning and may cause infections in an immunosuppressed individuals (Lin et al., 2001; Bateman et al., 2002). The results of this study were in good agreement with those found by Tournas and Katsoudas (2008). The study examined the microbiological quality of various medicinal herbal teas. The findings revealed that the most common fungal contaminants in the herbal teas were *Aspergillus niger*, *Penicillium* spp., *Eurotium rubrum*, *Eurotium chevalieri*, *Aspergillus flavus*, *Fusarium* spp., *A. alternata* and yeasts. Also, a South African study has reported contamination of herbal products with bacteria as well as fungi such as *Penicillium* and *Aspergillus* (Govender et al., 2006). Examination of pathogenic microorganisms in medicinal herbal drugs has equally shown that the most abundant fungi species were from *Fusarium*, *Aspergillus* and *Alternaria* according to Stevic et al. (2012). In addition, the fungal contamination of powdered herbal medicinal preparations sold in some parts of Nigeria was evaluated, the results showed that all of the herbal preparations had the presence of fungal contaminants with predominance of *Aspergillus* spp. and *Penicillium*

spp., while *Mucor* spp., *Candida* spp. and *Trichosporium* species were also present (Anyanwu, 2010). Similar to this study, is the findings of Zheng et al. (2017) which revealed that the surface of medicinal herbs are predominantly contaminated with species of *Aspergillus* and *Penicillium*.

Aspergillus is a group of moulds found in natural environment; it is an airborne fungus capable of causing Aspergillosis. Species of this genus are highly aerobic, possesses the ability to grow where high osmotic pressure exist and are found in oxygen rich environment. They are capable of growing at low water content; thus, to avoid their growth, quick drying of the herbs are highly important (Stevic et al., 2012). These species are detectable in the ground, air and in plants. *Aspergillus* does not normally cause infection except in an immunocompromised individual such as leukaemia, asthma, HIV/AIDS and in people with damaged lungs due to TB infection, thus causing severe pulmonary disease (WHO, 2011). According to WHO Bulletin, about one-third of TB patients develop cavities in their lungs, thus making them vulnerable to the infection (WHO, 2011). The presence of this *Aspergillus* in the remedies could be detrimental to the health of TB-patients considering their immunosuppressed status.

Species of *Fusarium* were also detected in four of the remedies. In a study carried out by Stevic et al. (2012), *Fusarium* was observed as the most dominant genus in most of the herbal drugs tested. The spores of this organism can survive drying conditions and remain dormant for several months. They are found abundantly in the soil and many of them are important plant pathogens causing various diseases such as crown rot, head blight and scab on cereal grains (Nelson et al., 1994). Some strains such as *Fusarium fumonisins* and *Fusarium trichothecenes* are toxins producers; however, they were not identified in these remedies. Several species of *Fusarium* have emerged as important opportunistic pathogens in humans causing a broad spectrum of infections such as hyalohyphomycosis, mycotic keratitis, onychomycosis, pneumonia, disseminated infections and sinusitis mostly in immunocompromised patients (Makowsky et al., 2005; Guarro, 2013). Species of *Fusarium* identified in this study are *Fusarium oxysporum*, *Fusarium verticillioides* and *Fusarium delphinoides*. Many studies have reported cases of Fusariosis in patients with acute myeloid and lymphoblastic leukemia (Jossi et al., 2010) however; there is dearth information on *Fusarium* infections associated with tuberculosis patients. Considering the immune-deficiency of the patients, the consumption of these remedies over a long period and prolong storage should be avoided in order to prevent the production of mycotoxins in the remedies.

Alternaria, *Candida* and *Ramularia* species were also found in all the tested remedies. *Alternaria* spp. are plant pathogens and field mycotoxin-producing moulds (Stevic et al., 2014). They are capable of producing tenuazonic

acid and other toxic metabolites which may be associated with diseases in humans or animals. These organisms have been reported as causative agents of subcutaneous phaeohyphomycosis, mycotic keratitis, hypersensitivity pneumonitis and extrinsic asthma (Crissey et al., 1995). They equally cause infections such as allergic bronchopulmonary mycosis and pheohyphomycotic lumbar spondylodiscitis in immunocompromised individuals. Therefore, it is important to maintain these remedies at the temperature or condition that would not support the growth this organism.

Candida spp. are yeasts and the most common cause of global fungal infections called Candidiasis (Manolakaki et al., 2010). They are commensals that colonize the skin, gastrointestinal and reproductive tracts. The species of *Candida* identified in this study are non-pathogenic strains but are capable of causing infections in immunocompromised patients. Species of these organisms have been reported as emerging pathogenic fungal in patients with pulmonary tuberculosis (Kali et al., 2013; Ndukwu et al., 2016). Since *Candida* spp can be found on the skin, coupled with the unsterile mode of preparation of these remedies, it could be said that this organism was introduced into the remedies by the handler during the process of preparation. The presence of these fungal contaminants in the polyherbal remedies can reduce or inactivate the therapeutic activity of the remedies and possess the potential to adversely affect the tuberculosis patients.

Conclusion

The findings of this study indicated that polyherbal medicines marketed in the study area are contaminated with fungi contaminants. Some of these organisms are capable of causing infections in immunocompromised patients while others are plant pathogens. The contamination of these polyherbal preparations could have resulted from contaminated soil, plants, inadequate drying, unhygienic mode of preparation and possibly prolonged storage. This could probably be a potential health risk to consumers. It is therefore suggested that quality-control measures and safe handling practices be established for medicinal herbs in the province.

ACKNOWLEDGEMENTS

The work was supported by National Research Foundation of South Africa. The authors also acknowledge the traditional healers for selling the polyherbal remedies to us.

CONFLICT OF INTERESTS

The authors have not declared any conflict of interests.

REFERENCES

- Adeleye IA, Okogi G, Ojo EO (2005). Microbial contamination of herbal preparations in Lagos, Nigeria. *Journal of Health, Population and Nutrition* 23(3):296-297.
- Anyanwu CU (2010). Fungal contaminants of powdered herbal drugs sold in parts of Enugu State, Southeast, Nigeria. *Plant Products Research Journal* 14(1):46-50.
- Bateman AC, Jones GR, O'Connell S, Clark FJ, Plummeridge (2002). Massive hepatosplenomegaly caused by *Penicillium marneffeii* associated with human immunodeficiency virus infection in a Thai patient. *Journal of Clinical Pathology* 55(2):143-144.
- Binu S (2008). Uses of pteridophytes among the tribals in pathanamthitta district, Kerala, India. *J. Non Timber Forest Products* 5(2):129-131.
- Buwa LV, Afolayan AJ (2009). Antimicrobial activity of some medicinal plants used for the treatment of tuberculosis in the Eastern Cape Province, South Africa. *African Journal of Biotechnology* 8(23):6683-6687.
- Chege IN, Okalebo FA, Guantai AN, Karanja S (2015). A Quality Control of Hypoglycemic Herbal Preparations in Nairobi, Kenya. *Journal of Pharmacognosy and Phytochemistry* 3(4):16-21.
- Crissey JT, Parish LC, Lang H (1995). *Manual of medical mycology*. Blackwell Scientific, Cambridge, Massachusetts P 263.
- Czech E, Kneifel W, Kopp B (2001). Microbiological status of commercially available medicinal herbal drugs-A screening study. *Planta Medica* 67:263-269.
- Dei-Tutuwa D, Amuna P, Rahman MA (2014). Rapid Detection of Microbial Contamination in Ghanaian Herbal Medicines by PCR Analysis. *Ghana Medical Journal* 48(2):106-111.
- Devi VM, Prasad PN, Kalirajan K (2010). Infrared spectral studies on Siddha drug-Pavala parpam. *International Journal of Pharma and Bio Sciences* 1(4):474-483.
- Dubey NK, Kumar A, Singh P, Shukla R (2008). Microbial contamination of raw materials: A major reason for the decline of India's share in the global herbal market. *Current Science* (00113891):95:717-718.
- Elujoba AA, Odeleye OM, Ogunyemi CM (2005). Traditional medicine development for medical and dental primary health care delivery system in Africa. *African Journal of Traditional, Complementary and Alternative Medicines* 2:46-61.
- Famewo EB, Clarke AM, Afolayan AJ (2016). Identification of bacterial contaminants in polyherbal medicines used for the treatment of tuberculosis in Amatole District of the Eastern Cape Province, South Africa, using rapid 16S rRNA technique. *Journal of Health, Population and Nutrition* 35(1):27.
- Famewo EB, Clarke AM, Afolayan AJ (2017). Ethno-medicinal documentation of polyherbal medicines used for the treatment of tuberculosis in Amathole District Municipality of the Eastern Cape Province, South Africa. *Pharmaceutical Biology* 55(1):696-700.
- Girridher P, Ready SM (1997). Incidence of mycotoxin producers on spices from Andhra Pradesh. *The journal of the Indian Botanical Society* 76:161-164.
- Govender S, Du Plessis-Stoman D, Downing TG, Van de Venter M (2006). Traditional herbal medicines: Microbial contamination, consumer safety and the need for standards. *South African Journal of Science* 102(5):253-255.
- Guarro J (2013). Fusariosis, a complex infection caused by a high diversity of fungal species refractory to treatment. *European Journal of Clinical Microbiology and Infectious Diseases* 32:1491-1500.
- Igbeneghu OA, Lamikanra A (2016). Assessment of the microbial quality of some oral liquid herbal medicines marketed in Ile-Ife, South-western Nigeria. *African Journal of Microbiology Research* 10(38):1618-1624.
- Jossi M, Ambrosioni J, Macedo-Vinas M, Garbino J (2010). Invasive fusariosis with prolonged fungemia in a patient with acute lymphoblastic leukemia: case report and review of the literature. *International Journal of Infectious Diseases* 14(4):354-356.
- Kali A, Charles MP, Noyal MJ, Sivaraman U, Kumar S, Easow JM (2013). Prevalence of *Candida* co-infection in patients with pulmonary tuberculosis. *The Australasian Medical Journal* 6(8):387.
- Kaume L, Foote JC, Gbur EE (2012). Microbial contamination of herbs marketed to HIV-infected people in Nairobi (Kenya). *South African*

- Journal of Science 108(9-10):1-4.
- Kozich JJ, Westcott SL, Baxter NT, Highlander SK., Schloss PD (2013). Development of a dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence data on the MiSeq Illumina sequencing platform. *Applied and Environmental Microbiology* 79(17):5112-5120.
- Kulshrestha R, Gupta CP, Shukla G, Kundu MG, Bhatnagar SP, Katiyar CK (2008). The effect of water activity and storage temperature on the growth of *Aspergillus flavus* in medicinal herbs. *Planta Medica* 74:1308-1315.
- Kuo HC, Su YL, Yang HL, Chen TY (2005). Identification of Chinese medicinal fungus *Cordyceps sinensis* by PCR-single-stranded conformation polymorphism and phylogenetic relationship. *Journal of Agricultural and Food Chemistry* 53(10):3963-3968.
- Lin SJ, Schranz J, Teutsch SM (2001). Aspergillosis case fatality rate: systematic review of the literature. *Clinical Infectious Diseases* 32(3):358-366.
- Louw CAM, Regnier TJC, Korsten L (2002). Medicinal bulbous plants of South Africa and their traditional relevance in the control of infectious diseases. *Journal of Ethnopharmacology* 82(2):147-154.
- Makowsky MJ, Warkentin DI, Savoie ML (2005). Caspofungin and amphotericin B for disseminated *Fusarium verticillioides* in leukemia. *Annals of Pharmacotherapy* 39:1365-1366.
- Manolakaki D, Velmahos G, Kourkoumpetis T, Chang Y, Alam HB, De Moya MM, Mylonakis E (2010). *Candida* infection and colonization among trauma patients. *Virulence* 1(5):367-375.
- Ndukwu CB, Mbakwem-Aniebo C, Frank-Peterside N (2016). Prevalence of *Candida* Co-Infections among Patients with Pulmonary Tuberculosis in Emuoha, Rivers State, Nigeria. *IOSR Journal of Pharmacy and Biological Sciences* 11(5):60-63.
- Nelson PE, Dignani MC, Anaissie EJ (1994). Taxonomy, biology, and clinical aspects of *Fusarium* species. *Clinical Microbiology Reviews* 7(4):479-504.
- Noor R, Huda N, Rahman F, Bashar T, Munshi SK (2014). Microbial contamination in herbal medicines available in Bangladesh. *Bangladesh Medical Research Council Bulletin* 39(3):124-129.
- Okunlola A, Adewoyin BA, Odeku AO (2007). Evaluation of pharmaceutical and microbial qualities of some herbal medicinal products in South Western Nigeria. *Tropical Journal of Pharmaceutical Research* 6:661-670.
- Osei-Adjei G, Hackman HK, Mills-Robertson FC, Tay SCK (2013). Quality assessment of aqueous herbal/medicinal products sold on the Ghanaian market. *Food Science and Quality Management* 20:13-18.
- Sago SK, Little CL, Greenwood M, Mithani V, Grant KA, McLauchlin J, de-Pinna E, Threlfall EJ (2009). Assessment of the microbiological safety of dried spices and herbs from production and retail premises in the United Kingdom. *Food Microbiology* 26(1):39-43.
- Schoch CL, Seifert KA, Huhndorf S, Robert V, Spouge JL, Levesque CA, Miller AN (2012). Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode marker for Fungi. *Proceedings of the National Academy of Sciences* 109(16):6241-6246.
- Sharma A (2001). Irradiation to decontaminate herbs and spices. In: Peter KV (eds) *Handbook of herbs and spices*. Cambridge, UK: Woodhead Publishing pp. 60-73.
- Siakrwar P, Mahajan S, Gupta A, Asthana M, Sharma R (2014). Isolation and Identification of Fungal Contamination in Stored Medicinal Plants. *American Journal of Pharmacology and Pharmacotherapeutics* 1(2):052-058
- Statistics South Africa (2017). TB Statistics for South Africa – National & provincial 2017. TBFACTS.ORG | Information about Tuberculosis, 2017. <https://www.tbfacts.org/tb-statistics-south-africa/> Accessed 2 November 2017.
- Stević T, Pavlović S, Stanković S, Šavikin K (2012). Pathogenic microorganisms of medicinal herbal drugs. *Archives of Biological Sciences* 64(1):49-58.
- Tassaneeyakul W, Razzazi-Fazeli E, Porasuphatana S, Bohm J (2004). Contamination of aflatoxins in herbal medicinal products in Thailand. *Mycopathologia* 158(2):239-244.
- Ting A, Chow Y, Tan W (2013). Microbial and heavy metal contamination in commonly consumed traditional Chinese herbal medicines. *Journal of Traditional Chinese Medicine* 33(1):119-124.
- Toma FM, Abdulla NF (2013). Isolation and identification of fungi from spices and medicinal plants. *Research Journal of Environmental and Earth* 5(3):131-138.
- Tournas VH, Katsoudas EJ (2008). Microbiological quality of various medicinal herbal teas and coffee substitutes. *Microbial Insights* 1:MBI-S943.
- Walther C, Marwa KJ, Seni J, Hamis P, Silago V, Mshana SE, Jande M (2016). Microbial contamination of traditional liquid herbal medicinal products marketed in Mwanza city: magnitude and risk factors. *Pan African Medical Journal* 23(1).
- White TJ, Bruns T, Lee SJWT, Taylor JW (1990). Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. *PCR protocols: a guide to methods and applications*. 18(1):315-322.
- World Health Organization (2016). *Global Tuberculosis Control*. Geneva, 2016 www.who.int/tb/en. Accessed 2 November 2017.
- World Health Organization (WHO) (2017). *Bending the curve - ending TB: Annual report 2017*. India: World Health Organization, Regional Office for South-East Asia; 2017. apps.who.int/iris/bitstream/10665/254762/1/978929022584-eng.pdf
- World Health Organization (WHO)/Bulletin (2011). *Tuberculosis survivors at risk of fungal infection*. <http://www.who.int/bulletin/releases/NFM1211EN.pdf?ua=1> Accessed 08/11/2017.
- Zheng RS, Wang WL, Tan J, Xu H, Zhan RT, Chen WW (2017). An investigation of fungal contamination on the surface of medicinal herbs in China. *Chinese Medicine* 12(1):2.

Related Journals:

