


OPEN ACCESS



Journal of
Dentistry and Oral Hygiene

June 2018
ISSN 2141-2472
DOI: 10.5897/JDOH
www.academicjournals.org



**ACADEMIC
JOURNALS**
expand your knowledge

ABOUT JDOH

The Journal of Dentistry and Oral Hygiene (JDOH) is published monthly (one volume per year) by Academic Journals.

Journal of Dentistry and Oral Hygiene (JDOH) is an open access journal that provides rapid publication (monthly) of articles in all areas of the subject such as endodontics, prosthodontics, Oral Pathology, dento-maxillofacial Radiology, periodontics etc.

The Journal welcomes the submission of manuscripts that meet the general criteria of significance and scientific excellence. Papers will be published shortly after acceptance. All articles published in JDOH are peer - reviewed.

Contact Us

Editorial Office: jdoh@academicjournals.org

Help Desk: helpdesk@academicjournals.org

Website: <http://www.academicjournals.org/journal/JDOH>

Submit manuscript online <http://ms.academicjournals.me/>

Editors

Dr. Denise Evans

*University of Johannesburg
4 Vlei street, Southcrest Alberton, 1449 Gauteng
South Africa.*

Prof. Azza A El-Housseiny

*Faculty of Dentistry, Alexandria university 1
Shampion St, Massarita, Alexandria. Egypt .*

Dr. Fawad Javed

*Karolinska Institutet
Box 4064, SE 14104 Huddinge. Stockholm.
Sweden*

Dr. Muneer Gohar Babar

*International Medical University
No. 126, Jalan 19/155B, Bukit Jalil, 57000 Kuala
Lumpur,
Malaysia.*

Dr. Janine Owens

*University of Sheffield, Department of Oral Health
and Development
Claremont Crescent, Sheffield S10 2TU
United Kingdom*

Dr. Satyabodh S.Guttal

*SDM College of Dental Sciences and Hospital
Sattur, Dharwad
India .*

Prof. E.J. Sauvetre

*Université Libre de Bruxelles (Faculty of Medicine) 10
rue bavaastro 06300 Nice France
France.*

Dr. Olfat Shaker

*Faculty of Medicine Cairo University
Department of Medical Biochemistry,
Egypt .*

Prof. Ayyaz Ali Khan

*Riphah International University IIDC&H, Off 7th Ave.,
G-7/4, Islamabad, Pakistan.*

Dr. Pei-Yi Chu

*Diagnostic and research pathologist,
Department of Surgical Pathology,
Changhua Christian Hospital/
135 Nan-Shiao Street, Changhua 500-
06, Taiwan.*

Dr. Abeer Gawish

*Al-Azhar University Faculty Of Dental Medicine
4 elsheikh Makhlof Street. Misr Elgedida,
Cairo Egypt.*

Dr. Murali Srinivasan

*Jebel Ali Hospital
PO Box 49207, Dubai,
UAE .*

Dr. Scardina Giuseppe Alessandro

*University of Palermo Department of Oral
Sciences "G. Messina"
Via del Vespro, 129 90127
Palermo Italy.*

Prof. Hussam M. Abdel-Kader

*Faculty of Dental Medicine, AlAzhar University, Cairo,
Egypt Madent Nasr 11884, Cairo,
Egypt.*

Dr. Mahmoud K. ALiri,

*BD-OmS, PhD, FDS RCS (England),
Jordanian Board, DCE (Ireland).
Associate Professor and Consultant
Faculty of Dentistry, The University of Jordan,
P.O. Box 710193, Amman 11171,
Jordan .*

Dr. Imtiaz Wani

*S.M.H.S Hospital
Srinagar, Kashmir,
India.
.*

Editorial Board

Dr. Santhosh Kumar

*Darshan Dental College and Hospital
Udaipur,
India .*

Prof. Sharon Struminger

*2350 Broadhollow Road Farmingdale, NY 11735/
Farmingdale State College of State
University of New York
USA.*

Dr. Saurab Bither

*Christain Dental College
C.M.C, Ludhiana. Punjab.
India.*

Dr. Ramesh Chowdhary

*HKE's S.Nijalingappa Institute of Dental Sciences and
Research
Ring Road, Gulbarga-585104,
India.*

Journal of Dentistry and Oral

Table of Contents: Volume 10 Number 1 June 2018

ARTICLE

Assessment of the soft tissue chins thickness with different skeletal vertical patterns in Pakistani adults

1

Anam Sattar, Imtiaz Ahmed and Taskeen Khan

Full Length Research Paper

Assessment of the soft tissue chin thickness with different skeletal vertical patterns in Pakistani adults

Anam Sattar*, Imtiaz Ahmed and Taskeen Khan

Department of Orthodontics, Dr. Irshat-Ul -Ebad Khan Institute of Oral Health Sciences, DOW University of Health Sciences, Karachi, Pakistan.

Received 26 February, 2016; Accepted 12 July, 2016

The aim of the study is to evaluate and compare the soft tissue chin thickness values among adult patients with different vertical growth patterns. A sample size of 180 adults patients (32 males and 148 females), with an average age of 21.42 years (range 17- 32 years), was selected. The sample was divided into three groups according to the vertical growth pattern using SNMP angle (hypodivergent $<27^\circ$, normodivergent $27-37^\circ$ and hyperdivergent $>37^\circ$). The soft tissue chin thicknesses in each group were measured at pogonion (Pog), gnathion (Gn), and menton (Me) and analyzed using the one-way analysis of variance and post-hoc Tukey test. The soft tissue chin thickness values were greater in hypodivergent group at Pog (12.71 ± 2.10 mm), Gn (9.72 ± 2.55 mm) and Me (9.13 ± 2.88 mm) and smallest in hyperdivergent group (Pog 8.05 ± 1.20 mm, Gn 6.07 ± 1.47 mm, Me 5.91 ± 1.21 mm). The soft tissue chin thicknesses were greater in men than women. Soft tissue chin thickness was greater in hypodivergent adults than those of normodivergent and hyperdivergent adults. In all the three groups, soft tissue thickness values were greater in men than women.

Key words: Vertical pattern, soft tissue, thickness, adult, chin.

INTRODUCTION

The vertical pattern plays a crucial role in diagnosis and treatment planning in both orthodontic growing and adult patients (Opdebeeck and Bell, 1978; Schendel et al., 1976). The development of vertical pattern has been related to several factors, like the growth of jaws, dentoalveolar development, eruption of the teeth, and function of the tongue and lips (Nielsen, 1991). Three basic types of vertical growth pattern are said to exist: Hypo-divergent (low angle), normo-divergent (average), and hyper-divergent (high angle) (Fields et al., 1984). Those with a hypo-divergent growth pattern have reduced vertical growth which is typically associated with

short face, deep anterior overbite, increased posterior to anterior facial height ratio, decreased lower facial height and reduced sella-nasion (SN)/mandibular plane (MP) angle (SN-MP) angle (Opdebeeck and Bell, 1978). The hyper-divergent growth pattern have increased vertical growth with long face, anterior open bite, decreased posterior to anterior facial height ratio, increased lower facial height and increased SN-MP angle (Schendel et al., 1976). Normo-divergent growth pattern lies between these two types.

Bony and soft tissue structures help in the determination of facial harmony and esthetics (Stephan

*Corresponding author. E-mail: anamsattar_88@yahoo.com.




Figure 1. Lateral cephalometric tracing showing angular (left) and linear measurements (right). Left: SN (sella to nasion representing anterior cranial base), FH (Frankfurt horizontal plane; porion to orbitale), PP (palatal plane; ANS to PNS), MP (mandibular plane; gonion to menton). Right: Pog-Pog'= length between bony pogonion (Pog) and its horizontal projection (Pog') over the vertical passing through soft tissue pogonion; Gn-Gn'= distance between bony gnathion (Gn) and soft tissue gnathion (Gn'); and Me-Me'= distance between bony menton (Me) and its vertical projection (Me') on the horizontal passing through soft tissue menton.

and Simpson, 2008); with the soft tissue structures has a major visual influence on facial esthetics (Dumont, 1986). Variation between the bony structures and its overlying soft tissue position may affect the treatment outcomes ranging from camouflage to orthognathic surgery (Kamak and Celikoglu, 2012; Ramos et al., 2005; Ramos et al., 2005).

Genioplasty is a method of recontouring the chin by changing its shape, or size, or both, in the horizontal direction, for esthetic purpose (Sarver et al., 2003; Rosen, 2007) therefore, precise soft tissue measurement is necessary for surgical outcomes (Cha, 2013). Different studies showed soft tissue thickness changes after orthognathic surgery in patients with thick and thin soft tissues (Shaughnessy et al., 2006; Reddy et al., 2011; Abeltins and Jakobson, 2011).

The aims and objectives of this study were to: (1) Evaluate and compare the soft tissue thickness values at different chin levels with different vertical growth patterns among the adult patients in a sample from Pakistani population and; (2) The difference in soft tissue chin thickness between men and women.

MATERIALS AND METHODS

The study was cross-sectional comparative, carried out on 180

pretreatment lateral cephalograms (32 male and 148 female) of adult Pakistani subjects. Mean age of subjects was 21.42 ± 3.178 years. Data was collected from patients coming to Orthodontics Department at Dr. Ishrat-ul-Ebad Khan Institute of Oral Health Sciences, DUHS for orthodontic treatment. Signed consent to use the radiographs was obtained from the patients, before the study was conducted.

The following inclusion and exclusion criteria were used for the selection of the subjects for this study: Inclusion criteria were Pakistani adults of age above 18 years, the lateral cephalometric radiograph of each subject was taken by the same operator using a single Cephalometer, Rotograph Plus at 80 kvp, 10 mA and 0.8 s exposure time using 8 × 10 inch Kodak green film. For exact calculation of mid-sagittal enlargement a scale of known dimensions was attached to the Cephalostat. All subjects were positioned in the cephalostat with the sagittal plane at a right angle to the path of the X-rays, the Frankfurt plane parallel to the horizontal, the teeth in centric occlusion, and the lips slightly closed with no lip strain, and well defined chin structures on the radiograph. Previous orthodontic and/ or orthognathic surgery treatment, presence of craniofacial anomaly like cleft lip and palate, TMJ abnormality, syndromes or presence of a noncontinuous soft tissue contour at the level of the chin indicating a chin strain were not included.

Tracing and measurement on cephalometric radiograph was done by a single investigator. Three linear and four angular measurements were measured on each radiograph. The landmarks were located and the following measurements were used (Figure 1), the angular measurements are:



Figure 2. Lateral cephalometric radiographs with different vertical skeletal pattern using SN-MP angle. Normodivergent group 27-37; hypodivergent group < 27; and hyperdivergent group >37.

Table 1. Description of the sample.

Parameter	Groups				P- value
	Total	Hypodivergent group MP/SN =<27	Normodivergent group 27<MP/SN <37	Hyperdivergent group MP/SN >37	
Total sample					
N	180	60	60	60	
Age, years (range)	21.42 (17-32)	21.02 (18-27)	21.97 (17-32)	21.27 (18-32)	NS
Men					
N	32	13	10	9	
Age, years (range)	20.91 (18-29)	19.85 (18-24)	22.70 (18-29)	20.44 (18-22)	NS
Women					
N	148	47	50	51	
Age, years (range)	21.53 (17-32)	21.34 (18-27)	21.82 (17-32)	21.41 (18-32)	NS

MP/SN, Mandibular plane to anterior cranial base. NS, not significant.

1. Mandibular plane to anterior cranial base (SN-MP).
2. Palatal plane to mandibular plane (MMA)
3. Mandibular plane to horizontal (FHMP), and
4. The ANB angle for the assessment of the sagittal relationship between the jaws.

The linear measurements for soft tissue thickness at chin were measured at three different levels:

1. Pog-Pog'= length between bony pogonion (Pog) and its horizontal projection (Pog') over the vertical passing through soft tissue pogonion;
2. Gn-Gn'= distance between bony gnathion (Gn) and soft tissue gnathion (Gn'); and
3. Me-Me'=distance between bony menton (Me) and its vertical projection (Me') on the horizontal passing through soft tissue menton.

Patients were divided into three groups based on vertical growth pattern using the SN-MP angle (Figure 2).

The sample included 60 patients in each group: Hyperdivergent group (9 men and 51 women; mean age, 21.27 ±3.34 years), hypodivergent group (13 men and 47 women; mean age,

21.02±2.37 years), and normodivergent group (10 men and 50 women; mean age, 21.97±3.64 years).

Statistical analysis

All statistical analyses were done using the SPSS software package (SPSS for Windows 98, version 16.0, SPSS Inc., Chicago, IL). Arithmetic mean and standard deviation were calculated for each variable. Shapiro-Wilk statistics showed that all data were normally distributed; thus, parametric tests were used for further comparisons. One-way analysis of variance (ANOVA) was used to compare soft tissue chin thickness and different cephalometric skeletal measurements among three facial divergence groups and post-hoc test (Tukey honestly significant difference) was used for multiple comparisons. Statistical significance level was set at P=0.05.

RESULTS

Table 1 shows the mean age and range of the age of the

Table 2. Comparison of the skeletal and soft tissue chin variables measurements among different vertical pattern.

Parameter	Groups						ANOVA (P)
	Hypodivergent group MP/SN = <27		Normodivergent group 27 < MP/SN < 37		Hyperdivergent group MP/SN > 37		
	Mean	SD	Mean	SD	Mean	SD	
Age, years	21.02	2.37	21.97	3.64	21.27	3.34	NS
Skeletal measurements							
SNMP	23.03	2.44	32.15	3.14	40.73	2.20	<0.001
FHMP	16.38	2.99	24.57	4.39	31.26	2.30	<0.001
MMA	18.02	3.93	25.04	4.71	30.75	4.58	<0.001
ANB	3.67	2.41	4.73	5.63	5.05	2.85	NS
Soft tissue measurements							
Pog-Pog'	12.71	2.10	9.70	1.62	8.05	1.20	<0.001
Gn-Gn'	9.72	2.55	7.03	1.46	6.07	1.47	<0.001
Me-Me'	9.13	2.88	7.13	1.66	5.91	1.21	<0.001

Groups	Comparison among groups		
	Hypodivergent-normodivergent	Hypodivergent-hyperdivergent	normodivergent-hyperdivergent
Age	NS	NS	NS
Skeletal measurements			
SNMP	<.001	<.001	<.001
FHMP	<.001	<.001	<.001
MMA	<.001	<.001	<.001
ANB	NS	NS	NS
Soft tissue measurements			
Pog-Pog'	<.001	<.001	<.001
Gn-Gn'	<.001	<.001	.017
Me-Me'	<.001	<.001	.004

MP/SN, Mandibular plane to anterior cranial base; SD, standard deviation; ANOVA, analysis of variance; NS, not significant; MMA, palatal plane to mandibular plane; FHMP, mandibular plane to horizontal; Pog-Pog', length between bony pogonion (Pog) and its horizontal projection (Pog') over the vertical passing through soft tissue pogonion; Gn-Gn', distance between bony gnathion (Gn) and soft tissue gnathion (Gn'); Me-Me', distance between bony menton (Me) and its vertical projection (Me') on the horizontal passing through soft tissue menton. NS, not significant.

patients in each group and the mean age of men and women within each of the three groups. Age was not statistically significantly different between men and women within each group and across the three groups.

Table 2 shows the comparison of the skeletal and soft tissue chin variables measurements among different vertical patterns. All skeletal and soft tissue chin thickness measurements except nasion-B point (ANB) shows statistically significant differences among the three groups.

Patients in hypodivergent group has thickest soft tissue thickness at Pogonion, Gnathion and Menton ($p < 0.001$), while the hyperdivergent group's patients has the thinnest soft tissue chin thickness at all the three points ($p < 0.001$).

The soft tissue chin thickness measurement of hypodivergent, normodivergent and hyperdivergent group of men and women are presented in Table 3 which shows men have thicker soft tissue chin thickness values as compared to the women.

DISCUSSION

The main purpose of this study is to determine the association between vertical skeletal pattern and soft tissue chin thickness in adults and to compare these values between men and women and also among different vertical skeletal groups. A previous study which evaluated the soft chin thickness in adult patients with various mandibular divergence pattern, showed that the values of soft tissue chin thickness was statistically significant only at point gnathion for both men and women (Macari and Hanna, 2013).

According to Celikoglu et al. (2015) the thickness values was statistically significant in women of high angle group at pogonion only.

In our study, we found that the soft tissue chin thickness values were the thickest in the hypodivergent group and thinnest in the hyperdivergent group for both men and women While in normodivergent group the

Table 3. Comparison of soft tissue thickness at chin between men and women among different vertical pattern

Sex	Soft tissue chin measurements	Groups						ANOVA (P)
		Hypodivergent group		Normodivergent group		Hyperdivergent group		
		MP/SN =<27		27<MP/SN <37		MP/SN >37		
		Mean	SD	Mean	SD	Mean	SD	
Men	Pog-Pog'	12.92	1.42	9.65	0.62	7.83	0.66	<0.001
	Gn-Gn'	10.07	2.83	7.05	1.25	6.33	0.66	<0.001
	Me-Me'	10.57	4.82	8.60	1.34	5.83	0.66	0.008
Women	Pog-Pog'	12.66	2.26	9.72	1.76	8.09	1.28	<0.001
	Gn-Gn'	9.62	2.49	7.03	1.51	6.02	1.57	<0.001
	Me-Me'	8.73	1.95	6.84	1.57	5.93	1.29	<0.001

MP/SN, Mandibular plane to anterior cranial base; SD, standard deviation; ANOVA, analysis of variance; NS, not significant; Pog-Pog', length between bony pogonion (Pog) and its horizontal projection (Pog') over the vertical passing through soft tissue pogonion; Gn-Gn', distance between bony gnathion (Gn) and soft tissue gnathion (Gn'); Me-Me', distance between bony menton (Me) and its vertical projection (Me') on the horizontal passing through soft tissue menton.

thickness values lies between these two groups. The statistically significant differences were found at pogonion ($P < 0.001$), gnathion ($P < 0.001$) for both men and women and at Menton ($P = 0.005$) and ($P < 0.001$) for men and women respectively. The difference between this study with previous studies (Macari and Hanna, 2013; Celikoglu et al., 2015) might be due to racial and ethnic groups. Hyperdivergent patients have deficient chin which required advancement genioplasty in comparison with the hypodivergent patients which do not require advancement genioplasty.

However, hypodivergent patients may require reduction genioplasty, or vertical augmentation genioplasty, or even sliding genioplasty for correction of any possible facial asymmetry. Soft tissue chin thickness is also correlated with sagittal plane discrepancies (Hoffelder et al., 2007). Soft tissue pogonion was influenced by the skeletal pogonion (Tanaka et al., 2011).

According to the soft tissue paradigm, more importance has been given to the soft tissue during diagnosis and treatment planning (Uysal et al., 2012). Soft tissue varies significantly with the difference in mandibular divergence pattern (Scheideman et al., 1980). Soft tissue should be considered before planning orthognathic surgery due to difference in soft tissue responses after surgery between the patients having thick and thin soft tissue (Abeltins and Jakobsone, 2011; Jakobsone et al., 2013; Wen-Ching et al., 2000)

Conclusions

The following conclusions were drawn:

1. Soft tissue chin thickness were greater in adults at pogonion, gnathion and menton in hypodivergent group.
2. These values are smallest in hyperdivergent adults.

3. In all the three groups, soft tissue thickness values were greater in men than women.

CONFLICT OF INTERESTS

The authors have not declared any conflict of interests.

REFERENCES

- Abeltins A, Jakobsone G (2011). Soft tissue thickness changes after correcting Class III malocclusion with bimaxillary surgery. *Stomatologija* 13:87-91
- Celikoglu M, Buyuk SK, Ekizer A, Sekerci AE, Sisman Y (2015). Assessment of the soft tissue thickness at the lower anterior face in adult patients with different skeletal vertical patterns using cone beam computed tomography. *Angle Orthodontist* 85:211-217.
- Cha KS (2013). Soft-tissue thickness of South Korean adults with normal facial profiles. *Korean Journal of Orthodontics* 43:178-185.
- Dumont RE (1986). Mid-facial tissue depths of white children: an aid in facial feature reconstruction *Journal of Forensic Sciences* 31:1463-1469.
- Fields HW, Proffit WR, Nixon WL, Phillips C, Stanek E (1984). Facial pattern differences in long-faced children and adults. *American Journal of Orthodontics* 85:217-223.
- Hoffelder LB, Lima EM, Martinelli FL, Bolognese AM (2007). Soft tissue changes during facial growth in skeletal Class II individuals. *American Journal of Orthodontics and Dentofacial Orthopedics* 131:490-495.
- Jakobsone G, Stenvik A, Espeland L (2013). Soft tissue response after Class III bimaxillary surgery. *Angle Orthodontist* 83:533-539.
- Kamak H, Celikoglu M (2012). Facial soft tissue thickness among skeletal malocclusions: is there a difference? *Korean Journal of Orthodontics* 42:23-31
- Macari AT, Hanna AE (2013). Comparisons of soft tissue chin thickness in adult patients with various mandibular divergence patterns. *Angle Orthodontist* 84:708-714.
- Nielsen IL (1991). Vertical malocclusions: etiology, development, diagnosis and some aspects of treatment. *Angle Orthod.* 61:247-260.
- Opdebeeck H, Bell WH (1978). The short face syndrome. *American Journal of Orthodontics* 73:499-511.
- Ramos AL, Sakima MT, Pinto AS, Bowman SJ (2005). Upper lip changes correlated to maxillary incisor retraction—a metallic implant

- study. *Angle Orthodontist* 75:499-505.
- Reddy PS, Kashyap B, Hallur N, Sikkerimath BC (2011). Advancement genioplasty-cephalometric analysis of osseous and soft tissue changes. *Journal of Oral and Maxillofacial Surgery* 10:288-295.
- Rosen HM (2007). Osseous genioplasty. In: Thorne CH (Eds.). *Grabb and Smith's Plastic Surgery*. Philadelphia, Lippincott Williams & Wilkins, a Wolters Kluwer Business. pp. 557-561.
- Sarver DM, Rousso DR, White RP Jr (2003). Adjunctive esthetic surgery. In: Proffit WR, White RP, Sarver DM (Eds.). *Contemporary Treatment of Dentofacial Deformity*. St Louis, Mo: Mosby pp. 394-415.
- Scheideman GB, Bell WH, Legan HL, Finn RA, ReischJS (1980). Cephalometric analysis of dentofacial normal. *American Journal of Orthodontics* 78:404-420.
- Schendel SA, Eisenfeld J, Bell WH, Epker BN, Mischelevich DJ (1976). The long face syndrome: vertical maxillary excess. *American Journal of Orthodontics* 70:398-408.
- Shaughnessy S, Mobarak KA, Høgevoid HE, Espeland L (2006). Long term skeletal and soft-tissue responses after advancement genioplasty. *American Journal of Orthodontics and Dentofacial Orthopedics* 130:8-17.
- Stephan CN, Simpson EK (2008). Facial soft tissue depths in craniofacial identification (part I): an analytical review of the published adult data *Journal of Forensic Sciences* 53:1257-1272.
- Tanaka O, Fabianski ST, Karakida LM, Knop LAH, Retamoso LB (2011). Changes in pogonion and nose according to breathing patterns. *Dental Press Journal of Orthodontics* 16(6):78-83.
- Uysal T, Baysal A, Yagci A, Sigler LM, McNamara JA Jr (2012). Ethnic differences in the soft tissue profiles of Turkish and European-American young adults with normal occlusions and well-balanced faces. *European Journal of Orthodontics* 34:296-301.
- Wen-Ching Ko E, Figueroa AA, Polley JW (2000). Soft tissue profile changes after maxillary advancement with distraction osteogenesis by use of a rigid external distraction device: a 1-year follow-up. *Journal of Oral and Maxillofacial Surgery* 58:959-969.

Related Journals:

