


OPEN ACCESS

A central image showing a complex, spiky biological structure, possibly a microorganism or a cell, rendered in shades of blue and white against a dark background. The structure has many thin, radiating filaments or spines. A horizontal teal band is overlaid across the middle of the image, containing the journal title.

# African Journal of **Microbiology Research**

28 October 2018  
ISSN 1996-0808  
DOI: 10.5897/AJMR  
[www.academicjournals.org](http://www.academicjournals.org)



**ACADEMIC  
JOURNALS**  
expand your knowledge

# ABOUT AJMR

**The African Journal of Microbiology Research (AJMR)** is published weekly (one volume per year) by Academic Journals.

**The African Journal of Microbiology Research (AJMR)** provides rapid publication (weekly) of articles in all areas of Microbiology such as: Environmental Microbiology, Clinical Microbiology, Immunology, Virology, Bacteriology, Phycology, Mycology and Parasitology, Protozoology, Microbial Ecology, Probiotics and Prebiotics, Molecular Microbiology, Biotechnology, Food Microbiology, Industrial Microbiology, Cell Physiology, Environmental Biotechnology, Genetics, Enzymology, Molecular and Cellular Biology, Plant Pathology, Entomology, Biomedical Sciences, Botany and Plant Sciences, Soil and Environmental Sciences, Zoology, Endocrinology, Toxicology. The Journal welcomes the submission of manuscripts that meet the general criteria of significance and scientific excellence. Papers will be published shortly after acceptance. All articles are peer-reviewed.

## Contact Us

**Editorial Office:** [ajmr@academicjournals.org](mailto:ajmr@academicjournals.org)

**Help Desk:** [helpdesk@academicjournals.org](mailto:helpdesk@academicjournals.org)

**Website:** <http://www.academicjournals.org/journal/AJMR>

**Submit manuscript online** <http://ms.academicjournals.me/>

## Editors

**Prof. Adriano Gomes da Cruz**  
*University of Campinas (UNICAMP),  
Brazil.*

**Prof. Ashok Kumar**  
*School of Biotechnology  
Banaras Hindu University Uttar Pradesh,  
India.*

**Dr. Mohd Fuat Abd Razak**  
*Infectious Disease Research Centre,  
Institute for Medical Research, Jalan  
Pahang, Malaysia.*

**Dr. Adibe Maxwell Ogochukwu**  
*Department of Clinical Pharmacy and  
Pharmacy Management,  
University of Nigeria  
Nsukka, Nigeria.*

**Dr. Mehdi Azami**  
*Parasitology & Mycology Department  
Baghaeei Lab.  
Isfahan, Iran.*

**Dr. Franco Mutinelli**  
*Istituto Zooprofilattico Sperimentale delle  
Venezie Italy.*

**Prof. Ebiamadon Andi Brisibe**  
*University of Calabar,  
Calabar,  
Nigeria.*

**Prof. Nazime Mercan Dogan**  
*Department of Biology  
Faculty of Science and Arts  
University Denizli Turkey.*

**Prof. Long-Liu Lin**  
*Department of Applied Chemistry  
National Chiayi University  
Chiayi County Taiwan.*

**Prof. Natasha Potgieter**  
*University of Venda  
South Africa.*

**Dr. Tamer Edirne**  
*Department of Family Medicine  
University of Pamukkale  
Turkey.*

**Dr Kwabena Ofori-Kwakye**  
*Department of Pharmaceutics  
Kwame Nkrumah University of Science &  
Technology  
Kumasi, Ghana.*

**Dr. Tülin Askun**  
*Department of Biology  
Faculty of Sciences & Arts  
Balikesir University Turkey.*

**Dr. Mahmoud A. M. Mohammed**  
*Department of Food Hygiene and Control  
Faculty of Veterinary Medicine  
Mansoura University Egypt.*

# Editors

**Dr. James Stefan Rokem**  
*Department of Microbiology & Molecular Genetics*  
*Institute of Medical Research Israel – Canada*  
*The Hebrew University – Hadassah Medical School Jerusalem, Israel.*

**Dr. Afework Kassu**  
*University of Gondar*  
*Ethiopia.*

**Dr. Wael Elnaggar**  
*Faculty of Pharmacy*  
*Northern Border University*  
*Rafha Saudi Arabia.*

**Dr. Maulin Shah**  
*Industrial Waste Water Research Laboratory*  
*Division of Applied & Environmental Microbiology, Enviro Technology Limited*  
*Gujarat, India.*

**Dr. Ahmed Mohammed**  
*Pathological Analysis Department*  
*Thi-Qar University College of Science*  
*Iraq.*

**Prof. Naziha Hassanein**  
*Department of Microbiology*  
*Faculty of Science*  
*Ain Shams University*  
*Egypt.*

**Dr. Shikha Thakur**  
*Department of Microbiology*  
*Sai Institute of Paramedical and Allied Sciences India.*

**Dr. Samuel K Ameyaw**  
*Civista Medical Center*  
*USA.*

**Dr. Anubrata Ghosal**  
*Department of Biology*  
*MIT - Massachusetts Institute of Technology*  
*USA.*

**Dr. Bellamkonda Ramesh**  
*Department of Food Technology*  
*Vikrama Simhapuri University*  
*India.*

**Dr. Sabiha Yusuf Essack**  
*Department of Pharmaceutical Sciences*  
*University of KwaZulu-Natal*  
*South Africa.*

**Dr. Navneet Rai**  
*Genome Center*  
*University of California Davis USA.*

**Dr. Iheanyi Omezuruike Okonko**  
*Department of Virology*  
*Faculty of Basic Medical Sciences*  
*University of Ibadan*  
*Ibadan, Nigeria.*

**Dr. Mike Agenbag**  
*Municipal Health Services,*  
*Joe Gqabi,*  
*South Africa.*

**Dr. Abdel-Hady El-Gilany**  
*Department of Public Health & Community Medicine, Faculty of Medicine*  
*Mansoura University*  
*Egypt.*

# **African Journal of Microbiology Research**

**Table of Content:    Volume 12    Number 40    28 October, 2018**

## **ARTICLES**

**Antibacterial activity of *Ocimum gratissimum* (scent leaf)  
on some pathogenic gastrointestinal bacteria**

Agholor Kin, Lucy M. Yaki, Idris Abubakar, Lucy F. Olusola and Rakiya Zubairu

**In vitro study on the anti-microbial efficacy of *Aloe vera* against  
*Candida albicans***

T. Jeevitha, K. Deepa and A. Michael

*Full Length Research Paper*

## **Antibacterial activity of *Ocimum gratissimum* (scent leaf) on some pathogenic gastrointestinal bacteria**

**Agholor Kin<sup>1\*</sup>, Lucy M. Yaki<sup>2</sup>, Idris Abubakar<sup>1</sup>, Lucy F. Olusola<sup>1</sup> and Rakiya Zubairu<sup>1</sup>**

<sup>1</sup>Department of Biological Science, Niger state Polytechnic, Zungeru, Niger State, Nigeria.

<sup>2</sup>Department of Microbiology, Kaduna State University, Kaduna, Kaduna State, Nigeria.

Received 14 March, 2018; Accepted 13 August, 2018

The emergence of antibiotic resistance as well as the recent undesirable side effect of some of the commercially available antibiotics has led to the screening of plant extract in order to discover new drug that could serve as alternative therapy for the treatment of various infections and diseases. Fresh leaf of *Ocimum gratissimum* (scent leaf) sample was collected, air-dried at room temperature and blended to powder using electric blender. The extraction was done using reflux extraction method and methanol as solvent. The phytochemical analysis and the antibacterial activity of *O. gratissimum* were determined to ascertain the different phytochemicals present in the plant extract. The extract was also tested against some selected Gram negative intestinal pathogenic bacteria; *Escherichia coli*, *Shigella* and *Salmonella* species, by reconstituting the extract in dimethyl sulphoxide (DMSO) to obtain different concentration (0.2, 0.1, 0.05 and 0.025 g/ml) and agar well diffusion techniques were used to evaluate the antibacterial susceptibility of the leaf extract. The qualitative phytochemical analysis of the extract revealed the presence of alkaloid, anthraquinone, flavonoid, glycoside, phenol, saponin, steroid and tannins. The result of antibacterial analysis showed that the extract of *O. gratissimum* has antibacterial activity against *E. coli*. This could be as a result of the presence of various phytochemicals or the interaction of one or more of the identified metabolites against the test organisms. However, there was no zone of inhibition (antibacterial effect) recorded on *Salmonella* and *Shigella* spp. as they were resistant to the extract. The results obtained from this research, suggest that *Escherichia coli* was susceptible to the leaf extract and the plant could be used as potential source of natural product for the treatment of infection.

**Key words:** Antibacterial activity, scent leaf, gastrointestinal bacteria, phytochemicals, plant extract.

### **INTRODUCTION**

Pathogenic gastrointestinal bacteria are bacteria that cause gastroenteritis (Okigbo and Igwe, 2007). They infect the gut leading to inflammation of the stomach and intestines (Ishiwu et al., 2014). This leads to vomiting, severe abdominal cramps, and diarrhea. They include

*Escherichia coli*, *Shigella* species, and *Salmonella* species. Bacterial gastroenteritis commonly occurs as a result of poor hygienic practices (Russell and Jarvis, 2011). However, infections can also occur after close contact with infected animals or consuming food or water

\*Corresponding author. E-mail: kinagholor@gmail.com. Tel: +2347062361650.

contaminated with bacteria or the toxic substances produced by bacteria (Opara et al., 2014).

*E. coli* are commonly found in fecaloid matters and can cause serious food poisoning in their hosts (Kotloff et al., 2013).

*Shigella* spp. causes shigellosis, commonly referred to as bacterial dysentery (Ram et al., 2008).

*Salmonella* spp. are facultative intracellular pathogens. They are two serotypes; the non-typhoidal and typhoidal serotypes. The non-typhoidal serotype invades only the gastrointestinal tract and cause *Salmonella* food poisoning while the typhoidal serotype spreads throughout the body, invades organs, and secretes endotoxins (Su and Chiu, 2007).

Antibacterial are forms of antimicrobial agent used especially against bacteria, for the treatment of bacterial infections (Prabhu et al., 2009). The discovery of antibiotics (a substance produced by microbes which inhibit or kill another microorganism at a very low concentration) has helped in the control of pathogenic bacteria until the recent development of resistance by most pathogens. Antibiotic resistance is becoming a worldwide problem posing danger to humanity as various diseases and infections that are formerly treated with this substance are now difficult to control (Suree and Pana, 2015).

In addition, some synthetic available antibacterial agents are becoming ineffective due to their side effects such as tendonitis, seizure, and Steven-Johnson syndrome (WHO, 2002). The emergence of antibiotic resistance as well as the recent undesirable side effect of some of the commercially available antibiotics has led to the screening of plant extracts in search for new drug that could serve as alternative therapy for the treatment of various infections and diseases (Effraim et al., 2013).

*Ocimum gratissimum* popularly referred to as scent leaf because of its aroma is commonly used as spices for food or soup preparation in Nigeria (Akinjogunla et al., 2009). It is a medicinal plant which has been used traditionally for the treatment of various infections (Abdullahi, 2012). The plant is cultivated in abundant in different part of Nigeria and it contains some bioactive substances such as tannis, saponins, alkaloids, glycosides, phenols and flavonoids, also referred to as phytochemicals.

These phytochemicals when consumed served as medicine for protection and treatment of human or animal disease (Abdullahi, 2012). The *in-vitro* antimicrobial screening of *O. gratissimum* against *Staphylococcus aureus*, *E. coli*, *Streptococcus fecalis*, *Pseudomonas aeruginosa* and *Lactobacilli* showed that the leaf extract is effective against human pathogens (Prabhu et al., 2009).

Hence, the 'Green' Movement in Western Society has established that naturally derived substances are safer and more desirable than synthetic chemicals products (Opara et al., 2014). Therefore, this study aimed at

determining the phytochemical constituent and antibacterial activity of *O. gratissimum* leaf extract on these pathogenic gastrointestinal bacteria; *Escherichia coli*, *Shigella* spp., and *Salmonella* spp.

## MATERIALS AND METHODS

### Collection and preparation of plant

Fresh samples of *O. gratissimum* leaves were collected from Niger State Polytechnic Staff Quarter, Wushishi Local Government, Niger State, on coordinate 9° 48'N 6° 9' E and elevation 149 m (489 ft) on 16 October, 2017. The plant was identified and authenticated in the Department of Biological Sciences, Niger State Polytechnic, Zungeru. The fresh leaves were dried completely for two weeks at room temperature (Figure 2) and were blended into powder form using an electric blender.

### Extraction process

The plant extraction was obtained using reflux extraction technique as described by Abdullahi (2012) and Sofowora (1993). 250 g of the blended plant leaf powder was weighed and wrapped in Whatman No. 1 filter paper and placed in the holding chamber of the reflux extractor. 600 ml of methanol was used for the extraction at 40°C for 48 h. Thereafter, the extract was concentrated by evaporating to dryness using water bath. The dark green coloured solid extract of the *O. gratissimum* was stored in an airtight container at 4°C in a refrigerator.

### Qualitative phytochemical analysis of the methanolic extract of *O. gratissimum*

The qualitative and quantitative phytochemical screening of the methanolic leaf extract were carried out following standard procedures according to the method described by Abdullahi (2012) and Okwu (2005) at the Centre for Genetic Engineering and Biotechnology Laboratory, Federal University of Technology Minna, Niger State.

### Determination of antibacterial activity of the plant extract

#### Reconstitution of the plant extract

The methanol plant extracts were reconstituted by weighing 0.2, 0.1, 0.050 and 0.025 g each into different sterile test tubes containing 1 ml of dimethyl sulphoxide (DMSO), respectively to obtain the following concentration of the extract: 0.2, 0.1, 0.05, and 0.025 g/ml.

#### Test for sterility of the plant extract

The plant extracts were tested for sterility by introducing 1 ml of the reconstituted plant extract into 5 ml of sterile nutrient broth and were incubated at 37°C for 24 h. The absence of turbidity of the broth (as compared to MacFarland standard) after incubation indicated that the extract was sterile (Preethi et al., 2010).

### Collection and confirmation of test organisms

The pure clinical isolates of some pathogenic gastrointestinal

**Table 1.** Qualitative and quantitative phytochemical constituent of the methanolic extract of *Ocimum gratissimum*.

Constituent	Qualitative analysis	Quantitative analysis
Saponin	+	52.48 ± 0.00
Flavonoid	+	5.33 ± 0.00
Steroid	+	N D
Terpenoids	-	N D
Glycoside	+	29.91 ± 0.03
Phenol	+	27.00 ± 0.01
Alkaloid	+	0.15 ± 0.03
Tannin	+	N D
Anthraquinones	+	N D

(+) = Positive, (-) = Negative; ND: Not determined.

**Table 2.** Antibacterial activity of methanolic extract of *Ocimum gratissimum* on the test organisms at different concentrations.

Test organism	Average zone of inhibition in diameters (mm)				Ciprofloxacin	DMSO
	0.20 g/ml	0.10 g/ml	0.05 g/ml	0.025 g/ml	10 mcg	
<i>Escherichia coli</i>	24	19	12	8	20	NI
<i>Salmonella</i> sp.	NI	NI	NI	NI	25	NI
<i>Shigella</i> sp.	NI	NI	NI	NI	23	NI

NI: No inhibition; Mcg: Micrograms.

bacteria belonging to the family Enterobacteriaceae: *Escherichia coli*, *Salmonella* spp. and *Shigella* spp. were obtained from the Department of Medical Microbiology, Kaduna State University, Kaduna State, Nigeria. All the isolates were checked for purity and confirmed by Gram staining and sub culturing them on selective media (*Salmonella shigella* Agar and MacConkey Agar); thus, observing the colony characteristics and morphology of the cells.

#### Antibacterial susceptibility assay

Agar well diffusion techniques were employed for the antimicrobial testing of the plant extracts. The 24 h old cultures were transferred into nutrient broth and incubated at 37°C for 5 h and standardized to Macfarland standard. Each of the test organisms from the broth cultures were streaked on 4 different Mueller Hinton agar plates under aseptic condition and were labeled accordingly.

Wells of approximately 5 mm in diameter were made on the surface of the inoculated agar medium using a sterile cork borer no. 1 and the wells labeled with a marker based on the concentration of the plant extract (0.2, 0.1, 0.05, and 0.025 g/ml) and the wells were filled with the different concentration of the extract. DMSO and ciprofloxacin were used as the negative and positive control, respectively as described by Lino and Deogracious (2006).

The plates were incubated at 37°C and the susceptibility of the test organisms to the plant extract were recorded after 24 h by measuring the average diameter of the clear zone of inhibition in millimeters (mm).

## RESULTS

The result of qualitative phytochemical analysis of methanolic extract of *O. gratissimum* leaves revealed the

presence of eight classes of secondary metabolites. However, the quantitative phytochemical analysis conducted on five classes of the identified metabolites showed that saponin has the highest phytochemical constituent (52.48 mg/g) of the extract, followed by tannin (29.91 mg/g), phenol (27.00 mg/g), flavonoid (5.33 mg/g) and alkaloid (0.15 mg/g) shown in Table 1. However, quantitative phytochemical analysis for anthraquinones, tannin, terpenoids and steroid were not determined.

The result of antibacterial activity of methanolic extract of *O. gratissimum* on *E. coli*, *Salmonella* spp. and *Shigella* spp. shown in Table 2, *E. coli* was sensitive to the extract while *Salmonella* spp. and *Shigella* sp. were resistant to the plant extract and there was no zone of inhibition of DMSO against the test organisms. However, the lower the concentration of the extract (from 0.20 to 0.025 g/ml), the lower the average zone of inhibition (from 24 to 8 mm) against *E. coli*.

The result of antibacterial activity of methanolic extract of *O. gratissimum* on *E. coli* shows decreased average zone of inhibition with decreased concentration of the extract (Figure 1), that is, the lower the concentration, the lower the antibacterial activity.

## DISCUSSION

The extraction of *O. gratissimum* leaf was obtained using methanol as solvent and the phytochemical analysis of the leaf extract revealed the presence of alkaloid,






Figure 1. Antibacterial activity of methanolic extract of *O. gratissimum* leaf on *E. coli*.



Figure 2. Collection and preparation of *O. gratissimum* leaf (scent leaf).



**Figure 3.** Antibacterial activity of *O. gratissimum* leaf extract on *E. coli*.



**Figure 4.** Antibacterial activity of *O. gratissimum* leaf extract on *Salmonella* sp.

anthraquinone, flavonoid, glycoside, phenol, saponin, steroid and tannins which agrees with the finding of Ladipo et al. (2010) that shows that *O. gratissimum* contains all the phytochemicals mentioned earlier.

The antibacterial analysis of the methanolic extract of *O. gratissimum* (Figure 3) revealed that the extract has antibacterial activity against *E. coli* shown in Table 2. This could be as a result of the presence of one of the phytochemicals or the interaction of two or more of the bioactive compounds against the test organism which agreed with the work of Abdullahi (2012) and also supported the traditional uses of the plant in the treatment of various bacterial enteric diseases such as

diarrhea, dysentery and other gastrointestinal infections (Nwinyi et al., 2009). Hence, from the result obtained, there was decreased in antibacterial activity with decreased in concentration of the extract as shown in Figure 1; as the concentration of the extract decreases from 0.20l to 0.025 g/ml, the average zone of inhibition also decreased from 24 to 8 mm. This also agreed with the finding of Ishiwu et al. (2014) who demonstrated that increase in the concentration of *O. gratissimum* extract reduces the number of viable *E. coli* from 36 to 5 cfu/ml.

However, there was no antibacterial effect of the extract on *Salmonella* (Figure 4) and *Shigella* spp. (Figure 5). This resistance may be due to the lipid content




Figure 5. Antibacterial activity of *O. gratissimum* leaf extract on *Shigella* spp.

on the membranes of these bacteria that prevented the permeability of the active phytochemicals into the cell or the low quantity of alkaloid ( $0.15 \pm 0.03$ ) extracted with methanol (Abdullahi, 2012). However, this disagreed with the work of Ladipo et al. (2010), who reported that the methanol extract of *O. gratissimum* leaf has antibacterial activity against *Salmonella*, *Shigella* and *Klebsiella* spp. But Justina and Solomon (2017) suggested that difference in phytochemical constituent of the leaf of *O. gratissimum* could be as a result of the planting location, seasonal and environmental variations. This could also have effect on the antibacterial activity of the leaf extract on *Salmonella* and *Shigella* spp. Therefore, the leaf extract of *O. gratissimum* could serve as natural antibacterial agent and herbal drug against gastroenteritis.

## Conclusion

The results obtained in this study, suggest that *Escherichia coli* was susceptible to the plant leaf extract and the methanolic extract of *O. gratissimum* leaf contains phytochemicals that possess antibacterial property and could be used as potential source of natural product in industrial manufacturing of drugs, for the treatment of infections/diseases caused by *E. coli*. However, the toxicity and the side effects of the plant leaf extract should be determined even if the plant is consumed locally.

## CONFLICT OF INTERESTS

The authors have not declared any conflict of interests.

## REFERENCES

- Abdullahi M (2012). Phytochemical constituents and antimicrobial and grain protectant activities of Clove Basil (*Ocimum gratissimum* L.) grown in Niger. *Journal of Plant Research* 2(1):51-58.
- Akinjogunla OJ, Adegoke AA, Udokang IP, Adebayo-Tayo B (2009). Antimicrobial potential of *Nymphaea lotus* (Nymphaeaceae) against wound pathogens. *Journal of Medicinal Plants Research* 3(3):138-141.
- Effraim KD, Jacks TW, Sopio OA (2013). Histopathological Studies on the Toxicity of *Ocimum Gratissimum* Leaves Extracts on Some Organs of Rabbits. *African Journal of Biomedical Research* 6:21-25.
- Ishiwu CN, Umenwanne CP, Obiegbuna JE, Uchegbu N (2014). *In vitro* assessment of antibacterial effect of extracts of *Ocimum gratissimum* and *carica papaya* leaves. *International Journal of Applied Science and Technology* 1(4).
- Justina YT, Solomon AM (2017). Proximate, Phytochemical, and In-Vitro Antimicrobial Properties of Dried Leaves from *Ocimum gratissimum*. *Nutrition and Food Science* 22(3):191-194.
- Kotloff KL, Nataro JP, Blackwelder WC (2013). Burden and aetiology of diarrhoeal disease in infants and young children in developing countries): a prospective, case-control study. *The Lancet* 382(9888):209-222.
- Ladipo MK, Doherty VF, Kanife UC (2010). Phytochemical Screening and antibacterial Investigation of The Extract of *Ocimum gratissimum* (Scent Leaf) On Selected *Enterobacteriaceae*. *PAT* 6(2):75-84.
- Lino A, Deogracious O (2006). The *in vitro* antibacterial activity of *Annonas negalensis*, *Sacuridecae longipendiculata* and *Steganotaema araliacea*. *The Journal of African Health Sciences* 6(1):31-35.
- Nwinyi OC, Chinedu NS, Ajani OO, Ikpo CO, Ogunniran KO (2009). Antibacterial effects of extracts of *Ocimum gratissimum* and *Piper guineense* on *Escherichia coli* and *Staphylococcus aureus*. *African Journal of Food Science* 3(3):071-081.
- Okigbo RN, Igwe M (2007). The antimicrobial effects of *Piperguineense uziza* and *Phyllantusamarus ebe-benizo* on *Candida albicans* and *Streptococcus faecalis*. *Acta microbiologica et immunologica Hungarica* 54(4):353-366.
- Okwu DE (2005). Chemical Composition of *Spondias Mombia* Linn Plant Parts. *Journal of Sustainable Agriculture and the Environment* 6:140-147.
- Opara A, Egbuobi R, Dike J, Ndudim E, Onyewuchi C, Nnodim J

- (2014). Antibacterial Activity of *Ocimum gratissimum* (Nchu-Anwu) and *Vernonia amygdalina* (Bitter-Leaf). *British Biotechnology Journal* 4(10):1115-1122.
- Prabhu KS, Lobo R, Shirwaikar AA, Shirwaikar A (2009). *Ocimum gratissimum*: A Review of its Chemical, Pharma-cological and Ethnomedicinal Properties. *The Open Complementary Medicine Journal* 1:1-15
- Preethi RM, Devanathan VV, Loganathan M (2010). Antimicrobial and antioxidant efficacy of some medicinal plants against food borne pathogens. *Advanced Biomedical Research* 4:122-125.
- Ram PK, Crump JA, Gupta SK, Miller MA, Mintz ED (2008). Analysis of Data Gaps Pertaining to *Shigella* Infections in Low and Medium Human Development Index Countries 1984–2005. *Epidemiology and Infection* 136(5):577-603.
- Russell JB, Jarvis GN (2011). Practical mechanisms for interrupting the oral-fecal life cycle of *Escherichia coli*. *Journal of Molecular Microbiology and Biotechnology* 3(2):265-272.
- Sofowora LA (1993). Medicinal plants and traditional Medicinal Medicine in African Spectrum books Ltd, Ibadan. pp. 55-71.
- Suree N, Pana L (2015). Antibacterial activity of crude ethanolic extracts and essential oil of spices against *Salmonella* and other *Enterobacteriaceae*. *Science Technology Journal* 5(3):527-538.
- Su L, Chiu CH (2007). *Salmonella*: Clinical Importance and Evolution of Nomenclature. *Chang Gung Medical Journal* 30(3):210-219.
- World Health Organization (WHO) (2002). *Traditional Medicine: Growing Needs and Potential. WHO Policy Perspectives on Medicines*. World Health Organization, Geneva pp. 1-6.

*Full Length Research Paper*

## ***In vitro* study on the anti-microbial efficacy of *Aloe vera* against *Candida albicans***

**T. Jeevitha<sup>1\*</sup>, K. Deepa<sup>2</sup> and A. Michael<sup>2</sup>**

<sup>1</sup>Department of Microbiology, Karpagam University, Coimbatore, India.

<sup>2</sup>Department of Microbiology, PSG College of Arts and Science, Coimbatore, India.

Received 18 June, 2015; Accepted 25 September, 2015

Oral *Candida* infections are common among the people who are immunocompromised, with poor oral hygiene, diabetics and advanced age. Even though several effective antifungal agents are available for treating oral *Candida* infections, on prolonged and frequent treatment with these drugs, they may result in development of drug resistant organisms. This leads to looking for an effective, inexpensive and simple alternative. Therapeutic products of plants play an important role in human health since ancient times. The present study was conducted to determine the bioactive components from the ethanol extract of *Aloe vera* by gas chromatograph-mass spectrometer (GC-MS) and to evaluate the antimicrobial effects of *A. vera* extracts on *Candida albicans*. GC-MS analysis revealed that ethanol extract of *A. vera* contains 26 bio active compounds. The results of *in vitro* antimicrobial activity of ethanol, methanol and aqueous extracts of *A. vera* against *Candida albicans* showed high inhibitory growth in yeast after treatment with ethanol extract followed by methanol extract with an inhibition zone of 23 and 12 mm, respectively. Aqueous extract did not show any zone of inhibition. The minimum inhibitory concentration (MIC) and the minimum fungicidal concentration (MFC) of the extracts were also determined. From this study, it is concluded that, *A. vera* has potent antifungal activity against *C. albicans*.

**Key words:** *Aloe vera*, gas chromatograph-mass spectrometer (GC-MS), antifungal, *Candida albicans*, minimum inhibitory concentration (MIC), the minimum fungicidal concentration (MFC).

### **INTRODUCTION**

The dimorphic fungus, *Candida albicans* is the normal inhabitant of oral microbiota in 30 to 50% of healthy people. The transition of *C. albicans* from normal flora to pathogenic organism is decided by various predisposing factors like suppression of immunity, poor oral hygiene, high carbohydrate diet, old age, infancy, dentures, nutritional deficiency, etc (Akpan and Morgan, 2002). Oral candidiasis can be treated with topical anti-

fungual drugs, such as fluconazole and amphotericin B. Topical therapy is given as an oral suspension which is washed around the mouth and then swallowed by the patient. Recently, *C. albicans* develops resistance against these antifungal agents on long-term treatment and leads to the development of drug-resistant *C. albicans* (Azmi and Tamer, 2014). So it is high time to look for an effective adjunct antifungal therapeutic

\*Corresponding author. E-mail: jeevithathangaraj@gmail.com.

strategy. Medicinal plants are the integral part of human life and rich source of novel drugs (Botelho et al., 2007). *Aloe vera* Linne or *Aloe barbadensis* Miller belongs to the family Liliaceae under which it comprises about 360 species. Historical evidence indicates that *A. vera* originated in the warm, dry climate of Southern and Eastern Africa, and was subsequently introduced into Northern Africa, the Arabian Peninsula, China, Gibraltar, the Mediterranean countries and the West Indies. It is perennial succulent xerophytes with thick leaves that supplies water for the plant to survive in dry areas for longer periods (Botelho et al., 2007). The Aloe plant has two parts, each of which produces substances that are completely varying in compositions and therapeutic properties. The inner portion is made up of parenchymal and forms the *A. vera* gel (or mucilage), a clear, thin, tasteless, jelly-like material. The other portion is made up of specialized cells known as the pericyclic tubules, which occur just beneath the outer green ring of the leaf. An exudate produced from that cells, consists of bitter yellow latex with powerful laxative-like actions (Hegggers and Kucukcelebi, 1996).

It has numerous therapeutic potentials and it is claimed to cure ailments like wounds, burns, immune modulation, inflammation and has antiseptic, antimicrobial, antifungal and antiviral properties (Richa et al., 2014). More than 75 active ingredients from inner gel have been identified including vitamins, minerals, enzymes, sugars, anthraquinones “phenolic compounds, lignin, saponins, sterols, amino acids and salicylic acid” (Nandal and Bhardwaj, 2012). Active ingredients include vitamin B<sub>12</sub>, magnesium, manganese, copper, zinc, calcium, potassium, folic acid, sodium, chromium, selenium and amino acids. It also contains lipases and proteases, which are enzymes that help with digestion; long-chain polysaccharides that boost the immune system and detoxify the body; fatty carboxylic acids that work as an anti-inflammatory; saponins that work as anti-viral, anti-fungal and anti-microbial agents; emodin and aloin that work as pain relievers as well as many other powerful medicinal agents (Nandal and Bhardwaj, 2012). In a study conducted by Bernardes et al. (2012), it was demonstrated that *A. vera* fresh leaves plant extract can inhibit both the growth and the germ tube formation by *C. albicans*.

Hence the present investigation was conducted to determine the bioactive components from the ethanol extract of *A. vera* by GC-MS and to determine the antifungal activity of *A. vera* extract against *C. albicans* and also the minimum inhibitory concentration (MIC) and the minimum fungicidal concentration (MFC) of whole leaf *A. vera* extract.

## MATERIALS AND METHODS

### Test microorganisms

The test organism, *C. albicans* (MTCC 227) was obtained from

Microbial Type of Culture Collection & Gene Bank (MTCC), Institute of Microbial Technology, Chandigarh. The test strain was maintained on Sabouraud's Dextrose agar plates (Hi-Media Laboratories Pvt. Limited, Mumbai, India) at 4°C and sub cultured on to yeast extract, peptone, dextrose (YPD) broth for 24 h prior to testing. This fungus served as test pathogen for antifungal activity assay.

### Preparation of *A. vera* extract

Fresh *A. vera* leaves were collected from Coimbatore, Tamilnadu, India. The whole leaf was chopped into pieces and air dried. All the dried parts of the leaves were ground into powdered form using mortar and pestle and solubilized with ethanol, methanol and water solvents at 0.2 g/ml concentration. These mixtures were maintained overnight in a shaker at 120 rpm for proper extraction of the active ingredients at (25°C) room temperature (Harbourne, 1991). The mixtures were then filtered using Whatman No. 1 filter paper (Pore size 0.47 µm). The solvents were evaporated using water bath at 40°C to ensure proper concentration and to avoid the influence of those solvents in the anti-fungal effect of *A. vera*.

### Phytochemical screening of the extracts

The phytochemical screening of the extract was carried out in order to ascertain the presence of its secondary metabolites such as saponins, alkaloids, flavonoids, steroids, terpenoids, anthraquinones, phlobatanins, tannins, cardiac glycosides, glycosides and phenols by gas chromatograph-mass spectrometer (GC-MS) analysis.

### GC-MS analysis

GC-MS analysis of this extract was performed using GC SHIMADZU QP 2010 system and gas chromatograph interfaced to a mass spectrometer (GC-MS) equipped with Elite-1 fused silica capillary column (Length : 30.0 m, Diameter : 0.25 mm, Film thickness : 0.25 µm composed of 100% dimethyl poly siloxane). For GCMS detection, an electron ionization energy system with ionization energy of 70eV was used. Helium gas (99.999%) was used as the carrier gas at a constant flow rate of 1.5 ml/min and an injection volume of 1 ml was employed (split ratio: 10). Injector temperature was 240°C and ion source temperature 200°C. The oven temperature was programmed from 70°C (isothermal for 2 min.), with an increase of 300°C for 10 min. Mass spectra were taken at 70 eV; a scan interval of 5 min with scan range of 40 – 1000 m/z. Total GC running time was 35 min. The relative percentage amount of each component was calculated by comparing its average peak area with the total areas.

### The antimicrobial screening of the extracts

The antimicrobial activity of extracts was determined *in vitro* in response to the *C. albicans*. The activities were measured using agar well diffusion method and broth dilution method, as previously described by the Clinical and Laboratory Standards Institute (CLSI; formerly known as the National Committee for Clinical Laboratory Standards) (NCCLS, 1997; Fothergill, 2011).

### Agar diffusion method

The organism was sub-cultured into prepared normal saline and incubated at 37°C for 30 min, the concentration of each organism

was increased to form a turbidity that matched with 0.5 McFarland's standard by visual comparison at which it was assumed that the number of cells was  $1.5 \times 10^8$  CFU/ml. Inoculum containing  $1.5 \times 10^8$  CFU/ml of fungal suspension to be tested was spread on Sabouraud's dextrose agar plates with a sterile swab moistened with the fungal suspension. Subsequently, wells were then made using sterile cork borer of 6 mm in diameter on the surface of pre seeded agar plates and filled with 100  $\mu$ l (0.2 g/ml) of each plant extracts and allowed to diffuse at room temperature for 2 h for proper diffusion. Wells containing the same volume of corresponding solvent served as negative control while standard antibiotic discs of fluconazole (50 mg/ml) used as the positive control. All the plates were incubated at 37°C for 24 h. The presence of zone of inhibition was regarded as the presence of antimicrobial action. From the inhibition zones seen, antimicrobial activity was expressed in terms of average diameter of the zones of inhibition measured.

#### Minimum inhibitory concentration (MIC)

The minimum inhibitory concentrations of the extracts were determined using broth micro dilution method approved by the National Committee for Clinical Laboratory Standards (M27- A) (NCCLS, 1997). The yeast were cultured overnight on Sabouraud's dextrose agar (SDA) and then resuspended in normal saline to obtain a final concentration of  $1.5 \times 10^8$  cfu/ml. 200 mg/ml of each of the extracts were reconstituted into Sabouraud's dextrose broth and serially diluted using Sabouraud's dextrose broth forming concentrations of 100, 50, 25, 12.5, 6.25 and 3.125 mg/ml. 0.1 ml of the cell suspension was inoculated into each of the tubes with varied concentrations. All the tubes were incubated at 37°C for 24 h. After incubation, the MIC was determined as the lowest concentration of extract for which there was no visible growth compared with the control (CLSI, 2008).

#### Minimum fungicidal concentration (MFC)

The MFC was determined by inoculating 0.1 ml of negative growth in MIC onto sterile SDA plates. The plates were incubated at 35°C for 48 h. The lowest concentration of plant extract that did not demonstrate growth of the tested organism was considered as the MFC; the negative control was a plate grown with media only (Ernst et al., 2002; Wiegand et al., 2008).

## RESULTS

The components present in the extract was identified based on the spectrum obtained using the data base of National Institute of Standard and Technology (NIST). The name of the compound, retention time, molecular formula and structure were determined. The area percentage of each component was calculated by comparing its average peak area with the total areas.

GC-MS analysis of *A. vera* revealed the presence of 35 compounds, of which 26 were found to have biological activities. These compounds were confirmed based on their peak area (%), retention time, molecular formula and molecular weight (MW) and are represented in Table 1 and Figure 1. Their reported biological activities are given in Table 2. The prevailing compounds of ethanol extract of *A. vera* were hexadecanoic acid, ethyl ester

(16.20%), octadecanoic acid, ethyl ester (13.62%), 4-3,5-di-tert-butyl-4-hydroxyphenyl)butyl acrylate (7.65%), 9-octadecenoic acid (z)-, ethyl ester (7.10%), 1-nonadecene (6.19%), ethyl (9z,12z)-9,12-octadecadienoate (4.49%), 1,4-dioxaspiro[4.14]nonadecane (3.56%), 1-pentadecene (3.25%), 1-hexadecene (3.19%), n-pentadecanol (3.03%), di-n-octyl phthalate (2.23), cyclopropane, 1-methyl-1-(1-methyl ethyl)-2-nonyl- (2.16%), 10-undecenoic acid, ethyl ester (2.13%), 1-heptacosanol (2.06%), cyclopropane, 1-methyl-1-(1- (1.96%), 13-hexyloxa-cyclotridec-10-en-2-one (1.94%), ricinoleic acid (1.72%), 1-undecanol (1.49%), n-nonadecanol-1 (1.48%), 3-eicosene, (e)- (1.35%), strychnidin-10-one, 2,3-dimethoxy- (1.26%), 1-undecanol (1.25%), tetraethyl silicate (1.00%), cyclopropane carboxylic acid (0.99%), 2-undecene, 3-methyl-, (z)- (0.97%), cyclohexanol, 3-methyl- (0.83%), propane, 1,1-diethoxy- (0.49%), 1-butanamine, 2-methyl-n-(2-methylb (0.78%), ethyl orthoformate (0.77%), furo[3,2-i][1]benzopyran decahydro-, [3as-(3a.alpha.,6a.beta.,10as\*)]- (0.41%), ethanone, 1-(6,6-dimethylbicyclo[3.1.0]hex-2-en-2-yl)- (0.08%) and eicosyl trifluoroacetate. *A. vera* is reported to contain carbohydrates, tannins, saponins, flavonoids and alkaloids (Yebpella et al., 2011). Phyto constituents have been found to inhibit bacteria, fungi, viruses and pests. The presence of phyto constituents in the *A. vera* extracts may be responsible for the antifungal activity of the plant (Cowan, 1999).

The antimicrobial activity of the ethanol, methanol and aqueous extracts against *C. albicans* had shown varying degrees of zones of inhibition and the results are depicted in Figure 2 and Table 3. The ethanol extracts exerted highest activity on *C. albicans* as compared to the methanol extract. The ethanol extract at the concentration of 0.2 g/ml showed 23 mm diameter zone of inhibition against *C. albicans*, methanol extract showed 12 mm diameter whereas aqueous extract showed no activity. The minimal inhibitory concentration of the ethanol, methanol and aqueous extracts against *C. albicans* was determined by broth dilution method and the results are depicted in Table 4. The MIC value of ethanol extract was found to be 50 mg/ml and methanol extract was 100 mg/ml. The MIC and the MFC were in the range of 25 to 100 mg/ml and 50 to 100 mg/ml, respectively. The growth of *C. albicans* in ethanol extract was inhibited at 25 mg/ml while the minimum concentration that showed no growth of *C. albicans* by methanol extract is 50 mg/ml.

## DISCUSSION

The present study on *A. vera* revealed the presence of medicinal active constituents. In the GC-MS analysis, 35 bioactive compounds were identified in the ethanol extract of *A. vera* based on the peak area, molecular

**Table 1.** Bioactive components in the ethanolic extract of *Aloe vera*.

S/N	RT	Name of the compound	Molecular formula	Molecular weight	Peak area (%)
1	6.067	Eicosyl trifluoroacetate	C22H41F3O2	394	-0.22
2	6.276	Furo[3,2-i][1]benzopyran, dec	C11H18O2	182	0.41
3	6.557	Cyclopropanecarboxylic a	C5H9NO2	115	0.99
4	8.035	Tetraethyl silicate	C8H20O4Si	208	1.00
5	9.578	Propane, 1,1-diethoxy-	C7H16O2	132	0.49
6	10.274	Cyclopentene, 1-(3-methylbutyl)-	C10H18	138	0.40
7	10.645	1-Butanamine, 2-methyl-N-(2-methylb	C10H21N	155	0.78
8	11.386	1-undecanol	C11H24O	172	1.49
9	13.481	Cyclohexanol, 3-methyl-	C7H14O	114	0.83
10	14.442	1-pentadecene	C15H30	210	3.25
11	19.776	1-hexadecene	C16H32	224	3.19
12	25.883	Cyclopropane, 1-methyl-1-(1-	C16H32	224	1.96
13	26.004	1-nonadecene	C19H38	266	6.19
14	26.304	2-undecene, 3-methyl-, (z)-	C12H24	168	0.97
15	27.727	Ethyl orthoformate	C7H16O3	148	0.77
16	29.248	Ethanone, 1-(6,6-dimethylbicyclo[3.1.0	C10H14O	150	0.08
17	29.467	Cyclopropane, 1-methyl-1-(1-methyle	C16H32	224	2.16
18	29.560	Hexadecanoic acid, ethyl ester	C18H36O2	284	16.20
19	29.782	4-(3,5-di-tert-butyl-4-hydrox	C21H32O3	332	7.65
20	30.651	13-hexyl-oxa-cyclotridec-10	C18H32O2	280	1.94
21	30.840	N-Nonadecanol-1	C19H40O	284	1.48
22	31.717	3-eicosene, (e)-	C20H40	280	1.35
23	31.850	Ethyl (9z,12z)-9,12-octadecadi	C20H36O2	308	0.94
24	31.933	9-octadecenoic acid (z)-, eth	C20H38O2	310	7.10
25	32.008	10-undecenoic acid, ethyl es	C13H24O2	212	2.13
26	32.238	Octadecanoic acid, ethyl es	C20H40O2	312	13.62
27	32.457	Ethyl (9z,12z)-9,12-octadecad	C19H34O2	294	3.25
28	32.988	Ethyl (9z,12z)-9,12-octadecadi	C19H34O2	294	4.49
29	34.051	1,4-Dioxaspiro[4.14]nonadecane	C17H32O2	268	3.56
30	34.203	Ricinoleic acid	C18H34O3	298	1.72
31	34.602	1-heptacosanol	C27H56O	396	2.06
32	37.000	Di-n-octyl phthalate	C24H38O4	390	2.23
33	37.383	Strychnidin-10-one, 2,3-dimet	C23H26N2O4	394	1.26
34	37.731	N-Pentadecanol	C15H32O	228	3.03
35	38.119	1-undecanol	C11H24O	114	1.25

weight and molecular formula. Of which 26 compounds were found to be biologically active. The prevailing compounds were hexadecanoic acid, ethyl ester (16.20%), octadecanoic acid, ethyl ester (13.62%), 13-hexyl-oxa-cyclotridec-10-en-2-one (1.94%), 9-octadecenoic acid (z)-, ethyl ester (7.10%), 1-nonadecene (6.19%) etc. These compounds were found to have anti-inflammatory (Langmead et al., 2004), anti-bacterial (Agarry et al., 2005), anti-fungal, anti-oxidant, immunomodulatory (Sun-A et al., 2010), analgesic (David, 1999) activity etc. Phytochemical compounds identified in plant material have been reported as having inhibitory action against *C. albicans* (Freeman and

Beattle, 2008). Below the minimum inhibitory concentrations, growth of the microbes was observed ranging from light growth to high growth (Table 4).

The growth of *C. albicans* was inhibited by the ethanol and methanol extracts of *A. vera* with varying zones of inhibition while aqueous extract did not show any inhibitory activity. Ethanol extracts had a better minimum inhibitory concentration as compared to methanol and aqueous extract used in this investigation as reported by Nejad et al. (2014). The negative control using the corresponding solvent had no effect on the test organism. The positive control (Fluconazole) had wide effect against the organism. Fluconazole is preferred as positive control







Figure 1. GC-MS analysis of *Aloe vera*.

Table 2. Biological activity of Identified phytochemical constituents in ethanolic extract of *Aloe vera*.

S/N	RT	Name of the compound	Peak area	Activity
1	6.067	Eicosyltrifluoroacetate	-0.22	Anti-microbial and anti-fungal activity <sup>29</sup>
2	6.557	Cyclopropanecarboxylic acid	0.99	Anti-fungal activity, Pesticide <sup>22</sup>
3	8.035	Tetraethyl silicate	1.00	Anti-aging activity, anti oxidant, immunostimulant <sup>19</sup>
4	11.386	1-undecanol	1.49	Anti-microbial, flavouring agent <sup>22</sup>
5	14.442	1-pentadecene	3.25	Insecticidal, pesticidal activity <sup>34</sup>
6	19.776	1-hexadecene	3.19	Anti-microbial, anti-oxidant <sup>23</sup>
7	25.883	Cyclopropane, 1-methyl-1-(1-	1.96	Anti-fungal, Insecticidal, herbicidal <sup>22</sup>
8	26.004	1-nonadecene	6.19	Anti-fungal, anti-tuberculosis activity <sup>22</sup>
9	29.467	Cyclopropane, 1-methyl-1-(1-methyle	2.16	Anti-fungal, Insecticidal, herbicidal <sup>22</sup>
10	29.560	Hexadecanoic acid, ethyl ester	16.20	Anti-microbial, anti-oxidant, Nematicide, Pesticide, anti-androgenic, hemolytic, flavor, hypocholesterolemic activity <sup>34</sup>
11	29.782	4-(3,5-di-tert-butyl-4-hydrox	7.65	Anti-inflammatory activity <sup>22</sup>
12	30.651	13-hexyl-oxa-cyclotridec-10	1.94	Effective against ESBL's and multi drug resistant organisms <sup>21</sup>
13	30.840	N-Nonadecanol-1	1.48	Cytotoxic properties <sup>9</sup>
14	31.717	3-eicosene, (e)-	1.35	Anti-microbial activity <sup>22</sup>
15	31.850	Ethyl (9z,12z)-9,12-octadecadi	0.94	Fragrance ingredient <sup>29</sup>
16	31.933	9-octadecenoic acid (z)-, eth	7.10	Plasticizer, food additive, vehicle for drug delivery <sup>29</sup>
17	32.008	10-undecenoic acid, ethyl es	2.13	Cosmetic ingredient <sup>22</sup>

Table 2. Contd.

18	32.238	Octadecanoic acid, ethyl es	13.62	Perturbs cell cycle and induce apoptosis in Hep G2 cells <sup>29</sup>
19	32.457	Ethyl (9z,12z)-9,12-octadecad	3.25	Fragrance ingredient <sup>29</sup>
20	32.988	Ethyl (9z,12z)-9,12-octadecadi	4.49	Fragrance ingredient <sup>29</sup>
21	34.203	Ricinoleic acid	1.72	Analgesic, anti-inflammatory activity, algicide, blood coagulant <sup>22</sup>
22	34.602	1-heptacosanol	2.06	Anti-microbial, anti-oxidant, anti, cancer, nematicidal, Anti-atherosclerosis <sup>27</sup>
23	37.000	Di-n-octyl phthalate	2.23	Plasticizer <sup>22</sup>
24	37.383	Strychnidin-10-one, 2,3-dimet	1.26	Pesticide <sup>22</sup>
25	37.731	N-Pentadecanol	3.03	Anti-microbial activity <sup>22</sup>
26	38.119	1-undecanol	1.25	Anti-fungal activity <sup>22</sup>

Figure 2. Antimicrobial activity of various *Aloe vera* extracts.

because it is a potent and selective inhibitor of fungal enzymes and also it does not have the hepatotoxicity as the imidazoles (Akpan and Morgan, 2002). A previous study conducted by Doddanna et al. (2013) demonstrated the antimicrobial effects of alcoholic extracts of tea leaves, onion leaves, onion bulb, *A. vera*, mint leaves and curry leaves against *C. albicans* and our results on ethanol extract was consistent with their results.

Oropharyngeal candidiasis (OPC) is an opportunistic infection of oral cavity. Oral candidiasis can appear as erythematous patches or white, scrapable lesions. Oral cavity is colonized by *C. albicans* or other *Candida* species in 40-60% of healthy persons. Many factors contribute to the development of oropharyngeal candidiasis (OPC) including malnutrition, poor oral hygiene, dental malocclusion, and immunosuppression.

In immune compromised patients, *Candida* species can trigger a variety of disease manifestations ranging from localized mild oral lesion to a disseminated candidiasis (Berberi et al., 2015). Oral candidiasis caused by *C. albicans* is generally managed by treatment with fluconazole and amphotericin B. Prolonged or frequent treatment with these azole drugs results in the rise of drug resistant organism. Use of medicinal plants is considered as an adjunct antifungal therapeutic strategy nowadays. According to the World Health Organization (WHO), as many as 80% of the world's people depend on traditional medicine for their primary healthcare needs. The development of indigenous medicines and the use of medicinal plants carry considerable economic benefits in the treatment of various diseases (Gunjan et al., 2013). Numerous studies have been carried out all over the world with *A. vera* and it has been reported that *A. vera* exhibits different levels of anti microbial property. The purified Aloe protein has been found to exhibit potent antifungal activity against *Candida parapsilosis*, *Candida krusei* and *C. albicans* (Das et al., 2011). Garnick et al (2008) evaluated a gel that combined allantoin, *A. vera* and silicon dioxide and its effects on aphthous ulcers of the oral cavity. *A. vera* tooth gel was as effective, and in some cases more effective than the commercial brands at controlling cavity-causing organisms (Namiranian and Serino, 2012). Sema and Suleyman (2009) reported that, a processed *A. vera* juice preparation showed inhibition of the growth of *C. albicans*.

## Conclusion

This study has revealed the presence of many secondary metabolites in the leaves of *A. vera* due to which it showed antifungal activity against *C. albicans*. Thus, based on the results obtained in the present study, ethanol extracts of *A. vera* may be useful as a better alternative for the treatment of antimycotic resistant *C. albicans* with respect to oral candidiasis. It is suggested that more *in vivo* and clinical research works should be done in the future.

**Table 3.** Antimicrobial activity of *Aloe vera* extracts against *Candida albicans*.

Organism	Solvent			Control	
<i>Candida albicans</i>	Ethanol	Methanol	Aqueous	Positive (fluconazole 50 mg/ml)	Negative (solvent)
Zone of inhibition in mm	23	12	-	36	-

**Table 4.** Minimum inhibitory concentration of *Aloe vera* extracts on *Candida albicans*.

Solvent	Concentration of extracts in mg/ml						MIC
	100	50	25	12.5	6.25	3.125	
Ethanol	-	-	0*	+	++	+++	25
Methanol	-	0*	+	++	++	+++	50
Aqueous	+	+	+	+	+	+	-

- = No growth (turbidity), 0\* = MIC, + = light growth, ++ = moderate growth, +++ = high growth.

### Conflict of interests

The authors have not declare any conflict of interest.

### ACKNOWLEDGEMENTS

The authors thank the Principal and the Management of Karpagam University, Coimbatore, India for providing facilities to carry out the work. They also thank the Principal and the Management of PSG College of Arts and Science, Coimbatore, India for their consistent support and help during this study.

### REFERENCES

- Agarry OO, Olaleye MT, Bello-Michael CO (2005). Comparative antimicrobial activities of *Aloe vera* gel and leaf. *African Journal of Biotechnology* 4:1413-1414.
- Akpan A, Morgan R (2002). Oral candidiasis. *Postgraduate Medical Journal* 78:455-459.
- Azmi MGD, Tamer AD (2014). What Makes Oral Candidiasis Recurrent Infection? A Clinical View. *Journal of Mycology Article ID 758394*.
- Berberi A, Noujeim Z, Aoun G (2015). Epidemiology of Oropharyngeal Candidiasis in Human Immunodeficiency Virus/Acquired Immune Deficiency Syndrome Patients and CD4+ Counts. *Journal of International Oral Health* 7(2):1-4
- Bernardes I, Felipe Rodrigues MP, Bacelli GK, Munin E, Alves LP, Costa MS (2012). *Aloe vera* extract reduces both growth and germ tube formation by *Candida albicans*. *Mycoses* 55:257-61
- Botelho A, NAP Nogueira, GM Bastos (2007). Antimicrobial activity of the essential oil from *Lippia sidoides*, carvacrol and thymol against oral pathogens. *Brazilian Journal of Medical and Biological Research* 40(3):349-356.
- CLSI (2008). Reference Method for Broth Dilution Antifungal Susceptibility Testing of Yeasts. 3<sup>rd</sup> Edn., Clinical and Laboratory Standards Institute, Wayne.
- Cowan MM (1999). Plant products as antimicrobial agents. *Clinical Microbiology Reviews* 12(4):564-582.
- Dalli AK, Saha. G, Chakraborty. U (2007). Characterization of Antimicrobial compounds from a common fern, *Pterin biaurita*. *Indian*

- Journal of Experimental Biology* (5):285-290.
- Das S, Mishra B, Gill K, Ashraf MS, Singh AK, Sinha M, Sharma S, Xess I, Dalal K, Singgh TP, Dey S (2011) Isolation and characterization of novel protein with anti-fungal and anti-inflammatory properties from *Aloe vera* leaf gel. *International Journal of Biological Macromolecules* 48:38-43.
- David U (1999). *Aloe vera-Nature's Gift: Aloe vera* in veterinary practice. Great Britain, Blackdown Publications.15.
- Doddanna SJ, Shilpa Patel, Madhusudan AS, Ravindra SV (2013). Antimicrobial activity of plant extracts on *Candida albicans*: An in vitro study. *Indian Journal of Dental Research* 24(4).
- Ernst EJ, Roling EE, Petzold CR, Keele DJ, Klepser ME (2002). *In vitro* activity of micafungin (FK-463) against *Candida* spp.: Microdilution, timekill and post antifungal-effect studies. *Antimicrobial Agents and Chemotherapy* 46:3846-3853.
- Fadipe LA, Haruna K, Mohammed I (2014). Antibacterial Activity of 1,2-Benzenedicarboxylic Acid, Dioctyl Ester Isolated from the Ethyl Acetate Soluble Sub-portion of the unripe Fruits of *Nauclea latifolia*. *International Journal of Pure & Applied Bioscience* 2(1):223-230.
- Fothergill AW (2011). Antifungal Susceptibility Testing: Clinical Laboratory and Standards Institute (CLSI) Methods. In: *Intractions of Yeast, Moulds and Antifungal Agents*, Hall, G.S. (Ed.), Humana Press, New York, ISBN-10: 1597451347, pp.65-74.
- Freeman BC, Beattle GA (2008). An Overview of Plant Defenses against Pathogens and Herbivores. *The Plant Health Instructor*. DOI: 10.1094/PHI-I-2008-0226-01.
- Garnick JJ, Singh B, Winkley G (2008). Effectiveness of a medicament containing silicon dioxide, aloe, and allantoin on aphthous ulcers. *Oral Surgery, Oral Medicine, Oral Pathology, and Oral Radiology* 86:550-556.
- Gunjan K, Md. Jalaluddin, Purnendu R, Rajat M, Dileep CL (2013). Emerging Trends of Herbal Care in Dentistry. *Journal of Clinical and Diagnostic Research* 7(8):1827-1829
- Harbourne JB (1991). *Phytochemical Methods. A Guide to Modern Techniques of Plant Analysis*. London: Chapman and Hall.
- Jurkić LM, Cepanec I, Pavelić SK, Pavelić K (2013). Biological and therapeutic effects of ortho-silicic acid and some ortho-silicic acid-releasing compounds: New perspectives for therapy. *Nutr. Metabolism* 10:2.
- Langmead L, Makins RJ, Rampton DS (2004). Anti-inflammatory effect of *Aloe vera* gel in human colorectal mucosa *in vitro*. *Aliment. Pharmacology & Therapeutics* 19:521-527
- Mohammed MP, Aysa OS, Dhamotharan R (2014). Insilico identification of potent inhibitor from andrographis paniculata for esbls. *International Journal of advances in Pharmacy, Biology and*

- Chemistry 3(3).
- Murugesan S, Vijayakumar R, Panneerselvam A (2011). Evaluation of Phytochemical Constituents from the Leaves of *Memecylon umbellatum* Burm.f. Research Journal of Pharmaceutical, Biological and Chemical Sciences 2(4).
- Muthumary J, Yogeswari S, Ramalakshmi S, Neelavathy R. 2012. Identification and Comparative Studies of Different Volatile Fractions from *Monochaetia kansensis* by GCMS. Global Journal of Pharmacology 6(2):65-71.
- Namiranian H, Serino G (2012). The effect of a toothpaste containing *Aloe vera* on established gingivitis. Swedish Dental Journal 36(4):179-185.
- Nandal U, Bhardwaj RL (2012). *Aloe vera* for human nutrition, health and cosmetic use -A review. International Journal of Plant Sciences 3(3):38-46.
- NCCLS (1997). Reference method for broth dilution antifungal susceptibility testing of yeasts. Approved standard M27-A. National Committee for Clinical Laboratory Standards, Wayne, Pa.
- Nejad BS, Mahsa R, Mamoudabadi AZ, Majid Z (2014). *In Vitro* Anti-*Candida* Activity of the Hydroalcoholic Extracts of *Heracleum persicum* Fruit against Pathogenic *Candida* Species. Jundishapur Journal of Microbiology 7(1):e8703.
- Nejatzadeh-Barandozi F (2013). Antibacterial activities and antioxidant capacity of *Aloe vera*. Organic and Medicinal Chemistry Letters 3:5.
- Richa W, Siraj DAAK, Gaurav S, Sabreena S (2014). *Aloe vera*: A Boon in Dentistry. International Journal of Pharmaceutical Sciences Review and Research 4(3):147-151.
- Sathyaprabha G, Kumaravel S, Ruffina D, Praveenkumar P (2010). A Comparative study on Antioxidant, Proximate analysis, Antimicrobial activity and phytochemical analysis of *Aloe vera* and *Cissus quadrangularis* by GC-MS. Journal of Pharmacy Research 3(12):2970-2973.
- Sema A, Suleyman AJ(2009). Investigation of in vitro antimicrobial activity of *Aloe vera* juice. Journal of Animal and Veterinary Advances. 8(1):99-102.
- Sun-A I, Young-Ran L, Young-Hee L, Myung-Koo L, Young IP, Sungwon L, Kyungjae K, Chong-Kil L (2010). *In Vivo* Evidence of the Immunomodulatory Activity of Orally Administered *Aloe vera* Gel. Archives of Pharmacal Research 33(3):451-456.
- Wei-seng ho, Zayed MZ, Ahmad FB, Pang S (2014). Gc-ms analysis of phytochemical constituents in leaf extracts of *neolamarckia cadamba* (rubiaceae) from malaysia. International Journal of Pharmacy and Pharmaceutical Sciences 6(9):123-127
- Wiegand I, Hilpert K, Hancock REW (2008). Agar and broth dilution methods to determine the Minimal Inhibitory Concentration (MIC) of antimicrobial substances. Nature Protocols 3:163-175.
- Yebpella GG, Adeyimi HMM, Hammuel C, Magomya AM, Agbaji AS, Okonkwo EM (2011). Phytochemical Screening and Comparative Study of Antimicrobial activity of *Aloe vera* various extracts. African Journal of Microbiology Research 5(10):1182-1187.

**Related Journals:**

