

OPEN ACCESS



International Journal of
Psychology and Counselling

January 2019
ISSN: 2141-2499
DOI: 10.5897/IJPC
www.academicjournals.org

 **ACADEMIC
JOURNALS**
expand your knowledge

ABOUT IJPC

International Journal of Psychology and Counselling (IJPC) will be published monthly (one volume per year) by Academic Journals.

International Journal of Psychology and Counselling (IJPC) is an open access journal that provides rapid publication (monthly) of articles in all areas of the subject counter transference, Clinical psychology, Professional practice of behaviour analysis, Behavioural model etc.

The Journal welcomes the submission of manuscripts that meet the general criteria of significance and scientific excellence. Papers will be published shortly after acceptance. All articles published in IJPC are peer-reviewed.

Contact Us

Editorial Office: ijpc@academicjournals.org

Help Desk: helpdesk@academicjournals.org

Website: <http://www.academicjournals.org/journal/IJPC>

Submit manuscript online <http://ms.academicjournals.me/>

Editors

Prof. Ahmed Awad Amin Mahmoud

*Faculty of Education and Higher Education
An-Najah National University,
Nablus.
Palestine.*

Dr. R. Joseph Ponniah

*Department of Humanities (English)
National Institute of Technology
Trichirappalli, Tamil Nadu
India.*

Dr. Kanwar Dinesh Singh

*# 3, Cecil Quarters,
Chaura Maidan, Shimla:171004 HP
India.*

Dr. S. D. Sindkhedkar

*Head, Department of English,
PSGVP Mandal's Arts, Science & Commerce
College,
Shahada: 425409, (Dist. Nandurbar), (M.S.),
India.*

Dr. Marta Manrique Gómez

*Middlebury College
Department of Spanish and Portuguese
Warner Hall, H-15
Middlebury, VT 05753
USA.*

Dr. Yanjiang Teng

*801 Cherry Lane, APT201
East Lansing
Michigan State University
MI 48824
USA.*

Prof. Radhakrishnan Nair

*SCMS-COCHIN
Address Prathap Nagar, Muttom, Aluva-1
India.*

Prof. Lianrui Yang

*School of Foreign Languages, Ocean University of
China
Address 23 Hongkong East Road, Qingdao,
Shandong Province, 266071 P
China.*

Editorial Board

Dr. Angeliki Koukoutsaki-Monnier

*University of Haute Alsace
IUT de Mulhouse
dep. SRC
61 rue Albert Camus
F-68093 Mulhouse
France.*

Dr. Martha Christopoulou

*Greek Ministry of National
Education & Religious Affairs
Xanthoudidou 2-4
Athens,
Greece.*

Dr. Zeynep Orhan

*Fatih University Hadımköy 34500 Istanbul
Fatih University Computer Engineering Department
Turkey.*

Dr. Tahar Labassi

*University of Tunis
94 Avenue 9 Avril, Tunis 1007
Tunisia.*

Table of Content

**Assessment of diabetes related distress among subjects
with type 2 diabetes in South India**

Hemavathi P., Satyavani K., Smina T. P. and Vijay V.

Full Length Research Paper

Assessment of diabetes related distress among subjects with type 2 diabetes in South India

Hemavathi P., Satyavani K., Smina T. P. and Vijay V.*

M.V. Hospital for Diabetes and Prof. M. Viswanathan Diabetes Research Centre, WHO Collaborating Centre for Research, Education and Training in Diabetes and IDF Centre for Excellence in Diabetes Care, No 4, West Madha Church Street, Royapuram, Chennai – 600 013, India.

Received 26 October, 2018; Accepted 18 December, 2018

Diabetes is a complex chronic disease that affects not only an individual's physical health, but can also have a profound impact on mental wellbeing. The aim of this study was to assess the prevalence of DRD among subjects with type 2 diabetes (T2DM) using Diabetes Distress Scale-17 (DDS-17) and to see its correlation with glycemic control and treatment modalities. A cross-sectional study of 400 T2DM subjects (200 men and 200 women) aged between 25 to 65 years who visited the tertiary care centre for diabetes in South India between April 2017 and May 2018 were included in this study. Subjects with T1DM, gestational diabetes mellitus (GDM) and psychiatric illness were excluded. The total score of DDS-17 was calculated by taking a sum of the 17 items' results and then dividing the total by 17. Clinical validation of the DDS suggests that the following thresholds of severity should be applied when interpreting the scores: little or no distress < 2.0, moderate distress = 2.0 - 2.9, and high distress ≥ 3.0. The mean age of men and women was 52.0 ± 8.4 and 51.7 ± 8.1 years, respectively. The mean score in women was 2.79 ± 1.52 as compared to men (1.62 ± 0.83) (p<0.001). The study findings highlighted that women had high levels of distress in managing diabetes as compared to men. Diabetes distress should be considered as a significant health problem and steps should be taken for effective management by lifestyle modifications, coping with their stress and diabetes.

Key words: Diabetes distress, HbA1c, emotional burden, type 2 diabetes (T2DM), South India.

INTRODUCTION

Diabetes is one of the most common chronic diseases, estimated to affect more than 400 million people worldwide. Prevalence is predicted to reach 642 million by 2040, and is anticipated to be the 7th leading cause of death by 2030. Diabetes affects not only an individual's physical health, but can also have a profound impact

on mental wellbeing (Mathers et al., 2006; Ogurtsova et al., 2017). It is a complex, chronic disease which requires patient self-management, involving daily decisions concerning diet, physical activity, blood glucose monitoring, and consistent medication adherence (Fowler, 2008). Diabetes complications are a significant cause of

*Corresponding author. E-mail: drvijay@mvdidiabetes.com; Tel: +9144 2595 4913.

increased morbidity and mortality among people. In addition, the substantial burden of impaired functioning and the demands of self management can contribute to significant emotional distress (Deshpande et al., 2008).

Diabetes distress has been defined across domains relating to: diabetic regimen like monitoring of blood glucose, dietary control, increasing physical activity, interpersonal or relational issues, referring to conflicts with careers, partners and friends, emotional burden such as feeling demotivated, unsupported emotionally, misunderstood and worrying about future complications (Polonsky et al., 2005). When an individual is diagnosed with diabetes, the coping resources they possess are insufficient to manage the illness threat, thus triggering emotional distress; that is, they experience high levels of emotional distress stemming from concerns and worries associated with their diabetes and its management (Fisher et al., 2012).

Diabetes distress refers to the unique, often hidden emotional burden and worries that are part of the spectrum of patient experience when managing a severe demanding chronic disease like diabetes (Gonzalez et al., 2011; Schetman et al., 2002). It may be sometimes misdiagnosed as depression and has links to problematic glycemic control and difficulties with self-care behaviors. It causes frustration with diabetes management and patients experience fear about potential complications with erratic blood glucose levels (Fisher et al., 2009, 2010).

Underlying causes of diabetes distress include feeling powerless about controlling diabetes, excessive worries about long-term complications, frustration with management tasks, fear that food constraints are controlling their life, poor confidence with regards to quickly identifying hypoglycemia, fear of embarrassment and potential risk of life, negative social perceptions, fear about being treated differently, less attractiveness to employers, family and friends over or under-involvement, treated as overly fragile and insufficient help or support (Gonzalez et al., 2011; Schetman et al., 2002). There is sparse data available on this topic, hence this study planned to assess the prevalence of DRD among subjects with type 2 diabetes mellitus (T2DM) using Diabetes Distress Scale-17 items (DDS-17) and to see its correlation with glycemic control and treatment modalities.

METHODOLOGY

In this cross-sectional study, 400 T2DM subjects (200 men and 200 women) aged between 25 to 65 years who visited the tertiary care centre for diabetes in South India between April 2017 and May 2018 were included. Patients with type 1 diabetes (T1D), gestational diabetes, cancer, psychiatric illness, and patients unwilling to participate were excluded from the study. DDS2 was first used for preliminary screening purpose to help the subjects familiarize with the content of DDS17. If a patient answered affirmatively to the DDS2 questions, then DDS 17 can be administered to define the extent of the distress.

Diabetes distress scale (DDS)

DDS is a validated 17- item self-report measure with each item scored on a Likert scale from 1 (no distress) to 6 (serious distress) concerning distress experienced over the last month (Fisher et al., 2012; Schetman et al., 2002). The scale yields four reliable subscales via item mean scores: emotional burden (e.g., "feeling angry, scared, and/or depressed when I think about living with diabetes," "feeling overwhelmed by the demands of living with diabetes"), physician-related distress (e.g., "feeling that my doctor does not take my concerns seriously enough," "feeling that my doctor does not give me clear directions on how to manage my diabetes"), regimen-related distress (e.g., "feeling that I am often failing with my diabetes routine," "not feeling confident in my day-to-day ability to manage diabetes"), and interpersonal distress (e.g., "feeling that friends or family do not appreciate how difficult living with diabetes can be," "feeling that friends or family are not supportive enough of self-care efforts"). The regimen distress scale assesses perceived problems with diabetes self-management. The total score is derived as the mean of all 17 items. Internal reliability of the total scale was excellent ($\alpha = 0.95$). All scales are treated as continuous variables. Clinical validation of the DDS suggests that the following thresholds of severity should be applied when interpreting scores: little or no distress < 2.0, moderate distress = 2.0-2.9 and high distress ≥ 3.0 (Fisher et al., 2012; Schetman et al., 2002).

Statistical analysis

Values reported are mean \pm standard deviation (SD) or number and percentages. Statistical comparison between different groups were made using independent samples t-test and Pearson correlation was done to find out the association between age, HbA1c and distress dimensions and total distress. SPSS version 20 was used for statistical analysis. A 'p' value of <0.05 was considered as statistically significant.

RESULTS

The mean age of men was 52.0 ± 8.4 and women 51.7 ± 8.1 years. Table 1 shows the distress levels among men and women. Analysis of DDS-17 results indicated that 77.5% of study subjects had moderate to high DRD based on the total score of the questionnaire. It was found that 22.5% of men and about 55% of women were screened positive for moderate to high levels of DRD on a DDS-17 scale with women having significantly higher DRD levels than men ($p < 0.001$).

Table 2 presents the DDS 17 individual domain scores by gender. In total, analysis of DDS-17 scale indicated that women had higher levels of distress (mean score of 2.79 ± 1.52) as compared to men (1.62 ± 0.83) ($p < 0.001$). HbA1c was significantly higher in women with high combined distress and high emotional distress compared to men. The mean scores for the total and individual components of DDS-17 were analyzed separately. It was found that women showed high distress in emotional burden domain as compared to men. In other domains like physician related, regimen related and interpersonal distress, women showed moderate distress, whereas men showed little or no distress.

Table 1. Over all distress levels among men and women.

Variable	Men [n (%)]	Women [n (%)]
Low distress (<2.0)	155 (77.5)	90 (45)
Moderate distress (2.0-2.9)	31 (15.5)	17 (8.5)
High distress (\geq 3.0)	14 (7)	93 (46.5)

Table 2. DDS-17 Domain score by gender.

Variable	Men (n=200)	Women (N=200)	Men Vs. Women (P-Value)
Emotional	1.78 \pm 0.97	3.16 \pm 1.55	<0.001*
Physician	1.40 \pm 0.70	2.21 \pm 1.46	<0.001*
Regimen	1.75 \pm 1.01	2.98 \pm 1.65	<0.001*
Interpersonal	1.46 \pm 0.90	2.66 \pm 1.6	<0.001*
Total	1.62 \pm 0.83	2.79 \pm 1.52	<0.001*

Values are mean \pm SD.


**Figure 1.** Individual DDS-17 domain scores by Gender. A. Gender wise details on emotional burden and physician related distress. B. Gender wise details on regimen and interpersonal related distress.

Figure 1A clearly depicts the gender-wise details on emotional burden and interpersonal related distress based on low, moderate and high distress levels among men and women. In emotional burden domain,

55.5% of women showed high distress, whereas it was only 13.5% in the case of men ($p < 0.0001$). In physician related distress, 6% of men expressed high distress levels whereas it was 37% in women ($p < 0.0001$). Figure 1B




Figure 2. Distress levels by age groups.

Table 3. Correlation of age and HBA1C with distress domain scores.

Variable		Emotional	Physician	Regimen	Interpersonal
AGE	Correlation	-0.15	-0.15	-0.16	-0.14
	P-Value	0.00	0.00	0.00	0.00
HBA1C	Correlation	0.95	0.91	0.98	0.94
	P-Value	0.00	0.00	0.00	0.00

shows the gender-wise details on regimen and interpersonal related distress based on low, moderate and high distress levels among men and women. Regimen related distress indicated that 11.5% of men and 49.5% of women showed high distress levels. In interpersonal related distress, 14% of men and 2% of women indicated moderate distress levels, whereas 44.5% of women and 7% men indicated high distress level.

Figure 2 shows the distress levels in relation to age groups. It was observed that the levels of diabetes distress decreased with increasing age. On examining the differences between the highly distressed and moderately/mildly distressed groups, significant relationship was noted between domain scores, age and HbA1c levels. In total, there was a strong statistically significant positive correlation between HbA1c and diabetes distress domain score; emotional burden ($r=0.95$, $p<0.001$), physician related distress ($r=0.91$, $p<0.001$), regimen related distress ($r=0.98$, $p<0.001$) and interpersonal distress ($r=0.94$, $p<0.001$). Age also was negatively correlated

with individual domain score; emotional burden ($r=-0.15$, $p<0.001$), physician related distress ($r=-0.15$, $p<0.001$), regimen related distress ($r=-0.16$, $p<0.001$) interpersonal distress ($r=-0.14$, $p<0.001$) (Table 3).

DISCUSSION

Diabetes distress is a common health problem which frequently co-exists with T2DM (Islam et al., 2013; Lloyd et al., 2018; Nanayakkara et al., 2018). The present study highlighted that 22.5% of men and about 55% of women were screened positive for moderate to high DRD on a DDS-17 scale. Another study using DDS-17 total scores (Aljuaid et al., 2018) also reported that T2DM subjects were more likely to have DRD especially women when they belonged to the low income, unemployed or had any diabetes complications. Similar results were also observed in our study which indicated high levels of diabetes distress in women.

An earlier study conducted in USA (Fisher et al., 2012) using the DDS-17 scale showed that 51.3% of the screened participants have moderate to high DRD. Similar distress proportions were shown from the studies conducted in Bangladesh (48.5%), China (43%) and Canada (39%) using DDS-17 (Islam et al., 2013; Wong et al., 2017; Zhou et al., 2017). Emotional burden was considered as the most important domain in measuring diabetes distress especially among women. This study finding is consistent with the study conducted in the Bangladesh (Islam et al., 2013) population.

The current study showed a positive correlation between both DRD total score and emotional distress with the glycemic control. It was noted that those who had high levels of DDS had poor glycemic control. Similar finding was also observed in other populations (Islam et al., 2013; Kuniss et al., 2017; Aljuaid et al., 2018). Diabetes related distress generally shows a closer association with glycemic control. This finding is consistent with another study conducted by Gonzalaz et al. (2011). DDS showed closer association with glycemic control than depression and appears to be more common and chronic than depression in adults with T2DM as shown by Fisher et al. (2010). International survey data also confirmed and suggested that emotional well-being is the domain of functioning most negatively affected by diabetes, second only to physical health (Nicolucci et al., 2013).

It was also observed in the current study that as the age increased the diabetes distress levels decreased. This finding is consistent with the earlier study conducted by Fisher et al. (2012) which documented the positive association of DD with age. This may be attributed to their gradual adjustments towards their diabetes life style modifications. The following are the limitations of the current study. Parameters like level of education, socioeconomic status, employment have not been evaluated in association with DDS. Secondly, a comparative level of distress was not assessed in subjects with oral antidiabetic drugs, insulin or combination. The factors associated with DRD should be explained and need to be further studied in depth to formulate proper guidance and empowerment in diabetes management plan.

Conclusion

This study highlights that women had high levels of distress in managing diabetes as compared to men. Diabetes distress should therefore be considered as a significant health problem and steps should be taken for effective management like lifestyle modifications as well as methods to cope with their stress and diabetes.

CONFLICT OF INTERESTS

The authors have not declared any conflict of interests.

ACKNOWLEDGEMENTS

The authors appreciate the help rendered by Sriram Ramachandran (Bio-statistician) for analyzing the data and all diabetes patients who participated in the study.

REFERENCES

- Aljuaid MO, Almutairi AM, Assiri MA, Dhifallah MA, Khaled A (2018). Diabetes related distress assessment among type 2 diabetes patients. *Journal of Diabetes Research* 1-10.
- Deshpande AD, Harris-Hayes M, Schootman M (2008). Epidemiology of diabetes and diabetes-related complications. *Physical Therapy* 88:1254-1264.
- Fisher L, Mullan JT, Areal P, Glasgow RE, Hessler D, Masharani U (2010). Diabetes distress but not clinical depression or depressive symptoms is associated with glycemic control in both cross-sectional and longitudinal analyses. *Diabetes Care* 33:23-28.
- Fisher L, Mullan JT, Skaff MM, Glasgow RE, Areal P, Hessler D (2009). Predicting diabetes distress in patients with type 2 diabetes: a longitudinal study. *Diabetes Medicine* 26:622-627.
- Fisher L, Hessler DM, Polonsky WH, Mullan J (2012). When is diabetes distress clinically meaningful? Establishing cut points for the Diabetes Distress Scale. *Diabetes Care* 35(2):259-264.
- Fowler MJ (2008). Microvascular and macrovascular complications of diabetes. *Clinical Diabetes* 26:77-82.
- Gonzalez JS, Fisher L, Polonsky (2011). Depression in diabetes: have we been missing something important? *Diabetes Care* 34:236-239.
- Islam RM, Karim MR, Habib SH, Yesmin K (2013). Diabetes distress among type 2 diabetic patients. *International Journal of Medicine and Biomedical Research* 2:113-124.
- Kuniss N, Rehtacek T, Kloos C, Müller UA, Roth J, Burghardt K, Kramer G (2017). Diabetes-related burden and distress in people with diabetes mellitus at primary care level in Germany. *Acta Diabetologica* 54(5): 471-478.
- Lloyd CE, Nouwen A, Sartorius N et al. (2018). Prevalence and correlates of depressive disorders in people with Type 2 diabetes: results from the International Prevalence and Treatment of Diabetes and Depression (INTERPRET-DD) study, a collaborative study carried out in 14 countries. *Diabetic Medicine* 35:760-769.
- Mathers ACD, Loncar D (2006). Projections of global mortality and burden of disease from 2002 to 2030. *PLoS Medicine* 3:e442.
- Nanayakkara N, Pease A, Ranasinha S, Wischer N, Andrikopoulos S, Speight J, de Courten B, Zoungas S (2018) Depression and diabetes distress in adults with type 2 diabetes: results from the Australian National Diabetes Audit (ANDA) 2016. *Scientific Report* 8:7846.
- Nicolucci A, Burns KK, Holt R, Comaschi M, Hermanns N, Ishii H, Kokoszka A, Pouwer F, Skovlund SE, Stuckey H, Tarkun I, Vallis M, Wens J, Peyrot M; DAWN2 Study Group (2013). Diabetes attitudes wished and needs second study (DAWN 2): cross-national benchmarking of diabetes-related psychosocial outcomes for people with diabetes. *Diabetic Medicine* 30:767-777.
- Ogurtsova KJD, Fernandes R, Huang Y, Linnenkamp U, Guariguata L, Cho NH, Cavan D, Shaw JE, Makaroff LE (2017). *IDF Diabetes Atlas: global estimates for the prevalence of diabetes for 2015 and 2040*, Diabetes Research and Clinical Practice 128:40-50.
- Polonsky WH, Fisher L, Earles J, Dudl RJ, Lees J, Mullan J, Jackson RA (2005). Assessing psychosocial Distress in diabetes: development of the Diabetes Distress Scale. *Diabetes Care* 28(3):626-631.
- Schetman JM, Nadkarni MM, Voss JD (2002). The association between diabetes metabolic control and drug adherence in an indigent population. *Diabetes Care* 25:1015-1021.
- Wong EM, Afshar R, Qian H, Zhang M, Elliott TG, Tang TS (2017). Diabetes distress, depression and glycemic control in a Canadian-based specialty care setting, *Canadian Journal of Diabetes* 41(4):362-365.
- Zhou H, Zhu J, Liu L, Li F, Fish AF, Chen T, Lou Q (2017). Diabetes-related distress and its associated factors among patients with type 2 diabetes mellitus in China. *Psychiatry Research* 252:45-50.

Related Journals:

