


OPEN ACCESS



30 March 2019
ISSN 1992-1950
DOI: 10.5897/IJPS
www.academicjournals.org

About IJPS

The International Journal of Physical Sciences (IJPS) is a peer reviewed journal. The journal publishes articles in all areas of physical sciences such as: Artificial intelligence, Neural processing, Nuclear and particle physics, Geophysics, Physics in medicine and biology, Plasma physics, Semiconductor science and technology Wireless and optical communications, Materials science, Energy and fuels, Environmental science and technology, Combinatorial chemistry, Geochemistry, Cement and concrete research, Metallurgy, Crystallography and Computer-aided materials design.

Open Access Policy

The International Journal of Physical Sciences is an Open Access journal. Abstracts and full texts of all articles published in this journal are freely accessible to everyone immediately after publication without any form of restriction.

Article License

All articles published by International Journal of Physical Sciences are licensed under the [Creative Commons Attribution 4.0 International License](#). This permits anyone to copy, redistribute, remix, transmit and adapt the work provided the original work and source is appropriately cited. Citation should include the article DOI. The article license is displayed on the abstract page the following statement:

This article is published under the terms of the [Creative Commons Attribution License 4.0](#)

Please refer to <https://creativecommons.org/licenses/by/4.0/legalcode> for details

about [Creative Commons Attribution License 4.0](#)

Article Copyright

When an article is published by in the International Journal of Physical Sciences, the author(s) of the article retain the copyright of article. Author(s) may republish the article as part of a book or other materials. When reusing a published article, author(s) should;

Cite the original source of the publication when reusing the article. i.e. cite that the article was originally published in the International Journal of Physical Sciences. Include the article DOI
Accept that the article remains published by the International Journal of Physical Sciences (except in occasion of a retraction of the article)

The article is licensed under the Creative Commons Attribution 4.0 International License.

A copyright statement is stated in the abstract page of each article. The following statement is an example of a copyright statement on an abstract page.

Copyright ©2016 Author(s) retains the copyright of this article.

Self-Archiving Policy

The International Journal of Physical Sciences is a RoMEO green journal. This permits authors to archive any version of their article they find most suitable, including the published version on their institutional repository and any other suitable website.

Please see <http://www.sherpa.ac.uk/romeo/search.php?issn=1684-5315>

Digital Archiving Policy

The International Journal of Physical Sciences is committed to the long-term preservation of its content. All articles published by the journal are preserved by [Portico](#). In addition, the journal encourages authors to archive the published version of their articles on their institutional repositories and as well as other appropriate websites.

<https://www.portico.org/publishers/ajournals/>

Metadata Harvesting

The International Journal of Physical Sciences encourages metadata harvesting of all its content. The journals fully supports and implement the OAI version 2.0, which comes in a standard XML format. [See Harvesting Parameter](#)

Contact

Editorial Office: ijps@academicjournals.org

Help Desk: helpdesk@academicjournals.org

Website: <http://www.academicjournals.org/journal/IJPS>

Submit manuscript online <http://ms.academicjournals.org>

Academic Journals
73023 Victoria Island, Lagos, Nigeria
ICEA Building, 17th Floor,
Kenyatta Avenue, Nairobi, Kenya.

Editors

Prof. Sanjay Misra

Editor-in-Chief

Department of Computer and Information Science
Covenant University
Nigeria.

Prof. Songjun Li

School of Materials Science and Engineering
Jiangsu University
Zhenjiang, China.

Prof. Xiao-Li Yang

School of Civil Engineering
Central South University
Hunan, China.

Editorial Board Members

Dr. G. Suresh Kumar

Biophysical Chemistry Division
Indian Institute of
Chemical Biology (IICB),
Kolkata, India.

Prof. Jr-Hau He

National Taiwan University
Taipei, Taiwan.

Dr. Sunil Kumar Yadav

Department of Mathematics
Alwar Institute of Engineering &
Technology
Alwar, India.

Prof. Zafar Iqbal

Department of Chemistry and Environmental Science
New Jersey Institute of Technology
Newark, USA.

Dr. Tomasz Baczek

Medical University of Gdansk
Gdansk, Poland.

Dr. Ricardo Martinho

Department of Informatics Engineering
School of Technology and Management
Polytechnic Institute of
Leiria Apartado, Portugal.

Dr. Jocenir Boita

Universidade Federal de
Santa Maria - UFSM
Campus Cachoeira do Sul
Cachoeira do Sul, RS,
Brazil.

Dr. Omar Abu Arqub

Mathematics Department
Al Balqa Applied
University Jordan.

Prof. H. M. Srivastava

Department of Mathematics and Statistics
University of Victoria,
Victoria, Canada.

Prof. Liangchi Zhang

School of Aerospace Mechanical
and Mechatronic Engineering,
The University of Sydney
Australia.

Prof. Awrejcewicz Jan

Department of Automatics
and Biomechanics Łódź,
Lodz University of
Technology Poland.

Prof. Nazmul Islam

Department of Basic Sciences
& Humanities/Chemistry,
Techno Global-Balurghat
Mangalpur, India.

Prof. Ismail Musirin

Centre for Electrical
Power Engineering
Studies (CEPES),
Faculty of Electrical
Engineering,
Universiti Teknologi
Mara,
Selangor, Malaysia.

Dr. Mohammadreza Saeidi

Department of Physics
Faculty of Basic Science
Shahed University
Tehran, Iran.

Prof. N. V. Sastry

Department of Chemistry
Sardar Patel University
Gujarat, India.

Dr. Luigi Maxmilian Caligiuri

Department of Chemistry and Chemical
Technology

University of Calabria
and Foundation of Physics Research
Center (FoPRC)
Italy.

Dr. Walid Mohamed

Physics Department
King Saud University
Saudi Arabia.

Dr. Masroor Hassan Shah Bukhari

Physics Department
Jazan University
Saudi Arabia.

Dr. Premkumar Thathan

Department of Chemistry
Sungkyunkwan University
South Korea.

Dr. Bidyut Saha

Chemistry Department
Burdwan University
WB, India.

Dr. Pouya Derakhshan-Barjoei

Electrical and Computer Engineering
Islamic Azad University
Naein, Iran.

Dr. Pooran Koli

Department of Chemistry,
Jai Narain Vyas University,
Jodhpur-342001,
Rajasthan,
India

Table of Content

Magnetic source basement depth determination of Ikogosi Warm Water Spring South Western Nigeria and the environ using aeromagnetic data
Ayobami I. Ojoawo and Muideen I. Lateef

Full Length Research Paper

Magnetic source basement depth determination of Ikogosi Warm Water Spring South Western Nigeria and the environ using aeromagnetic data

Ayobami I. Ojoawo* and Muideen I. Lateef

Department of Physics, Faculty of Science, University of Ibadan, Ibadan, Nigeria.

Received 2 December, 2018; Accepted 19 February, 2019

Structural features of Ikogosi Warm Spring and its environs have been delineated using qualitative and quantitative interpretation methods. Euler-Deconvolution and local wave number methods were adopted for quantitative and qualitative interpretation of aeromagnetic data sheet 243 by Geological Survey Agency of Nigeria (GSN) in 2008, respectively. Euler solution revealed depth range of -2.98 to -290 m to the magnetic sources, while local wave number method revealed depths to magnetic sources/basements ranging from -2.1 to -1311.4 m of the geologic lineaments in Ikogosi Warm Spring and its environs. Both methods revealed outcrop features characterized by their positive depth values above the mean ground level. Both methods proved efficient way of resolving shallow magnetic source depth giving a very plausible depth of -2.1 m for local wave number and -2.98 m for Euler solution in the area of study.

Key words: Warm-Spring, Ikogosi, lineament.

INTRODUCTION

Ikogosi Warm Spring at Ekiti, South Western Nigeria is believed by many to be a viable geothermal energy source for the generation of electricity which is a major need of the nation for economic growth and all other forms of development which directly or indirectly depend on electricity. This belief is attributed to the fact that natural water is supposed to be cool and not warm or hot as the Ikogosi Warm Spring is. There is need to scientifically study the substructures of the area to understand the structural setting of the area. This information about these substructures and their depths to the surface can be used to suggest the existence of any potential reservoir or fault and fractures in the area.

Location of the study area

The Ikogosi Warm Spring is located in the southwestern part of Ekiti State of Nigeria. It is situated between lofty steep-sided and heavily wooded, north-south trending hills about 17 miles (approximately 27.4 km) east of Ilesha, and about 6.5 miles (approximately 10.4 km) southeast of Efon Alaye (Rogers et al., 1969). It lies on the geographic latitude of 7°35'N and longitude 5°00'E (Figure 1) within the central region of the area covered by this study. Located within the Precambrian basement complex of South Western Nigeria, it is at an altitude of 450 to 500 m (Adegbuyi and Abimbola, 1997).

*Corresponding author. Ojoawo@yahoo.com. Tel: +2348059350482.




Figure 1. Geological map of Nigeria showing the location of Ikogosi Warm Spring (blue arrow) in the basement complex of Nigeria. Source: Modified from Obaje (2009).

The dominant geology of Nigeria is made mainly by crystalline (Precambrian basement complex) and sedimentary rocks (Cretaceous recent sediments).

The area covered by this study lies approximately between geographic latitudes 7°30'N and 8°00'N and geographic longitude 4°30'E and 5°00'E within the Precambrian of South Western Nigeria. The area is covered by the aeromagnetic map sheet 243 (Ilesha).

Geological setting of the study area

The warm spring temperature is 38°C near the foot of the eastern slope of the north-south trending ridge from a thin quartzite unit within a belt of quartzite which includes quartz-mica schist and granulitic migmatite east of Ilesha. The Okemesi quartzite member is characterized by a North-South trending ridge called the Effon ridge (Elueze, 1988; Oyinloye, 2011). The quartzitic rocks (as shown in Figure 2) are composed of dominant quartz with muscovite, chlorite and sericite occurring in minor proportions (Adegbuyi and Abimbola, 1997). It was suggested that the source of springs in the Effon Psammite formation is associated with a faulted and fractured quartzite band sandwiched between schists

(Rogers et al., 1969). Chemical data of Ikogosi shows that quartzite is largely metamorphosed sandstones containing minor arkosic intercalations (Elueze, 1988). On the basis of petrology, a medium pressure Barrovian and low medium pressure types of metamorphism had been suggested for the Precambrian basement rocks in South Western Nigeria (Oyinloye, 2011). It is believed that the intersections of the NNE-SSW epeirogenic belts with the NW-SE fracture trends in Nigeria coincide with the centers of warm springs like the Wikki (Bauchi State) and Ikogosi (Ekiti State) springs (Mbonu, 1990). The issue of the springs is controlled by permeability developed within the quartzite as a result of intergranular pore spaces coupled with fracturing of the relatively competent quartzite (Rogers, 1969).

THEORETICAL BACKGROUND

Geomagnetic field is produced by electric currents induced within the conductive liquid outer core as a result of slow convective movements within it. The behavior of these fields can be characterized by a vector quantity known as magnetization, M . The magnetic induction B , which is composed of the ambient earth's field and




Figure 2. Geology map of Ikogosi Warm Spring and environ. Source: Adegbuyi and Abimbola (1997).

magnetic material in the subsurface is related to M by:

$$B(R) = \frac{-\mu_0}{4\pi} \int M(R_0) \cdot \nabla_0 \frac{1}{R - R_0} dV_0 \quad (1)$$

where μ is known as permeability of free space and R and R_0 are the observation and source locations, respectively.

The Total Magnetic Field B represents the sum of the magnetizing field strength and the magnetization of the medium:

$$B = \mu_0(1 - k)H = \mu\mu_0H \quad (2)$$

where μ_0 is magnetic permeability of free space (4×10^{-7} H/m), B is also called the magnetic flux density or magnetic induction.

Earth's magnetic field varies between 20,000 nT in the equator and 60,000 nT at the pole (Reynolds, 1997).

Euler De-Convolution

The Euler De-convolution is an interpretation tool in potential field for locating anomalous sources and the determination of their depths by De-convolution using Euler's homogeneity relation (Reid et al., 1990).

The Euler's homogeneity equation relates the magnetic field and its gradient components to the location of the source of an anomaly, with the degree of homogeneity expressed as a structural index (Reynolds, 1997). Euler's homogeneity relationship can be written for magnetic data as:

$$(x - x_0) \frac{dT}{dx} + (y - y_0) \frac{dT}{dy} + (z - z_0) \frac{dT}{dz} = N(B - T) \quad (3)$$

where

$$\frac{dT}{dx}, \frac{dT}{dy} \text{ and } \frac{dT}{dz} \quad (4)$$

represent first order derivative of the magnetic field along x-, y-, and z-directions, respectively. Where (x_0, y_0, z_0) is the location of a magnetic source, whose total field magnetic anomaly at the point (x, y, z) is T and B is the regional field. N is structural index and is a measure of the rate of decay of field with distance and assumes different values for different types of magnetic source. The structural index, S.I, depends on the geometry of the source. For a homogeneous point source $N = 3$, a linear source (line of dipoles or poles, and for a homogeneous cylinder (rod, etc.) $N = 2$, for extrusive bodies (thin layer, dike, etc.) $N = 1$, for a contact, vertex of a block and a pyramid with a big height $N = 0$.




Figure 3. Residual anomaly contour map showing the profile lines in the direction NE-SW.

Local wave number method

This method is based on extension of complex analytical signal to estimate magnetic depths. The method works for 2-D slopping contact or 2-D dipping thin-sheet. For dipping contacts, the maxima of K are located directly over the isolated field contact edges and are independent of the magnetic field inclination, declination, dip, strike, and any remanent magnetization.

Depth can also be estimated without assumptions about the thickness of the source bodies. For magnetic field M, the local wave number (Salako, 2014) is given by:

$$K = \frac{\frac{\partial^2 M}{\partial x \partial z} \frac{\partial M}{\partial x} - \frac{\partial^2 M}{\partial x^2} \frac{\partial M}{\partial z}}{\left(\frac{\partial M}{\partial x}\right)^2 + \left(\frac{\partial M}{\partial z}\right)^2} \tag{5}$$

The depth is estimated at the source edge from the reciprocal of the local wave number (K).

MATERIALS AND METHODS

A high resolution aeromagnetic data which are part of the airborne geophysical data of Nigeria acquired by Fugro Exploration in 2008 and published by the Geological Survey Agency of Nigeria (GSN) on a total magnetic intensity map scale of 1:100,000 were processed and interpreted to yield the set objectives. The data were

leveled, cultural-edited, high frequency noise filtered before further processing. Error correction, spike removal, gridding were carried out prior to other filtering operations to yield the proposed target.

Subsequently, the general description of the survey results and the explanation of the major features called anomalies which can be geological formation and/or structures were discussed. The qualitative interpretation was performed using grids of residual field of the total magnetic intensity, reduction to the equator, upward continuation, horizontal derivatives and CET (Centre for Exploration Targets) analysis maps. The aeromagnetic data were input into the Geosoft Oasis Montaj software that processed them into grids and maps after which the gridded data was contoured on Surfer. The output results were then interpreted qualitatively using the Euler De-convolution and quantitatively using Local Wave Number Method.

RESULTS AND DISCUSSION

Residual anomaly contour map

The residual anomaly contour map was digitized into ten profiles as shown in Figure 3 and profile data were analyzed.

Most of the profiles have both relatively low negative and high positive magnetic anomaly field strengths. Zones with low negative magnetic anomalies are suspected to be the embedded features (faults/fractures) and those with positive anomalies are suspected to be massive quartzite/basement complex embedding the embedded features. The prominent sudden change in the contour over an appreciable distance that trends




Figure 4. Lineament map of Ikogosi Warm Spring and its environs.

frequently in the south-west direction of the study area implies discontinuity in depth. This could be subsurface major faults.

CET analysis map (Lineament Map)

The map (Figure 4) shows geological features (faulted zones, fractured zones) that are elongated in NE-SW direction. This shows that the majority of the fracture zones and faults are NE-SW in direction although they are of different depths and width. The lineaments bounded from (7°35', 4°55'), (7°35', 5°) to (7°40', 4°55'), (7°40', 5°) (Figure 4) are sandwiched in the quartzite formation of the basement complex of the study area (Figure 2).

Euler-Deconvolution solution

The Euler solution revealed depth range of -290 to 47 m for sources located below and above the mean ground level. The positive depths coincide with regions where sources/basement rocks outcrop on the surface while the negative depths coincide with depth to magnetic sources/

basement rocks.

The Ikogosi Warm Spring region with location (7°35'30"N,4°58'45"E) and (7°35'45"N,4°58'54"E) has features with depth range of -2.98 to -290 m with the shallowest anomaly source (basement) having a depth of -2.98 m.

Prominent clustering of lineaments was obtained on the Lineament map (Figure 4) between (7°30'N, 4°40'E) and (7°40'N,5°E) showing definite magnetic trends within the Ikogosi Warm Spring and its environs. These linear features trending NE-SW are suspected fractured/faulted zones with depths in between -2.98 and -290 m if it is superimposed on the depth contour plot (Figure 6).

With the aforementioned information and the range of depth values obtained, it could be suggested that the linear trending features (fracture or fault or both) rimmed by basement complex existing in the shallow and deep basements of Ikogosi Warm Spring and its immediate neighbourhood interface the spring outlet and act as channels for movement of warm ground water from profound depths to the spring outlet (Figure 5).

The Euler depth solution grid was digitized at the Ikogosi Warm Spring location {(7°35'30"N,4°58'45"E) and (7°35'45"N,4°58'54"E)} on the map. The digitized depth data were krigged to give Figure 6. This plot shows



Figure 5. Ikogosi Warm Spring source.




Figure 6. Depth contour plot from Euler-Deconvolution grid of Ikogosi Warm Spring.

that the shallowest depth is between 0 and -5 m which is inline with the shallowest depth of this region earlier suggested.

Local wave number solution

The Local wave number solution revealed depths range




Figure 7. Depth contour plot from local wave number grid of Ikogosi Warm Spring.

of -1311.4 to 2403 m for sources located below and above the mean ground level. The positive depths coincide with region where basement rocks outcrop on the surface while the negative depths coincide with depth to magnetic sources/basement rocks

The local wave number solution revealed depths to magnetic sources/basements ranging from -2.1 to -1311.4 m. So the shallowest magnetic source in the area of the study has a depth of -2.1 m and the deepest-seated magnetic source has a depth of -1311.4 m.

The local wave number depth solution grid was digitized at the Ikogosi Warm Spring location $\{(7^{\circ}35'30''N, 4^{\circ}58'45''E)$ and $(7^{\circ}35'45''N, 4^{\circ}58'54''E)\}$ on the map. The digitized depth data were grided to give Figure 7. This plot shows that the shallowest depth is between 0 and -100 m which encompasses the shallowest depth as suggested.

This finding revealed shallowest depths to magnetic sources compared to depths to magnetic sources (quartzite) which ranged from 50.9 to 227.7 m suggested by Ojo et al. (2011).

The depths suggested in other authors' works on this area presented curie depths associated with the intense temperature that causes surface manifestation of heat.

Conclusion

The analysis of aeromagnetic data from this region revealed lineaments which could be fracture or fault or both have been delineated with trends mapped out to be elongated in the NE-SW directions. Both methods used have proved efficient for resolving well for shallow magnetic source depth giving a very plausible depth of -2.1 m for local wave number and -2.98 m for Euler solution in the area of study. But local wave number solution resolved very well for deep-seated magnetic sources by yielding the least depth value of -1311.4 m. Both methods also have revealed heights of outcrop features characterized by their positive depth values above the mean ground level.

These estimated relatively shallow magnetic sources in the area of study could be an indication of magma intrusion which solidified to form a basement rock and thus has several implications on geothermal resources and tectonic activities in the area. The heat flow in the tectonically active region is enough to cause surface manifestation of geothermal resource.

It is also suspected that the probable fractured/faulted zones mapped out may have acted as channels for movement of warm deep water from profound depths to

the surface and the spring outlet is on a lineament interface.

CONFLICT OF INTERESTS

The authors have not declared any conflict of interests.

REFERENCES

- Adegbuyi O, Abimbola AF (1997). Energy resource potential of Ikogosi Warm Spring Area, Ekiti State, Southwestern Nigeria. *African Journal of Science* 1(2):111-117.
- Elueze AA (1988). Geology of the Precambrian schist belt in Ilesha area, southwestern Nigeria. *Precambrian geology of Nigeria*. Geological Survey of Nigeria pp. 77-89.
- Kearey P, Brooks M, Hill I (2002). *An Introduction to Geophysical Exploration*, Blackwell Publishing, 3rd Edition.
- Keating and Pilkington (2004). Euler deconvolution of the analytic signal and its application to magnetic interpretation. *Geophysical prospecting* 52(3):165-182. <https://doi.org/10.1111/j.1365-2478.2004.00408.x>
- Mbonu WO (1990). The trends and geologic consequences of belts of epeirogeny in Nigeria. *Book of Abstracts, Nigeria Min. Geoscience Conference* P 141.
- Obaje NG (2009). Geology and mineral resources of Nigeria. In: *Lecture Notes in Earth Sciences* 120:221.
- Ojo JS, Olorunfemi MO, Faletiba DE (2011). An appraisal of the geologic structure beneath the Ikogosi warm spring in South Western Nigeria using integrated surface geophysical methods. *Earth Sciences Research Journal* 15(1):27-34.
- Olaniyi OE, Atalor NO (2018). Land use/ Land cover dynamics around ecotourism attractions and support facilities in Ikogosi Warm Spring Resorts, Nigeria. *Journal of Forestry Research and Management*, 15(1):196-220.
- Oyinloye AO (2011). Geology and geotectonic setting of the basement complex rocks in South Western Nigeria: implications on provenance and evolution. In *Earth and Environmental Sciences*. Intech Open.
- Parasnis DS (1986). *Principles of Applied Geophysics*, Chapman Hall, London pp. 43-44.
- Ravat D (1996). Analysis of the Euler method and its applicability in environmental magnetic investigations. *Journal of Environment Engineering and Geophysics* 1:229-238.
- Reid AB, Allsop JM, Grauser H, Millet AJ, Somerton IN (1990). Magnetic interpretation in three dimensions using Euler deconvolution. *Geophysics* 55:80-91.
- Reynolds JM (1997). *An Introduction to Applied and Environmental Geophysics*. John Wiley and Sons, England. pp. 118-121.
- Roest WR, Verhoef J, Pilkington M (1992). Magnetic interpretation using the 3-D analytic signal. *Geophysics* 57(1):116-125
- Rogers AS, Imevbore AMA, Adegoke OS (1969). Physical and chemical properties of Ikogosi Warm Spring, western Nigeria. *Nigeria Journal of Mining Geology* 4:1-2.
- Salako KA (2014). Depth to Basement Determination Using Source Parameter Imaging (SPI) of Aeromagnetic Data: An application to Upper Benue Trough and Borno Basin, Northeast, Nigeria. *Academic Research International* 5(3):10-29
- Thompson DT (1982). EULDPH: A new technique for making computer-assisted depth estimates from magnetic data. *Geophysics* 47(1):31-37.

Related Journals:

