

OPEN ACCESS



International Journal of
**Biotechnology and Molecular
Biology Research**

August 2019
ISSN 2141-2154
DOI: 10.5897/IJBMBR
www.academicjournals.org



**ACADEMIC
JOURNALS**
expand your knowledge

ABOUT IJBMBR

The **International Journal for Biotechnology and Molecular Biology Research (IJBMBR)** (ISSN 2141-2154) is published Monthly (one volume per year) by AcademicJournals.

International Journal for Biotechnology and Molecular Biology Research (IJBMBR) provides rapid publication (monthly) of articles in all areas of the subject such as Green energy from chemicals and bio-wastes, Studies in the graft copolymerization of acrylonitrile onto cassava starch by ceric ion induced initiation, Antimutagenic activity of aqueous extract of Momordica charantia, Ethnomedicinal plants and other natural products with anti-HIV active compounds and their putative modes of action etc.

The Journal welcomes the submission of manuscripts that meet the general criteria of significance and scientific excellence. Papers will be published shortly after acceptance. All articles published in IJBMBR are peer-reviewed.

Contact Us

Editorial Office: ijbmb@academicjournals.org

Help Desk: helpdesk@academicjournals.org

Website: <http://academicjournals.org/IJBMBR>

Submit manuscript online <http://ms.academicjournals.me/>

Editors

Prof Atagana, Harrison

*Institute for Science and Technology Education
University of South Africa*

Prof. UC Banerjee

*Department of Pharmaceutical Technology
(Biotechnology)
National Institute of Pharmaceutical Education and
Research
Punjab, INDIA*

Dr. Y. Omid

*Faculty of Pharmacy,
Research Center for Pharmaceutical Nanotechnology,
School of Advanced Biomedical Sciences,
Tabriz University of Medical Sciences,
Tabriz, Iran.*

Prof. Mohamed E. Wagih

*University of New Brunswick (UNB-SJ),
Saint John College, NB,
E2L 4L5, Canada*

Dr. Sripada M. Udupa

*ICARDA-INRA Cooperative Research Project,
International Center for Agricultural Research in the
Dry Areas(ICARDA), B.P. 6299,
Rabat Instituts, Rabat, Morocco*

Dr. Amjad Masood Husaini

*Sher-e-Kashmir University of Agricultural Sciences &
Technology
Bohlochipora, Dr. Ali Jan Road,
Nowshera, Srinagar, J&K-190011, India*

Dr. Om Prakash Gupta

*Directorate of Wheat Research (ICAR)
Post Box-158, A
grasain Marg, Karnal-132001,
Haryana, India*

Editorial Board

Dr. Amro Hanora

*Suez Canal University, Department of Microbiology
and Immunology,
Faculty of Pharmacy, Suez Canal University,
Box 41522 Ismailia, Egypt*

Dr. C. Rajasekaran

*VIT University
School of Bio-Sciences & Technology (SBST)*

Dr. Yasar Karadag

*Gaziosmanpasa University
Faculty of Agriculture,
Department of Field Crops, Tokat-Turkey*

Dr. Ahmet Tutus

*KSU (Kahramanmaras Sutcu Imam University)
Faculty of Forestry,
Department of Forest Industrial Engineering,
Kahramanmaras 46100 Turkey*

Dr. Vinod Joshi

*Desert Medicine Research Centre,
Indian Council of Medical Research
New Pali Road, Jodhpur, India*

Dr. Eshrat Gharaei Fathabad

*K.M.18 Khazarabad road,
Sari, Mazandaran, Iran*

Dr. Shashideep Singhal

*121 Dekalb Ave, Brooklyn,
NY 11201, New York, USA*

Dr Masayoshi Yamaguchi

*101 Woodruff Circle, 1305 WMRB,
Atlanta, Georgia 30322-0001, USA*

Dr. Okonko Iheanyi Omezuruike

*Department of Virology,
Faculty of Basic Medical Sciences,
College of Medicine,
University College Hospital,
Ibadan, Nigeria*

Dr. S. M. Shahid

*University of Karachi,
Karachi-75270, Pakistan*

Prof. Reda Helmy Sammour

*Botany Department,
Faculty of Science,
Tanta University,
Tanta, Egypt*

Dr. Premendra D. Dwivedi

*Food Toxicology Division,
Room No 303,
P.O. Box 80,
M. G. Road, Lucknow-226001, UP, India*

Dr. Amro Hanora

*Microbiology and Immunology department,
Faculty of Pharmacy,
Suez Canal University,
Box 41522 Ismailia, Egypt*

Dr. Tamilnadu

*1501 N. Campbell Ave
Tucson, AZ 85724 India*

Dr. Yadollah Omid

*Faculty of Pharmacy,
Tabriz University of Medical Sciences,
Daneshghah St., Tabriz, Iran*

Dr. Mohsen Selim Asker

*National Research Centre, Dokki, Cairo,
Egypt*

Dr. Fanuel Lampiao

P.O.Box 360, Blantyre, Malawi

Prof. Mohamed E. Wagih

Saint John, NB, E2L 4L5, Canada

Dr. Santosh Kumar Singh

*Centre of Experimental Medicine and Surgery,
Institute of Medical Sciences,
Banaras Hindu University, Varanasi-221005, India*

Dr. Zhanhuan Shang

*No.768, Jiayuguan West Road,
Lanzhou City, Gansu Province, China*

Dr. Worlanyo Eric Gato

*Southern Illinois University – Carbondale,
1245 Lincoln Dr, 144 Neckers, Carbondale IL 62901*

Dr. Chun Chen

*College of Life Sciences,
China Jiliang University
Xueyuan Street, Xiasha, Hangzhou,
Zhejiang Province, PR China*

Dr. Efthimios J.Karadimas

*LGI, Leeds NHS Trust 10th Timoleontos Vassou str,
11521, Athens Greece*

Dr. Samuel Toba Anjorin

University of Abuja, Abuja, Nigeria

Dr. Rupali Agnihotri

*Department of Periodontics,
Manipal college of Dental Sciences,
Manipal,576104. Karnataka. India*

Dr. Mahbuba Begum

*Tuber Crops Research Centre,
Joydebpur, Gajipur-1701, Bangladesh*

Prof. S. Mohan Karuppaiyl

*School of Life Sciences
Srtm University
Nanded. MS. India*

Dr. Neveen Helmy Aboelsoud

*Complementary Medicine Researches and Application
Department
National Research Center, Cairo
Egypt.*

Dr. D.E. Chandrashekar Rao

*National Research Council,
Plant Biotechnology Institute
Canada*

Dr. Nikolaos Papanas

*Democritus University of Thrace
G. Kondyli 22, Alexandroupolis, Greece*

Dr. Sivakumar Swaminathan

*Iowa State University
USA*

Dr. El Sayed Hussein El Sayed Ziedan

*National Research Centre,
Plant Pathology Department
Tahrir St.,Dokki Cairo,
Egypt*

Dr. Chethan Kumar M

*Post Graduate Departments of Bio-technology and Biochemistry,
Ooty Road, Mysore - 570 025,
Karnataka,
India*

Dr. M. Sattari

*Rice Research Ins. of Iran
Iran*

Dr. Zaved Ahmed Khan

*VIT University
India*

Dr. Subbiah Poopathi

*Vector Control Research Centre
Indian Council of Medical Research
(Ministry of Health & Family Welfare,
Govt. of India)
Medical Complex, Indira Nagar
India*

Dr. Reyazul Rouf Mir

*International Crops Research Institute for the Semi-Arid Tropics (ICRISAT),
Patancheru - 502 324, Greater Hyderabad,
India*

Dr. Prasanna Kumar S

*Virginia Commonwealth University,
USA*

Dr. Naseem Ahmad

*Plant Biotechnology Laboratory
Department of Botany
Aligarh Muslim University
Aligarh- 202 002, (UP)
India*

Dr. Zhen-Xing Tang

*Food Bioengineering institute, Hangzhou Wahaha Co.
Ltd, Hangzhou,
Zhejiang,
China*

Dr. Jayanthi Abraham

*VIT (Vellore Institute of Technology) University,
Tamilnadu,
India*

Dr. Gobianand Kuppannan

*National Institute of Animal Science
South Korea*

Dr. R. Harikrishnan

*Jeju National University
South Korea*

Dr. Asit Ranjan Ghosh

*Vellore Institute of Technology (VIT) University,
School of Bio Sciences & Technology,
Medical Biotechnology Division, Vellore-632014,
India*

Dr. Kamal Dev

*Shoolini University of Biotechnology and Management
Sciences (SUBMS)
India*

Dr. Wichian Sittiprapaporn

*Maharakham University
Thailand*

Dr. Vijai Kumar Gupta

*Molecular Glycobiotechnology Group, Department of Biochemistry,
School of Natural Sciences,
National University of Ireland, Galway,
Ireland*

Dr. Jeffy George

*Department of Microbiology and Immunology
F. Edward Hébert School of Medicine
Uniformed Services University of the Health Sciences
4301 Jones Bridge Road,
Bethesda, MD 20814
USA.*

Dr. Gyanendra Singh

*Stanley S. Scott Cancer Center,
School of Medicine,
Louisiana State University Health Sciences Center
New Orleans, LA 70112,
USA.*

Dr. Anupreet Kour

*1620 Chevy Chase Dr.
Champaign, IL 61821
USA.*

Dr. Arun Sharma

*Institute for Plant Genomics and Biotechnology
(IPGB)
Borlaug Center,
TAMU 2123
Texas A&M University
College Station, TX 77843
USA.*

Dr. Mohsen Asker

*Microbial Biotechnology Dept.
National Research Centre
Cairo,
Egypt.*

Dr. Elijah Miinda Ateka

*Department of Horticulture,
Jomo Kenyatta University of Agriculture and
Technology (JKUAT)
Kenya.*

Dr. Jozélio Freire De Carvalho

*Faculdade de Medicina Da USP, Reumatologia
Av. Dr. Arnaldo, 455 - 3ª andar – Sala 3133.
São Paulo - SP
Brazil*

Dr. Premendra Dhar Dwivedi

*Food Toxicology Division
Industrial Institute of Toxicology Research,
Post Box No: 80, Mahatma Gandhi Marg,
Lucknow 226001,
India*

Dr. Muhammad Abd El-Moez El-Saadani

*Universities and Research Center District,
New Borg El-Arab,
P.O.Box: 21934 Alexandria,
Egypt.*

Dr. Donald J. Ferguson

*Advanced Orthodontic Training Program,
Nicolas & Asp University College
Dubai,
UAE*

Dr. Kalyan Goswami

*Department of Biochemistry & JB Tropical Disease
Research Centre,
Mahatma Gandhi Institute of Medical Sciences,
Sevagram, Wardha-442102*

Dr. A.K. Handa

*National Research Centre for Agroforestry,
Gwalior Road, JHANSI-284003 UP
India.*

Dr. Amjad M.Husaini

*Metabolic Engineering & Biotechnology Laboratory
Division of Plant Breeding & Genetics
Sher-e-Kashmir University of Agricultural Sciences &
Technology of Kashmir
J&K-191121,
India*

Dr. Vinod Joshi

*Laboratory of Virology & Molecular Biology,
Desert Medicine Research Centre,
Pali Road, Jodhpur-342 005,
India*

Dr. T. Kalaivani

*D/O S. Thiagarajan
B-43, Rajaram Nagar,
Salem - 636 007,
Tamil Nadu, India*

Dr. Priya Kalia

*Orthopaedic Research Unit,
Department of Surgery,
Cambridge University, Cambridge,
UK*

Dr. Patricia Khashayar

*Tehran University of Medical Sciences
Endocrinology and Metabolism Research Center
Shariati Hospital*

Dr. Zaringhalam Moghadam

*Shahid Beheshti Medical University (M.C)
Tehran,
Iran*

Dr. Okeke Ikechukwu Linus

*Department of Surgery, University of Ibadan
Nigeria.*

Dr. Rajesh Kumar Patel

*Centre for Analysis and Learning in Livestock and Food
(CALF)
National Dairy Development Board (NDDB)
Anand- 388 001 (Gujarat)
INDIA*

Dr. Pooja Ralli-Jain

*Department of Pathology and Laboratory Medicine
University of California Irvine,
Irvine, California,
U.S.A.*

Dr. Meltem Sesli

*College of Tobacco Expertise,
Turkish Republic, Celal Bayar University 45210,
Akhisar, Manisa,
Turkey*

Dr. Reda H. Sammour

*Tanta University, Faculty of Science, Tanta,
Egypt*

Dr. Seyed Soheil Saeedi Saravi

*Mazandaran University of Medical sciences, Sari,
Iran*

Dr. R. Senthil Kumar

*St. Matthew's University, School of Medicine
Grand Cayman
Cayman Islands*

Dr. Mohammad Reza Shakibaie

*Kerman University of Medical Sciences, Kerman,
Iran*

Dr. Srividya Shivakumar

*Dept of Microbiology,
CPGS, Jain university,
Bangalore*

Dr. Shashideep Singhal

*The Brooklyn Hospital Center
NewYork-Presbyterian Healthcare System
Brooklyn,
NY.*

Dr. Sripada M. Udupa

*International Center for Agricultural Research in the
Dry Areas (ICARDA),
B.P. 6299, Rabat Instituts, Rabat,
Morocco.*

Dr. Wei Wu

*Institute for Biocomplexity and Informatics
Department of Bio Science
The University of Calgary
Canada*

Dr. Xiao-Bing Zhang

*Molecular Regeneration Laboratory, MC1528B
11234 Anderson Street
Loma Linda, CA 92350*

Prof. Dr. Ozfer Yesilada

*Inonu University
Faculty of Arts and Sciences
Department of Biology
44280 Malatya
Turkey*

Dr. Edson Boasquevisque

*Universidade do Estado do Rio de Janeiro- UERJ
Av 28 de setembro, 87, fundos (LMMC-IBRAG).
Vila Isabel, city: Rio de Janeiro/ RJ
Brasil*

Dr. Abhilash M.

*The Oxford College of Engineering
Hosur Road, Bangalore - 560068*

Dr. Nasar Uddin Ahmed

*Department of Genetics and Plant Breeding
Patuakhali Science and Technology University
Dumki, Patuakhali-8602
Bangladesh*

Dr. Mervat Morsy EL- Gendy

*Chemistry of Natural and Microbial Products
Department,
National Research Center, Dokki, Cairo,
Egypt*

Dr. Gjurmakch Aliev

*Health Science and Healthcare Administration
Program,
University of Atlanta, Atlanta, Georgia,
USA*

Dr. Muhammad Asgher

*Department of Chemistry and Biochemistry,
University of Agriculture,
Faisalabad,
Pakistan*

Dr. Anand Bharatkumar

*Parul Institute of Pharmacy, Limda, Waghodia,
Vadodara*

Dr. Chinmoy Kumar Bose,
*Netaji Subhash Chandra Bose Cancer Research
Institute
16A, Park Lane, Park Street, Kolkata 700 016,
India.*

Dr. Mousumi Debnath
*Jaipur Engineering College and Research Centre
(JECRC) Department of Biotechnology,
Shri Ram ki Nangal, Via Vatika ,Tonk Road , Jaipur-
303905 ,
India*

Dr. Dolan C. Saha
*Dept. of Biochemistry and Molecular Biology,
Faculty of Medicine,
University of Calgary,
Canada*

Dr. Ramasamy Harikrishnan
*Department of Aquatic Biomedical Sciences
School of Marine Biomedical Science
College of Ocean Sciences
Jeju National University
Jeju city, Jeju 690 756,
South Korea*

Dr. Abdul Haque
*Health
Biotechnology division, nibge,
Faisalabad,
Pakistan*

Dr. Kuvalekar Aniket Arun
*Interactive Research School for Health Affairs
(IRHSA),
Bharati Vidyapeeth University, Pune, Maharashtra,
India*

Dr. Asit Ranjan Ghosh
*School of Bio Science & Technology,
Division of Medical Biotechnology,
Vellore Institute of Technology (VIT) University,
Vellore-632014,
India*

Dr. Prasanna Kumar Santhekadur
*Department of Human and Molecular Genetics
Virginia Commonwealth University
Richmond,
VA*

Dr. Majid Sattari
*Rice Research Institute of Iran
Iran*

Dr. Mihael Cristin Ichim
*National Institute Research and Development for
Biological Sciences /
"Stejarul" Research Centre for Biological Sciences
Alexandru cel Bun St., 6, Piatra Neamt, 610004,
Romania*

Dr. Sailas Benjamin
*Enzyme Technology Laboratory
Biotechnology Division
Department of Botany
University of Calicut
Kerala - 673 635
India*

Dr. Sreeramanan Subramaniam
*School of Biological Sciences,
Universiti Sains Malaysia (USM),
Minden Heights, 11800, Penang,
Malaysia*

Dr. Vijai Kumar Gupta,
*Department of Biochemistry, NUI, Galway,
Ireland*

Dr. Vitor Engrácia Valenti
*Universidade Federal de São Paulo
Rua Napoleão de Barros, 715, Térreso
São Paulo, SP
Brazil.*

Dr. Ravindra Pogaku
*School of Engineering and IT
Universiti Malaysia Sabah
88999 Kota Kinabalu
Sabah,
Malaysia*

Dr. Ahmed Eid Abdel-Hamid Eweis Fazary
*School of Pharmacy,
College of Medicine,
National Taiwan University, Taipei 100,
Taiwan.*

Dr. Mohammad Hashemi
*Dept. of Clinical Biochemistry,
School of Medicine,
Zahedan University of Medical Sciences,
Zahedan,
Iran*

Dr. Hesham, Abd El-Latif

*Genetics Department,
Assiut University, Assiut 71516,
Egypt.*

Prof. Jia-ying Xin

*College of Food Engineering
Harbin University of Commerce
138 Tongda Road
Daoli District
Harbin 150076, Heilongjiang
P.R.China*

Dr. Kabir Mohammad Humayun

*Plant Molecular Biotech Lab
Department of Medical Biotechnology
College of Biomedical Science
Kangwon National University
Kangwon-do, Chuncheon, 200-701
South Korea*

Dr. Kalpesh Gaur

*Geetanjali College of Pharmaceutical Studies Manwa
Khera,
Udaipur- 313002. Rajasthan,
India*

Dr. Meganathan, Kannan

*Center for Biologics Evaluation and Research (CBER),
U.S. Food and Drug Administration (FDA),
Bldg. NIH 29A, Room 2C-10,
8800 Rockville Pike,
Bethesda, MD 20892.
USA.*

Assist. Prof. Ali Karadeniz

*Department of Physiology,
Faculty of Veterinary Medicine,
University of Atatürk 25240 ERZURUM
Turkey*

Dr. Matthew Kostek

*Department of Kinesiology
University of Connecticut
Storrs CT*

Dr. Tansu Kucuk

*Gulhane School of Medicine
Department of Obstetrics and Gynecology
Etlik 06018 Ankara,
Turkey*

Dr. Kuo-Sheng Hung

*Department of Neurosurgery
Taipei Medical University - Wan Fang Medical Center
111 Section 3, Hsing-Long Rd,
Taipei 116,
Taiwan*

Dr. V. Manju

*Department of Biochemistry,
Periyar University,
Salem -11.*

Dr. Mbagwu Ferdinand Nkem

*Department of Plant science and Biotechnology,
Faculty of Science,
Imo State University
Nigeria.*

Dr. Anand Pithadia

*Parul Institute of Pharmacy
Vadodara, Gujarat,
India*

Dr. Radhakrishnan Ramaraj

*Department of Internal Medicine
University of Arizona
Tucson 85724
AZ*

Dr. M. Rasool

*School of Bio Sciences and Technology,
VIT University,
Vellore-632104, Tamil Nadu,
India*

Dr. Reda A.I. Abou-Shanab

*Genetic Engineering & Biotechnology Research
Institute (GEBRI)
Mubarak City for Scientific Research and Technology
Applications
New Burg El-Arab City, Universities and Research
Institutes
Zone, P.O. 21934, Alexandria,
Egypt.*

Dr. MR. Pravin Babarao Suruse

*Department of Pharmaceutics
Sharad Pawar College of Pharmacy
Wanadongri, Hingna Road
Nagpur- 441 110. (M. S.)*

Dr. Jan Woraratanadharm

*GenPhar, Inc.,
Mount Pleasant,
SC*

Dr. Serap Yalin

*Mersin University Pharmacy Faculty
Department of Biochemistry, Mersin
Turkey*

Dr. YongYong Shi

*Bio-X Center,
Shanghai Jiao Tong University,
Hao Ran Building, 1954 Hua Shan Road,
Shanghai 200030,
PR China*

Dr. Jyotdeep Kaur

*Department of Biochemistry,
Post Graduate Institute of Medical Education and
Research (PGIMER),
Chandigarh*

Dr. Rajkumar

*Dept. Of Radiation Biosciences,
Institute of Nuclear Medicine and Allied Sciences
Brig. S.K. Mazumdar Road, Timarpur,
Delhi-110054
India*

Dr. Meera Sumanth

*Visveswarapura Institute of Pharmaceutical Sciences,
22nd Main, 24th Cross, B.S.K II stage,
Bangalore-560070
Karnataka,
India.*

Dr, Jai S. Ghosh

*Department of Microbiology,
Shivaji University,
Kolhapur 416004,
India*

Prof. Dr. Alaa H. Al-Charrakh

*Babylon University, College of Medicine.
Dept. of Microbiology
Hilla, Iraq*

Table of Content

**First detection of the BES-type Extended-Spectrum β -Lactamase
produced by Enterobacteria at Saint Camille Hospital
of Ouagadougou (Burkina Faso)**

1

Sougué Serge, Mètuor-Dabiré Amana, Zohoncon Théodora Mahoukèdè,
Tiemtoré Rahimatou Yasmine Wend-kuni, Bangré Yasmine Aminata,
Zongo Jacob Koudbi and Simporé Jacques

Full Length Research Paper

First detection of the BES-type Extended-Spectrum β -Lactamase produced by Enterobacteria at Saint Camille Hospital of Ouagadougou (Burkina Faso)

Sougué Serge^{1*}, Mètuor-Dabiré Amana², Zohoncon Théodora Mahoukèdè³, Tiemtoré Rahimatou Yasmine Wend-kuni¹, Bangré Yasmine Aminata¹, Zongo Jacob Koudbi² and Simporé Jacques¹

¹Department of Biochemistry and Microbiology, Faculty of life and earth Science, University Joseph KI-ZERBO of Ouagadougou, National Public Health Laboratory of Ouagadougou, Burkina Faso.

²Laboratory of Biology and Molecular Genetics (LABIOGENE), Pietra Annigoni Biomolecular Research Center (CERBA), University of Dédougou, Burkina Faso.

³Saint Camille Hospital of Ouagadougou, Saint Thomas D'Aquin University of Ouagadougou, Burkina Faso.

Received 21 May, 2019 ; Accepted 2 August, 2019.

Several studies have been reported on the *bla*_{TEM}, *bla*_{CTX-M} and *bla*_{SHV} genes in Extended-spectrum β -lactamase (ESBL) producing Enterobacteria, however very few studies reported in the literature are related to *bla*_{BES} in ESBL producing Enterobacteria. This study concerns the molecular epidemiology of the *bla*_{BES} gene in Enterobacteria identified from in-patients and out-patients at Saint Camille hospital of Ouagadougou (Burkina Faso). The study was first involved microbiological identification of Enterobacteria that are implicated in antibiotic resistance using API 20 E system; the antibiotics susceptibility test was secondly performed by the diffusion method and the molecular characterization was finally made by PCR to detect the *bla*_{BES} gene. Data were entered and analyzed using Excel 2013 and EPI Info version 6.0 software. A p-value < 0.05 was considered as significant. A total of 60 isolates of ESBL-producing Enterobacteria were found: 21 (35%) *Escherichia coli*; 18 (30%) *Klebsiella pneumoniae*; 6 (10%) *Enterobacter cloacae*; 4 (7%) *Proteus mirabilis*; 4 (7%) *Serratia marcescens*; 3 (5%) *Citrobacter freundii*; 1 (1.6%) *Enterobacter aerogenes*; 1 (1.6%) *Citrobacter brakii*; 1 (1.6%) *Citrobacter youngae* and 1 (1.6%) *Salmonella arizonae*. Molecular characterization revealed the presence of the *bla*_{BES} gene in 38 (63.3%) of bacterial isolates carried by patients. The presence of *bla*_{BES} gene in ESBL producing Enterobacteria at Saint Camille Hospital in Ouagadougou was therefore established in this study for the first time in Burkina faso.

Key words: Enterobacteria, Extended-spectrum β -lactamase (ESBL), *bla*_{BES}, gene, hospital, resistance, Ouagadougou.

INTRODUCTION

Antimicrobial resistance constitutes an increasingly human health hazard worldwide, but the hospital has always been known to be the greatest risk (Bradford, 2001). Thereby, the first antimicrobial resistance surveillance data published by the World Health

Organization (WHO, 2018) showed high levels of resistance to several serious bacterial infections in both high and low income countries. Antimicrobial resistance is responsible for about 700,000 deaths a year worldwide and has huge implications for the cost of healthcare

(Jasovsky et al., 2016). The production of Extended-Spectrum β -lactamases (ESBLs) by Enterobacteria is the main mechanism of the antimicrobial resistance. Several studies have been conducted on the major genes involved in the production of ESBLs. The most common ESBLs are the Temoneira (TEM), Variable sulfhydryl (SHV) and Cefotaximase-Munich (CTX-M) types (Sadeeq et al., 2018).

The first plasmid TEM-1-type β -lactamase was isolated in 1965 in Greece from a strain of *Escherichia coli* isolated in a patient named Temoneira hence the name (Zubair et al., 2015). The SHV-types ESBLs are derived by punctual mutations from the original SHV-1 enzyme. The origin of these enzymes is probably a variant of the *Klebsiella pneumoniae* chromosomal penicillinase *bla_{SHV}* gene (Brisse and Verhoef, Haeggman et al., 2004). Currently, more than 180 SHV-types ESBLs variants have been described (Liakopoulos et al., 2016). CTX-M ESBLs were initially described in 1986 in Japan, Germany and France in 1989 (CTX-M-1) and have since spread widely around the world (Thomson and Moland, 2000). CTX-M is the most prevalent ESBLs worldwide (Anna et al., 2014).

Besides the so-called major ESBLs there are minor types ESBLs such as BES-type (*Brazilian extended spectrum*) which are less studied (Cattoir, 2008).

It has less wide distribution and is characterized by a high level of resistance to Cefotaxime and sometimes to Aztreonam rather than Cefotaxime (Arlet and Philippon 2003; Bradford, 2001). The prevalence of the *bla_{BES}* gene in ESBL-producing Enterobacteria is rare. Investigations on ESBL-producing Enterobacteria in Burkina faso is relatively recent and the *bla_{BES}* gene is not concerned yet.

To fill the gap on the molecular epidemiology of the *bla_{BES}* gene, this study was undertaken to detect the *bla_{BES}* gene in ESBLs-producing Enterobacteria at Saint Camille Hospital of Ouagadougou (Burkina Faso).

MATERIALS AND METHODS

Type of study

This is a cross-sectional study conducted at Saint Camille Hospital of Ouagadougou (Burkina faso) from September to November 2018. Samples collected consisted of stool samples, urine samples and vaginal swab samples from hospitalized patients or out-patients. 250 samples were collected from 250 patients. Samples are inoculated on ordinary media like Uri Select 4 medium Bio-Rad (Marnes-la-Coquette, France), Hektoen Enteric Agar medium (Taden, France) and BD Salmonella Shigella (SS) Agar medium (Heidelberg, Germany) to allow the growth of Enterobacteria and then incubated for 24 h at 37°C. Subsequently, Enterobacteria who grew on the previous media were subcultured on a BD Mueller-Hinton (MH) Agar medium (Heidelberg, Germany) and then

incubated for 24 h at 37°C for antimicrobial assays.

Antimicrobial susceptibility test

Isolates were identified using Analytical Profile Index API 20 E (BioMérieux S. A., Marcy l'Etoile, France) identification method. Antibiotics susceptibility/resistance test was carried out on Mueller-Hinton (MH) medium with pure colonies of Enterobacteria according to the recommendations of the Committee of antibiogram of the French Society of Microbiology (CASFM/EUCAST, 2018). The antibiotic discs used were: Amoxicilline + Clavulanic acid (Augmentin) (30 μ g), Cefotaxime (30 μ g), Cefotaxidime (30 μ g), Ceftriaxone (30 μ g) and Aztreonam (30 μ g). All Enterobacteria resistant to Augmentin and at least one third generation cephalosporin were considered as ESBL-producing Enterobacteria for the study.

Molecular characterization of ESBLs

Extraction of bacterial DNAs

The boiling method was used to extract DNAs from bacteria (Ribeiro Junior et al., 2016). The strains were reactivated by culture on the MH medium for 18 to 24 h. An isolated colony is taken from the Petri dishes and suspended in 200 μ l of distilled water previously aliquoted in labeled eppendorf tubes. Then immersion in a water bath at 100°C for 15 min was followed to release the genetic material. After immersion in a water bath, the suspension was centrifuged at 12,000xg for 10 min and the supernatant containing the DNA released is transferred to a new eppendorf tube. The concentration of the DNA was then determined using the Nanodrop type spectrophotometer (Biodrop®, Holliston, USA).

Molecular analysis

The reaction medium for PCR was constituted by a volume of 25 μ l composed of the Master Mix, the DNA and the primers. PCR amplification of *bla_{BES}* gene was carried out with the following specific primers: BES-F 5' TAATAACCCTGACCAAGCCTA 3' and BES-R 5' CCCTTTCAAAGTCATAATC 3' were used for *bla_{BES}* with 879 bp as molecular weight (Bonnet et al., 2000) and using thermocycler Gene Amp 9700 PCR System (Applied Biosystems, Californie, États - Unis). PCR program consisted of an initial denaturation at 95°C for 5 min followed by 30 cycles (Denaturation 95°C/59 s, Annealing 50°C/59 s, Elongation 68°C/59 s) and a final Extension at 68°C for 5 min.

Electrophoresis on agarose gel

Agarose gel (1%) for electrophoresis was prepared with 1X TBE buffer with addition of Ethidium bromide (BET) which allowed visualization of the bands in the UV light. An electrophoretic migration at 110 millivolts for 30 min was performed on the PCR products using a molecular weight marker (1 kb). The fragments were visualized under UV light (Gene Flash) and the images were recorded.

*Corresponding author. E-mail: sergesougue@yahoo.fr. Tel: 0022670414258.

Frequency of bacterial species




Figure 1. Distribution of enterobacterial isolates by size.

Data processing

The clinical data was entered on Excel 2013 and then analyzed with the Standard Statistical Package for Social Sciences (SPSS) version 17.0 for Windows and the EPI Info version 6.0 software. All tests of significance were considered statistically significant at P-value < 0.05.

RESULTS AND DISCUSSION

In this study, out of 250 strains isolated, 60 (24.09%) of Enterobacteriaceae were resistant to at least one third generation cephalosporin or azthreonam and Amoxicillin + Clavulanic acid. The distribution of strains involved in bacterial resistance by ESBL production is presented by Figure 1.

Results for the sensitivity/resistance of the 60 enterobacterial isolates to the different antibiotics tested showed that 46 strains (76.6%) were resistant to Cefotaxime, 45 strains (75%) were resistant to Ceftriaxone, 42 strains (70%) were resistant to Aztreonam and 35 strains (58.3%) were resistant to Ceftazidime. Resistance profile by bacterial strain were shown in Table 1. It was noticed that all strains were resistant to Amoxicillin + Clavulanic acid (Augmentin®).

Molecular characterization of the ESBLs by PCR revealed that 38 (63.3%) strains carried the *bla*_{BES} gene as shown by the electrophoresis bands (Figure 2). The distribution of the *bla*_{BES} gene according to the bacterial species is presented in Table 2.

These results are comparable to those found in some studies, particularly those found at the Laghouat Hospital (Algeria): 43% *Escherichia coli*, 30% *Klebsiella pneumoniae* and 20% *Enterobacter cloacae* (Lagha, 2015). Other studies that was done at Charles De Gaulle Paediatric Teaching Hospital (CHUP/CDG) of Ouagadougou (Burkina faso), showed 47.22% of *Escherichia coli*; 15.55% *Klebsiella pneumoniae* and 3.33% *Klebsiella oxytoca* (Metuor-Dabire, 2014). These prevalences of ESBLs produced by *Escherichia coli* and *Klebsiella pneumoniae* were also described in South America (45.4 to 51.9%) (Villegas et al., 2008) and Saudi Arabia (55%) (Al-Agamy et al., 2009).

These results confirm that the overall prevalence of ESBL production by Enterobacteria fluctuates considerably according to the geographical zones, to the countries and to different hospitals. However, the bacterial strains mainly concerned by antibiotic resistance were *E. coli* and *K. pneumoniae* with the high level of ESBL production.

The antibiotic susceptibility profile of the 60 strains tested showed resistance to most of β -lactams antibiotics (Table 1): 46 strains (76.6%) were resistant to Cefotaxime, 45 strains (75%) were resistant to Ceftriaxone, 42 strains (70%) were resistant to Aztreonam and 37 strains (58.3%) were resistant to Ceftazidime. These levels of antibiotic resistance in the study could be explained by the misuse of antibiotics in animal and human health or because of the poor living and hygiene conditions observed in developping

Table 1. Resistance profile of different isolates to antibiotics used.

Bacterial strain	ATM	CAZ	CTX	CTR	Synergy image
<i>Escherichia coli</i>	13	12	15	15	4
<i>Klebsiella pneumoniae</i>	12	8	11	12	0
<i>Enterobacter cloacae</i>	6	4	4	4	0
<i>Proteus mirabilis</i>	0	0	2	2	2
<i>Serratia marcescens</i>	3	3	3	3	0
<i>Enterobacter aerogenes</i>	0	1	0	0	0
<i>Citrobacter freundii</i>	1	2	3	1	0
<i>Citrobacter brakii</i>	0	0	0	1	0
<i>Citrobacter youngae</i>	0	0	0	0	1
<i>Salmonella arizonae</i>	1	0	1	0	0
Total	36	30	39	38	7

ATM = Aztreonam, CTX = Cefotaxime, CTR =Ceftriaxone, CAZ = Ceftazidime.




Figure 2. Agarose gel image showing PCR products of *bla*_{BES} gene in identified isolates. Lane (M) = Molecular Weight Marker (DNA Ladder (1kb)); Lane (1-8) = Samples; Lane (9) = Negative control; Lane (3) = Positive to *bla*_{BES} gene.

Table 2. Prevalence of the *bla*_{BES} gene according to bacterial species.

Bacterial species	BES gene
<i>Escherichia coli</i>	13
<i>Klebsiella pneumoniae</i>	10
<i>Enterobacter cloacae</i>	5
<i>Proteus mirabilis</i>	3
<i>Serratia marcescens</i>	2
<i>Enterobacter aerogenes</i>	1
<i>Citrobacter freundii</i>	2
<i>Citrobacter brakii</i>	1
<i>Citrobacter youngae</i>	1
<i>Salmonella arizonae</i>	0
Total	38

countries (Ouédraogo, 2017). It is currently admitted that the use of antibiotics, including third-generation cephalosporins for therapeutic purposes is the most important risk factor in the development of bacterial resistance (Metuor-Dabire, 2014).

Others types of resistance mechanisms could explain these levels of antibiotic resistance like the modification of the membrane permeability, the modification of the antibiotic target, the metabolic pathway change or the efflux phenomena.

The molecular characterization of the 60 bacterial strains by PCR revealed the BES gene in 38 (63.3%). Few studies have evaluated the prevalence of the *bla*_{BES} gene in ESBL-producing Enterobacteria. The *bla*_{BES} gene was first described in *Serratia marcescens* in Brazil in 1996 (Bonnet et al., 2000). This strain had a very high level of resistance to Aztreonam; distinctly high to Cefotaxime than to Ceftazidime. In this study, 21 strains, more than 55% of the strains carried the BES gene were resistant to Aztreonam. Philippon found that BES-1 ESBLs, characterized by a high level of resistance to Ceftazidime and sometimes to Aztreonam rather than to Cefotaxime, have a less wide distribution than the CTX-M group (Philippon and Arlet, 2006). The presence of the BES gene in our study could be explained by the variation of ESBLs frequencies between different geographical areas or by the genetic support of BES gene. If it was plasmids or integrons, genes could be easily transmitted between several bacteria. It could also be explained by the spread of this gene in West Africa and at Saint Camille Hospital in Ouagadougou).

CONFLICT OF INTERESTS

The authors have not declared any conflict of interests.

REFERENCES

- Al-Agamy MH, Shibl AM, Tawfik AF (2009). Prevalence and molecular characterization of extended-spectrum beta-lactamase-producing *Klebsiella pneumoniae* in Riyadh, Saudi Arabia. *Annals of Saudi Medicine* 29(4):253-257.
- Anna O, Soderquist B, Persson K, Molling P (2014). Characterization of CTX-M-producing *Escherichia coli* by repetitive sequence-based PCR and real-time PCR-based replicon typing of CTX-M-15 plasmids 122(11):1136-1143.
- Arlet G, Philippon A (2003). The new β -lactamases at the dawn of the third millennium. *French review of laboratories* 352:41-55.
- Bonnet R, Sampaio JL, Chanal C, Sirot D, De Champs C, Viillard J, Labia R, Sirot J (2000). A novel class A extended-spectrum β -lactamase (BES-1) in *Serratia marcescens* isolated in Brazil. *Antimicrobial Agents Chemotherapy* 44(11):3061-3068.
- Bradford PA (2001). Extended-spectrum β -lactamases in the 21st century: characterization, epidemiology and detection of this important resistance threat. *Clinical Microbiology Reviews* 14(4):933-951.
- Brisse S, Verhoef J (2001). Phylogenetic diversity of *Klebsiella pneumoniae* and *Klebsiella oxytoca* clinical isolates revealed by randomly amplified polymorphic DNA, *gyrA* and *parC* genes sequencing and automated ribotyping. *International Journal of Systematic and Evolutionary Microbiology* 51(3):915-924.
- CASFM/EUCAST (2018). Antibiotic Committee of the French Microbiology Society, Recommendation 2018. <http://www.sfm.asso.fr, 09/09/2018>
- Cattoir V (2008). Les nouvelles β -lactamases à spectre étendu (BLSE). <http://www.mapar.org/article/1/Communication%20MAPAR/qm6segij/Les%20nouvelles%20b%20C3%AAta-lactamases%20C3%A0%20spectre%20C3%A9tendu%2028BLSE%29.pdf>
- Haeggman S, Lofdahl S, Paauw A, Verhoef J, Brisse S (2004). Diversity and evolution of the class A chromosomal beta-lactamase gene in *Klebsiella pneumoniae*. *Antimicrobial Agents Chemotherapy* 48(7):2400-2408.
- Jasovsky D, Littmann J, Zorzet A, Cars O (2016). Antimicrobial resistance a threat to the world's sustainable development. *Upsala Journal of Medical Sciences* 121(3):159-164.
- Lagha N (2015). Antibiotic resistance study of extended spectrum β -lactamase-producing Enterobacteriaceae (ESBL) isolated from Laghouat Hospital: Alger. www.dspace.univ-tlemcen.dz/bitstream/112/7203/1/LAGHA-n.pdf
- Liakopoulos A, Mevius D, Ceccarelli D (2016). A Review of SHVExtended-Spectrum β -Lactamases:Neglected Yet Ubiquitous. *Frontiers in Microbiology* 7:1374.
- Metuor-Dabiré A (2014). Molecular and kinetic characterization of broad-spectrum β -lactamase (ESBL) types of bacterial strains collected at Charles De Gaulle Paediatric Teaching Hospital (CHUP/CDG): Ouagadougou. www.labiogene.org/IMG/pdf/these_metuor_dabire_amana_corrige2.pdf
- Ouédraogo A (2017). Prévalence, circulation et caractérisation des bactéries multirésistantes au Burkina faso: Hal Id: tel-01476152 Spécialité/ Biologie Santé.
- Philippon A, Arlet G (2006). β -Lactamases of Gram négatif bacteria: Never-ending clockwork. *Annals of Clinical Biology* 64(1):37-51.
- Ribeiro Junior JC, Tamanini R, Soares BF, Marangon de Oliveira A, Silva F (2016). Efficiency of boiling and four other methods for genomic DNA extraction of deteriorating spore-forming bacteria from milk. *Agricultural Sciences* 37(5):3069-3078.
- Sadeeq UR, Tariq A, Ijaz A, Nazir AK, Bo H, Jian G (2018). The Growing Genetic and Functional Diversity of Extended Spectrum Beta-Lactamases. *BioMed Research International* 2018:9519718.
- Thomson KS, Moland ES (2000). The new β -lactamases of Gram negative bacteria at the dawn of the new millennium. *Microbes Infection* 2(10):1225-1235.
- Villegas MV, Kattan JN, Quinteros MG, Casellas JM (2008). Prevalence of extended-spectrum β -lactamases in South America. *Clinical Microbiology and Infection* 14:154-158.
- WHO (2018). High levels of antibiotic resistance found worldwide, new data shows. <https://www.who.int/news-room/detail/29-01-2018>
- Zubair M, Malik A, Ahmad J (2015). Microbiology of Diabetic Foot Ulcer with Special Reference to ESBL Infections. *American Journal of Clinical and Experimental Medicine* 3(1):6-23.

Related Journals:

