

Full Length Research Paper

Evaluation of yield performance of *Pleurotus sajor-caju* on different agro-based wastes

Shauket Ahmed Pala*, Abdul Hamid Wani and Riyaz Ahmed Mir

Department of Botany, Section of Mycology and Plant Pathology, University of Kashmir, Hazratbal, Srinagar-190006, India.

Accepted 14 June, 2013

Pleurotus sajor-caju a nutrient rich mushroom was cultivated on four different substrates, viz. Paddy straw, wheat straw, Apple leaf and Chinar leaf substrates. It was observed that its yield or biological efficiency was maximum on paddy straw followed by wheat straw, apple leaves and Chinar leaves. It was also observed that *P. sajor-caju* gave the maximum yield in the first flush followed by second, third and fourth flush except in case of Chinar leaves where the yield obtained in second flush was comparatively higher than first flush.

Key words: *Pleurotus sajor-caju*, cultivation, biological efficiency, substrate.

INTRODUCTION

Food production in large quantity is a challenge but safe disposal of crop residues is a great problem. Edible fungi are natural recycler which converts lignocelluloses wastes into protein rich health food. Malnutrition is a problem in developing third world countries. The Food and Agriculture Organization have recognized mushrooms as food contributing protein nutrition to the countries depending largely on cereals. Mushrooms with their flavour, texture, nutritional value and high productivity per unit area have been identified as an excellent food source (Eswaran and Ramabadrhan, 2000). *Pleurotus sajor-caju* commonly known as Dhingri is an important edible mushrooms gaining popularity in recent years because of its high nutritional value and ability to grow on diverse agricultural wastes. The genus is characterized by its high protein content 30 to 40% on dry weight basis (Sharma and Madan, 1993) which is twice that of vegetable. Dhingri mushroom can help in solving the problems of malnutrition and disease. Poppe (2000) reported that there are about 200 kinds of waste in which edible mushrooms can be produced. Various agricultural

wastes rich in cellulose are being used as substrates for cultivation of Dhingri mushrooms (Thomas et al., 1998). *Pleurotus* species require a temperature of 20 to 30°C both for its vegetative growth and reproductive phase in natural habitat (Chang and Miles, 2004). Since the climate of Kashmir is quite conducive for the growth of *Pleurotus* species, therefore its cultivation can be carried out on large scale easily which in turn can empower the economy of farmers. Also in Kashmir a lot of agricultural residues rich in lignocelluloses are generated every year, which can be used as base material for cultivation of mushrooms *P. sajor-caju*. The present study aimed to examine the biological efficiency or yield of *P. sajor-caju* on different agro-based wastes used for its cultivation.

MATERIALS AND METHODS

The pure mycelial culture was maintained on potato-dextrose agar (pH=7) containing 20% potato extract; 2% dextrose; 2% agar as recommended by earlier workers (Das and Mukherjee, 1996). For preparing the pure spawn wheat grains were used as a nutrient

*Corresponding author. E-mail: sapala29@gmail.com / ahamidwani@yahoo.com.

Table 1. Days for completion of spawn running, pinhead formation and fruiting body formation of *P. sajor-caju* on different substrates.

Substrates	Spawn running	Pinhead formation	Fruiting body formation
	(Days)		
Paddy straw	17-19	21-23	25-27
Wheat straw	22-24	28-30	32-34
Apple leaves	25-28	31-36	42-44
Chinar leaves	30-34	40-44	47-49

Table 2. Yield performance of fresh mushrooms.

Substrates	Yield (g)/500 g dry substrate					Biological efficiency (%)
	first flush	second flush	third flush	fourth flush	Total	
Paddy straw	280.4±4.3	265.7±3.9	120.6±2.7	80.4±2.1	747.1±3.2	149.4
Wheat straw	225.1±2.6	220.4±2.9	113.5±3.1	64.7±2.8	623.7±2.8	124.7
Apple leaves	169.3±3.1	146.5±2.4	101.2±2.7	61.1±2.3	478.1±2.1	95.62
Chinar leaves	115.4±2.7	145.2±3.2	110.4±2.9	55.8±2.9	426.8±2.9	85.3

The values are mean of three replicates ± S.D.

capsule. Wheat grains were mixed with gypsum and chalk powder (to maintain a neutral pH and avoid clumping of grains) in a definite proportion (1 Kg wheat grains, 20 g of gypsum and 5 g of calcium carbonate). The grains were filled in bottles and plugged with non absorbent cotton and subjected to sterilization by autoclaving at 21 lb pressure for half an hour followed by the inoculation with pure mycelia. For initiation of mycelial growth, inoculated bottles were kept in an incubator at 25°C till the mycelium spreads homogeneously. When all the grains in the bottles were run over by mycelium they were used as spawn for mass cultivation of mushrooms on different substrates.

The substrate materials viz. Paddy straw, Wheat straw, Apple leaves and Chinar leaves were cut into 4 to 5 cm pieces followed by overnight (12 h) dip in water. After this the substrate was washed in a formalin solution (36 parts of water and 1 part formaldehyde) followed by another solution containing 1 g carbendazim in 10 L of water. Now the drained substrates was put in polythene bags of 35 × 50 cm size to 3/4th of its capacity, perforated with holes all over the surface to allow free exchange of gases. Inoculation of the bags, that is, spawning was carried out through multilayered spawning.

The inoculated bags were kept in the cropping room in dark at the temperature of 25±2°C till the cottony growth proliferates. When the substrate was completely covered by the white cottony mycelia growth, the polythene bags were removed and white light was switched on. For the initiation and subsequent development of fruiting bodies the temperature and relative humidity inside the cropping room was maintained between 20 to 22°C and 80 to 85% respectively.

Observations on period of spawn run, appearance of pinhead, maturation of fruiting bodies were recorded upto forth flush. Fresh weights of mature fruit bodies were also recorded upto forth flush to calculate the total yield and corresponding biological efficiency. Total yield was calculated as the fresh weight of mushrooms harvested upto forth flush per 500 g of dry substrate used for its cultivation. Biological efficiency (B.E.) was determined by the ratio of total fresh weight (g) of mushrooms obtained from four flushes to dry weight (g) of substrate and expressed as percentage.

$$\text{Biological efficiency} = \frac{\text{Fresh weight(g) of mushrooms harvested}}{\text{Dry weight (g) of substrate}} \times 100$$

RESULTS

The analysis of yield on all the four substrates (Paddy straw, Wheat straw Apple leaves and Chinar leaves) used for the cultivation of *P. sajor-caju* gave more or less significant results. However, there was a quite momentous variation in the time interval needed for completion of spawn running, pinhead formation and fruiting body formation on different substrates. In all the cases the time duration for the formation of fruiting bodies was longer in case of Chinar leaf substrate (47-49 days), followed by Apple leaf substrate (42-44), Wheat straw (32-34 days) and Paddy straw substrate (25-27 days) as shown in Table 1. The trend was same for the spawn running and pinhead formation that is, more time was consumed in case of Chinar leaf substrate followed by apple leaf substrate, wheat straw and paddy straw substrate. While analyzing the total yield of *P. sajor-caju* on the above mentioned substrates the trend was opposite, where the highest yield was found on Paddy straw substrate (747.1 g/500 g dry weight), followed by Wheat straw (623.7/500 g dry weight), Apple leaf (478.1/500 g) and Chinar leaf substrate (426.8/500 g dry weight) as shown in Table 2. The biological efficiency of the mushroom was 149.4, 124.7, 95.62 and 85.3 on Paddy straw, Wheat straw, Apple leaf and Chinar leaf substrate respectively. While examining the yield achieved in each flush it was observed that *P. sajor-caju*



Figure 1. Mature fruiting bodies of *P. sajor-caju* on a) Chinar leaf s, b) Paddy straw, c) Wheat straw and d) Apple leaf substrates.

gave the maximum yield in the first flush followed by second, third and fourth flush except in case of Chinar leaf substrate where the yield obtained in second flush was comparatively higher than first flush (Table 2). It is also quite evident that the Paddy straw proved the most efficient while leaf Chinar leaves proved least efficient substrate both in terms of yield and time consumption (Figure 1).

DISCUSSION

Since *P. sajor-caju* can easily grow on the by-products or lignocellulosic residues and lignin, therefore a large number of agricultural, forests, and agro-industrial by-products can be used for its cultivation. In the present study *P. sajor-caju* was cultivated under *in-vitro* conditions on different substrates, viz. Paddy straw, Wheat straw, Apple leaves and Chinar leaves. The study revealed that highest biological efficiency *P. sajor-caju* on Paddy straw, followed by Wheat straw and Chinar leaf substrates. Zhang et al. (2002) cultivated *P. sajor-caju* on

rice straw and wheat straw and observed 10% higher yield in case of rice straw under the same cultivation conditions. Madan et al. (1986) cultivated *P. sajor-caju* on leaves of *Morus alba* and *Ricinus communis*. Several species of *Pleurotus* are known to be cultivated on different substrates in India (Suman and Sharma, 1990). Ragnathan et al. (1996) cultivated three species of *Pleurotus*, viz. *P. sajor-caju*, *P. platypus* and *P. citrinopileatus*, on various agro-residues such as Paddy straw, maize stover, sugarcane bagasse, coir pith and a mixture of these wastes, where the maximum yield was obtained from *P. sajor-caju* cultivated on Paddy straw. According to Chang and Miles (2004)) nutrient content of substrates affects the growth and formation of fruit bodies of *Pleurotus* species. Banik and Nandi (2004) observed that yield of *P. sajor-caju* can be increased significantly when grown on a lignocellulosic crop residue-rice straw supplemented with biogas residual slurry manure in 1:1 ratio as substrate. Khan et al. (2011) studied the impact of various sterilization methods using different substrates on the yield of *Pleurotus* sps. In Kashmir there is a lot of potential for growing mushrooms from Paddy straw as

Paddy cultivation takes place on 140970 ha of land generating 245993 tons of Paddy straw annually (Anonymous, 1999-2000). Only 5706 quintals of mushrooms were produced during 2001-2002 in Kashmir (Munshi and Ghani, 2003). Large scale cultivation of *P. sajor-caju* using suitable substrates can help people in rural areas improve their income. Mushroom cultivation is not just an agribusiness but also a noteworthy means for restoration, replenishment and remediation of earths overburden ecosphere, thereby benefiting all the inhabitants of the planet earth.

ACKNOWLEDGEMENT

The authors are highly thankful to Dr. Nazir Ahmed Munshi, Associate professor SKUAST-K for their guidance during the course of work.

REFERENCES

- Anonymous (1999-2000). Economic Survey, Directorate of Economics and Statistics, planning and Development Department, Government of Jammu and Kashmir, P. 147.
- Banik S, Nandi R (2004). Effect of supplementation of rice straw with biogas residual slurry manure on the yield, protein and mineral contents of Oyster mushroom. *Ind. Crops Prod.* 20(3):311-319.
- Chang ST, Miles PG (2004). *Mushrooms: Cultivation, Nutritional Value, Medicinal Effect, and Environmental Impact* (11nd edition), CRC Press, New York.
- Das N, Mukerje M (1996). Preparation and regeneration of mycelial protoplasts of *Pleurotus florida* and *Pleurotus ostreatus*. *Folia Microbiol.* 41:208-210.
- Eswaran A, Ramabadrhan R (2000). Studies on some physiological, cultural and post harvest aspects of Oyster mushroom, *Pleurotus ostreatus*. *Trop. Agric. Res.* 12:360-374.
- Khan NA, Abbas A, Rehman A, Imran ul H, Hanan A (2011). Impact of various sterilization methods using different substrates for yield improvement of *Pleurotus* spp. *Pak. J. Phytopathol.* 23(1):20-23.
- Madan P, Vasudevan P, Sharma S (1986). Cultivation of *P. sajor-caju* on different wastes. *Biol. Wastes* 22(4):241-250.
- Munshi NA, Ghani MY (2003). Mushroom industry in Kashmir valley-present status future prospects and problems. *SKUAST J. Res.* 5:1-19.
- Poppe J (2000). Use of the agricultural waste materials in the cultivation of mushrooms. *Mushroom Sci.* 15:3-23.
- Ragunathan R, Gurusamy R, Palaniswamy M, Swaminathan K (1996). Cultivation of *Pleurotus* spp. on various agro-residues. *Food Chem.* 55(2):139-144.
- Sharma S, Madan M (1993). Microbial protein from leguminous and non-leguminous substrates. *Acta Biotechnol.* 13:131-139.
- Suman BC, Sharma S (1990). Compact bag method. A new method of increasing the yield of *P. sajor-caju*. *Ind. J. Mushrooms* 17:11-13.
- Thomas GV, Prabhu SR, Reeny MZ, Bopaiah BM (1998). Evaluation of lignocellulosic biomass from cocoonut palm as substrate for cultivation of *Pleurotus sajor-caju*. *World J. Microbiol. Biotechnol.* 14:879-882.
- Zhang R, Li X, Fadel JG (2002). Oyster mushroom cultivation with rice and wheat straw. *Bioresour. Technol.* 82(3):277-284.