

Full Length Research Paper

Impact of allelopathy of sunflower (*Helianthus annuus* L.) roots extract on physiology of wheat (*Triticum aestivum* L.)

Javed Kamal

Department of Plant Sciences, Faculty of Biological Sciences, Quaid-I-Azam University, Islamabad, Pakistan. E-mail: javed1743@yahoo.com. Tel: +923046886850.

Accepted 2 August, 2011

Petri-dishes grown seedlings of wheat cvs. Margalla 99 and Chakwall 97 were treated with distilled water and with different extracts of sunflower roots. Data on physiology of wheat seedlings including germination rate, protein, proline, sugars, DNA, peroxidase, superoxide dismutase and chlorophyll contents were recorded; all increased with allelochemicals treatments when compared with the controls.

Key words: Allelopathy, chlorophyll contents, DNA, peroxidase, protein, proline, sunflower, sugar, superoxide dismutase, wheat.

INTRODUCTION

There is increasing emphasis on sustainable agriculture and concerns about the adverse effects of extensive use of synthetic chemicals, such as contamination of the environment, greater plant resistance to herbicides and high costs. Consequently, research attention is now focused on reducing the dependence upon synthetic herbicides and finding alternative strategies for weed management. Allelopathy is one promising strategy, which can be put to good use in several ways in agroecosystems. Sunflower (*Helianthus annuus* L.) in general and cv. Hysun 38 in particular is increasingly recognized as an important crop in several areas of Islamabad, (Pakistan) given the suitability of the crop to local agroclimatic conditions, its importance as source of edible oil and protein, resistance to drought and its short duration, which makes it a suitable crop if sowing is delayed. However, yields of some crops following sunflower are lower than normal, possibly because of inadequate nutrition and chemical inhibition. Sunflower is often grown when rainfall is marginal and depletion of soil moisture by sunflower may be a factor, although, this remains unproven. Both sunflower and the crops that follow it receive routinely specified amounts of fertilizers and there is no evidence that nutrient deficiency is the cause of lower yields. Sunflower is known to actively influence the growth of surrounding plants because of its high allelo-

pathic potential and > 200 natural allelopathic compounds have been isolated from sunflower cultivars. Most of these known allelochemicals affect seed germination (kamal and Bano, 2008). Wheat (*Triticum aestivum* L.) is the staple food of Pakistan. Various factors lower the productivity of wheat, such as delayed sowing, inadequate doses of fertilizers, water shortage, non-availability of improved seed, diseases and drought and now weed infestation has emerged as a serious problem. Weeds compete with the crop for nutrients, water, space, light and carbon dioxide and also interfere with its normal growth by secreting biomolecules into the rhizosphere. Thus, the objective of the present study was to assess the allelopathic effects of sunflower root extracts on germination, contents of chlorophyll, protein, proline, deoxyribonucleic acid, sugar, superoxidase dismutase (SOD) and peroxidase (POD) in wheat.

MATERIALS AND METHODS

Sunflower plants (cv. Hysun 38) were grown in pots in the Department of plant sciences, Quaid-i-Azam University, Islamabad. 3 seeds were sown in each pot and were given a basal fertilizer dose of 2 g diammonium phosphate, 1 g urea, and 1 g potash. When the plants reached the vegetative stage (40 d after sowing), they were uprooted and separated into leaves, stems and roots. They were then washed thoroughly with distilled water, dried,

pulverized in a mill and stored in a cool place along with anhydrous CaCl_2 to maintain dryness. Allelopathic extracts of sunflower at different concentrations were prepared as described by Bogatek et al. (2005). The solution was centrifuged at low speed (3000 rpm) for 15 min and the supernatant filtered through one layer of Whatman No. 42 filter paper. The extracts were stored at $< 5^\circ\text{C}$ until use. The extracts were determined to be free of fungal contamination. Seeds of 2 wheat varieties cvs. Margalla 99 and Chakwall 97 were sown in pots (3 seeds per pot) and a basal fertilizer dose of 1 g urea and 1 g diammonium phosphate applied to soil in each pot at the time of sowing through either plain water (control) or water mixed with extracts of sunflower leaves, stems or roots (1 g of extract mixed with 9 ml of water). Contents of chlorophyll, protein, proline, sugar, SOD and POD were determined. The experiment had 3 replications.

Plant material and growth conditions

Allelopathic extracts of sunflower at different concentrations were prepared as described by Bogatek et al. (2005). The solution was centrifuged at low speed (3000 rpm) for 15 min and the supernatant filtered through one layer of Whatman No. 42 filter paper. The extracts were stored below 5°C until use. It was ascertained that the extracts were free of fungal contamination. Seeds of two wheat varieties, namely Margalla 99 and Chakwall 97 were germinated in pots. The experiment consisted of 3 replications.

Leaf protein contents

Protein contents of leaves were determined following the methods of Lowry et al. (1951) using BSA as the standard. Phosphate buffer (Stock solution), Monobasic sodium phosphate (27.6 g) was dissolved in 1000 ml of distilled water.

Chlorophyll content of leaves

The fresh leaves of wheat were collected at 50% flowering for extraction of chlorophyll. The chlorophyll estimation of leaves followed the method of Arnon (1949) and Kirch (1968). The crude preparation (1 ml) was mixed with 4 ml of 80% (v/v) acetone and allowed to stand in the dark at room temperature for 10 min. It was centrifuged at 2000 rpm for 5 min to clear the suspension. The supernatant, which contained soluble pigment, was used to determine chlorophyll. Absorbance of the solution was read at 645 nm for chlorophyll a and at 663 nm for chlorophyll b on a spectrophotometer against 80% (v/v) acetone blank. Total chlorophyll was determined with the equation of Arnon (1949):

$$\text{Total chlorophyll (mg/L)} = (20.2 \times a \text{ 645.A645}) + (8.02 \times b \text{ 663.A663})$$

Sugar content of leaves

Sugar content of wheat leaves at flowering stage was estimated by the method of Dubo et al. (1956) as modified by Johnson et al. (1966). Fresh plant material (0.5 g) was homogenized with 10 ml of distilled water in a clean mortar. It was centrifuged at 3000 rpm for 5 min. Then to 0.1 ml of supernatant, 1 ml of 5% (v/v) phenol was added. After 1 h incubation at room temperature, 5 ml of concentrated H_2SO_4 was added. The absorbance of each sample was recorded at 420 nm. The concentration of unknown samples was calculated with reference to a standard curve made using glucose.

Proline content of leaves

Proline content of leaves was estimated at flowering stage by using the following method (Bates et al., 1973). Fresh plant material (0.1 g) was homogenized with 5 ml of 3.0% sulfosalicylic acid in a mortar. Samples were centrifuged at room temperature at 3000 rpm for 5 min. Supernatant was adjusted to 5 ml with distilled water, to which 5 ml of glacial acetic acid and 5 ml of ninhydrin (1.25 g of ninhydrin dissolved in a solvent prepared by mixing 30 ml of glacial acetic acid, 8 ml of orthophosphoric acid and 12 ml distilled water). The reaction mixture was shaken and the contents placed in tubes and heated in a boiling water bath for 1 h. Then tubes were cooled and the mixture extracted with 10 ml of toluene in a separating funnel. The absorbance of the toluene layer was recorded at 520 nm. A standard curve was constructed from a dilution series of 10 to 100 g of proline at increments of 10 g. The concentration of the unknown sample was calculated with reference to the standard curve.

Peroxidase of leaves

To 3 ml of assay buffer (20 mM mixed phosphate buffer, Ph 7.0 to 7.5 at 30°C) was added 100 μl of 100 mM guaiacol and 300 μl of partially purified enzyme preparation. The mixture was incubated for 1 min at 30°C and the reaction initiated by addition of 1 μmol H_2O_2 (30% v/v). Activity was measured colorimetrically at 436 nm and calculated using an extinction coefficient of 6.39 mM/cm for the guaiacol dehydrogenation product (Putter, 1974).

SOD of leaves

SOD (EC 1. 15. 1.1) activity was assayed by monitoring the inhibition of photochemical reduction of nitroblue tetrazolium (NBT) according to the method of Beauchamp and Fridovich (1971) taking into account the notes by Beyer and Fridovich (1987). Leaf samples were homogenized in 4 volumes (w/v) of an ice-cold buffer containing 0.1 M Tris-HCL (pH 7.8), 0.1 mM EDTA and 0.05% Triton X- 100. The homogenates were filtered through 4 layers of cheesecloth and centrifuged at 4°C for 30 min at 15,000 rpm. The crude extracts were dialyzed for 24 h against half-strength extraction buffer without Triton X-100, centrifuged at 4°C for 30 min at 15,000 rpm and the supernatants were used for SOD assay. The reaction mixture contained 50 mM phosphate buffer (pH 7.8), 0.053 Mm NBT, 10 mM methionine, 0.0053 mM riboflavin and an appropriate aliquot of enzyme extract. The reaction was started by switching on the light and allowed to run for 7 min. One unit of SOD activity was defined as the amount of enzyme required to cause 50% inhibition of NBT reduction as monitored at 560 nm.

DNA analysis

Total DNA was extracted according to Gumin and estimated using an UV spectrophotometer (Spectrophotometer 601) as adopted by Ogur and Rosen genomic DNA was extracted from 60 mg of frozen tissue (young leaves ground in liquid nitrogen) using Genomicprep cells and tissue DNA Isolation Kit (Amersham Pharmacia Biotech Inc.) according to the instruction manual. The DNA concentration in samples was measured by UV-absorption spectroscopy at 260 nm at which DNA gives a maximum absorption at a concentration of approximately 50 $\mu\text{g/g}$ of double-stranded DNA. DNA was treated with 4 restriction enzymes (Bam HI, EcoRI, HinfI and SmaI at 5 U per reaction) according to manufacturer's instructions. The reaction mixture was incubated for 1 h at 37°C except SmaI (25°C). The restriction fragments were size-fractionated using 2% agarose gel

Table 1. Effect of sunflower root extract on germination (%) of wheat varieties Margalla 99 and Chakwall 97.

Treatment	1st year		2nd year	
	V ₁	V ₂	V ₁	V ₂
T ₁	13.67 ^A	12.00 ^A	14.00 ^A	12.33 ^A
T ₂	10.33 ^C	9.00 ^C	9.333 ^C	6.667 ^C
T ₃	11.67 ^{BC}	9.667 ^C	10.67 ^{BC}	7.667 ^C
T ₄	12.00 ^{ABC}	10.67 ^B	11.00 ^B	9.333 ^B
T ₅	12.67 ^{AB}	11.33 ^A	11.67 ^B	9.667 ^{AB}

Values followed by the same letter within a column are not significantly different.

Table 2. Effect of root extract of sunflower on root length (cm) of seedlings of wheat varieties Margalla 99 Chakwall 97.

Treatment	1st year		2nd year	
	V ₁	V ₂	V ₁	V ₂
T ₁	13.00 ^A	17.50 ^A	12.57 ^A	17.00 ^A
T ₂	9.633 ^C	13.20 ^B	8.283 ^C	11.88 ^C
T ₃	10.50 ^{BC}	13.67 ^B	9.950 ^B	13.68 ^B
T ₄	10.90 ^B	14.20 ^B	11.65 ^A	12.01 ^C
T ₅	12.40 ^A	16.96 ^A	12.30 ^A	13.98 ^B

Values followed by the same letter within a column are not significantly different.

Table 3. Effect of root extract of sunflower on shoot length (cm) of seedlings of wheat varieties Margalla 99 and Chakwall 97.

Treatment	1st year		2nd year	
	V ₁	V ₂	V ₁	V ₂
T ₁	18.30 ^A	17.50 ^A	18.43 ^A	17.00 ^A
T ₂	11.86 ^D	12.50 ^C	11.64 ^D	12.92 ^B
T ₃	12.66 ^{CD}	14.20 ^B	13.03 ^{CD}	13.17 ^B
T ₄	13.19 ^C	12.13 ^C	13.40 ^C	13.70 ^B
T ₅	14.63 ^B	14.70 ^B	15.35 ^B	16.32 ^A

Values followed by the same letter within a column are not significantly different. T₁ = Control (distilled water; DW); T₂ = undiluted extract (9 g extract + 10 ml DW); T₃ = 75% extract + 25% DW (7.5 g + 10 ml); T₄ = 50% extract + 50% DW (5 g + 10 ml); T₅ = 25% DW (2.5 g + 10 ml).

electrophoresis. The agarose gels were stained with ethidium bromide and photographed under UV light. The results were documented by Image analyzer gel Doc 2000 (Bio Rad).

Statistical analysis

The data were analysed statistically using MStatC. A completely randomized design was followed.

RESULTS

Effect of sunflower root extract on germination (%) of wheat varieties Margalla 99 and Chakwall 97

The data regarding germination rate under the effect of sunflower roots extract are presented in Table 1. A

perusal of the data revealed that in case of both wheat varieties, Margalla 99 and Chakwall 97, the T₁ (control) showed maximum germination, followed by T₅ (25% extract), whereas minimum germination was shown by T₂ (undiluted extract). Similar pattern was followed in the 2nd year).

Effect of root extract on root length and shoot length (cm) of seedlings of wheat varieties Margalla 99 and Chakwall 97

Root length and shoot length indicates the productive efficiency of a crop. The higher the root and shoot length, the greater the efficiency and vice versa. The data given in Tables 2 and 3 revealed that in case of both wheat

Table 4. Effect of root extract of sunflower on fresh weight (g) of seedlings of wheat varieties Margalla 99 and Chakwall 97.

Treatment	1st Year		2nd year	
	V1	V2	V1	V2
T ¹	0.8900 ^A	0.8900 ^A	0.9100 ^A	0.8300 ^A
T ²	0.2333 ^E	0.7500 ^B	0.1900 ^E	0.7100 ^B
T ³	0.3200 ^D	0.8300 ^A	0.2900 ^D	0.7900 ^A
T ⁴	0.6333 ^C	0.8667 ^A	0.6000 ^C	0.8233 ^A
T ₅	0.8300 ^B	0.8733 ^A	0.7500 ^B	0.8400 ^A

Values followed by the same letter within a column are not significantly different.

Table 5. Effect of root extract of sunflower on dry weight (g) of seedlings of wheat varieties Margalla 99 Chakwall 97.

Treatment	1st year		2nd year	
	V ₁	V ₂	V ₁	V ₂
T ₁	0.8600 ^A	0.8600 ^A	0.8933 ^A	0.8333 ^A
T ₂	0.1833 ^E	0.5433 ^C	0.1633 ^E	0.5167 ^D
T ₃	0.2833 ^D	0.6300 ^B	0.2633 ^D	0.6133 ^C
T ₄	0.5833 ^C	0.6700 ^B	0.5567 ^C	0.6467 ^{BC}
T ₅	0.7833 ^B	0.6833 ^B	0.7533 ^B	0.6700 ^B

Values followed by the same letter within a column are not significantly different. T₁ = Control (distilled water; DW); T₂ = undiluted extract (9 g extract + 10 ml DW); T₃ = 75% extract + 25% DW (7.5 g + 10 ml); T₄ = 50% extract + 50% DW (5 g + 10ml); T₅ = 25% DW (2.5 g + 10 ml).

Table 6. Effect of root extract of sunflower on GA ($\mu\text{g g}^{-1}$) content of leaves in wheat varieties Margalla 99 and Chakwall 97.

Treatment	1st year		2nd year	
	V ₁	V ₂	V ₁	V ₂
T ₁	500.0 ^A	490.00 ^A	503.0 ^A	492.00 ^A
T ₂	404.00 ^D	351.00 ^C	402.00 ^D	348.7 ^C
T ₃	418.7 ^C	388.3 ^{BC}	416.7 ^C	386.0 ^{BC}
T ₄	527.0 ^C	394.3 ^{BC}	424.7 ^C	391.7 ^{BC}
T ₅	480.0 ^B	437.0 ^{AB}	477.0 ^B	434.0 ^{AB}

Values followed by the same letter within a column are not significantly different. T₁ = Control (distilled water; DW); T₂ = undiluted extract (9 g extract + 10 ml DW); T₃ = 75% extract + 25% DW (7.5 g + 10 ml); T₄ = 50% extract + 50% DW (5 g + 10 ml); T₅ = 25% DW (2.5 g + 10 ml).

varieties Margalla 99 and Chakwall 97, T₁ (control) showed the maximum root and shoot length, followed by T₅ (25% extract) < T₄ (50% extract) < T₃ (75% extract), and T₂ (undiluted extract), in the second year, the results were the same.

Effect of sunflower stem extract on fresh weight and dry weight (g) of seedlings of wheat varieties Margalla 99 and Chakwall 97

The data in Tables 4 and 5 revealed that in the case of both first and second year, wheat variety Margalla 99 showed better response. In the two wheat varieties, T₁

(control) showed maximum response, followed by T₅ (25% extract), whereas T₂ (undiluted extract) showed minimum response. In the case of Margalla 99, T₄ was at par with T₃ (75% extract), while in the case of Chakwall 97, T₄ (50% extract) was at par with T₂ (undiluted extract).

Effect of root extract of sunflower on gibberellic acid ($\mu\text{g g}^{-1}$) contents of leaves and roots in wheat varieties Margalla 99 and Chakwall 97

The data on the GA content in leaves are given in Tables 6 and 7. The maximum GA contents in case of wheat

Table 7. Effect of root extract of sunflower on GA ($\mu\text{g g}^{-1}$) content of roots in wheat varieties Margalla 99 and Chakwall 97.

Treatment	1st year		2nd year	
	V ₁	V ₂	V ₁	V ₂
T ₁	505.3 ^A	422.3 ^A	407.0 ^A	419.7 ^A
T ₂	392.3 ^B	318.0 ^C	301.0 ^E	313.0 ^E
T ₃	399.0 ^B	360.0 ^B	349.7 ^D	349.0 ^D
T ₄	411.7 ^B	364.7 ^B	364.3 ^C	362.0 ^C
T ₅	482.3 ^A	417.0 ^A	417.0 ^A	412.3 ^B

Values followed by the same letter within a column are not significantly different.

Table 8. Effect of root extract of sunflower on IAA ($\mu\text{g g}^{-1}$) content of leaves in wheat varieties Margalla 99 and Chakwall 97.

Treatment	1st year		2nd year	
	V1	V2	V1	V2
T ₁	172.7 ^A	131.0 ^A	190.0 ^A	132.0 ^A
T ₂	100.0 ^E	102.0 ^E	113.0 ^E	96.00 ^D
T ₃	127.0 ^D	107.0 ^D	127.0 ^D	106.0 ^C
T ₄	139.3 ^C	114.0 ^E	139.0 ^C	107.0 ^C
T ₅	160.0 ^B	122.0 ^B	149.0 ^B	121.0 ^B

Values followed by the same letter within a column are not significantly different. T₁ = Control (distilled water; DW); T₂ = Undiluted extract (9 g extract + 10 ml DW); T₃ = 75% extract + 25% DW (7.5 g + 10 ml); T₄ = 50% extract + 50% DW (5 g + 10 ml); T₅ = 25% DW (2.5 g + 10 ml).

Table 9. Effect of root extract of sunflower on IAA ($\mu\text{g g}^{-1}$) content of roots in wheat varieties Margalla 99 and Chakwall 97.

Treatment	1st year		2nd year	
	V1	V2	V1	V2
T ₁	193.0 ^A	164.0 ^A	164.0 ^A	134.0 ^A
T ₂	117.0 ^D	94.00 ^E	96.00 ^E	93.30 ^D
T ₃	129.0 ^C	120.0 ^D	125.7 ^D	109.0 ^C
T ₄	135.0 ^C	139.0 ^C	140.3 ^C	113.0 ^C
T ₅	155.7 ^B	153.0 ^B	155.0 ^B	123.0 ^B

Values followed by the same letter within a column are not significantly different.

variety Margalla 99 were given by T₅ (25% extract) which was at par with T₁ (control), the minimum amount of GA was found in T₂ (undiluted extract), while in the second year T₁ (control) showed the maximum amount of GA and T₂ (undiluted extract) showed the minimum amount of GA. In the wheat variety Chakwall 97, in both years, T₁ (control) showed the maximum and T₂ (undiluted extract) showed the minimum quantity of GA in wheat leaves.

Effect of root extract of sunflower on indole-3-acetic acid content of leaves and roots in wheat varieties Margalla 99 and Chakwall 97

The data regarding IAA contents in leaves are presented in Tables 8 and 9. A perusal of the tables revealed that in

the case of both wheat varieties Margalla 99 and Chakwall 97, the maximum IAA contents in leaves samples were calculated in T₁ (control) and the least amount were found in T₂ in the first year as well as second year.

Effect of root extract of sunflower on abscisic acid ($\mu\text{g g}^{-1}$) content of leaves and root in wheat varieties Margalla 99 and Chakwall 97

The data regarding ABA contents in leaves presented in Tables 10 and 11 revealed that in both the wheat varieties Margalla 99 and Chakwall 97, maximum amount of ABA was recorded in T₂ (undiluted extract) and the minimum in T₁ (control). Similar result was found in the

Table 10. Effect of root extract of sunflower on ABA ($\mu\text{g g}^{-1}$) contents of leaves in wheat varieties Margalla 99 and Chakwall 97.

Treatment	1st year		2nd year	
	V1	V2	V1	V2
T ₁	68.00 ^E	90.30 ^E	66.33 ^E	91.61 ^E
T ₂	177.3 ^A	177.3 ^A	180.3 ^A	180.0 ^A
T ₃	156.0 ^B	159.0 ^B	159.0 ^B	162.3 ^B
T ₄	133.7 ^C	132.7 ^C	136.3 ^C	135.0 ^C
T ₅	98.00 ^D	102.0 ^D	100.7 ^D	104.0 ^D

Values followed by the same letter within a column are not significantly different. T₁ = Control (distilled water; DW); T₂ = undiluted extract (9 g extract + 10 ml DW); T₃ = 75% extract + 25% DW (7.5 g + 10 ml); T₄ = 50% extract + 50% DW (5 g + 10 ml); T₅ = 25% DW (2.5 g + 10 ml).

Table 11. Effect of root extract of sunflower on ABA ($\mu\text{g g}^{-1}$) content of roots in wheat varieties Margalla 99 and Chakwall 97.

Treatment	1 st year		2 nd year	
	V ₁	V ₂	V ₁	V ₂
T ₁	100.0 ^C	81.33 ^D	98.67 ^C	82.33 ^D
T ₂	151.0 ^A	113.7 ^A	153.0 ^A	116.3 ^A
T ₃	130.0 ^B	106.0 ^B	132.0 ^B	108.0 ^B
T ₄	105.7 ^C	96.67 ^C	107.7 ^C	98.67 ^C
T ₅	100.3 ^C	84.00 ^D	102.7 ^C	86.00 ^D

Values followed by the same letter within a column are not significantly different.

Table 12. Effect of root extract of sunflower on DNA (mg /100 g F. wt) contents of leaves in wheat varieties Margalla 99 and Chakwall 97.

Treatment	1st year		2nd year	
	V ₁	V ₂	V ₁	V ₂
T ₁	512.9 ^A	485.3 ^A	515.0 ^A	487.0 ^A
T ₂	150.4 ^D	101.3 ^E	148.7 ^D	99.33 ^E
T ₃	239.0 ^C	206.7 ^D	237.7 ^D	204.7 ^D
T ₄	272.3 ^C	269.3 ^C	269.0 ^C	266.7 ^C
T ₅	398.3 ^B	396.7 ^B	395.3 ^B	393.7 ^B

Values followed by the same letter within a column are not significantly different. T₁ = Control (distilled water; DW); T₂ = undiluted extract (9 g extract + 10 ml DW); T₃ = 75% extract + 25% DW (7.5g + 10ml); T₄ = 50% extract + 50% DW (5 g + 10 ml); T₅ = 25% DW (2.5 g + 10 ml).

second year.

Effect of root extracts of sunflower on deoxyribonucleic acid content of leaves in wheat varieties Margalla 99 and Chakwall 97

The data regarding DNA is presented in Table 12. The data pertaining to the DNA contents in leaves of wheat varieties were also affected significantly by leaf, stem and root extract of sunflower. In both wheat varieties Margalla 99 and Chakwall 97, the maximum DNA was observed in T₁ (control) and the minimum in T₂ (undiluted extract) in

the first as well as second year of experiment under the effect of leaves, stem and root extract of sunflower.

Effect of root extracts of sunflower on chlorophyll content (mg/g) of leaves in wheat varieties Margalla 99 and Chakwall 97

The data regarding effect of leaf stem and root extract on chlorophyll contents are presented in Table 13. From the table, it is revealed that the maximum chlorophyll contents under the effect of leaf, stem and root extract of sunflower were recorded in T₁ (control) and the minimum

Table 13. Effect of root extract of sunflower on chlorophyll (mg/ 100 g F. wt) content of leaves in wheat varieties Margalla 99 and Chakwall 97.

Treatment	1st year		2nd year	
	V1	V2	V1	V2
T ₁	109.7 ^A	107.0 ^A	111.9 ^A	108.0 ^A
T ₂	87.34 ^D	84.00 ^D	86.66 ^D	83.00 ^D
T ₃	94.34 ^{CD}	93.00 ^C	93.00 ^{AB}	92.00 ^C
T ₄	100.00 ^{BC}	98.67 ^{BC}	99.00 ^{AB}	97.67 ^{BC}
T ₅	105.3 ^{AB}	103.3 ^{AB}	104.3 ^A	102.3 ^{AB}

Values followed by the same letter within a column are not significantly different.

Table 14. Effect of root extract of sunflower on proline (mg/ 100 g F. wt) contents of leaves in wheat varieties Margalla 99 and Chakwall 97.

Treatment	1st year		2nd year	
	V1	V2	V1	V2
T ₁	82.70 ^E	65.06 ^D	81.00 ^E	64.63 ^D
T ₂	130.0 ^A	119.7 ^A	131.3 ^A	121.0 ^A
T ₃	121.0 ^B	112.4 ^A	123.0 ^B	113.4 ^A
T ₄	114.7 ^C	101.3 ^B	116.7 ^C	102.3 ^B
T ₅	96.00 ^D	91.67 ^C	97.00 ^D	92.67 ^C

Values followed by the same letter within a column are not significantly different. T₁ = Control (distilled water; DW); T₂ = undiluted extract (9 g extract + 10 ml DW); T₃ = 75% extract + 25% DW (7.5 g + 10 ml); T₄ = 50% extract + 50% DW (5 g + 10 ml); T₅ = 25% DW (2.5 g + 10 ml).

Table 15. Effect of root extract of sunflower on sugar (mg/ 100 g F. wt) content of leaves in wheat varieties Margalla 99 and Chakwall 97.

Treatment	1st year		2nd year	
	V1	V2	V1	V2
T ₁	175.0 ^C	180.0 ^{CD}	174.0 ^C	169.0 ^C
T ₂	235.0 ^A	226.0 ^A	222.0 ^A	212.3 ^A
T ₃	215.0 ^B	210.0 ^B	202.7 ^B	197.0 ^B
T ₄	190.0 ^C	184.3 ^C	178.0 ^C	175.3 ^C
T ₅	182.0 ^E	177.7 ^D	176.0 ^C	172.0 ^C

Values followed by the same letter within a column are not significantly different.

chlorophyll content was found in T₂ (undiluted extract) in both wheat varieties Margalla 99 and Chakwall 97. In the second year, results were found similar.

Effect of root extract of sunflower on proline content (mg/g) of leaves in wheat varieties Margalla 99 and Chakwall 97 Data presented in Table 14 revealed that proline contents of leaves under the effect of leaf, stem and root extract were affected significantly by sunflower extract.

A perusal of the tables revealed that in both the wheat varieties, T₂ (undiluted extract) showed the maximum proline contents under the effect of leaf, stem and root extract and T₁ (control) showed the minimum proline

contents. Results were found similar in the second year of experiment.

Effect of root extract of sunflower on sugar content (mg/g) of leaves of wheat varieties Margalla 99 and Chakwall 97

Effect of root extract of sunflower on sugar content of leaves in wheat varieties is given in Table 15. A perusal of the table showed that in the case of wheat variety Margalla 99, T₂ (undiluted extract) showed the maximum while T₅ (25% extract) showed the minimum amount of

Table 16. Effect of root extract of sunflower on protein (mg/ 100 g F. wt) activity of leaves in wheat varieties Margalla 99 and Chakwall 97.

Treatment	1st year		2nd year	
	V ₁	V ₂	V ₁	V ₂
T ₁	1685 ^D	1766 ^B	1753 ^D	1675 ^C
T ₂	1723 ^A	1785 ^A	1783 ^A	1725 ^A
T ₃	1720 ^A	1775 ^{AB}	1775 ^B	1760 ^B
T ₄	1712 ^B	1772 ^B	1768 ^B	1755 ^A
T ₅	1695 ^C	1765 ^B	1761 ^A	1692 ^C

Values followed by the same letter within a column are not significantly different. T₁ = Control (distilled water; DW); T₂ = undiluted extract (9 g extract + 10 ml DW); T₃ = 75% extract + 25% DW (7.5 g + 10 ml); T₄ = 50% extract + 50% DW (5 g + 10 ml); T₅ = 25% DW (2.5 g + 10 ml).

Table 17. Effect of root extract of sunflower on superoxidase (mg/ 100 g F. wt) of leaves in wheat varieties Margalla 99 and Chakwall 97.

Treatment	1st year		2nd year	
	V ₁	V ₂	V ₁	V ₂
T ₁	2.670 ^C	3.000 ^B	3.000 ^B	3.000 ^{BC}
T ₂	6.000 ^A	5.000 ^A	5.330 ^A	5.000 ^A
T ₃	5.000 ^{AB}	4.000 ^{AB}	5.000 ^A	4.330 ^{AB}
T ₄	3.000 ^C	2.670 ^B	4.670 ^A	3.670 ^{ABC}
T ₅	3.330 ^{BC}	2.330 ^B	3.000 ^B	2.330 ^C

Values followed by the same letter within a column are not significantly different.

sugar contents; this is in par with T₁ (control), and T₁ (control) showed the minimum values in the Chakwall 97. Similar result was found in the second year of experiment.

Effect of root extract of sunflower on protein content (mg/g) in leaves of wheat varieties Margalla 99 and Chakwall 97

Effect of leaf and root extract of sunflower on protein content in leaves of wheat varieties is shown in Table 16. A perusal of the table revealed that in the case of both wheat varieties Margalla 99 and Chakwall 97, T₂ (undiluted extract) showed the maximum contents of protein under the effect of leaf and root extract, which was followed by T₃ (75% extract), and T₁ (control) showed minimum amount of protein.

Effect of root extract of sunflower on superoxide dismutase (mg/100 g fresh weight) of leaves in wheat varieties Margalla 99 and Chakwall 97

From Table 17, it is revealed that in the case of both wheat varieties Margalla 99 and Chakwall 97, T₂ (undiluted extract) showed the maximum superoxide

dismutase (SOD) content under the effect of sunflower leaf and stem extract, followed by T₃ (75% extract), and T₄ (50% extract) whereas T₁ (control) showed minimum amount of SOD. Similar results were found in the second.

Effect of root extract of sunflower on peroxidase (POD) content (mg/100 g fresh weight) of leaves in wheat varieties Margalla 99 and Chakwall 97

From Table 18, it is revealed that in the case of both wheat varieties Margalla 99 and Chakwall 97, T₂ (undiluted extract) showed the maximum POD content under the effect of leaf and stem extract of sunflower, which was followed by T₃ (75% extract) and T₄ (50% extract), whereas T₁ (control) showed minimum amount of POD. Similar result was shown in the second year.

DISCUSSION

The allelopathic phenomenon has received much attention, as shown by numerous reports on the subject (Anaya, 1999; Khan et al., 2005; Narwal et al., 1998; Reigosa et al., 2006; Singh et al., 2001; Weston and Duke, 2003; Kamal and Bano, 2009; Kamal, 2011). The present study has shown that sunflower (*H. annuus* L.)

Table 18. Effect of root extract of sunflower on peroxidase (mg/ 100 g F. wt) activity of leaves in wheat varieties Margalla 99 and Chakwall 97.

Treatment	1st year		2nd year	
	V1	V2	V1	V2
T ₁	8.000 ^C	7.000 ^B	7.670 ^C	6.00 ^B
T ₂	14.67 ^A	13.00 ^A	15.33 ^A	14.00
T ₃	13.67 ^{AB}	11.00 ^A	14.67 ^A	11.67 ^A
T ₄	12.33 ^{AB}	10.67 ^A	13.33 ^{AB}	11.33 ^A
T ₅	10.67 ^{BC}	10.33 ^A	11.00 ^B	8.00 ^C

Values followed by the same letter within a column are not significantly different. T₁ = Control (distilled water; DW); T₂ = undiluted extract (9 g extract + 10 ml DW); T₃ = 75% extract + 25% DW (7.5 g + 10 ml); T₄ = 50% extract + 50% DW (5 g + 10 ml); T₅ = 25% DW (2.5 g + 10 ml).

extracts (leaves, stems, and roots) affect seed germination; so, aqueous extracts of sunflower parts, like leaves, stems, and roots, significantly affected germination. This indicates that reduction in these parameters (fresh weight, dry weight, root length, and shoot length) might have been the result of water-soluble allelochemicals in the extracts and their inhibitory effect or phytotoxicity on the measured parameters. Results show that sunflower plant parts varied in their allelopathic activity against wheat seedlings extracts of leaves, followed by roots which showed a higher allelopathic potential on the test plants as compared to stems. These results are in accordance with previous studies reporting that allelopathy may vary among plant parts (Turk and Tawaha, 2002; Peng et al., 2004). For all extracts, allelopathic activity increased with increases in extract concentrations. This is in agreement with the finding that under certain conditions, the rate of an elementary reaction is positively related to the reactant concentration. Previous studies have shown that the phytotoxicity of extracts was significantly increased as their concentration increased (Rice, 1984; Putnam, 1994; Sinkkonen, 2001; Peng et al., 2004; kamal and Bano, 2009). Wheat seed germination was reduced by all extracts of sunflower as compared with the untreated control. The observed inhibition of the two wheat varieties, namely Margalla 99 and Chakwall 97, for germination of seeds could be attributable to a contribution of allelochemicals released from the sunflower extracts of leaves, stems, and roots, as the allelochemicals are water-soluble and can accumulate upon release in seeds in direct contact with bioactive concentrations. There are several reports from literature showing that addition or incorporation of plant residues into the growth environment of another plant can result in growth inhibition (Patterson, 1981; Qasem, 1994; Chung and Miller, 1995; Al-Khatib et al., 1997). This study observed seed chlorosis on wheat seed at high rates of sunflower extracts; this may be due to a sunflower phenolic allelochemical effect on Mg-chelatase activity and chlorosis of seeds. The effect of higher concentration of extracts of sunflower suggests that

reduction in measured parameters may have the result of either or both the osmotic potential of the extracts or the presence of allelochemicals in the extracts. Beside possible allelochemicals, higher chemical concentrations in the extracts might possibly cause an osmotic stress during seed germination and seedling growth. The inhibition of seed germination and seedling growth was concentration-dependent; the present results are similar to those obtained by other workers, who observed inhibition of seed germination and seedling growth on other crops and weeds by sunflower extracts (Batish et al., 2002; Bogatek et al., 2005; Anjum et al., 2005; kamal and Bano, 2008). The allelochemicals inhibit germination and seedling growth, probably by affecting the cell division and elongation process that are very important at this stage or by interfering with enzymes involved in the mobilization of nutrients necessary for germination. Levizou et al. (2002) found that mitosis in root apex of lettuce retarded with leaf extracts. Growth hormones, like GA and IAA, are very important in agriculture. Growth-inhibiting compounds of agricultural importance have recently received considerable research attention (Siquira et al., 1991; Inderjit, 1996). These compounds which affect phytohormones are phenols with allelopathic characteristics (Schenk et al., 1999; Callaway and Aschhoug, 2000). Allelopathy is the plant harm to other plants in their vicinity or is an adverse influence of one plant on another (Rice, 1984). In plants exposed to sunflower extracts, changes in ABA were also recorded. In the leaves, the ABA content increased. In the case of roots, levels of ABA are very low in control plants and in treated plants they were high. This was probably due to the fact that in roots, as being very affected by allelochemicals early, cell death would occur. In roots with gradually dying cells, most of the already synthesized ABA is probably transported to the shoot, as a stress signal messenger. The increase of ABA in leaves correlates well with changes described earlier in the plant water status; that is, loss of turgor, lowered osmotic potentials and RWC, and the increase in leaf diffusive resistance. The involvement of ABA in plants coping with

abiotic and biotic stresses, especially those dealing with cell water deficit is well documented in the literature (Davies and Jones., 1991). This study clearly shows that ABA also plays an important role in the defense processes against allelopathic stress. An increase of ABA levels in seedlings of wheat exposed to allelopathy stress was observed in another study (Bernat et al., 2003). However, in this study, conversely to this work, the increase was recorded not only in leaves but also in roots. Different patterns of ABA accumulation in these two growth stages by wheat plants due to the same stress suggest that there are differences in mechanisms of coping with allelopathy stress or in sensitivity to this stress. Results presented in this work demonstrated that plants of sunflower cv. Hysun 38 are the donors of allelopathic compounds which affect acceptor plants via disorder in many physiological processes and that these changes, being mostly statistically significant showed a dose- and time-dependent manner. Recently, some progress has been made in the study of molecular processes involved in morphological and physiological adaptation of plants exposed to allelopathic chemicals of other plants. The present research was carried out to study the allelopathic effects of leaf stem and root on morphological, biochemical, and molecular criteria of wheat leaves. These extracts can be used in biological control as natural herbicides to reduce the risk of manufactured herbicides. With respect to the total soluble protein and the nucleic acids contents, there was a highly significant increase in the level of DNA in all treated samples of wheat seedlings. Wheat seedlings treated with sunflower extracts showed a highly significant increase in the level of DNA. These results were confirmed partially by (Duhan et al., 1995) who revealed a drastic increase in the level of nucleic acids and decrease in the level of soluble proteins in legume crops in response to plant extracts. They attributed this effect to the enzymes of synthetic pathways. In accordance with these results, Baziramakenga et al. (1997) reported that many phenolic acids reduced the incorporation of phosphorus into DNA in soybean. The present results are in controversy with those obtained by (Padhy et al., 2000), who reported that chlorophyll synthesis in leaves, as well as protein, carbohydrate, and nucleic acid (DNA and RNA) contents, in both shoots and roots of seedling was also decreased with increases in extracts concentrations. It was found that the activities of SOD and POD of many plants were affected by (allelochemicals) drought. Numerous studies have shown that the degree of injury caused by allelochemicals was negatively correlated to the increase of activities of SOD and POD (Dhindsa and Matow, 1981; Chowdhury and Choudhuri, 1985). But some results were the same as this; for example, Sun et al. (2010a) found that SOD and POD activities increased under allelochemical water stress (Wang et al., 2002); water stress and allelochemicals are correlated with each other because more allelochemicals

are produced during stress conditions. Some others got the similar results (Iturbe-ormatexe et al., 1998). In this study, it was found that SOD and POD activities in roots and leaves increased with allelochemicals. These differences may result in different treatment methods. In this study, the allelochemicals were applied during the whole growth period. Allelochemicals probably acted directly on roots, whereas leaves could reduce water transpiration by curling and stoma regulation. Furthermore, the degree of leaf injury caused by allelochemicals could be lessened through morphological adaptation and regulation of stem and sheath water content, stem diameter, or plant height. This showed that the ability of crops to resist allelochemicals was connected to the activities of protective enzymes and their defensive function. These mechanisms may be the main physiological action and defense from injury of crops under allelochemicals. There were significant differences in endogenous concentrations of GA and IAA in both leaves and roots of two wheat varieties during their growth when different concentrations of extracts of sunflower were applied. The GA and IAA concentration in leaves as well as roots decreased when applied different concentrations of extracts of sunflower were applied, while a higher reduction in both IAA and GA concentration was found when more concentrations of extracts of sunflower were applied in 2 wheat varieties. This decrease in leaves' GA and IAA may be attributed to slow transport of growth-promoting hormones from vegetative organs. Phytohormones may also enhance root development under growing conditions. Kuroha et al. (2002) reported that plant hormones stimulate adventitious root formation. Similar findings have been reported previously (Oda et al., 2003). The ability of the plants to absorb nutrients efficiently can be attributed to a higher number of adventitious root formations, and GA promotes this (Wahyuni et al., 2003; Kono, 1995; Watanabe, 1997). Root initiation and early development of root are also stimulated by auxin (Bellamine et al., 1998; Pan and Tian, 1999). There were significant decreases in endogenous concentrations of GA and IAA in the T1 and T5 treatments in both varieties (Margalla 99 and Chakwall 97). In controlling stomatal resistance, the most important signal is considered to be the concentration of plant hormones in xylem sap (Borel et al., 1997). Root-produced organic compounds in the xylem sap, such as hormones and amino acids, are considered to be vital for plant development (Oda et al., 2003). Extracts of sunflower (leaves, stems, and roots) decreased IAA and GA and increased ABA concentration in both leaves and roots compared to the control. Plant responses to allelopathy (stress) can also be determined by the variations in IAA, GA, and ABA concentration (Naqvi, 1999). A rapid decrease in leaf ABA concentration is observed in low concentrations of allelochemicals and in control plant. The GA concentration was decreased in plants under water stress conditions (Aharaoni, 1979; Guinn-Brummet,

1988; Yang et al., 2001; Xie et al., 2003). The increase in ABA may possibly be attributed to an induction in ABA synthesis. Abscisic acid contents in seedlings of leaves treated with extracts of sunflower were greater than those of control (Yang et al., 2001, 2004). Allelochemicals increased in the concentrations of ABA in leaves and in roots. This may be due to a higher concentration of growth-promoting hormones. During the present investigation, it was found that extracts of sunflower significantly increased the accumulation of soluble sugars in both wheat varieties. In these experiments, extracts of leaves, followed by root extracts and stem extracts, significantly increased the soluble sugar contents of wheat varieties Margalla 99 and Chakwall 97. Allelochemical treatment (T1) markedly increased sugar content in leaves. This increase may be due to the positive effect of ABA on assimilate translocation. Assimilate translocation to the developing seeds is reported to be under the control of ABA (Brenner and Cheikh, 1995; Yang et al., 1999, 2004). Allelochemicals decreased the accumulation of sugars markedly in leaves. The allelochemical treatments also showed an increase in sugar contents both in leaves. Ahmadi and Bakicer (1999) reported that ABA is involved in osmolyte regulation under moisture stress conditions. Mahajan and Tuteja (2005) reported that under severe drought, growth was inhibited by high concentrations of ABA and sugar, whereas low concentrations promote growth. Increased rates of photosynthesis and higher chlorophyll content might cause accumulation of sugars due to ABA or allelochemical treatments (Dong et al., 1995; Ndung et al., 1997). Drought tolerance in plants is enhanced by ABA caused by allelochemical treatment, possibly due to the accumulation of osmolytes, such as sugars. The present investigation indicated that application extracts of sunflower parts, like leaves, stems, and roots, caused accumulation of protein in both wheat varieties, Margalla 99 and Chakwall 97. Nevertheless, the intensity of increase was higher in Margalla 99 and Chakwall 97. The application of extracts of sunflower significantly increased the protein accumulation in both wheat varieties Margalla 99 and Chakwall 97. In these experiments, the extracts of leaves have more of an effect, followed by root extracts, and the lowest effect was observed in the case of stem extract. It was observed that allelochemicals increased protein content in leaves. This may be due to a positive role of ABA on protein accumulation. Guerrero and Mullet (1986) and Schmitz et al. (2000) reported that protein synthesis in developing seeds is induced by ABA. Zhang et al. (2001) reported that protein phosphorylation is enhanced under water stress due to increased concentration of ABA. Bartels and Sunkar (2005) and Ingram and Bartels (1996) investigated late embryo-genesis-abundant (LEA) proteins induced in vegetative tissues of plants in response to osmotic stress that may interact with carbohydrates to prevent cellular damage during dehydration. The recovery of leaf protein levels in

drought-stressed plants is more rapid at panicle initiation. In the present study, it is shown that allelopathy of sunflower caused accumulation of proline in both wheat varieties Margalla 99 and Chakwall 97. The intensity of increase was higher in Margalla 99. The application of extracts of sunflower significantly increased the proline accumulation in both wheat varieties. However, a significantly higher increase was recorded in Margalla 99. The efficiency of sunflower leaves, followed by root extracts, to increase proline contents was higher than root and stem extracts. Applications of allelochemical treatments markedly increased proline content in leaves of two wheat varieties, Margalla 99 and Chakwall 97. This magnitude of the increase was similar to that under drought stress. Under drought stress, in addition to its role as an osmoregulator, proline like other soluble organic compounds, may also act as osmo-protectants (Kamelie and Lose, 1995). ABA was considered to be involved in the accumulation of proline (Hose et al., 2000; Trotel Aziz et al., 2000; Nayyar and Walia, 2003), carbohydrates (Ahmadi and Baker, 2001), and other osmolytes in plants (Popova et al., 2000). Allelochemical-caused stress increased the accumulation of proline significantly in leaves. ABA enhanced the accumulation of proline contents in leaves. This increase may be due to the role of ABA, which may stimulate proline accumulation under water deficit conditions. Allelochemicals in T2, followed by T3, T4, and T5, caused a significant increase in proline levels compared to control (Bajji et al., 2001). Root length, shoot length, fresh weight, and dry weight of wheat seedling were significantly different in both the varieties with respect to different concentrations of extracts of sunflower; leaf extract showed more of an effect, followed by root extract, and the lowest effect was found in the case of stem extracts of sunflower.

ACKNOWLEDGEMENT

The author is thankful to Higher Education Commission of Pakistan for giving him a scholarship to complete his research work leading to doctoral degree.

Conclusions

From the experiments, these were concluded: allelochemicals secreted by sunflower inhibited germination and lowered the level of hormones, GA, IAA, shoot length, root length, fresh weight, dry weight, chlorophyll contents, and DNA, while they increased the values of ABA, protein, proline, sugar, SOD, and POD contents in wheat seedlings.

REFERENCES

Aharoni N, Blummenfeld A, Richmond AE (1979). Hormonal activity in

- detached lettuce leaves as affected by leaf water content. *Plant Physiol.*, 59: 1169-1173.
- Ahmadi A, Baker DA (1999). Effects of abscisic acid (ABA) on grain filling processes in wheat. *Plant Growth Regul.*, 28: 187-197.
- Ahmadi A, Baker DA (2001). The effect of water stress on the activities of key regulatory enzymes of the sucrose to starch pathway in wheat. *Plant Growth Regul.*, 35: 81-91.
- Al-Khatib K, Libbey C, Boydston R (1997). Weed suppression with Brassica green manure crops in green pea. *Weed Sci.*, 45: 439-445.
- Anjum T, Bajwa R, Javaid A (2005). Biological Control of *Parthenium* L: Effect of *Imperata cylindrica* L. on distribution, germination and seedling growth of *Parthenium hysterophorus* L. *Int. J. Agric. Biol.*, 7(3): 448-450.
- Anaya AL (1999). Allelopathy as a tool in the management of biotic resources in agrosystems. *Crit. Rev. Plant Sci.*, 18: 697- 739.
- Anjum T, Stevenson P, Hall D, Bajwa R (2005). Allelopathic potential of sunflower (*Helianthus annuus* L.) as a natural herbicide. In proceedings of the 4th World Congress on Allelopathy. Harper, J. D. I., M. An, H. Wu and J.H. Kent (Eds), 26 August 2005, International Allelopathy Society.
- Bajji M, Lutts S, Kinet J (2001). Water deficit effects on solute contribution to osmotic adjustment as a function leaf aging in three durum wheat (*Triticum durum* Desf.) cultivars performing differently in arid conditions. *Plant Sci.*, 160: 699-681.
- Bartels D, Sunkar R (2005). Drought and salt tolerance in plants. *Crit. Rev. Plant Sci.*, 24: 1-36.
- Borel C, Simonneau T, This D, Tardieu F (1997). Stomatal conductance and ABA concentration in the xylem sap of barley lines and constricting genetic origins. *Aust. J. Plant Physiol.*, 24: 607-615.
- Brenner ML, Chiekh N (1995). The role of hormones in photosynthate partitioning and seed filling.. In P.J. Davies (ed.) *Plant hormones, physiology, biochemistry and molecular biology*. Kluwer Academic Publ., Dordrecht, the Netherlands., pp. 649-670.
- Bogatek R, Gniazdowska A, Zakrzewska W, Oracz K, Gawroski SW (2005). Allelopathic effects of sunflower extracts on mustard seed germination and seedling growth. *Biol. Plant.*, 50 (1):156-158. Batish DR, Singh HP, Saxena DB, Kohil RK (2002). Weed suppressing ability of parthenin – A sesquiterpene lactone from *Parthenium hysterophorus* L.. *New Zealand Plant Prot.*, 55: 218-221.
- Bernat W, Gawaronska H, Janowiak F, Gawronski SW (2003). The effect of sunflower allelopathics on germination and seedling vigor of winter wheat water and mustard. Abstract of Fifth International Conference, Ecophysiological Aspects of plants responses to stress factor, Cracow, Cracow, Poland. *Acta Physiol. Plant.* 25 (Supplement): pp. 24- 25.
- Baziramakenga R, Heroux GD, Simard RR, Nadeau P (1997). Allelopathic effects of phenolic acids on nucleic acids and protein levels in soybeans seedlings. *Can. J. Bot.*, 75: 445- 450.
- Chung IM, Miller DA (1995). Natural herbicide potential of alfalfa residues on selected weed species. *Agron. J.*, 87: 920-925.
- Chowdhury RS, Chowdhuri MA (1985). Hydrogen peroxide metabolism as index of water stress tolerance in jute. *Phylogica Plant.*, 65: 503-507.
- Callaway RM, Aschehoug ETM (2000). Invasive plant versus their new and old neighbours: a mechanism for exotic invasion. *Science.*, 290: 521-523.
- Dong B, Rengel Z, Graham RD (1995). Root morphology of wheat genotypes differing in Zn efficiency. *J. Plant Nutr.* 18: 2761-2773.
- Davies WJ, Jones HG (1991). *Abscisic acid, Physiology and Biochemistry*. Bios Scientific Publishers Limited. St Thomas House, Backet Street, Oxford OX1 1SJ, UK: 63: 77, 137-149, 189-197: 201-225.
- Dhindsa RS, Matsow EW (1981). Drought tolerance in two mosses: correlated with enzyme defence against lipid peroxidation. *J. Exp. Bot.*, 32: 79- 91.
- Guerrero F, Mullet JE (1986). Increased abscisic acid biosynthesis during plant dehydration requires transcription. *Plant Physiol.*, 80: 588-591.
- Guinn G, Brummett DL (1988). Abscisic acid in young cotton furts and their abscission zones in relation to fruit retention during and after moisture stress. *Plant Physiol.*, 86: 28-31.
- Hose E, Studle E, Hartung W (2000). Abscisic acid and hydraulic conductivity of maize roots: a study using cell-n and root pressure probes. *Planta.*, 211: 874-882.
- Iturbe-ormatexe I, Escredo PR, Arrese- Igor C, Becana M (1998). Oxidative damage in pea plants exposed to water deficit or paraquat. *Plant Physiol* 116: 173-181.
- Inderjit A (1996). Plant phenolics in allelopathy. *Botanical Rev.* 62., 182-202.
- Ingram J, Bartels D (1996). The molecular basis of dehydration tolerance in plants. *Annu. Rev. Plant Physiol. Plant Mol. Biol.*, 47: 377- 403.
- Kamal J (2011). Quantification of alkaloids, phenols and flavonoids in sunflower (*Helianthus annuus* L.) *Afr. J. Biotechnol.*, 10(16): 3149-3151.
- Kamal J, Bano A (2008). Allelopathic potential of sunflower (*Helianthus annuus* L.) on soil metals and its leaves extracts on physiology of wheat (*Triticum aestivum* L.) seedlings. *Afr. J. Biotechnol.*, 7(18): 3261-3265.
- Kamal J, Bano A (2008). Effects of sunflower (*Helianthus annuus* L.) extracts on wheat (*Triticum aestivum* L.) and physicochemical characteristics of soil. *Afr. J. Biotechnol.*, 7(22): 4130-4135.
- Kamal J, Bano A (2008). Potential allelopathic effects of sunflower (*Helianthus annuus* L.) on microorganisms. *Afr. J. Biotechnol.*, 7 (22): 4208-4211.
- Kamal J, Bano A (2008). Efficiency of allelopathy of sunflower (*Helianthusannuus* L.) on physiology of wheat (*Triticum aestivum* L.). *Afr. J. Biotechnol.*, 8(15): 3555-3559.
- Kamal J, Bano A (2009). Efficiency of allelopathy of sunflower (*Helianthusannuus* L.) on physiology of wheat (*Triticum aestivum* L.). *Afr. J. Biotechnol.*, 8(15): 3555-3559.
- Kamelie A, Losel DM (1995). Contribution of carbohydrates and solutes to osmotic adjustment in wheat leaves under water stress. *Plant Physiol.*, 145: 363-366.
- Kono M (1995). Physiological aspects of lodging In: Mastuo T, Kumazawa K, Ishii R, Ishihara K, Hirata H (eds.). *Science of the Rice. Vol. II. Physiology*. Food and Agricultur Policy Research Center, Tokyo, Japan.
- Khan TD, Chung MI, Xuan TD, Tawanta S (2005). The exploitation of crop allelopathy in sustainable agriculture production. *J. Agron. Crop Sci.*, 188: 19- 24.
- Kuroha T, Kato H, Asami T, Yoshida S, Kamada H, Satoh S (2002). A transzeatin riboside in root xylem sap negatively regulates adventitious root formation on cucumber hypocotyls. *J. Exp. Bot.*, 53: 2193- 2200.
- Levizou E, Karageorgou P, Psaras GK, Manetas Y (2002). Inhibitory effects of water soluble leaf leachates from *Dittrichia viscosa* on lettuce root growth, statocyte development and graviperception. *Flora*, 197: 152- 157.
- Mahajan S, Tueja N (2005). Cold, salinity and drought stresses: An overview. *Arch. Biochem. Biophys.* 444: 139-158.
- Naqvi SSM (1999). Plant hormones and stress phenomena. In: Pressarakli, M. (ed.): *Handbook of Plant and Crop Stress*. Marcel Dekker, New York-Basel., pp. 709-730.
- Nayyar H, Walia DP (2003). Water stress induced praline accumulation in contrasting wheat genotypes as affected by calcium and abscisic acid. *Biol. Plant.*, 46: 275-279.
- Ndungu CK, Shimizu M, Okamoto G, Hirano K (1997). Abscisic acid, carbohydrates, and nitrogen contents of Kyoho grapevines in relation to bud break induction by water stress. *Am.J. Enol. Vitic.*, 48: 115-120.
- Narwal SS, Hoagland RE, Dilchey RH, Reigosa MJ (1998). Allelopathy in Ecological Agriculture and Forestry. In proceedings of the 3rd International Congress on Allelopathy in Ecological Agriculture and Forestry, Dharawad, India. pp. 18-21.
- Oda A, Sakuta C, Masuda S, Mizoguchi T, Kamoda H, Satoh S (2003). Possible involvement of leaf gibberellins in the clock-controlled expression of XSP 30, a gene encoding a xylem sap lectin, in cucumber root. *Plant Physiol.*, 133: 1779- 1790.
- Pan R, Tian X (1999). Comparative effect of IBA, BSAA, and 5,6-C1AA-Me on the rooting of hypocotyls in mungbean. *Plant Growth Regul.*, 27: 91-98.

- Popova LP, Outlaw WH, Aghoram K, Hite DRC (2000). Abscisic acid and intra leaf water-stress signal. *Physiol. Plant.*, 108, 376-381.
- Peng SL, Wen J, Guo QF (2004). Mechanism and active variety of allelochemicals. *Act. Bot. Sinica.*, 46: 757-766.
- Putnam AR, De Frank J, Barnes JP (1994). Exploitation of allelopathy for weed control in annual and perennial cropping system. *J. Chem. Ecol.*, 9; 1001-1010.
- Patterson DT (1981). Effects of allelopathic chemicals on growth and physiological responses of soybean (*Glycine max* L.). *Weed Sci.*, 29: 53-59.
- Padhy BB, Patanaik PK, Tripathy AK (2000). Allelopathic potential of *Eucalyptus* leaf litter leachates on germination and seedling growth of finger millet. *Allel. J.*, 7: 69- 78.
- Qasem JR (1994). Allelopathic effect of white top (*Lepidium draba* L.) on wheat and barley. *Allel. J.*, 1: 29-40.
- Reigosa MJ, Pedrol M, Gonzales L, Narwal SS (2006). Allelopathy in Ecological Sustainable Agriculture. In: *Allelopathy: A physiological Process with Ecological Implications*. Reigosa, M. J., M. Pedrol and L. Gonzales (eds.), Springer, Netherlands, pp. 537-564.
- Reigosa MJ, Pedrol N, Sanchez-Moreiras AM, Gonzales L (2002). Stress and allelopathy. In: *Allelopathy, from Molecules to Ecosystems*, Reigosa MJ and Pedrol N, Eds. Science Publishers, Enfield, New Hampshire.
- Rice EL (1984). *Allelopathy*, 2nd ed. New York: Academic Press.
- Schmitz N, Abrams SR, Kermode AR (2000). Changes in the abscisic acid content and embryo sensitivity to (+)-abscisic acid during the termination of dormancy of yellow cedar seed. *J. Exp. Bot.*, 51, 1159-1162.
- Singh HP, Kohli RK, DR Batish (2001). Allelopathy in agrosystems: An overview. *J. Crop Prod.*, 4: 1- 41.
- Sinkkonen A (2001). Density-dependent chemical interference-an extension of the biological response model. *J Chem Ecol.*, 27: 1513-1523.
- Siqueira JO, Safir GR, Nair MG (1991). Stimulation of vesicular-arbuscular mycorrhiza formation and growth of white clover by flavonoid compounds. *New Phytol.*, 118, 87-93.
- Schenk HJ, Callaway RM, Mahal BE (1999). Spatial root segregation: are plants territorial. *Adv. Ecol. Res.*, 28: 146-180.
- Trotel-Aziz P, Niogret MF, Larher F (2000). Proline level is partly under the control of abscisic acid in canola leaf discs during recovery from hyper-osmotic stress in the adult maize leaf. *J. Exp. Bot.*, 54, 2177-2186.
- Turk MA, Tawaha AM (2002). Effect of sowing rates of weeds control method on winter wheat under Mediterranean environment. *Pak. J. Agron.*, 1: 25-27.
- Weston LA, Duke SO (2003). Weed and crop allelopathy. *Crit. Rev. Plant Sci.*, 22: 367- 389.
- Wang J, Li D, Guld Q (2002). The response to water stress of the antioxidant system in maize seedling root with different drought resistance. *Acta Botanica Boreali-Occidentalia Sinica.*, 22, 285- 290.
- Wahyuni S, Sinniah UR, Amarthalingam R, Yusop MK (2003). Enhancement of seedling establishment in rice by selected growth regulators as seed treatment. *Jurn Penelitian Pertanian Pangan.*, 22: 51-55.
- Watanabe T (1997). Lodging resistance., In: Futsuhara TY, Kikuchi F and Yamaguchi H (eds.). *Science of the rice Plant. Vol. III. Genetics. Food and Agriculture Policy Research Center, Tokyo, Japan.* pp. 567-577.
- Xie Z, Jiang D, Cao W, Dai T, Jing Q (2003). Relationships of endogenous plant hormones to accumulation of grain protein starch in winter wheat under different post-anthesis soil water statuses. *Plant Growth Regul.*, 41: 117-127.
- Yang J, Zhang J, Wang Z, Zhu Q, Wang W (2001). Hormonal changes in the grains of rice subjected to water stress during grain filling. *Plant physiol.*, 127: 315-323.
- Yang JC, Zhang JH, Ye YX, Qang ZQ, Zhu QS, Liu LJ (2004). Involvement of abscisic and ethylene in the responses of rice grains to water stress during filling. *Plant Cell Environ.*, 27: 1055-1064.
- Zhang X, Miao YC, An GY, Zhou Y, Shangguan ZP, Gao JF, Song CP (2001). K⁺ channels inhibited by hydrogen peroxide mediate abscisic acid signalling in guard cells. *Cell Res.*, 11: 195-202.