

## Review

# Advances in genetic engineering for plants abiotic stress control

Tchouopou Lontchi Josine<sup>1,2</sup>, Jing Ji<sup>2\*</sup>, Gang Wang<sup>2</sup> and Chun Feng Guan<sup>2</sup>

<sup>1</sup>School of Chemical Engineering and Technology, Department of Bioengineering, Tianjin University, Tianjin 300072, People's Republic of China.

<sup>2</sup>School of Agriculture and Bioengineering, Department of Biotechnology, Tianjin University, Tianjin 300072, People's Republic of China

Accepted 8 April, 2011

**Agricultural productivity is highly influenced by abiotic stresses, known as the most harmful factor concerning the growth and productivity of crops worldwide. Furthermore, industrial crops are nowadays highly influenced by abiotic stress; these include extremes in temperature, drought, salinity, heavy metals and radiation. Typical studies were discussed by many researchers about the control of abiotic stress in plants by the expression, over-expression or switching off abiotic stress-related genes. Despite the rapid evolution of the research, some crops are still expected to decline by 15 to 32% in the next fifty years. Consequently, engineering genes that protect and maintain the function and structure of cellular components can enhance tolerance to stress. This review presents principal methods adapted in the control of plants abiotic stresses including recent advances in using transgenes for the improvement of abiotic stress tolerance in plants. Specified analysis of recent advances in abiotic stress control could describe trehalose as a better compound in the control of plant abiotic stresses. Therefore, studies of genes-related trehalose metabolism and associated patterns could not only provide an improved plant metabolism, phenotypes and texture, but in fact, the plants become highly resistant to abiotic stress.**

**Key words:** Abiotic stresses, crops, expression, over-expression, switching off, trehalose, genes-related.

## IMPACT OF ABIOTIC STRESS IN PLANTS

Abiotic stress is known as the most harmful factor concerning the growth, productivity of crops and worldwide agricultural loss, posing a substantial challenge in the face of an ever increasing world population (Gao et al., 2007). In the recent years, many researchers have drawn their investigations in biotic and abiotic stress in plants (Boyer, 1982; Bray et al., 2000; Mittler, 2006; Brown et al., 2003). Abiotic stressors are at their most harmful stage when they occur together, in combinations of abiotic stress factors (Mittler, 2006). Although some biotic stresses can be controlled by germicide, pesticides, antibiotic treatment and/or plant metabolism, it is urgent to control abiotic stress, which can possibly affect response to biotic stress control with detrimental effect on plant growth and yields. Therefore, solving few threats such as the depletion of the

the ozone layer, high risk of exposure to ultra violet (UV) and infra-red radiation, environmental warming, increasing soil salinity, drinking polluted underground water, production of toxic chemicals and contaminated food materials which were not visible till 1960 is highly required. Moreover, cold weather crops like rye, oats, wheat and apples are expected to decline by about 15% in the next fifty years and strawberries will drop as much as 32% simply because of projected climate changes of a few degrees (Lane and Jarvis, 2007). Presently, industrial needs are highly influenced by abiotic stress. Improved production efficiency can be achieved by developing stress tolerant crops (Ribaut et al., 1997), which necessitates research on plant adaptation mechanisms against adverse environmental conditions. Activation of stress-response transcription factors has been found to enhance plant tolerance to various kinds of stress in transgenic plants (Kasuga et al., 1999; Vogel et al., 2005; Villalobos et al., 2004; Gu et al., 2002)

Salinity and drought are the most commonly observed

\*Corresponding author. E-mail: [jjing@tju.edu.cn](mailto:jjing@tju.edu.cn), Tel/fax: 0086-22-87401878

abiotic stress; they significantly reduce yield and affect almost every aspect of the physiology and biochemistry of plants. Great efforts have been devoted to understanding the physiological aspects of response in plants to salinity and drought-tolerant genotypes (Cuartero et al., 2006). Studies on tomato cultivar Liaoyuan Duoli (*Solanum lycopersicum* L.) indicated that, soluble sugar and hexoses accumulated to higher levels and the contents of sucrose and starch were lower in mature fruit under the two stress treatments. Salt and water stresses can enhance the invertase and sucrose synthase activities of tomato fruit in a long period of time (45 to 60 days after anthesis) and elevate the expression of soluble acid invertase mRNA. Two different stresses could also regulate the soluble acid invertase activity by controlling its gene expression. The activity of sucrose synthase was linked to the changes in soluble sugar levels but not with transcript levels (Lu et al., 2009). However, little is still known about the function of other components of the plant transcription machinery during stress.

## **METHODS DEVELOPED AND ADVANCES IN THE CONTROL OF PLANTS ABIOTIC STRESS**

### **Maintenance of soil biodiversity**

Healthy environment (pollution free, habitat conservation, climate variability and some other changes) as well as biologically diverse soil are necessary for productivity concerns. The healthy soil community keeps pests and diseases under control. Several microorganisms have been used in the control of soil biodiversity; the fungi *Beauveria brongniartii* and *Metarhizium anisopliae* already in commercial formulas to control pests *Glomus clarum* and to grow maize and shorgumin in *Striga hermonthica* infected soils.

Resilience to abiotic stress and soil rehabilitation used some microorganisms for restoration of soil quality; for example, arbuscular mycorrhizal fungi (AMF) is used to reduce potentially toxic elements (Cd, Cr, Mn) from the soil commonly known as bioremediation. Some use mycorrhizal fungi to enable plants to cope with abiotic stress by alleviating nutrient deficiencies, improving drought tolerance, overcoming the detrimental effects of salinity, improving the adaptation of sterile micro-propagated plants to unsterile substrates and to field conditions. Although these researches have provided sufficient information in the maintenance of plants' good environment, they are still not enough to understand the impact of soil biodiversity in the control of abiotic stress and agricultural production (Barrios, 2007; Wall and Moore, 1999; Bloem et al., 2003; Swift et al., 2004).

### **Positive interactions between plant species**

Positive interactions between plant species are wide-

spread in natural communities and have been acknowledged as a major driver of plant community dynamics and ecosystem processes (Bruno et al., 2003). There have been no large-scale experiments designed to examine the generality of positive interactions in plant communities and their importance relative to competition. Ragan et al. (2002) found that, competition generally but not exclusively dominate interactions at lower elevations where conditions are less physically stressful. In contrast, at high elevations where abiotic stress is high the interactions among plants are predominantly positive. Furthermore, across all high and low sites, positive interactions are more important at sites with low temperatures in the early summer and competition prevails at warmer sites. Facilitative and competitive interactions are thought to be of great importance in arid and semi-arid areas (Whitford, 2002). However, only a few studies have evaluated how changes in abiotic conditions modify the relative importance of facilitation and competition in these environments (Tielborger and Kadmon, 2000; Pugnaire and Luque, 2001; Maestre et al., 2003). Furthermore, none of these studies were conducted at multiple sites across a gradient of abiotic stress. A shift from facilitation to competition under high abiotic stress conditions is likely to occur when the levels of the most limiting resource are so low that the benefits provided by the facilitator cannot overcome its own resource uptake (Fernando et al., 2004). Positive interactions between different species of plants, including cross-pollination or mutualistic actions were also studied (Maestre et al., 2007).

### **Conventional breeding to genetic engineering**

Success in breeding for better adapted varieties to abiotic stresses depends upon the concerted efforts by various research domains including plant and cell physiology, molecular biology, genetics and breeding (Pooja et al., 2008; Muhammad and Nudrat, 2009). Common bean (*Phaseolus vulgaris* L.) because of its wide genetic variability, is one of the crops that received attention for more than 20 years in breeding program for abiotic stress control such as drought, tolerance to low soil phosphorus (P) and nitrogen (N) availability and resistance to high aluminium (Al) in acid soils (Lynch and Beebe, 1995; Thung and Rao, 1999; Rao, 2002; Beebe et al., 2008). Meanwhile, the use of modern molecular biology tools for elucidation of the control mechanisms of abiotic stress tolerance and for engineering stress tolerant crops is based on the expression of specific stress-related genes. Instead of relying on sexual recombination to thoroughly stir the parental genes, genetic engineering preserves the integrity of the parental genotype, inserting only a small additional piece of information that controls a specific trait. Genetic engineering allows the direct transfer, removing or switching off (Tables 1 and 2) a particular or just a few genes between either closely or distantly related

**Table 1.** Illustration of some recent genes expressed and response to abiotic stress.

Gene/molecule used and origin	Plant tested	Abiotic stress controlled	Associated pattern	Reference
<i>Arabidopsis DREB2C</i>	<i>Arabidopsis</i>	Heat stress	Transactivation of the DRE-dependent transcription factor HsfA3	Huan et al. (2010)
<i>Arabidopsis ftl1-1D/ddf1</i>	<i>Arabidopsis</i>	Enhances tolerance to cold, drought, heat and salt stresses	The induced gene encoded an AP2 transcription factor of the <i>CBF/DREB1</i> subfamily	Hong-Gyu et al. (2011)
Rice <i>OsABF2</i>	Rice	Induced by different types of abiotic stress treatments such as drought, salinity, cold, oxidative stress, and ABA.	The group A <i>bZIP</i> transcription factors play important roles in the ABA signaling pathway in <i>Arabidopsis</i> . In yeast experiments, <i>OsABF2</i> was shown to bind to ABA-responsive elements ( <i>ABREs</i> ) and its N-terminal region found to be necessary to transactivate a downstream reporter gene.	Md. Amir et al. (2010)
<i>Hevea brasiliensis HbMT2</i>	<i>Escherichia coli</i>	Strongly induced by ethephon and hydrogen peroxide (H <sub>2</sub> O <sub>2</sub> ) stress. Overproduction induce more tolerance on Cu <sup>2+</sup> and Zn <sup>2+</sup> , and the recombinant <i>HbMT2</i> scavenge the reactive oxidant species (ROS) in vitro.		Jiahong et al. (2010)
Apple <i>cyMDH</i>	Apple callus and tomato	Involved in plant and cell growth as well as the tolerance to cold and salt stresses.		Yu-Xin et al. (2011)
<i>Panax ginseng</i> PgCat1	<i>Panax ginseng</i>	Different stresses, heavy metals, plant hormones, osmotic agents, high light irradiance, abiotic stresses, triggered a significant induction of PgCat1.		Munkhbaatar et al. (2010)

Table 1. Continue.

Emmer wheat ( <i>Triticum turgidum</i> ssp. dicoccoides) <i>TdicDRF1</i>	Wheat; <i>Arabidopsis</i> <i>thaliana</i> .	Drought resistance	Analysis of the AP2/ERF DNA-binding domain of TdicDRF1 as a GST-fusion protein and its binding to DRE by electrophoretic mobility shift assay (EMSA) indicate functional differences between wheat <i>DREBs</i> and those characterized in <i>Arabidopsis thaliana</i> .	Stuart et al. (2011)
<i>Citrus unshiu</i> <i>CitERF</i>	<i>Citrus unshiu</i>	Play an important role in some biological processes during fruit ripening and in improving tolerance to drought, low temperature and salt stress.		Xiao-Yan et al. (2011)
<i>Trichoderma harzianum</i> hsp70	<i>Arabidopsis</i>	Enhanced tolerance to heat stress, osmotic, salt and oxidative stresses	Increased transcript levels of the Na <sup>+</sup> /H <sup>+</sup> exchanger1 ( <i>SOS1</i> ) and ascorbate peroxidase1( <i>APX1</i> ) genes	Marta et al. (2009) and Hervé and Ary (2010)
Rice <i>OsAP2LP</i>	Rice	Plays important roles when plants response to cold, salt and drought stresses through ABA and gibberellin pathways.		Ma et al. (2010)
<i>Brassica juncea</i> $\gamma$ -TMT	<i>Arabidopsis</i>	Enhanced tolerance to salt, heavy metal and osmotic stress		Mohd et al. (2010)
Cauliflower mosaic virus (CaMV) 35S promoter	Tobacco	Acute single dose of UVC, UVB or X-ray radiation and exposure to high temperature respectively decreased and increased the transgene expression level		Alex et al. (2010)
<i>Oryza sativa</i> miRNAs	Rice	Control in some extend cold, dehydration, salinity and abscisic acid (ABA)		Xinyu et al. (2010)

**Table 1.** Continue.

L. temulentum ( <i>eEF- 1a</i> , <i>ACT11</i> and <i>UBQ5</i> )	L. temulentum	Drought, UV stress, Heat stress, Wounding, Salt stress, Osmotic stress, Cold stress	<i>eEF-1a</i> and <i>UBQ5</i> were the most stable and <i>ACT11</i> was the least stable in stress Condition	James and Ruth (2009)
Maize ( <i>Zea mays</i> L.) <i>ZmCOI6.1</i>	<i>Arabidopsis</i>	Cold, drought and salinity as well as the signaling molecules abscisic acid and salicylic Acid		Orlene et al. (2009)
<i>CaBI-1</i> from the cDNA library of heat-stressed hot pepper ( <i>Capsicum annuum</i> )	Tobacco	High or low temperatures, drought, high salinity, flooding and heavy metal stresses, and ABA		Mohammad et al. (2009)
L. maritime <i>AtbZIP24</i> <i>Tamarix hispida</i> <i>ThbZIP1</i>	<i>Arabidopsis</i> tobacco	Decreased <i>AtbZIP24</i> expression activated the expression of stress-inducible genes: the Na <sup>+</sup> transporter <i>AtHKT1</i> , the Na <sup>+</sup> /H <sup>+</sup> antiporter <i>AtSOS1</i> , the aquaporin <i>AtPIP2.1</i> , and a glutamine synthetase. overexpression of <i>ThbZIP1</i> can enhance the activity of both peroxidase (POD) and superoxide dismutase (SOD), and increase the content of soluble sugars and soluble proteins under salt stress conditions.	Target genes of <i>AtbZIP24</i> with functions in plant growth and development were identified such as an argonaute (AGO1)-related protein and cyclophilin <i>AtCYP19</i> . The salt tolerance in transgenic plants correlated with reduced Na <sup>+</sup> accumulation in leaves	Oksoon et al. (2009) and Yucheng et al. (2010)
Hot peppers <i>PepRSH</i>	<i>Escherichia coli</i>	<i>PepRSH</i> transcripts were highly accumulated in non-host resistance response-induced leaves and in leaves following induction with salicylic acid, methyl jasmonate, wounding, hydrogen peroxide, and ultraviolet- B. The expression of <i>PepRSH</i> was also influenced by abiotic stresses, such as flooding and high salinity	<i>PepRSH</i> has (p)ppGpp synthetase activity only in the (p)ppGpp synthetase domain	Tae-Ho et al. (2009)
Cypress seedlings ( <i>Cupressus simper- virens</i> L.) <i>cyplp009</i> ; <i>cyplp005</i> ; <i>cyplp006</i> ; <i>cyplp017</i> ; <i>cyplp010</i> ; <i>cyplp020</i> ; <i>cyplp012</i> ; <i>cyplp002</i> ; <i>cyplp007</i> ; <i>cyplp019</i> ; <i>cyplp003</i> ; <i>cyplp016</i> ; <i>cyplp008</i> ; <i>cyplp015</i> ; <i>cyplp021</i>	Cypress canker	Control of cold		Luca et al. (2009)

Table 1. Continue.

Wheat <i>TaVP1</i> , <i>TaVP2</i> and <i>TaVP3</i>	Wheat	Salinity stress	<i>TaVP3</i> transcripts were only detected in Developing seeds, and no <i>TaVP2</i> transcripts were found in germinating seeds. <i>TaVP2</i> was mainly expressed in shoot tissues and down-regulated in leaves under dehydration. Its expression was up-regulated in roots under high salinity. <i>TaVP1</i> was relatively more ubiquitously and evenly expressed than <i>TaVP2</i>	Yuezhi et al. (2009)
Wheat <i>TaSTRG</i>	Rice	Expression induced by salt, polyethylene glycol (PEG), abscisic acid (ABA), and cold. Overexpression showed higher salt and drought tolerance	Under salt stress, the transgenic rice had a lower intracellular Na <sup>+</sup> /K <sup>+</sup> ratio Under salt and PEG treatments, these <i>TaSTRG</i> overexpressing rice plants had higher survival rate, fresh weight and chlorophyll content, accumulated higher proline and soluble sugar contents, and had significantly higher expression levels of putative proline synthetase and transporter genes	Wei et al. (2009)
<i>Arabidopsis CRT/DRE</i> binding factor 1 ( <i>CBF1</i> )	Tobacco	Enhancing low temperature tolerance of plants	The photochemical efficiency of PSII(Fv/Fm) and the photo-oxidizable P700 in the transgenic lines overexpressing <i>CBF1</i> were higher. A higher NPQ and qf, lower Fv/Fm, higher activity of SOD, and lower content of MDA Higher expression levels of <i>Nicotiana tabacum</i> copper/zinc superoxide dismutase ( <i>Cu/Zn SOD</i> ) were also detected in the transgenic lines	Jia-Sen et al. (2010)
<i>Arabidopsis AVP1</i>	Alfalfa ( <i>Medicago sativa</i> )	Enhanced salt and drought tolerance	Transgenic plants accumulate more Na <sup>+</sup> , K <sup>+</sup> and Ca <sup>2+</sup> in leaves and roots. Moreover, the leaves of transgenic plants retain more water during drought stress	Ai-Ke et al. (2009)
Bruguiera Gymnorhiza Bg70 and cyc02	<i>Arabidopsis</i>	Increased tolerance to NaCl		Shota and Yuichi (2009)

Table 1. Continue.

Cotton <i>GhTIP1;1</i> tonoplast intrinsic protein (TIP)	<i>Xenopus laevis</i> oocytes	Involved in cell freezing-tolerance; salt, heavy metals and nutrient deficiency), abscisic acid (ABA), gibberellic acid (GA) osmotic stress and cold.	<i>GhTIP1;1</i> facilitated water transport to the cells <i>GhTIP1;1</i> transcripts were predominantly accumulated in roots and hypocotyls. Trans-activating ability tests in yeast demonstrate that <i>RAP2.6</i> could act as a transactivator. <i>RAP2.6</i> is able to bind to the <i>GCC</i> and <i>CE1</i> cis-elements	Deng-Di et al. (2009), Ayalew et al. (2011) and Qiang et al. (2010)
Barley <i>HvTIP1;2</i> , <i>HvTIP2;1</i> , <i>HvTIP2;2</i> , <i>HvTIP2;3</i> and <i>HvTIP4;1</i>	Barley ( <i>Hordeum vulgare</i> )			
<i>Arabidopsis RAP2.6</i>	<i>Arabidopsis</i>			
Rice ( <i>Oryza rufip-ogon</i> ) <i>OrbHLH2</i>	<i>Arabidopsis</i>	Increased tolerance to salt and osmotic stress	The stress-responsive genes <i>DREB1A/CBF3</i> , <i>RD29A</i> , <i>COR15A</i> and <i>KIN1</i> were up regulated in transgenic plants	Jing et al. (2009)
<i>Brassica juncea</i> AnnBj1	Tobacco	Significant tolerance to dehydration (mannitol), salt (NaCl), heavy metal ( <i>CdCl2</i> ) and oxidative stress (H <sub>2</sub> O <sub>2</sub> ) at the seedling stage	Retain higher chlorophyll levels in response to the studied stresses	Sravan et al. (2008)
Cassava <i>CuZnSOD</i> and pea <i>APX</i>	Fescue	Functionally expressed under methyl viologen (MV), H <sub>2</sub> O <sub>2</sub> , heavy metals copper, cadmium, and arsenic		Sang-Hoon et al. (2007)
Maize kernel <i>SAT41</i> and <i>Arabidopsis MBF1a</i>	Maize and <i>Arabidopsis</i>	Maize clone ( <i>SAT41</i> ) seemed to confer elevated salt tolerance		Min-Jung et al. (2007)
Rice <i>OsUGE-1</i>	<i>Arabidopsis</i>	Tolerance to salt, drought and freezing stress	No alteration of plant morphology. Transgenic plants showed a higher level of the soluble sugar raffinose	Hui-li et al. (2007)
Rice <i>AtNDPK2</i>	Rice	<i>AtNDPK2</i> plays a regulatory role in chilling and antioxidant signaling in plants	Accumulation of transcripts of pBC442 and pBC601, is related to chilling stress. <i>OsAPX1</i> -, <i>OsAPX2</i> -, and <i>OsSodB</i> -scavenging free oxygen radicals, such as superoxide eO <sub>2</sub> _T and hydrogen peroxide (H <sub>2</sub> O <sub>2</sub> ), could be induced in <i>AtNDPK2</i> -overexpressed rice plants.	Eun et al. (2007)
Pepper <i>CaXTH3</i>	<i>Arabidopsis</i>	Resistance to drought, high salinity and cold temperature	Abnormal leaf morphology	Seok et al. (2006)
Yeast <i>YCF1</i>	<i>Arabidopsis</i>	Enhanced in salt or xenobiotic chemical (1-chloro-2, 4-dinitrobenzene,CDNB) tolerance	<i>COR15A</i> and <i>KIN1</i> were less strongly induced in the transgenic plants	Eun-Ji et al. (2006)

**Table 2.** Illustration of advanced trehalose genes expressed in the control of abiotic stress.

Gene/molecule used and origin	Plant tested	Abiotic stress controlled	Associated pattern	References
Exogenous trehalose	<i>Arabidopsis</i>	Promotes non-structural carbohydrate accumulation; Induces chemical detoxification and stress response proteins	Increased starch and decreased ATP levels, glutamate, glutamine, hexoses and chlorophyll were unaffected. Phi glutathione S-transferase 2 ( <i>AtGSTF2</i> ), flavin mononucleotide-binding flavodoxin-like quinone reductase 1 ( <i>FQR1</i> ), cytosolic dehydroascorbate reductase 1 ( <i>DHAR1</i> ) and S-adenosylmethionine synthetase 2 ( <i>SAMS2</i> ), were involved in either detoxification or stress responses.	Hanhong et al. (2005)
Yeast trehalose-6-phosphate synthase ( <i>TPS1</i> )	Tomato	Improved tolerance under drought, salt and oxidative stress	Thick shoots, rigid dark-green leaves, erected branches and an aberrant root development. Higher chlorophyll and starch content	Carolina and Francisco (2005)
Fungal entomopathogen <i>Beauveria bassiana</i> Neutral trehalase(NTH1)	<i>E. coli</i>	Crucial stress-responsive elements; NaCl and thermal stress	BbT4 is controlled by the shortest NthP4. that is, this promoter fragment is crucial to initiating high-level <i>BbNTH1</i> expression in <i>B. bassiana</i> under oxidative stress.	Qian et al. (2010)

organisms. Crop improvement can be achieved in a shorter time compared with conventional breeding. Hence, genetic engineering for developing stress tolerant plants, based on the introgression of genes that are known to be involved in stress response and putative tolerance, might prove to

be a faster track towards improving crop varieties. Nevertheless, the task of generating transgenic cultivars is not only limited to the success in the transformation process, but also proper incorporation of the stress tolerance genetic material. Evaluation of the transgenic plants under

stress conditions and understanding the physiological effect of the inserted genes at the whole plant level is still the major challenges to overcome (Pooja et al., 2008). Following plant breeding for salt stress tolerance in plants, genetic engineering seems to be at the moment, the only



method for efficient improvement in abiotic stress control. Some recent advances in genetic engineering for abiotic stress control in plants are represented in Table 1. Genes responsible for stress control were cloned from a wide variety of organism and studied for expression or over-expression in either model plant like *Arabidopsis*, tobacco or some other plants like rice, maize and barley as the most studied ones. Tae-Ho et al. (2009) studied the expression of *PepRSH* gene in *Escherichia coli* and provided wide information about its controlled activity due to the easy manipulation and short life cycle of *E. coli* (Tables 1 and 2) cells. Information about the abiotic stress control and interaction that may happen in the cell is important before gene transformation and analysis in a targeted higher plant. Meanwhile, multiple genes were successfully transformed to the control of abiotic stress in plant with substantial information on their mechanisms; this include the *Trichoderma harzianum* hsp70, *L. maritima* *AtbZIP24*, wheat *TaVP1*, *TaVP2* and *TaVP3*, *Arabidopsis* *CRT/DRE* binding factor 1 (CBF1), *Arabidopsis* *AVP1*, 1 tonoplast intrinsic protein (TIP) *GhTIP1*, rice *OrbHLH2*, rice *AtNDPK2*, pepper *CaXTH3*, *Arabidopsis* *DREB2C*, *Arabidopsis* *ftl1-1D/ddf1*, rice *OsABF2*, emmer wheat (*Triticum turgidum* ssp. *dicoccoides*) *TdicDRF1*, barley *HvTIP1;2*, *HvTIP2;1*, *HvTIP2;2*, *HvTIP2;3* and *HvTIP4;1*, *Arabidopsis* *RAP2.6* and yeast *YCF1* genes. The studied associated genetic patterns provides bright insight useful for the application of the gene in plant abiotic stress control and improved metabolism for a better stress-resistance leading to a large productivity. Some other genes: *Brassica juncea*  $\gamma$ -TMT, (CaMV) 35S promoter cauliflower mosaic virus, *Oryza sativa* miRNAs, maize (*Zea mays* L.) *ZmCO16.1*, *CaBI-1* from the cDNA library of heat-stressed hot pepper (*Capsicum annuum*), cypress seedlings (*Cupressus simper-virens* L.) *cyplp009*; *cyplp005*; *cyplp006*; *cyplp017*; *cyplp010*; *cyplp020*; *cyplp012*; *cyplp002*; *cyplp007*; *cyplp019*; *cyplp003*; *cyplp016*; *cyplp008*; *cyplp015*; *cyplp021*, *Bruguiera* Bg70 and *cyc02* *Gymnorhiza*, pea *APX* and cassava *CuZnSOD*, maize kernel *SAT41* and *Arabidopsis* *MBF1a*, *Hevea brasiliensis* *HbMT2*, apple *cyMDH*, *Panax ginseng* *PgCat1*, *Citrus unshiu* *CitERF*, rice *OsAP2LP* and *Tamarix hispida* *ThbZIP1*, were extensively studied, but still provided less information on the patterns that could affect their expression and/or overexpression. Furthermore, research on abiotic control genes overexpression is gaining an advantage on the study of expressed genes. This suggests that, over-expression of genes under the control of abiotic stress may efficiently control the stresses, although, it is still of huge interest to bring out discovery of abiotic stress-related gene expression in organisms. One specific molecule, trehalose (Kidd and Devorak, 1994; Herdeiro et al., 2006) is present in a wide range of organisms and expressed under stress conditions. Some genes involve in its metabolism were studied for their overexpression and found to over-

produce trehalose in the control of abiotic stress.

## TREHALOSE FOR ABIOTIC STRESS CONTROL IN PLANT

All major crops are sensitive to abiotic stress and sophisticated approaches for solving crops exposed to abiotic stress are therefore needed. These will require the unraveling of additional stress-associated gene resources, from crop plants, model plant and even microorganism species that are highly stress-tolerant. Most of the recent molecular studies on plant stress tolerance have used *Arabidopsis*, rice, maize, tobacco, Cypress canker, fescue, barley, Alfalfa, etc, and the microorganism *E. coli* (in addition to a limited number of microorganism). All these wide range of organisms are viewed as a potential source of trehalose. It is one specific multifunctional molecule commercially produced and that has also been successfully applied for the control of abiotic stress in yeast, bacteria and fungi as well as in plant metabolism and is still considered to be a great challenge in the field of genetic engineering (Li et al., 2008; nez-Esparza et al., 2009; Rohan et al., 2009). Trehalose expression and activity have been successfully studied in yeast as well as many other microorganisms (Herdeiro et al., 2006; Qian et al., 2010) in the control of stress response. Despite its application in industries as active stabilizer of enzymes, proteins, biomasses, pharmaceutical preparations, organs for transplantation, agri-food and cosmetic, and because of its interesting properties (Kidd and Devorak, 1994), researchers went further to study the effect of the expression of trehalose genes in plants. *otsA*, *otsB*, *TPS*, *TPS1* and *TPP* from *E. coli* and yeast were transformed in tobacco, potato and rice with improved growth under stress conditions (Goddijn et al., 1997; Paul et al., 2001; Holmstrom et al., 1996; Garg et al., 2002; Goddijn and Van, 1999; Romero et al., 1997). Culture media supplemented with trehalose or sucrose presented different growth abilities; protocorm-like body (PLB) derived from *Phalaenopsis* and *Doritaenopsis* cultivars, were grown on solidified trehalose amended new phalaenopsis (NP) medium and showed higher proliferation rate than on NP medium containing sucrose. *Phalaenopsis* and *Doritaenopsis* protocorm-like-bodies proliferation rates on solidified trehalose media are almost two times higher than those on sucrose media after 8-week culture (Tsu-Hwie et al., 2006).

Recently, only few reports up till now have developed the application of trehalose in plants abiotic stress control. Exogenous trehalose functioned as an elicitor of genes involved in biotic and abiotic stress. Carbohydrate alterations produced by trehalose biosynthesis can be linked to the stress response. Carolina and Francisco (2005) found that, the yeast *TPS1* gene transformed in tomato plants exhibited pleiotropic changes such as thick shoots, rigid dark-green leaves, erected branches and an

aberrant root development, features commonly missed in transformed plant. Additionally, leaves of transgenic TPS1 tomato plants showed that chlorophyll and starch content were higher than the wild-type plants, which was also observed in some transgenic plants in Table 2. Under drought, salt and oxidative stress TPS1 tomato plants improved tolerance than wild type, suggesting that, carbohydrate alterations produced by trehalose biosynthesis may be linked to the stress response. The TPS1 overexpression activity emerged the feasibility of engineering tomato for increased tolerance of abiotic stress, without decreasing productivity, under both stress and non-stress conditions through trehalose biosynthesis (Table 2). Some plants were transformed with the coding sequence of a gene for trehalose-6-phosphate synthase fused to a non-constitutive plant promoter, which allows for temporal, topological or stress-induced control over the expression of the gene. This protects plants against drought, high salinity or temperature extremes and improves the storage properties of harvested plants, including green food stuffs, picked fruits and ornamental plants. Although the interaction of genes are still not well understood, trehalose control genes could even be more useful in plant tissue culture as it is not only a good carbohydrate source, but also good in regulating plant growth and metabolism during stress conditions.

## THE IMPORTANT SUGAR MOLECULE ALPHA-D-GLUCOPYRANOSYL-ALPHA-D-GLUCOPYRANOSIDE

Abiotic stress is the primary cause of plant loss worldwide. Therefore, resolutions from plant biotechnology discussions aiming at overcoming severe environmental stresses need to be quickly and fully implemented, with intensive molecular assisted genetic engineering. Knowledge of how to alleviate this loss is still limited due to the complexity of both the stress condition and plant responses. A number of reports provide novel and more detailed information concerning the involvement of genes and mediated signaling in stress responses. Despite many other plant stress response factors, genes and key transcription factors involved in stress tolerance can induce the specific functions of trehalose (alpha-D-glucopyranosyl-alpha-D-glucopyranoside), its degrading enzyme trehalase and associated genes that have been studied extensively and appear to be species dependent. They are present in a wide variety of organisms, including bacteria, yeast, fungi, insects, invertebrates and plants. The fundamental role of trehalose is to protect proteins and cellular membranes from inactivation or denaturation caused by a variety of stress conditions. In addition, it also has a pivotal role in regulating sugar metabolism and signal transduction pathways involved in cell proliferation and differentiation. Trehalose increases the thermostability and thermoactivity of reverse transcriptase,

proteins, biomasses, pharmaceutical preparations, organs for transplantation, agri-food, cosmetic, plant tissue culture and improving the length of cDNA synthesis. It has an *in vivo* antioxidant activity and also has a potential role as part of a therapeutic strategy for treating Alzheimer's disease. Hopefully, Parkinson's disease, Huntington's disease and amyotrophic lateral sclerosis (ALS) (also referred to as Lou Gehrig's disease, in rats) can also have possible issue of treatment through trehalose effect investigations. Hence, genes involved in the regulation of its metabolism could be viewed as an important tool in genetic engineering for stress control on one hand and disease treatment on the other hand.

In addition, several important comestibles, cosmetics, ornamental plants and many other industrially used crops are affected by abiotic stress. The overall impact of this article is to stimulate many more researchers to utilize a genetic approach, especially with interesting and promising genes combined with a better understood connection of the four complimenting areas of physiology and biochemistry to the genetic component that controls abiotic stress. This could possibly find an answer in trehalose gene control, regulation and expression. Studies of gene-related trehalose metabolism and associated patterns did not only improve the plant metabolism, phenotypes and texture (one of the problems encountered in plants genetically transformed), but in fact, provide plants that are highly resistant to abiotic stress.

## ACKNOWLEDGEMENT

The authors thank Dr. Jiang Wuof the school of Agriculture and Bioengineering, Tianjin University for helpful literacy criticism, discussion and suggestions.

## REFERENCES

- Ai-Ke B, Suo-Min W, Guo-Qiang W, Jie-Jun X, Jin-Lin Z, Chun-Mei W (2009). Overexpression of the *Arabidopsis* H<sup>+</sup>-PPase enhanced resistance to salt and drought stress in transgenic alfalfa (*Medicago sativa* L.). *Plant Sci.* 176: 232–240.
- Alex B, Jean M, Waine C, Andre L, Igor K (2010). Acute but not chronic exposure to abiotic stress results in transient reduction of expression levels of the transgene driven by the 35S promoter. *New Biotech.* 27(1): 70–77.
- Ayalew L, Maki K, Mineo S, Gemechis D (2011). Abiotic stresses modulate expression of major intrinsic proteins in barley (*Hordeum vulgare*). *C. R. Biolog.* 334: 127–139.
- Barrios E (2007). Convergence in Agriculture of Some Asian Countries. ADBI Discussion Paper Asian Development Bank Institute, Tokyo, Japan, ([http://www.adbi.org/files/dp\\_71\\_Agricultural.Growth.Convergence.pdf](http://www.adbi.org/files/dp_71_Agricultural.Growth.Convergence.pdf)). p. 71.
- Beebe SE, Rao IM, Cajiao C, Grajales M (2008). Selection for drought resistance in common bean also improves yield in phosphorus limited and favorable Environments. *Crop Sci.* 48: 582–592.
- Bloem S, Carpenter JE, Hofmeyr JH (2003). Radiation biology and inherited sterility in false codling moth (Lepidoptera: Tortricidae). *J. Econ. Ent.* 96: 1724–1731.
- Boyer JS (1982). Plant productivity and environment. *Sci.* 218: 443-448.
- Bray EA, Bailey-Serres J, Weretilnyk E (2000). Responses to abiotic stresses. In: Gruijsem W, Buchannan B, Jones R (Eds.).

- Biochemistry and Molecular Biology of Plants, American Society of Plant Physiologists. Rockville, MD, pp. 1158–1249.
- Brown RL, Kazan K, McGrath KC, Maclean DJ, Manners JM (2003). A role for the GCC-box in jasmonate-mediated activation of the PDF1.2 gene of *Arabidopsis*. *Plant Physiol.* 132: 1020–1032.
- Bruno JF, Stachowicz JJ, Bertness MD, (2003) Inclusion of facilitation into ecological theory. *Trends in Ecol.Evol.* 18: 119–125.
- Carolina C, Francisco ACM (2005). Tomato abiotic stress enhanced tolerance by trehalose biosynthesis. *Plant Sci.* 169: 75–82.
- Cuartero J, Bolarin MC, Asins MJ, Moreno V (2006). Increasing salt tolerance in the tomato. *J. Exp. Bot.* 5: 1045–1058.
- Deng-Di L, Fu-Ju T, Ze-Ting Z, Yang L, Yong Z, Yan-Feng W, Xue-Bao L (2009). A cotton gene encodes a tonoplast aquaporin that is involved in cell tolerance to cold stress. *Gene*, 438: 26–32.
- Eun SS, Jia G, Young HK, Joon-Hyeong C, Chun KL, Jang HH, Myeong-Hyeon W (2007). Regulations of marker genes involved in biotic and abiotic stress by overexpression of the AtNDPK2 gene in rice. *Bioch. Biophys. Res. Commun.* 363: 126–132.
- Eun-Ji K, Won-Yong S, Youngsook L, Kyoung HK, Kideok K, Namhyun C, Kwang-Won L, Suk-Whan H, Hojoung L (2006). Expression of yeast cadmium factor 1 (YCF1) confers salt tolerance to *Arabidopsis thaliana*. *Plant Sci.* 170: 534–541.
- Fernando TM, Jordi C (2004). Do positive interactions increase with abiotic stress? A test from a semi-arid steppe. *The Royal Soc.* 271: 331–333.
- Garg AK, Kim JK, Owens TG, Ranwala AP, Choi YD, Kochian LV, Wu RJ (2002). Trehalose accumulation in rice plants confers high tolerance levels to different abiotic stresses. *Proc. Nat. Acad. Sci. USA.* 99: 15898–15903.
- Goddijn OJM, Theo CV, Eline V, Ronny WHH, Krutwagen, Peggy THM De G, Juul P, Kees VD, Anne SP, Brigitte D, Jan P (1997). Inhibition of trehalase activity enhances trehalose accumulation in transgenic plants. *Plant Physiol.* 113: 181–190.
- Goddijn OJM, Van DK (1999). Trehalose metabolism in plants. *Trends Plant Sci.* 4: 315–319.
- Gu YQ (2002). Tomato transcription factors Pti4, Pti5, and Pti6 activate defense Responses when expressed in *Arabidopsis*. *Plant Cell* 14: 817–831.
- Hanhong B, Eliot H, Richard S (2005). Exogenous trehalose promotes non-structural carbohydrate accumulation and induces chemical detoxification and stress response proteins in *Arabidopsis thaliana* grown in liquid culture. *Plant Sci.* 168: 1293–1301.
- Herdeiro RS, Pereira MD, Panek AD, Eleutherio ECA (2006). Trehalose protects *Saccharomyces cerevisiae* from lipid peroxidation during oxidative stress. *Biochim. Biophys. Acta*, 1760: 340–346.
- Hervé C, Ary H (2010). Gene and protein expression of *Drosophila* Starvin during cold stress and recovery from chill coma. *Insect Biochem. Mol. Biol.* 40: 425–428.
- Holmstrom KO, Mantyla E, Welin B, Mandal A, Paiva ET, Tunnela OE, Londesborough J (1996). Drought tolerance in tobacco. *Nat.* 379: 683–684.
- Hong-Gyu K, Joonki K, Bohwa K, Hana J, Sun HC, Eun KK, Hyo-Yeon L, Pyung OL (2011). Overexpression of FTL1/DDF1, an AP2 transcription factor, enhances tolerance to cold, drought, and heat stresses in *Arabidopsis thaliana*. *Plant Science*. article in press. (Provide page number).
- Huan C, Jung EH, Chan JL, Dool YK, Sang YL, Chae OL (2010). *Arabidopsis* DREB2C functions as a transcriptional activator of HsfA3 during the heat stress response. *Biochem. Biophys. Res. Commun.* 401: 238–244.
- Hui-li L, Xiao-yan D, Yun-yuan X, Kang C (2007). Over-expression of OsUGE-1 altered raffinose level and tolerance to abiotic stress but not morphology in *Arabidopsis*. *J. Plant Physiol.* 164: 1384–1390.
- James ED, Ruth CM (2009). Evaluation of reference genes for quantitative RT-PCR in *Lolium temulentum* under abiotic stress. *Plant Sci.* 176: 390–396.
- Jiahong Z, Quanqi Z, Rui W, Zhili Z (2010). HbMT2, an ethphon-induced metallothionein gene from *Hevea brasiliensis* responds to H<sub>2</sub>O<sub>2</sub> stress. *Plant Physiol. Biochem.* 48: 710–715.
- Jia-Sen Y, Ran W, Jing-Jing M, Yu-Ping B, Ping-Li X, Feng G, Shu-Bo W, Qi-Wei H, Xin-Guo L (2010). Overexpression of *Arabidopsis* CBF1 gene in transgenic tobacco alleviates photo inhibition of PSII and PSI during chilling stress under low irradiance. *J. Plant Physiol.* 167: 534–539.
- Jing Z, Fei L, Jin-lan WG, Yun M, Kang CG, Yun-yuan X (2009). Basic helix-loop-helix transcription factor from wild rice (OrbHLH2) improves tolerance to salt-and-osmotic stress in *Arabidopsis*. *J. Plant Physiol.* 166: 1296–1306.
- Kasuga M, Liu Q, Miura S, Yamaguchi-Shinozaki K, Shinozaki K (1999). Improving plant drought salt, and freezing tolerance by gene transfer of a single stress-inducible transcription factor. *Nat. Biotech.* 17: 287–291.
- Kidd G, Devorak J (1994). Trehalose is a sweet target for agbiotech. *Bio/Tech.* 12: 1328–1329.
- Lane A, Jarvis A (2007). Changes in Climate will modify the Geography of Crop Suitability: Agricultural Biodiversity can help with Adaptation. *J.* 4: 1–12.
- Li BQ, Zhou ZW, Tian SP (2008). Combined effects of endo- and exogenous trehalose on stress tolerance and biocontrol efficacy of two antagonistic yeasts. *Biol. Cont.* 46: 187–193.
- Luca P, Paolo B, Ari MH, Nicola LP (2009). Genotype-specific regulation of cold-responsive genes in cypress (*Cupressus sempervirens* L.). *Gene*, 437: 45–53.
- Lynch JP, Beebe SE (1995). Adaptation of beans (*Phaseolus vulgaris* L.) to low phosphorus availability. *Horticult. Sci.* 30: 1165–1171.
- Ma Hl, Zhou Hl, Zhang HY, Zhao J (2010). Cloning and Expression Analysis of an AP2/ERF Gene and Its Responses to Phytohormones and Abiotic Stresses in Rice. *Rice Sci.* 17(1): 1–9.
- Maestre FT, Bautista S, Cortina J (2003). Positive, negative, and net effects in grass–shrub interactions in Mediterranean semiarid grasslands. *Ecology*, 84: 3186–3197.
- Maestre FT, Jordi C, Susana B (2007). Mechanisms underlying the interaction between *Pinus halepensis* and the native late-successional shrub *Pistacia lentiscus* in a semi-arid plantation. *Ecography*, 27: 776–786.
- Marta MB, Rosa H, Rosa EC, Santiago G, Carlos N, Enrique M (2009). Transgenic expression of the *Trichoderma harzianum* hsp70 gene increases *Arabidopsis* resistance to heat and other abiotic stresses. *J. Plant Physiol.* 167(8): 659–665.
- Md Amir H, Jung-Il C, Muho H, Chul-Hyun A, Jong-Seong J, Gynheung A, Phun BP (2010). The ABRE-binding bZIP transcription factor OsABF2 is a positive regulator of abiotic stress and ABA signaling in rice. *J. Plant Physiol.* 167: 1512–1520.
- Min-Jung K, Gah-Hyun L, Eun-Seon K, Chang-Beom K, Kwang-Yeol Y, Jin-An J, Myung-Chul L, Cheol SK (2007). Abiotic and biotic stress tolerance in *Arabidopsis* overexpressing the Multiprotein bridging factor 1a (MBF1a) transcriptional coactivator gene. *Bioch. Biophys. Res. Commun.* 354: 440–446.
- Mittler R (2006). Abiotic stress, the field environment and stress combination. *Trends In Plant Sci.* 11: 15–19.
- Mohammad IL, Naheed ZI, Seong RK, Choo BH (2009). A BAXinhibitor-1 gene in *Capsicum annuum* is induced under various abiotic stresses and endows multi-tolerance in transgenic tobacco. *J. Plant Physiol.* 166: 1685–1693.
- Mohd AY, Deepak K, Ravi R, Reto JS, Merope T-M, Govindjee, Neera BS (2010). Overexpression of  $\gamma$ -tocopherol methyl transferase gene in transgenic *Brassica juncea* plants alleviates abiotic stress: Physiological and chlorophyll a fluorescence measurements. *Biochim. Biophys. Acta.* 1797(8): 1428–1438.
- Munkhbaatar P, Yeon-Ju K, Myung KK, Rama KP, Deok-Chun Y (2010). Isolation of a novel catalase (Cat1) gene from *Panax ginseng* and analysis of the response of this gene to various stresses. *Plant Physiol. Biochem.* 48: 451–460.
- Nez-Esparza MM, Nez-Vicente EM, Lez-Parraga P, Ros JM, Arrubia PA, Elles J (2009). Role of trehalose-6Pphosphatase (TPS2) in stress tolerance and resistance to macrophage killing in *Candida albicans*. *Int. J. Med. Microbiol.* 299: 453–464.
- Oksoon Y, Olga VP, Ulrike S, Ines L, Karl-Josef D, Dortje G (2009). The *Arabidopsis* basic leucine zipper transcription factor AtbZIP24 regulates complex transcriptional networks involved in abiotic stress resistance. *Gene*, 436: 45–55.
- Orlene GP, Ha TN, Peter S, Leipner J (2009). ZmCOI6.1, a novel, alternatively spliced maize gene, whose transcript level changes under abiotic stress. *Plant Sci.* 176: 783–791.

- Paul M, Pellnyk T, Goddijn O (2001). Enhancing photosynthesis with sugar signals. *Trends Plant Sci.* 6: 197–200.
- Pooja BM, Vadez V, Kiran KS (2008). Transgenic approaches for abiotic stress tolerance in plants: retrospect and prospects. *Plant cell reports*, 27: 411–24.
- Pugnaire FI, Luque MT (2001). Changes in plant interactions along a gradient of environmental stress. *Oikos*, 93: 42–49.
- Qian L, Sheng-Hua Y, Ming-Guang F (2010). Characterization of *Beauveria bassiana* neutral trehalase (BbNTH1) and recognition of crucial stress-responsive elements to control its expression in response to multiple stresses. Article in press.
- Qiang Z, Jiantao Z, Xiaoshu G, Jianhua T, Langtao X, Wenbin L, Hongxia Z (2010). The *Arabidopsis* AP2/ERF transcription factor RAP2.6 participates in ABA, salt and osmotic stress responses. *Gene*, 457: 1–12.
- Ragan MC, Brooker RW, Philippe C, Zaal K, Christopher JL, Richard M, Leonardo P, Francisco IP, Beth N, Erik TA, Cristina A, David K, Bradley JC (2002). Positive interactions among alpine plants increase with Stress. *Nature*, 417: 844–848.
- Rao IM (2002). Role of physiology in improving crop adaptation to abiotic stresses in the tropics: The case of common bean and tropical forages. In: M. Pessarakli (ed). *Handbook of Plant and Crop Physiol.*, Marcel Dekker, Inc., New York, USA, pp. 583–613.
- Ribaut JM, Jiang C, Gonzalez-de-Leon D, Edmeades GO, Hosington DA (1997). Identification of quantitative trait loci under drought conditions in tropical maize, 2. Yield components and marker assisted selection strategies. *Theor. Appl. Genet.* 94: 887–896.
- Rohan GTL, Maryn L, Kasia R, Robert DT, Richard PO, Peter SS (2009). Trehalose biosynthesis is involved in sporulation of *Stagonospora nodorum*. *Fungal Genet. Biol.* 46: 381–389.
- Romero C, Belles JL, Vaya JL, Serrano R, Culiñez-Macia FA (1997). Expression of the yeast trehalose-6-phosphate synthase gene in transgenic tobacco plants: pleiotropic phenotypes include drought tolerance. *Planta*, 201: 293–297.
- Sang-Hoon L, Nagib A, Ki-Won L, Do-Hyun K, Dong-Gi L, Sang-Soo K, Suk-Yoon K, Tae-Hwan K, Byung-Hyun L (2007). Simultaneous overexpression of both CuZn superoxide dismutase and ascorbate peroxidase in transgenic tall fescue plants confers increased. *J. Plant Physiol.* 164: 1626–163.
- Seok KC, Jee EK, Jong AP, Tae JE, Woo TK (2006). Constitutive expression of abiotic stress-inducible hot pepper CaXTH3, which encodes a xyloglucan endotransglucosylase/hydrolase homolog, improves drought and salt tolerance in transgenic *Arabidopsis* plants. *FEBS Letters*, 580: 3136–3144.
- Shota E, Yuichi T (2009). Identification of salt tolerance genes from the mangrove plant *Bruguiera gymnorhiza* using *Agrobacterium* functional screening. *Plant Sci.* 176: 272–278.
- Sravan K, Greg BC, Swathi AT, Craig H, Stanley JR, Pulugurtha BK (2008). Ectopic expression of an annexin from *Brassica juncea* confers tolerance to abiotic and biotic stress treatments in transgenic tobacco. *Plant Physiol. Biochem.* 46: 1019–1030.
- Stuart L, Emel D, Bala AA, Hikmet B (2011). The drought response displayed by a DRE-binding protein from *Triticum dicoccoides*. *Plant Physiol. Biochem.* 49: 346–351.
- Tae-Ho K, Sung HO, Donghern K, Seok-Cheol S, Myung OB, Jeong SS (2009). Molecular characterization of a biotic and abiotic stress resistance-related gene RelA/SpoT homologue (PepRSH) from pepper. *Plant Sci.* 176: 635–642.
- Thung M, Rao IM (1999) Integrated management of abiotic stresses. In: S. P. Singh (ed.), *Common Bean Improvement in the Twenty-First Century*. Kluwer Academic Publishers, Dordrecht. The Netherlands. pp. 331–370.
- Tielborger K, Kadmon R (2000). Temporal environmental variation tips the balance between facilitation and interference in desert plants. *Ecology*, 81: 1544–1553.
- Tsu-Hwie AL, Jing-Jing L, Rey-Yuh W (2006). The effects of using trehalose as a carbon source on the proliferation of *Phalaenopsis* and *Doritaenopsis* protocorm-like bodies. *Plant Cell, Tiss. Organ Cult.* 86: 125–129.
- Villalobos MA, Bartels D, Iturriaga G (2004). Stress tolerance and glucose insensitive phenotypes in *Arabidopsis* overexpressing the CpMYB10 transcription factor gene. *Plant Physiol.* 135: 309–324.
- Vogel JT, Zarka DG, Van Buskirk HA, Fowler SG, Thomashow MF (2005). Roles of the CBF2 and ZAT12 transcription factors in configuring the low temperature transcriptome of *Arabidopsis*. *Plant J.* 41: 195–211.
- Wall DH, Moore JC (1999). Interactions underground. *BioSci.* 49: 109–117.
- Wei Z, Ying L, Bao-Cun Z, Rong-Chao G, Yin-Zhu S, Gang W, Zhan-Jing H (2009). Overexpression of TaSTRG gene improves salt and drought tolerance in rice. *J. Plant Physiol.* 166: 1660–1671.
- Whitford WG (2002). *Ecology of desert systems*, New York, NY: Academic Press.
- Xiao-Yan Y, Jin-Xia X, Xiao-Peng, Yong-Zhong L, Shu-Ang P (2011). Isolation of a citrus ethylene-responsive element binding factor gene and its expression in response to abiotic stress, girdling and shading. *Sci. Hortic.* 127: 275–281.
- Xinyu J, Lin Z, Guanglin L, Liang Z, Xiujie W, Xiaofeng C, Xiaohua F, Fan C (2010). Identification of novel stress-regulated microRNAs from *Oryza sativa* L. *Genomics*, 95: 47–55.
- Yucheng W, Caiqiu G, Yanan L, Chao W, Chuanping Y, Guifeng L (2010). A novel *bZIP* gene from *Tamarix hispida* mediates physiological responses to salt stress in tobacco plants. *J. Plant Physiol.* 167: 222–230.
- Yuezhi W, Haibin X, Guangxiang Z, Huilan Z, Lixia Z, Zhengzhi Z, Caiqin Z, Zhengqiang M (2009). Expression and responses to dehydration and salinity stresses of V-PPase gene members in wheat. *J. Genet. Genomics*, 36: 711–720.
- Yu-Xin Y, Qing-Long D, Heng Z, Chun-Xiang Y, Yu-Jin H (2011). The functions of an apple cytosolic malate dehydrogenase gene in growth and tolerance to cold and salt stresses. *Plant Physiol. Biochem.* 49: 257–264.