

Full Length Research Paper

Molecular cloning and characterization of a cytoplasmic cyclophilin gene in sugarcane

Youxiong Que¹, Jinxian Liu^{1,2}, Liping Xu^{1*}, Jinrong Guo¹, Jingsheng Xu¹ and Rukai Chen¹

¹Key Lab of Sugarcane Genetic Improvement, Ministry of Agriculture, Fujian Agriculture and Forestry University, Fuzhou 350002, Fujian Province, China.

²Department of Tea Science and Biology, Wuyi University, Wuyishan 354300, Fujian Province, China.

Accepted 20 May, 2011

Cyclophilins are ubiquitous proteins with an enzymatic activity of peptidyl-prolyl cis-trans isomerase (PPIase), which play important roles in a variety of stress responsiveness. In this study, we reported the cloning and characterization of a full-length cytoplasmic cyclophilin gene in sugarcane. Sequence analysis showed the cDNA of this gene (GenBank accession number:GQ246462), termed as *Sc-CyP*, was 904 bp long, including a 519 bp complete ORF, the 5' UTR of 74 bp and 3'UTR of 311 bp, plus a typical AATAA motif and poly (A) tail. It encoded the 172 amino acid polypeptide with a molecular weight of 18.4 KD and the isoelectric point of 8.68. The *Sc-CyP* encoding protein had the conserved site Trp128 (W128) ubiquitous of all cyclophilins in eukaryotes and the KSGKPLH48-54 region specific to cytoplasmic cyclophilins in plants. SDS-PAGE analysis and PPIase assay revealed that the expression product, with PPIase activity, was a fusion protein with a molecular weight about 25 and 18.4 kD of *Sc-CyP* plus 7 kD of His · Tag peptides. In real-time qPCR analysis, the *Sc-CyP* gene showed induced expression under PEG, NaCl, SA and H₂O₂ stresses, indicating it a stress-related gene for drought and salt stress, signal transduction and disease resistance response in sugarcane.

Key words: Sugarcane (*Saccharum officinarum*), cyclophilin, PPIase, real-time quantitative PCR.

INTRODUCTION

Immunophilins, including cyclosporin A, FK506 and rapamycin are defined as receptors for immunosuppressive drugs. FK506- and rapamycin-binding proteins were abbreviated as FKBP, while the cyclosporin A was referred to as cyclophilins (CyPs) (He et al., 2004). Cyclophilins constituted a family of abundant and highly conserved proteins which appeared to be ubiquitous in organisms, ranging from bacteria to plants and animals (Galat, 1999). The first cyclophilin was identified as the specific target of the immunosuppressant cyclosporin A (CsA) in mammalian-cells (Handschumacher

T et al., 1984). Since then, a series of cyclophilins had been identified and annotated in higher plants, especially in monocotyledonous plants (Opiyo and Moriyama, 2009). These proteins belonged to the cluster of immunophilins which possessed an endogenous enzymatic activity. They exhibited peptidyl-prolyl cis-trans isomerase (PPIase or rotamase) activity, which catalyzed cis-trans isomerization of the amide bond between a proline residue and the preceding amino acid residue and they could also participate in the rate-limiting step in protein folding whose process could be accelerated by the rotamase of cyclophilin (Brandts et al., 1975).

To date, the diverse functions of cyclophilins have been reported (He et al., 2004; Opiyo and Moriyama, 2009). In animal cells, the CyP/CsA complex was found to inhibit the Ca²⁺-dependent phosphatase activity of the calcineurin protein (Schreiber, 1991). In yeast, genetic analysis of CyP-deficient mutants revealed that cyclophilin was critical for the recovery from heat shock (Sykes et al., 1993). Besides, cyclophilins were reported to play a role in a wide variety of processes, including cell

*Corresponding author. E-mail: queyouxiong@yahoo.com.cn or xlpmail@yahoo.com.cn. Tel: 0086-591-83772604. Fax: 0086-591-83851742.

Abbreviations: PPIase, Peptidyl-prolyl cis-trans isomerase; IPTG, isopropyl-beta- D-thiogalactopyranoside; ORF, open reading frame; UTR, untranslated region; HR, hypersensitive response; SAR, systemic acquired resistance.

division (Faure et al., 1998), transcription regulation (Rycyzyn and Clevenger, 2002), signaling (Brazin et al., 2002) and so on. In plants, cyclophilin genes existed as a large gene family, with 29 members in *Arabidopsis thaliana* (Romano et al., 2004). Plant cyclophilins required various proteins to execute different kinds of physiological processes, with a striking feature of various expression modes in response to different abiotic or biotic stresses (Godoy et al., 2000; Sharma and Singh, 2003; Opiyo and Moriyama, 2009). Marivet et al. (1994) found that the transcription of a bean cyclophilin gene was up-regulated due to environmental or pathogenic stresses, indicating this gene may be important for signal transduction or protein folding under these stresses (Marivet et al., 1994). Research on a cyclophilin gene in maize revealed that it expressed at a basal level in all tissues and the expression could be enhanced by treatment of salicylic acid (SA), which acted as an important signal molecule responsible for induction of plant defense response (Marivet et al., 1995). It was also demonstrated that CyP/CsA complex may reactivate Ca^{2+} -inactivated K^+ channel by inhibiting phosphatase activity in guard cells of *Vicia faba*, leading to changes in guard cell turgor, which suggested that a conserved Ca^{2+} -dependent signaling pathway was also existed in plants (Luan et al., 1993).

To our knowledge, there is not any report on the isolation and characterization of a sugarcane cyclophilin gene, which highlights the meanings of the ongoing exploitation of cyclophilin genes in sugarcane. In this study, we reported the cloning and characterization of a cyclophilin gene in sugarcane, aimed to lay the foundation for further research and application of this gene in sugarcane molecular breeding through genetic engineering.

MATERIALS AND METHODS

Materials and treatment

Sugarcane varieties FN 22, sugarcane stem full-length cDNA library, *E. coli* DH5 α and BL21 (DE3) and the prokaryotic expression vector pET29a (+) were provided by key lab of sugarcane genetic improvement, Ministry of Agriculture, P. R. China. *EcoR* I, *Sal* I, T₄ DNA ligase, Taq enzyme, DNA and protein molecular marker, SYBR[®]Premix Ex Taq[™] were from TaKaRa (Japan); Hepes, Triton X100, chymotrypsin, IPTG (isopropyl-beta-D-thiogalactopyranoside) and reverse transcription kit from Promega Corporation (USA); FK506 and rapamycin from Sigma (USA) and ABI PRISM7500 real-time PCR system was used.

Strong and evenly growing sugarcane stalks were soaked and sterilized in water with 0.1% potassium permanganate for 24 h and then planted into autoclaved fine sand in 10 trays, 40 shoots in each tray. These trays were then placed into the illumination incubator where each day they were kept under light for 12 h at 29°C and dark for 12 h at 26°C with a light intensity of 440 mol m⁻² s⁻¹. When growing to the height of 15 cm, they were transferred into the greenhouse for cultivation and sprayed with Hongland nutrient solution (Zhang and Qu, 2003). In the six leaves period, evenly growing stalks were selected and the sand was removed. After

rehydration for 2 days, the stalks were cultivated in Hongland nutrient solution for one week and then treated as follows. For the first group, 5.0 m mol·L⁻¹ SA and 10 mmol·L⁻¹ H₂O₂ were sprayed onto the leaves, respectively (Que et al., 2009a). For the second group, the sugarcane roots were soaked in PEG8000 solution (25%) and NaCl solution (250 mmol·L⁻¹), respectively (Zhang, 2007). The sampling times were 0, 12, 24, 36, 48, 60 and 72 h for SA and H₂O₂ treatment, 0, 3, 6, 12, 24 and 72 h for PEG treatment and 0, 3, 18, 24, 48 and 72 h for NaCl treatment. All the samples collected were immediately fixed in liquid nitrogen and stored in a refrigerator at -85°C until RNA extraction.

Obtaining and sequence analysis of a full-length cDNA sequence of *Sc-CyP* gene

The large-scale sequencing and bioinformatics analysis were conducted for sugarcane stem full-length cDNA library (Private bulletin). Those clones which showed high homology to cyclophilin genes in NCBI database were selected and sequenced completely to obtain the full-length cDNA sequence of a cyclophilin gene in sugarcane.

For the full-length cDNA sequence of the cyclophilin gene, ORF was predicted with the online tool ORF Finder from NCBI (<http://www.ncbi.nlm.nih.gov/gorf/gorf.html>). The BLAST program in NCBI was used for the homology analysis. Besides, the ProtParam (<http://cn.expasy.org/tools/protparam.html>) was applied to analyze the basic property of the encoding protein. Multiple sequence alignment and phylogenetic analysis were performed with DNAMAN software (version 6.0) and MEGA3.1. InterProScan (<http://ebi.ac.uk/InterProScan>) and SMAR (<http://smart.embl-heidelberg.de>) were used to analyze the putative domain of the encoding protein. The gene sub-cellular localization was carried out with SubLoc V1.0 (<http://www.bioinfo.tsinghua.edu.cn/SubLoc/>) and the signal peptide prediction was performed with SignalP3.0 (<http://www.cbs.dtu.dk/services/SignalP/>). SOPMA was adopted for the secondary protein structure prediction (http://npsa-pbil.ibcp.fr/cgi-bin/npsaautomat.pl?page=npsa_sopma.html) and the analysis of the repetitive sequence of the amino acid was performed with the REP searcher (<http://www.embl-heidelberg.de/~andrade/papers/rep/search.html>).

Construction of prokaryotic expression vector of *Sc-CyP* gene

pET29a (+) was used as the prokaryotic expression vector and the specific primer pairs amplifying the ORF region of *Sc-CyP* gene were designed. The primer sequences were as follows, of which the underlined parts were the restriction endonuclease sites of *EcoR* I and *Sal* I, respectively. *Sc-CyP*-F: 5'-TATGGATCCATGGCGA ACCCGCG-3'; *Sc-CyP*-R: 5'-TTAGTTCGACGCTGAGCTGGCCGC-3'.

The primers were synthesized by Shanghai Sangon Biological Engineering Technology and Services CO., Ltd. With plasmid DNA extracted from the library clone containing *Sc-CyP* gene as the template, the PCR amplification was carried out. The PCR reaction was composed of 5.0 μ l 10 \times PCR buffer, 4.0 μ l 2.5 mmol·L⁻¹ dNTPs; 2.0 μ l 10 μ mol·L⁻¹ forward and reserves primers respectively; 2.0 μ l plasmid DNA; 0.25 μ l Taq enzyme (5 U· μ l⁻¹); ddH₂O was added as the supplement and the final volume was 50 μ l. PCR amplification program was pre-denaturation for 5 min at 94°C; denaturing for 30 s at 94°C, annealing for 30 s at 55°C and extending for 1 min at 72°C, 30 cycles; followed by finally extending for 10 min at 72°C. When the reaction finished, 1% agarose gel electrophoresis was performed and the target PCR product was recovered. After receiving the double digestion with *EcoR* I and *Sal* I, the obtained ORF sequence was ligated by T4 DNA ligase into pET29a (+) which had also executed the double digestion with *EcoR* I and *Sal* I. The

recombinant was transferred into *E. coli* DH5 α . Plasmid DNA of clones selected by blue/white colony screening was extracted. The positive clone validated by PCR and double digestion was just the prokaryotic expression vector which termed as pET29a- ScCyP (Que et al., 2009b).

SDS-PAGE analysis of prokaryotic expression product and enzyme activity assay

pET29a-ScCyP and empty pET29a(+) were both transformed into *E. coli* BL21 (DE3). The single colony was inoculated into the LB medium containing the ampicillin (100 $\mu\text{g}\cdot\text{ml}^{-1}$) for overnight shake culture at 150 r/min at 37°C. In the following day, 1% of the medium was inoculated into the LB medium containing of 100 $\mu\text{g}\cdot\text{ml}^{-1}$ ampicillin and shake cultured. When OD600 reached 0.4 to 0.6, 1.0 ml of LB liquid medium was collected as the control and the remaining medium was added with IPTG with the final concentration 1.0 $\text{mmol}\cdot\text{L}^{-1}$. The LB medium with pET29a-ScCyP was induced for 2, 4 and 6 h, respectively at 37°C and 1.0 ml medium was collected at each time point; the LB medium with empty pET29a (+) was induced with IPTG for 6 h and 1.0 ml corresponding medium was collected. The medium collected was used for the SDS-PAGE with 25 μl sample loading and also for peptidyl-prolyl-cis-trans-isomerase (PPIase) assay according to Blecher et al. (1996). When the electrophoresis finished, the gel was colored with Coomassie brilliant blue and imaged (Que et al., 2009b).

Expression profile of Sc-CyP gene under various exogenous stresses

25S rRNA was selected as the control gene in the Real-time qPCR analysis (Que et al., 2009c). According to the sequence of Sc-CyP gene and 25S rRNA (BQ536525), two pairs of real-time qPCR primers were designed with GenScript online PCR primers designs tool (http://www.genscript.com/cgi-bin/tools/primer_genscript.cgi). The forward and reverse primers of Sc-CyP gene were 5'-ACTTCTCGCCGTAGATGGAC-3' and 5'-CACTACAAGGGC TCCACCTT-3', respectively. For the 25S rRNA, they were 5'-GCAGCCAAGCGTTCATAGC-3' and 5'-CC TATTGGTGGGTG AACAAATCC-3', respectively.

The 20 μl reverse transcription system contained 4.0 μl 25 $\text{mmol}\cdot\text{L}^{-1}$ MgCl₂, 2.0 μl 5 \times RT buffer solution, 2.0 μl 10 $\text{mmol}\cdot\text{L}^{-1}$ dNTPs, 0.5 μl Rnase inhibitor, 0.5 μl random primers (0.5 μg), 15 U AMV reverse transcriptase and 4.0 μl total RNA (1.0 μg) as the template and RNase free H₂O as the supplement. Reverse transcription conditions were as follows: kept at room temperature for 10 min, then incubated at 42°C for 15 min and 95°C for 5 min and then kept at 5°C for 5 min. In real-time qPCR analysis, ABI PRISM7500 real-time PCR system was used. For real-time qPCR amplification, the RT-PCR product was chosen as the template and the volume of the reaction system was 25 μl , including SYBR Primix Ex TaqTM (2 \times), 12.5 μl ; Rox reference dye II, 0.5 μl ; forward and reverse primers (10 $\mu\text{mol}\cdot\text{L}^{-1}$), 0.5 μl , respectively; cDNA template, 2.5 μl and sterile water, 8.5 μl . Three replicas were set for each sample. The PCR reaction conditions were pre-denaturation at 95°C for 10 s and 40 cycles with 94°C 5 s, 60°C 25 s. When the reaction was completed, the melting curve was analyzed. The method of $2^{-\Delta\Delta\text{CT}}$ was adopted to analyze the real-time qPCR results (Livak and Schmittgen, 2001; Que et al., 2009d).

RUSULTS

Cloning and sequence analysis Sc-CyP gene

According to the results of large-scale sequencing and

bioinformatics analysis, one EST sequence which had 95% homology to the sorghum cyclophilin gene (EU722309) was obtained from the sugarcane stem full-length cDNA library. The corresponding library clone of this EST was then completely sequenced and the full-length cDNA sequence of a cyclophilin gene in sugarcane was obtained and termed as Sc-CyP. It had been submitted to the GenBank with the accession number of GQ246462. Sequence analysis showed that the Sc-CyP gene had a full length of 904 bp, ORF (open reading frame) length of 519 bp, 5' UTR (un-translated region) of 74 bp and 3'UTR of 311 bp and the typical AATAA motif and poly(A) tail could also be found in the 3'UTR (Figure 1).

The primary structure of the predicted Sc-CyP protein is shown in Figure 1. The ORF encoded 172 aa with the molecular weight of 18.4 kD and isoelectric point of 8.68. The Sc-CyP encoding protein in sugarcane had a highly conserved functional site Trp128 (W128) present in all cyclophilins in eukaryotes (Liu et al., 1991), and the 7 amino acid residues region of KSGKPLH48-54, which was specific to cytoplasmic cyclophilins in plants, was also found (Lippuner et al., 1994). Protein domain prediction indicated that Sc-CyP encoding protein had the conserved structure domain of cyclophilin (Figure 2). With SubLoc v1.0, it was located in the cytoplasm.

Sequence homology analysis of Sc-CyP gene

The blastn analysis indicated that the homology of the sugarcane cyclophilin gene (Sc-CyP) with the sorghum cyclophilin gene (EU722309) and with the corn peptide prolyl cis-trans isomerase gene (PPI) (EU967603) was 88% (818/920) and 88% (757/856), respectively. It also showed in blastn analysis that the homology of the Sc-CyP gene with that of the cyclophilin genes from sorghum bicolor (ACD93011), *Zea mays* (ACG35201), *Dasypyrum villosum* (ABU56008), *Triticum aestivum* (AAS17067), *Oryza sativa* (AAA57045), *Gerbera* (ABV26711), *Ricinus communis* (CAC80550) and *Picea abies* (CAC81066) was 95% (165/172), 95% (165/172), 87% (149/171), 86% (148/171), 86% (149/172), 83% (144/172), 83% (143/171) and 84% (145/172), respectively. Multiple alignment analysis of amino acid sequences of the cyclophilins encoded by Sc-CyP gene and those from other plant species was performed using DNAMAN software and the results are shown in Figure 3. It indicated that the amino acid sequences of all these cyclophilins had the similar length. What is more, the Sc-CyP encoding protein had rather high homology with the cyclophilins from other plant species, especially in the region KSGKPLH, which should be the key for its enzymatic activity.

The phylogenetic tree of the cyclophilin encoded by Sc-CyP gene and the cyclophilins of other 15 plant species was constructed using the MEGA3.1 software. Figure 4 shows that the Sc-CyP and the cyclophilin of *P. abies*, *Z.*

```

1      ACCAGATCAG ATCACACAAC AGCCGAAGGA AAATTTCCCC CAAAACCCTC CCCAAACCCT
61     AGCTTCGGAT CCCGATGGCG AACCCGCGCG TCTTCTTCGA CATGACCGTC GGCGGCGCCC
          M A N P R V F F D M T V G G A P
121    CGGCGGGGCG GATCGTGATG GAGCTGTACG CCAACGAGGT GCCCAAGACG GCCGAGAACT
          A G R I V M E L Y A N E V P K T A E N F
181    TCCGCGCGCT GTGCACGGGC GAGAAGGGCG TGGGGAAATC GGGGAAGCCG CTCCACTACA
          R A L C T G E K G V G K S G K P L H Y K
241    AGGGCTCCAC CTTTCACCGC GTCATCCCGC AGTTCATGTG CCAGGGCGGC GACTTCACCC
          G S T F H R V I P Q F M C Q G G D F T R
301    GCGGCAACGG CACGGGAGGG GAGTCCATCT ACGGCGAGAA GTTCCCCGAC GAGAAGTTCC
          G N G T G G E S I Y G E K F P D E K F V
361    TGC GCAAGCA CACGGGGCCC GGGGTGCTCT CCATGGCCAA CGCGGGGCCC AACACCAACG
          R K H T G P G V L S M A N A G P N T N G
421    GCTCCCAGTT CTTCGTCTGC ACCGTCGAGA CCCCCTGGCT CGACGGCAAG CACGTCGTCT
          S Q F F V C T V E T P W L D G K H V V F
481    TCGGCCAGGT CGTCGAGGGC ATGGACGTCG TCAAGGCCAT CGAGAAGGTG GGCTCCCGCA
          G Q V V E G M D V V K A I E K V G S R S
541    GCGGCTCCAC CTCCAAGGAG GTC AAGGTCG CTGACTGCGG CCAGCTCAGC TAGATCTCTG
          G S T S K E V K V A D C G Q L S *
601    GTCGTCTCGT CTCGTCTGTC CGGCCGCCCT CCCCTCCGTC ATCGTCGACT CCGCCTGCGT
661    CCGTCCCTTT TCCATTTGCT TCGATCTGAA TAAGATGATG GTGATCTGAG TGGTGGTCTG
721    TGTTGAGTCG TTCAATCATG TGT TGTGTC GCTCGTGGTT TAATTTAGAG GTTTATTTAG
781    GCATGGATCT GCGAATCCCA TGGCGCCTCT GCTTACTTCC TGTTTCACCT GTTATGTTAT
841    GAGATCCATC TCAAGATAAT GCTATGAGAA CTGAAATTCT GTTTAAAAAA AAAAAAAAAA
|901    AAAA

```

Figure 1. Nucleotide sequence of *Sc-CyP* gene and primary structure of *Sc-CyP* protein. Capital and small letters represent amino acid and nucleotide, respectively; *shows stop codon; the underline shows the polyadenylation signal AATAA; the big frame shows the conservative sequence of the plant cytoplasmic cyclophilin; the small frame shows conservative amino acid of cyclophilin in all kinds of organisms.




Figure 2. Predicted domain of *Sc-CyP* protein.

mays, *Sorghum bicolor*, *O. sativa*, *T. aestivum* and *D. villosum* belongs to the same category, indicating that these cyclophilins may evolve from a common ancestor

through different pathways. Besides, the *Sc-CyP* and the cyclophilin of *Z. mays* were classified into one group and that from *D. villosum* and *T. aestivum* were in another




Figure 3. Multiple sequence alignment of cyclophilins isolated from different plant species. Frame shows conservative sequence of the plant cytoplasmic CyP protein; ▲ shows conservative amino acid of CyP protein *S. bicolor* (ACD93011); *T. aestivum*, (AAS17067); *Z. mays*, (ACG35201); *D. villosum*, (ABU56008); *G. hybrid_cultiva*, (ABV26711); *O. sativa*, (AAA57045); *P. abies*, (CAC81066); *R. communis*, (CAC80550).

group, which suggested that the evolutionary relationship between the two within one group was the closest.

Construction of the *Sc-CyP* gene prokaryotic expression vector

After the PCR product of the ORF amplification of *Sc-CyP* gene was recovered, it received the *BamH* I and *Sal* I digestion. The target fragment was then recovered and used to ligate into the pET29a (+) which also received double digestion of *BamH* I-*Sal* I and a positive recombinant was transformed into the *E. coil* DH5a competent cell. Through PCR amplification and double

digestion with *BamH* I+*Sal* I, the fragment with the same size as that of the target fragment could be obtained from the positive clone pET29a-*ScCyP*. The results of the agarose gel electrophoresis are shown in Figure 5, indicating the construction of the prokaryotic expression vector pET29a-*ScCyP* of the *Sc-CyP* gene was successful.

SDS-PAGE analysis of prokaryotic expression product and enzyme activity assay

As shown in Figure 6, the target protein began to express in the second hour with the molecular weight of about 25 kD, which was basically consistent with the deduced




Figure 4. Phylogenetic tree of cyclophilins isolated from different plant species.




Figure 5. Validation of pET29a-ScCyP by PCR and double digestion with *Bam*H I+*Sal* I. A: M, DL2000; Lane 1, negative control; Lane 2, PCR product; B: M, DL2000; 1, digested product with *Bam*H I+*Sal* I.

molecular weight of the fusion protein of 25.4 kD, that was 18.4 kD of Sc-CyP plus 7 kD of His • Tag peptides. Enzyme activity assay demonstrated that the purified prokaryotic expression protein exhibited PPLase activity and this kind of PPLase activity could be inhibited by

FK506 and rapamycin. Though the *Sc-CyP* gene had the successful expression within *E. coli* BL2, this target protein could not be expressed in the empty vector, while no wrong coding or reading-frame shift of amino acid sequences were found, all of which further confirmed the




Figure 6. Expression of pET29a-ScCyP in *E. coli* BL21. M, protein maker; Lane 1-3, pET29a-ScCyP induction for 6, 4 and 2 h respectively; Lane 4, pET29a-ScCyP without induction; Lane 5, pET29a (+) induction for 6 h; Lane 6, pET29a (+) without induction; Lane 7, BL21 without induction.

successful construction of this prokaryotic expression vector.

Expression profile of *Sc-CyP* gene under various exogenous stresses

Real-time qPCR was used to examine the expression profile of *Sc-CyP* gene under different kinds of exogenous stresses. Figure 7 shows that the *Sc-CyP* gene expression was induced during the whole period under all the stresses of PEG, NaCl, SA and H₂O₂, with various expression profiles under different kinds of exogenous stresses. Under PEG stress, the induced expression of *Sc-CyP* gene had the trend of “increasing-declining-increasing-declining”. At the time point of 3 h, the *Sc-CyP* gene expression was strongly induced, reaching more than 24 times that of the control at 0 h. Then the expression began to decrease, which lasted until the time point of 6 h. From the time point of 6 h on, the expression of *Sc-CyP* gene increased again. The peak was found at the time point of 24 h, which was more than 72 times that of the control, and after that it began to drop again. But even at the time point of 72 h, the expression was still 12 times that of the control. Under NaCl stress, the induced expression of *Sc-CyP* gene was strong and just at the time point of 3 h, more than 40 times expression that of the control at 0 h was observed. It reached the peak at 18 h at which the expression was more than 630 times that of the control. Although, it began to decrease later, at the following three time points, it was still more than 97 times that of the control. Under the stress of SA, the expression of *Sc-CyP* gene had the “increasing-declining-increasing” trend. At the time point of 12 h, the induced expression of *Sc-CyP* gene was not very strong. However, it was rather strong at the time point of 24 h with the expression 5.7 times that of the control at 0 h. Although, the expression

began to decrease after the time point of 24 h, it was still more than twice that of the control. It reached the peak at 72 h, which was more than 6 times that of the control. Under the stress of H₂O₂, the induced expression of *Sc-CyP* gene had the trend of “increasing-declining-increasing-declining”. At the time point of 3 h, the *Sc-CyP* gene expression was more than 2.5 times that of the control at 0 h. The peak was found at the time point of 24 h which was more than 6 times that of the control. Although, the expression was induced during the period of 48 to 72 h, it began to decrease after that which was to some extent equal to that of the control at 0 h. Based on the real-time qPCR analysis earlier, it could be concluded that the *Sc-CyP* gene showed induced expression under all the four kinds of exogenous stresses, PEG, NaCl, SA and H₂O₂. This indicated that the *Sc-CyP* gene cloned in this study was stress-related and it may play the protective roles of resistance in sugarcane.

DISCUSSION

Cyclophilin is a kind of intracellular binding protein which exists widely in a variety of biological cells and has high affinity with the cyclosporine A (CsA). According to their subcellular localization, plant cyclophilins could be divided into several types, such as cytoplasmic, chloroplast and mitochondrial (Chou and Gasser, 1997). So far, a lot of cyclophilin genes had already been cloned in *A. thaliana*, *G. max* and several other crops and also received in-depth studies (He et al., 2004; Opiyo and Moriyama, 2009). Previous studies of cytoplasmic cyclophilins in mammals showed that they played a role in drug-induced immunosuppression by inhibiting calcineurin phosphatase activity in Ca²⁺-dependent signaling pathway (Crabtree and Clipstone, 1994). In addition, there were at least five kinds of cytoplasmic cyclophilins




Figure 7. Expression profiles of *Sc-CyP* gene under various exogenous stresses.

in *A. thaliana*, indicating that cytoplasmic cyclophilins may play even more diverse roles in plants than previously suspected (Chou and Gasser., 1997). However, there is not any report about studies on cyclophilin genes

in sugarcane.

In this study, a full-length cytoplasmic cyclophilin gene in sugarcane was obtained (GenBank accession number: GQ246462). Using the PET-29a (+)/BL21 (DE3) expres-

sion system, the prokaryotic expression vector pET29a-*ScCyP* with the *Sc-CyP* gene was successfully constructed. The prokaryotic expression product was a fusion protein of 25.4 kD, that was 18.4 kD of *Sc-CyP* plus 7 kD of His • Tag peptides. The PPIase activity of the prokaryotic expression product and its inhibition by the immunosuppressive drugs FK506 and rapamycin further demonstrated that the *Sc-CyP* gene obtained was indeed a kind of cyclophilin in sugarcane. Since cyclophilin comprised the PPIase activity that catalyzed the rate limiting step in protein folding, it was hypothesized that one possible function of *Sc-CyP* gene may be to act as “chaperone-like” protein to facilitate the folding of the stress-related proteins or to protect these proteins from proteolytic degradation or aggregation under exogenous stresses, as it had been speculated for some other cyclophilins (Ferreira et al., 1996). Due to the existence of one lysine residues, this KSGKPLH48-54 region would carry a positive charge, while the conservation of this region between the distantly related monocotyledonous and dicotyledonous plants indicated the strong selective pressure for maintenance of the sequence in this region (Figures 3 and 4). It was predicted by SubLoc v1.0 that the protein encoded by *Sc-CyP* gene was located in the cytoplasm and thus, it was further confirmed that the *Sc-CyP* gene identified in this study should be classified into the cytoplasmic cyclophilin gene.

Many studies showed that cyclophilins were the proteins in response to adverse stress (Godoy et al., 2000; He et al., 2004; Opiyo and Moriyama, 2009) and their expression and accumulation could be regulated by various environmental stress factors, mainly consisting of the abiotic or biotic stress factors such as mercuric chloride, abscisic acid (ABA), ethylene, salicylic acid (SA), methyl jasmonate (MJ), salt stress, cold and heat, light, drought, injury and fungal infection (Godoy et al., 2000; Nuc et al., 2001; He et al., 2004; Romano et al., 2004; Opiyo and Moriyama, 2009). Kullertz et al. (1999) revealed that when the expression level of cyclophilin in *Grecian foxglove* was enhanced, the freezing tolerance of the plant under low temperature was increased. Godoy et al. (2000) found that the potato cyclophilin responded to SA and bacterial stimulation. Chen et al. (2007) demonstrated that the cyclophilin expression in *Lepidium sativum* could be induced by salt, ABA, H₂O₂ and heat shock stress and also by fungal infection. Previous studies revealed that PEG and NaCl could produce osmotic stress and the exogenous PEG or NaCl treatment could thus be used as the simulated condition of drought stress (Jia et al., 2008), while SA and H₂O₂ were proved to play important roles in signal transduction and resistance response of plant at early stages, such as hypersensitive response (HR) and systemic acquired resistance (SAR) (Wu et al., 1995; Scott et al., 1999). In this study, it was demonstrated in real-time qPCR analysis that when sugarcane received the simulated drought or salt stress, PEG and NaCl stresses, the

expression of *Sc-CyP* gene was greatly induced and thus, it was inferred that the *Sc-CyP* gene played the protective role under the drought and salt stress. What is more, the expression of *Sc-CyP* gene could also be improved significantly under the stresses of H₂O₂ and SA, which suggested that the *Sc-CyP* gene played a certain role under exogenous stresses similar to the H₂O₂ and SA as well, mostly probably a role in signal transduction and disease resistance response in sugarcane.

This study reported the molecular cloning and characterization of a full-length cyclophilin gene in sugarcane and the results should be helpful in the ongoing exploitation of cyclophilin genes in sugarcane.

ACKNOWLEDGEMENTS

This research was funded by the earmarked fund for Modern Agro-industry Technology Research System (CARS-20); Research Fund for the Doctoral Program of Higher Education (20103515120006); Natural Science Foundation of Fujian province (2010J01078) and type A Research Fund of Fujian Education Department (JA10115).

REFERENCES

- Brazin KN, Mallis RJ, Fulton DB, Andreotti AH (2002). Regulation of the tyrosine kinase Itk by the peptidyl-prolyl isomerase cyclophilin A. *Proc. Natl. Acad. Sci. USA.*, 99: 1899-1904.
- Blecher O, Erel N, Callebaut I, Aviezer K, Breiman A (1996). A novel plant peptidyl-prolyl-cis-trans-isomerase (PPIase): cDNA cloning, structural analysis, enzymatic activity and expression. *Plant Mol. Biol.*, 3: 493-504.
- Brandts JF, Halvorson HR, Brennan M (1975). Consideration of the possibility that the slow step in protein denaturation reactions is due to cis-trans isomerism of proline residues. *Biochemistry*, 14: 4953-4963.
- Chen AP, Wang GL, Qu ZL, Lu CX, Liu N, Wang F, Xia GX (2007). Ectopic expression of ThCYP1, a stress-responsive cyclophilin gene from *Thellungiella halophila*, confers salt tolerance in fission yeast and tobacco cells. *Plant Cell Rep.*, 2: 237-245.
- Chou IT, Gasser CS (1997). Characterization of the cyclophilin gene family of *Arabidopsis thaliana* and phylogenetic analysis of known cyclophilin proteins. *Plant Mol. Biol.*, 35: 873-892.
- Crabtree GR, Clipstone NA (1994). Signal transmission between the plasma membrane and nucleus of T lymphocytes. *Annu. Rev. Biochem.*, 63: 1045-1083.
- Faure JD, Gingerich D, Howell SH (1998). An *Arabidopsis* immunophilin, AtFKBP12, binds to AtFIP37 (FKBP interacting protein) in an interaction that is disrupted by FK506. *Plant J.* 15: 783-789.
- Ferreira PA, Nakayama TA, Pak WL, Travis GH (1996). Cyclophilin-related protein RanBP2 acts as chaperone for red/green opsin. *Nature*, 383: 637-640.
- Godoy AV, Lazzaro AS, Casalongue CA, San Segundo B (2000). Expression of a *Solanum tuberosum* cyclophilin gene is regulated by fungal infection and abiotic stress conditions. *Plant Sci.* 152: 123-134.
- Galat A (1999). Variations of sequences and amino acid compositions of proteins that sustain their biological functions: an analysis of the cyclophilin family of proteins. *Arch. Biochem. Biophys.*, 371: 149-162.
- He ZY, Li LG, Luan S (2004). Immunophilins and parvulins. Superfamily of peptidyl prolyl isomerases in *Arabidopsis*. *Plant Physiol.* 4: 1248-1267.
- Handschumacher RE, Harding MW, Rice J, Drugge RJ, Speicher DW

- (1984). Cyclophilin: a specific cytosolic binding protein for cyclosporin A. *Science*, 226: 544-546.
- Kullertz G, Liebau A, Rucknagel P, Schierhorn A, Dietrich B, Fischer G, Luckner M (1999). Stress-induced expression of cyclophilins in proembryonic masses of *Digitalis lanata* does not protect against freezing/thawing stress. *Planta*, 4: 599-605.
- Livak KJ, Schmittgen TD (2001). Analysis of relative gene expression data using Real-time quantitative PCR and the $2^{-\Delta\Delta CT}$ method. *Methods*, 25: 402-408.
- Luan S, Li W, Rusnak F, Assmann SM, Schreiber SL (1993). Immunosuppressants implicate protein phosphatase regulation of K⁺ channels in guard cells. *Proc. Natl. Acad. Sci. USA*. 90: 2202-2206.
- Marivet J, Frendo P, Burkard G (1995). DNA sequence analysis of a cyclophilin gene from maize: developmental expression and regulation by salicylic acid. *Mol. Gen. Genet.*, 2: 222-228.
- Nuc K, Nuc P, Stomaski R (2001). Yellow lupine cyclophilin transcripts are highly accumulated in the nodule meristem zone. *Mol. Plant Microbe Interact.* 12: 1384-1394.
- Opiyo SO, Moriyama EN (2009). Mining the Arabidopsis and rice genomes for cyclophilin protein families. *Int. J. Bioinform. Res. Appl.*, 3: 295-309.
- Que YX, Yang ZX, Xu LP, Chen RK (2009a). Isolation and identification of differentially expressed genes in sugarcane infected by *Ustilago scitaminea*. *Acta Agron. Sin.*, 3: 452-458.
- Que YX, Jiang KQ, Xu LP, Guo Y, Lin JW, Chen RK (2009b). Prokaryotic expression and purification of CP gene of sugarcane mosaic virus strain E and preparation of polyclonal antibody. *Lett. Biotechnol.*, 20(1): 69-71, 102.
- Que YX, Xu LP, Xu JS, Zhang JS, Zhang MQ, Chen RK (2009c). Selection of control genes in Real-time qPCR analysis of gene expression in sugarcane. *Chin. J. Trop. Crops*, 3: 274-278.
- Que YX, Xu LP, Lin JW, Chen RK (2009d). Isolation and characterization of NBS-LRR resistance gene analogs from sugarcane. *Acta Agron. Sin.*, 4: 631-639.
- Romano PGN, Horton P, Gray JE (2004). The Arabidopsis cyclophilin gene family. *Plant Physiol.* 134: 1268-1282.
- Rycyzyn MA, Clevenger CV (2002). The intranuclear prolactin/cyclophilin B complex as a transcriptional inducer. *Proc. Natl. Acad. Sci. USA*. 99: 6790-6795.
- Sharma AD, Singh P (2003). Effect of water stress on expression of a 20 kD cyclophilin-like protein in drought susceptible and tolerant cultivars of Sorghum. *J. Plant Biochem. Biotechnol.*, 12: 77-80.
- Scott IM, Dat JF, Lopez-Delgado H, Foyer CH (1999). Salicylic acid and hydrogen peroxide in abiotic stress signaling in plants. *Phyton (Austria)*. 3: 13-17.
- Sykes K, Gething MJ, Sambrook J (1993). Proline isomerases function during heat shock. *Proc. Natl. Acad. Sci. USA*. 90: 5853-5857.
- Schreiber SL (1991). Chemistry and biology of immunophilins and their immunosuppressive ligands. *Science*, 251: 283-287.
- Wu G, Shortt BJ, Lawrence EB, Levine EB, Fitzsimmons KC, Shah DM (1995). Disease resistance conferred by expression of a gene encoding H₂O₂-generating glucose oxidase in transgenic potato plants. *Plant Cell*. 7: 1357-1368.
- Zhang JS (2007). Expression profile of *Erianthus arundinaceus* cDNA microarray under drought-stress and key drought-resistant genes cloning. Fuzhou: Fujian Agriculture and Forestry University.
- Zhang ZL, Qu WJ (2003). The experimental guide for plant physiology. Beijing: Higher Education Press, p: 23-25.