

Full Length Research Paper

First observation of excision and integration in Class 1 integron in *staphylococci*

Shengjie Guo^{1†}, Yanmei Li^{2†}, Jin Chu³, Li Wang⁴, Xihong Zhao^{5*}, Nanjing Zhong⁶, Shaopeng Qiu¹, Junxing Chen^{1*} and Yanyan Li^{7,8}

¹Department of Urology, the First Affiliated Hospital of Sun Yat-sen University, Guangzhou, 510080, China.

²Guangzhou Women and Children's Medical Center, Guangzhou 510623, China.

³School of Food Science and Nutrition, Faculty of Mathematics and Physical Sciences, University of Leeds, Leeds LS2 9JT, United Kingdom.

⁴Food Safety Key Laboratory of Guangdong Province, College of Food Science, South China Agricultural University, 510642 Guangzhou, China.

⁵Key Laboratory for Green Chemical Process of Ministry of Education, School of Chemical Engineering and Pharmacy, Wuhan Institute of Technology, Wuhan 430073, China.

⁶Guangdong Pharmaceutical University, Guangzhou 510006, China.

⁷State Key Laboratory of Food Science and Technology, Jiangnan University, Wuxi 214122, China.

⁸Department of Microbial Pathogenesis, Dental School, University of Maryland, Baltimore, MD 21201, USA

Accepted 24 August, 2011

Class 1 integron had been well-known as a mobile genetic element involved in the antibiotic resistance of various bacteria. Up to date, most of the relevant publications concentrated on gram-negative bacteria, with limited exceptions. However, in recent studies, class 1 integron had been detected in methicillin-resistant *Staphylococcus aureus* (MRSA) and methicillin-resistant coagulase negative staphylococci (MRCNS). So in this study, we tested in *S. aureus*, the class 1 integron mediated excision and integration. We first asked 8 plasmids from previous studies, then established some transformants and perform the excision and integration reaction. As the results revealed, we observed positive excision assay, which had been confirmed by further integration assays. This observation may raise the public attention of integron as a novel antibiotic resistance determinant in gram-positive bacteria, especially in staphylococci.

Key words: *Staphylococcus aureus*, class 1 integron, mobile genetic element, excision and integration, antibiotic resistance.

INTRODUCTION

In recent years, the role of integrons and gene cassettes in the dissemination of resistance genes has been well established, which is responsible for the facile spread of resistance genes and the rapid evolution of resistance to a wide range of unrelated antibiotics among diverse bacteria (Tauch et al., 2002; Yang et al., 2004; Yu et al., 2003). An integron comprises three elements including the integrase gene (*intI*) encoding an integrase, a recom-

bination site, *attI* and a promoter gene. Several classes of integrons have been reported on the basis of the *intI* gene and classes 1 to 3 are so-called multi-resistant integron. Of these, the class 1 integron platform is the most ubiquitous among resistant clinical isolates of gram-negative bacteria and is found to be associated with the Tn21 transposon family, while the class 2 integrons are associated with the Tn7 transposon family (Arakawa et al., 1995; Barlow et al., 2004; Correia et al., 2003; Hall and Stokes, 1993; Nandi et al., 2004; Nesvera et al., 1998; Nield et al., 2001; Sun et al., 2002). Class 1 integron can capture the gene cassettes, which also contain *attC* site for recombination, via a site-specific recombination event between *attI* and *attC*. Its 3' conserved segment (3'CS) possesses *qacE*_{Δ1} and *sulI*

*Corresponding authors. E-mail: xhzhao2006@gmail.com, chenjunxing@hotmail.com; Tel: +86-13419663487; Fax: +86-13419663487; Tel: +86-20-87333300; Fax: +86-20-87333300.

†These authors contributed equally to this work.

genes, encoding resistance to quaternary ammonium salts and sulfonamide, respectively (Recchia and Hall, 1997). Although, the role of class 1 integron in the spread of antibiotic resistance genes in gram-negative bacteria is clear, little is known about the prevalence of class 1 integron in gram-positive bacteria, especially in *Staphylococcus aureus* (Clark et al., 1999; Nandi et al., 2004). Increasing antibiotic resistance in gram-positive bacteria has become a great concern.

In the past decades, methicillin-resistant *S. aureus* (MRSA) has become one of the most prevalent pathogens that cause nosocomial infections throughout the world. Since it can spread easily by direct or indirect contact between patients and environment or through patients or medical personnel, MRSA is an important risk factor for nosocomial infection and MRSA mediated nosocomial outbreaks are common in hospitals. As consequence, MRSA mediated nosocomial infections continue to be a challenge for clinicians, hospital epidemiologists and administrators since such infections may lead to a serious problem for therapy of patients and infection control. MRSA shows resistance to practically all β -lactam antibiotics, which is caused by *mecA* gene. The *mecA* gene is carried by a mobile genetic element, designated by staphylococcal cassette chromosome *mec* (SCC*mec*), which contains the *mec* gene complex (the *mecA* gene and its regulators) and the *ccr* gene complex encoding site-specific recombinases responsible for the mobility of SCC*mec* (Ito et al., 2001; Kuroda et al., 2001; Xu et al., 2007, 2008a,b, 2010c, 2011a and b).

In recent years, class 1 integron had been found commonly existed in MRSA and methicillin-resistant coagulase negative staphylococci (MRCNS) strains (Xu et al., 2007, 2008a and b, 2009, 2010a and b). In this study, we investigated class 1 integron-mediated recombination in staphylococci, including class 1 integron mediated excision of different gene cassettes with 2 *att* sites and 3 arrays of cassettes and further integration of them by site-specific recombination.

MATERIALS AND METHODS

Bacteria and plasmids

Plasmid-free *S. aureus* strain RN4220 was used as recipient bacterial, with storage, culturing and processing performed as described previously (Hauschild et al., 2003). *S. aureus* strain RN4220 was routinely cultured in Luria-Bertani (LB) medium or agar supplemented with or without appropriate antibiotics as ampicillin (100 μ g/ml), chloramphenicol (25 μ g/ml), sulfamethoxazole (25 μ g/ml) or trimethoprim (25 μ g/ml). Plasmids R388 (a 33 kb IncW plasmid containing In3 with cassettes *dfxB2-orfA*, antibiotic phenotype including Tpr (trimethoprim resistant) Sur (sulfamethoxazole resistant) and pSU2056 (1,176 bp *RsaI*-*Bam*HI fragment of In2 in pUC9, antibiotic phenotype including Apr (ampicillin resistant)) had been acquired from Hall et al. (1991). PLQ425 (with *bla_{imp}* as gene cassette and GTTAGAG as associated *attC*), PLQ431 (with *pse-1* as gene cassette and GTTAGCC as associated *attC*), PLQ438 (with *bla_{imp}* as gene cassette and

GTTAGAA as associated *attC*), PLQ440 (with *aacA1a-orfG* + *orfH* as gene cassette and GTTAGGG as associated *attC*), PLQ443 (with *aadA1* as gene cassette and GTTAAAC as associated *attC*) and PLQ444 (with *aacA1a-orfG* + *orfH* as gene cassette and GTTAGGGG as associated *attC*) were obtained from Leon and Roy (2003).

Electroporation of cassettes carried plasmids into RN4220

To establish different transformants for the excision and integration assay, the aforementioned plasmids had been transformed into RN4220 by electroporation, which had been performed as described previously (Hauschild et al., 2003). In brief, 100 μ l of the competent cells and 1 or 2 μ g of each plasmid were subjected to electroporation, 200 μ l aliquots were then streaked on plated supplemented with proper antibiotics according to relevant phenotypes.

Excision and integration reaction

In vivo excision and integration tests were performed by double transformants (pSU2056 with one of the cassette containing PLQ clones) and triple transformants (R388, pSU2056 with one of the cassette containing PLQ clones), respectively. Growth condition and induction of *Int11* in transformants were performed as described previously (Leon and Roy, 2003; Martinez and de la Cruz, 1990). In brief, transformants were grown overnight at 37°C in blood medium supplemented with selective antibiotics (ampicillin 50 g/ml, chloramphenicol 50 g/ml) and 2% of the overnight culture was then inoculated in 15 ml of new medium (without antibiotic). Isopropyl- β -D-thiogalactopyranoside (IPTG) was added (with final concentration of 0.4 mM for overnight induction) when the optical density at 600 nm reached 0.6. Plasmid isolation was performed as described previously (Schwarz et al., 1989). Excision events were determined by polymerase chain reaction (PCR) amplification and further confirmed by excision-integration tests, with different primer pairs as indicated (Table 1). All experimental procedure had been performed as described previously with slight changes (Leon and Roy, 2003). In brief, for excision assays, sixty-four colonies had been randomly selected and further subjected to PCR amplification for each transformant. For integration assays, ten colonies recovered had been selected to sequencing. In detail, PCR products were purified from agarose gels and the DNA fragments were ligated with the pGEM-T easy vector (Promega, Madison, WI, USA). The ligation mixture was transformed into *E. coli* DH5 α strain and the recombinants were selected on Luria agar containing ampicillin (100 μ g/ml). Recombinant plasmid DNA was purified by standard method and subjected to sequencing and further analyses. The nucleotide sequences of the insert DNA was determined using the BigDye terminator cycle sequencing FS ready reaction kit on an ABI PRISM 310 genetic analyzer (Perkin-Elmer Japan Applied Biosystems, Tokyo, Japan). The nucleotide sequences were analyzed and compared using GenBank database by using the BLAST algorithm, which is available through the National Center for Biotechnology Information (NCBI) website (<http://www.ncbi.nlm.nih.gov>). Definition of excision and integration efficiency was according to previous report (Leon and Roy, 2003).

RESULTS

Transformants in this study

Twelve different types of transformants had been

Table 1. Primers used for the different mobile gene cassettes constructs.

Primer	Nucleotide sequence (5' - 3')	Temperature (°C)	Function	Reference
pACYC184-5'	TGTAGCACCTGAAGTCAGCC	55	Excision test	Leon and Roy (2003)
pACYC184-3'	AGTGATCGAAGTTAGGCTG			
ATTINEU	AGGCGTGATAGATTCTC	48		Leon and Roy (2003)
AADA1	TGCATGACGCCAACTAC	48		
BLA _{IMP}	TGCGGTAGCAATGCTGC	48	Integration test	Leon and Roy (2003)
AACA1a	TAATTGCTGCATTCCTCCGC	48		
PSE-1	CGGATGGTATTA AAAAGC	44		

Table 2. List of transformants and its characteristics.

Transformant	Plasmid	Phenotype
RN4220-PLQ425	pSU2056 and PLQ425	Ap ^r , Cm ^r
RN4220-PLQ431	pSU2056 and PLQ431	Ap ^r , Cm ^r
RN4220-PLQ438	pSU2056 and PLQ438	Ap ^r , Cm ^r
RN4220-PLQ440	pSU2056 and PLQ440	Ap ^r , Cm ^r
RN4220-PLQ443	pSU2056 and PLQ443	Ap ^r , Cm ^r
RN4220-PLQ444	pSU2056 and PLQ444	Ap ^r , Cm ^r
RN4220-R388-PLQ425	pSU2056 and PLQ425	Ap ^r , Cm ^r , Su ^r , Tr ^r
RN4220-R388-PLQ431	pSU2056 and PLQ431	Ap ^r , Cm ^r , Su ^r , Tr ^r
RN4220-R388-PLQ438	pSU2056 and PLQ438	Ap ^r , Cm ^r , Su ^r , Tr ^r
RN4220-R388-PLQ440	pSU2056 and PLQ440	Ap ^r , Cm ^r , Su ^r , Tr ^r
RN4220-R388-PLQ443	pSU2056 and PLQ443	Ap ^r , Cm ^r , Su ^r , Tr ^r
RN4220-R388-PLQ444	pSU2056 and PLQ444	Ap ^r , Cm ^r , Su ^r , Tr ^r

constructed in this study (Table 2), including RN4220-PLQ425 (Ap^r, Cm^r, with pSU2056 and PLQ425), RN4220-PLQ431 (Ap^r, Cm^r, with pSU2056 and PLQ431), RN4220-PLQ438 (Ap^r, Cm^r, with pSU2056 and PLQ438), RN4220-PLQ440 (Ap^r, Cm^r, with pSU2056 and PLQ440), RN4220-PLQ443 (Ap^r, Cm^r, with pSU2056 and PLQ443), RN4220-PLQ444 (Ap^r, Cm^r, with pSU2056 and PLQ444), RN4220-R388-PLQ425 (Ap^r, Cm^r, Su^r, Tr^r, with R388, pSU2056 and PLQ425), RN4220-R388-PLQ431 (Ap^r, Cm^r, Su^r, Tr^r, with R388, pSU2056 and PLQ431), RN4220-R388-PLQ438 (Ap^r, Cm^r, Su^r, Tr^r, with R388, pSU2056 and PLQ438), RN4220-R388-PLQ440 (Ap^r, Cm^r, Su^r, Tr^r, with R388, pSU2056 and PLQ440), RN4220-R388-PLQ443 (Ap^r, Cm^r, Su^r, Tr^r, with R388, pSU2056 and PLQ443), RN4220-R388-PLQ444 (Ap^r, Cm^r, Su^r, Tr^r, with R388, pSU2056 and PLQ444).

Excision and integration reaction

According to the results, excision efficacy ranged from 7.8 to 62.5%, which was thoroughly lower than previous reports when the assays induced in bacteria other than *S. aureus* (Leon and Roy, 2003). Two major factors affecting the excision included *att* site and gene cassettes. As the results showed, greater efficacy had been observed when

attC site on the left-hand neighbour comparing with *attI*, based on 7.8% (5/64) for RN4220-PLQ425 with 31.2% (20/64) for RN4220-PLQ438, 25% (16/64) for RN4220-PLQ440 with 53.1% (34/64) for RN4220-PLQ444. In addition, transformants containing *aadA1* and *aacA1a-orfG+orfH* obtained higher efficacy than *pse-1* and *bla_{IMP}*, as excision efficacy was 15.6% (10/64) for RN4220-PLQ431 and 31.25% for RN4220-PLQ438, comparing with 53.1% for RN4220-PLQ444 and 62.5% (40/64) for RN4220-PLQ443. These results were similar to those published previously (Hansson et al., 1997; Leon and Roy, 2003) and had also been further confirmed by subsequent integrations of excised cassettes into relevant *attI* site. Amplicons obtained were around expected 200 bp and sequencing amplicons confirm that, the excised cassettes were integrated specifically into the G/TTRRRY consensus site of the *attI1* or *attC* site.

DISCUSSION

Since the first description of class 1 integron in 1989 (Stokes and Hall, 1989) integrons have been identified as a primary source of resistance genes and were suspected to serve as reservoirs of antimicrobial resistance genes within microbial populations (Xu et al., 2007,

2009). However, most of these studies were done in gram-negative bacteria with only a few exceptions. While in recent years, integrons had been identified as a novel and unexpected antibiotic determinant in gram-positive organism, especially in MRSA and MRCNS (Xu et al., 2007, 2008a and b, 2009, 2010). The Integron system had been well-known as a mobile genetic element, playing key role in antibiotic resistance in various bacteria (Rowe-Magnus and Mazel, 2001). In this study, we tested and investigated the class 1 integron-mediated excision and integration in *S. aureus* RN4220 and positive result had been observed for both assays. Up to date, no report is available on the dissemination of class 1 integron in any species of staphylococci strain and we are convinced that, the observation of the class 1 integron-mediated excision and integration in *S. aureus* may definitely raise the attention of integron as the novel antibiotic mechanism in gram-positive bacteria, especially in staphylococci, which had been neglected throughout the time.

ACKNOWLEDGEMENTS

We sincerely thank Dr. Stokes and Dr. Roy for providing us plasmids [Stokes et al., 1989; Leon et al., 2003], and Xu et al [Xu et al., 2007] for technical support and kind assistance in writing for this study. This work was supported by National Natural Science Foundation (30872584), Guangdong Natural Science Foundation (8251008901000018), Science Research Fund of Wuhan Institute of Technology (10113201), China; Innovative Experiment Projects of Experimental Teaching Demonstration Center for Environmental and Chemical Engineering Clean Production of Wuhan Institute of Technology (2011), China; College young teacher in-depth business project of Hubei Provincial, Department of Education (XD20100620) Hubei province, China and State Scholarship Fund of China Scholarship Council (2009679002).

REFERENCES

- Arakawa Y, Murakami M, Suzuki K, Ito H, Wacharotayankun R, Ohsuka S, Kato N, Ohta M (1995). A novel integron-like element carrying the metallo- β -lactamase gene blaIMP. *Antimicrob. Agents Chemother.* 39: 1612-1615.
- Barlow RS, Pemberton JM, Desmarchelier PM, Gobius KS (2004). Isolation and characterization of integron-containing bacteria without antibiotic selection. *Antimicrob. Agents Chemother.* 48: 838-842.
- Chang S, Sun C, Yang L, Luh K, Hsieh W (1997). Increasing nosocomial infections of methicillin-resistant *Staphylococcus aureus* at a teaching hospital in Taiwan. *Int. J. Antimicrob. Agents*, 8: 109-114.
- Clark NC, Olsvik Ø, Swenson JM, Spiegel CA, Tenover FC (1999). Detection of a streptomycin/spectinomycin adenyltransferase gene (*aadA*) in *Enterococcus faecalis*. *Antimicrob. Agents Chemother.* 43: 157-160.
- Correia M, Boavida F, Grosso F, Salgado MJ, Lito ML, Cristino JM, Mendo S, Duarte A (2003). Molecular characterization of a new class 3 integron in *Klebsiella pneumoniae*. *Antimicrob. Agents Chemother.* 47: 2838-2843.
- Hall RM, Brookes DE, Stokes HW (1991). Site-specific insertion of genes into integrons: role of the 59-base element and determination of the recombination cross-over point. *Mol. Microbiol.* 5: 1941-1959.
- Hall RM, Stokes HW (1993). Integrons: novel DNA elements which capture genes by site-specific recombination. *Genetica*, 90: 115-132.
- Hansson K, Skold O, Sundstrom L (1997). Non-palindromic *attI* sites of integrons are capable of site-specific recombination with one another and with secondary targets. *Mol. Microbiol.* 26: 441-453.
- Hauschild T, Kehrenberg C, Schwarz S (2003). Tetracycline Resistance in Staphylococci from Free-living Rodents and Insectivores. *J. Vet. Med.* 50: 443-446.
- Ito T, Katayama Y, Asada K, Mori N, Tsutsumimoto K, Tiensasitorn C, Hiramatsu K (2001). Structural comparison of three types of staphylococcal cassette chromosome *mec* integrated in the chromosome in methicillin-resistant *Staphylococcus aureus*. *Antimicrob. Agents Chemother.* 45: 1323-1336.
- Kuroda MT, Ohta I, Uchiyama T, Baba H, Yuzawa I, Kobayashi L, Cui A, Oguchi K, Aoki YN, Lian J, Ito T, Kanamori M, Matsumaru H, Maruyama A, Murakami H, Hosoyama A, Mizutani-Ui Y, Takahashi NK, Sawano T, Inoue R, Kaito C, Sekimizu K, Hirakawa H, Kuhara S, Goto S, Yabuzaki J, Kanehisa M, Yamashita A, Oshima K, Furuya K, Yoshino C, Shiba T, Hattori M, Ogasawara N, Hayashi H, Hiramatsu K (2001). Whole genome sequencing of methicillin-resistant *Staphylococcus aureus*. *Lancet*, 357: 1225-1240.
- Leon G, Roy PH (2003). Excision and Integration of Cassettes by an Integron Integrase of *Nitrosomonas europaea*. *J. Bacteriol.* 185: 2036-2041.
- Martinez E, de la Cruz F (1990). Genetic elements involved in Tn21 site-specific integration, a novel mechanism for the dissemination of antibiotic resistance genes. *EMBO J.* 9: 1275-1281.
- Nandi S, Maurer JJ, Hofacre C (2004). Gram-positive bacteria are a major reservoir of class 1 antibiotic resistance integrons in poultry litter. *Proc Natl. Acad. Sci. USA.* 101: 7118-7122.
- Nesvera J, Hochmannova J, Patek M (1998). An integron of class 1 is present on the plasmid pCG4 from gram-positive bacterium *Corynebacterium glutamicum*. *FEMS Microbiol. Lett.* 169: 391-395.
- Nield BS, Holmes AJ, Gillings MR, Recchia GD, Mabbutt BC, Nevalainen KM, Stokes HW (2001). Recovery of new integron classes from environmental DNA. *FEMS Microbiol. Lett.* 195: 59-65.
- Recchia GD, Hall RM (1997). Origins of the mobile gene cassettes found in integrons. *Trends Microbiol.* 5: 389-394.
- Rowe-Magnus DA, Mazel D (2001). Integrons: Natural tools for bacterial genome evolution. *Curr. Opin. Microbiol.* 4: 565-569.
- Schwarz S, Cardoso M, Blobel H (1989). Plasmid-mediated Chloramphenicol Resistance in *Staphylococcus hyicus*. *J. Gen. Microbiol.* 135: 3329-3336.
- Stokes HW, Hall RM (1989). A novel family of potentially mobile DNA elements encoding site-specific gene-integration functions: integrons. *Mol. Microbiol.* 3: 1669-1683.
- Sun XG, Wang ZQ, Chen L, Luo SY, Jin XY, Zhang WH (2002). Etiological analysis on bacterial ocular disease in northern China (1989-1998). *Chin. Med. J.* 115: 933-935.
- Tauch A, Gotker S, Puhler A, Kalinowski J, Thierbach G. (2002). The 27.8-kb R-plasmid pTET3 from *Corynebacterium glutamicum* encodes the aminoglycoside adenylyltransferase gene cassette *aadA9* and the regulated tetracycline efflux system Tet 33 flanked by active copies of the widespread insertion sequence IS6100. *Plasmid*, 48: 117-129.
- Xu Z, Shi L, Zhang C, Zhang L, Li X, Cao Y, Li L, Yamasaki S (2007). Nosocomial infection caused by class 1 integron-carrying *Staphylococcus aureus* in a hospital in South China. *Clin. Microbiol. Infect.* 13: 980-984.
- Xu Z, Shi L, Alam MJ, Li L, Yamasaki S (2008). Integron-bearing methicillin-resistant coagulase-negative staphylococci in South China, 2001-2004. *FEMS Microbiol. Lett.* 278: 223-230.
- Xu Z, Li L, Alam MJ, Yamasaki S, Shi L (2008). First confirmation of integron-bearing methicillin-resistant *Staphylococcus aureus*. *Curr. Microbiol.* 57: 264-268.
- Xu Z, Li L, Shirliff ME, Alam MJ, Yamasaki S, Shi L (2009). Occurrence and characteristics of class 1 and 2 integrons in *Pseudomonas*

- aeruginosa* isolates from patients in Southern China. J. Clin. Microbiol. 47: 230-234.
- Xu Z, Li L, Shirliff ME, Peters BM, Peng Y, Alam MJ, Yamasaki S, Shi L (2010a). First report of class 2 integron in clinical *Enterococcus faecalis* and class 1 integron in *Enterococcus faecium* in South China. Diag. Microbiol. Infect. Dis. 68: 315-317.
- Xu Z, Li L, Shirliff ME, Peters BM, Li B, Peng Y, Alam MJ, Yamasaki S, Shi L (2010b). Resistance class 1 integron in clinical methicillin-resistant *Staphylococcus aureus* strains in Southern China, 2001-2006. Clin. Microbiol. Infect. 17: 714-718.
- Xu Z, Li L, Shi L, Shirliff ME (2010c). Class 1 integron in staphylococci. Mol. Biol. Rep. (DOI: 10.1007/s11033-011-0676-7).
- Xu Z, Li L, Chu J, Peters BM, Harris M, Li B, Shi L, Shirliff ME (2011a). Development and application of loop-mediated isothermal amplification assays on rapid detection of various types of staphylococci strains. Food. Res. Int. (DOI:10.1016/j.foodres.2011.04.042).
- Xu Z, Li L, Zhao X, Chu J, Li B, Shi L, Su J, Shirliff ME (2011b). Development and application of a novel multiplex polymerase chain reaction (PCR) assay on rapid detection of various types of staphylococci strains. Afr. J. Microbiol. Res. 5: 1869-1873.
- Yang H, Chen S, White DG, Zhao S, McDermott P, Walker R, Meng J (2004). Characterization of multiple antimicrobial resistant *Escherichia coli* isolates from diseased chickens and swine in China. J. Clin. Microbiol. 42: 3483-3489.
- Yu HS, Lee JC, Kang HY, Ro DW, Chung JY, Jeong YS, Tae SH, Choi CH, Lee EY, Soel SY, Lee YC, Cho DT (2003). Changes in gene cassettes of class 1 integrons among *Escherichia coli* isolates from urine specimens collected in Korea during the last two decades. J. Clin. Microbiol. 41: 5429-5433.