

## Full Length Research Paper

# Screening of *in vitro* derived mutants of banana against nematodes

A. Ramesh Kumar\*, N. Kumar, K. Poornima and K. Soorianathasundaram

Department of Fruit Crops, HC and RI, Tamil Nadu Agricultural University- 641 003, India.

Accepted 5 September, 2008

Investigations were carried out to screen the *in vitro* derived mutants of banana cv. Robusta (Cavendish-AAA) and Rasthali (Silk- AAB) by using certain bio-chemical parameters including some enzyme activities. The mutants tested were Ro Im V<sub>4</sub> 6-1-1, Ro Im V<sub>4</sub> 6-1-2, Ro Im V<sub>4</sub> 6-2-1, Si Im V<sub>4</sub> 10-5-3, Si Im V<sub>4</sub> 6-2-5 along with respective susceptible checks (Robusta and Rasthali), tolerant check (Anaikomban-AA) and resistant check (Pisang Lilin- AA). Various biochemical assays used were total phenol, tannin content, lignin content, peroxidase, polyphenol oxidase, phenyl alanine ammonia lyase and ascorbic acid oxidase. The results revealed that the mutants namely Ro Im V<sub>4</sub> 6-1-1 and Si Im V<sub>4</sub> 10-5-3 were found to be resistant while the mutant Ro Im V<sub>4</sub> 6-2-1 was moderately resistant. The rest of the mutants namely Ro Im V<sub>4</sub> 6-1-2 and Si Im V<sub>4</sub> 6-2-5 were found to be susceptible to nematodes. The resistant and moderately resistant mutants of banana could be further used in breeding programmes as well as being recognized as potential cultivars of commerce.

**Key words:** Banana, nematode, resistance, biochemical parameters, enzymes, screening.

## INTRODUCTION

*Musa* production is threatened by pest and disease pressure, which has been increasing during the past 20 years. Most alarming has been the spread of more virulent types of nematodes like burrowing nematode (*Radopholus similis*), root lesion nematode (*Pratylenchus* sp), root knot nematode (*Meloidogyne incognita*) apart from the serious threat by diseases like banana bunchy top disease, sigatoka leaf spots (*Mycosphaerella* sp) and *Fusarium* wilt (*Fusarium oxysporum* f. sp. *cubense*). Crop losses caused by nematodes are very high, with 20% annual yield losses worldwide (Sundararaju and Cannayane, 2002). The existing practice of chemical control of nematodes leaves lot of residues causing much threat to the environment. Hence, there is a need to develop commercially acceptable types of banana with resistance/tolerance to this biotic stresses. In response to these production constraints, efforts aimed at the genetic improvement of *Musa* have gained renewed interest to generate resistant cultivars. Classical breeding consisting of recombination and selection is difficult for banana.

Polyploidy and sterility are both serious handicaps in

the genetic improvement of *Musa* cultivars. An alternative procedure to synthesis nematode resistant cultivars would be to induce mutants under *in vitro* conditions as vegetatively propagated crops like banana are usually heterozygous and the genetic nature of *Musa* is suitable for the application of mutation breeding.

## MATERIALS AND METHODS

The mutants (Table 1) derived from the *in vitro* mutation studies from two commercial cultivars viz., Robusta (Cavendish group-AAA) and Rasthali (Silk group- AAB) were screened for nematode resistance along with Pisang Lilin (AA) (resistant check) and Anaikomban (AA) (tolerant check). Original Robusta and Rasthali were used as susceptible check. These mutants were selected based on their performance for yield and quality parameters during the preliminary screening trials.

### Inoculation with nematodes

The suckers with rhizome weight of approximately 1.5 kg were selected, pared and planted in earthen pots containing four kilograms of sterilized pot mixture (red soil : sand : FYM in the ratio 2:1:1 v/v) at one sucker per pot. The experiment was conducted in a Completely Randomized Design (CRD) with two replications each. Banana mutants maintained in the pots were inoculated with

\*Corresponding author. E-mail: rameshort@yahoo.com.

**Table 1.** List of *in vitro* derived mutants used for the study.

S/N	Name	Genome
1	Ro Im V <sub>4</sub> 6-1-1 (Derived from Robusta)	Cavendish group-AAA
2	Ro Im V <sub>4</sub> 6-1-2 (Derived from Robusta)	Cavendish group-AAA
3	Ro Im V <sub>4</sub> 6-2-1 (Derived from Robusta)	Cavendish group-AAA
4	Robusta -Susceptible check	Cavendish group-AAA
5	Si Im V <sub>4</sub> 10-5-3 (Derived from Rasthali)	Silk group- AAB
6	Si Im V <sub>4</sub> 6-2-5 (Derived from Rasthali)	Silk group- AAB
7	Rasthali -Susceptible check	Silk group- AAB
8	Pisang Lilin - Resistant check	Unique- AA
9	Anaikomban - Tolerant check	Unique- AA

mixed population of nematodes; infective juveniles of root-lesion nematode *Pratylenchus coffeae* (1000 no.s/pot) and burrowing nematode *R. similis* (400 no.s/pot). The nematodes were extracted by modified Baermann funnel technique. The nematode suspension was then poured in the holes made around the rhizosphere of the plants after the emergence of the roots i.e. at 45 days after planting. After inoculation the soil was lightly watered.

#### Estimation of total phenols

The total phenol content was estimated by Folin Ciocalteu method. From the ethanol extract, one ml was taken in two different test tubes and the volume was made up to three ml with distilled water. One ml of Folin Ciocalteu reagent was added to each test tube followed by addition of 2 ml of 20% sodium carbonate solution. The tubes were kept in boiling water bath for a minute, cooled and diluted to 10 ml with distilled water. The intensity of deep blue colour developed was measured at 660 nm wavelength in a Spectrophotometer (Spies, 1955). From the standard graph, amount of phenol present was calculated and expressed as  $\mu\text{g/g}$  of fresh weight.

#### Estimation of tannins

The tannin content in the leaves of banana was quantified by following vanillin hydrochloride method (Robert, 1971)

#### Estimation of lignin

The lignin content of banana leaf tissue was gravimetrically estimated following the method of Chesson (1978). One gram of the banana leaf or root tissue was shaken in a mixture of 5 ml of conc.  $\text{H}_2\text{SO}_4$  and 50 ml of HCl for 16 h at 25°C in a shaker. The mixture was then transferred into a flask with 450 ml of distilled water. After boiling for 20 min, the content of the flask was filtered through a Geena G<sub>3</sub> glass filter. The acid residue was washed to neutrality with distilled water, dried at 105°C and weighed. The results were expressed in terms of percent lignin content on dry weight basis of the tissues.

#### Enzymes assay

Activity of the enzymes such as polyphenol oxidase, peroxidase, phenyl alanine ammonia lyase (PAL) and ascorbic acid oxidase was estimated in leaves at 90 days after nematode inoculation. The

banana leaf samples were taken and homogenized at the rate of one gram per five ml of 0.1 M phosphate buffer (pH 6.5). The homogenate was centrifuged for 20 min at 10,000 rpm at 4°C. Borate buffer 0.2M (pH 8.7) was used for the extraction of PAL. The supernatant was used as the enzyme extract and the activities were recorded as units  $\text{min}^{-1} \text{g}^{-1}$  fresh weight

Peroxidase activity was analysed spectrophotometrically (Hartec, 1955). The reaction mixture was prepared with 0.05 ml of 20 mM guaiacol, 3 ml of phosphate buffer, 0.1 ml of enzyme extract and 0.03 ml of  $\text{H}_2\text{O}_2$ . The changes in absorbance of the reaction mixture at 420 nm were recorded at every 30 s interval for 3 min.

Polyphenol oxidase was assayed using the modified method of Mayer et al. (1965). Standard reaction mixture contained 1.5 ml of 0.1 M phosphate buffer (pH 6.5), 0.5 ml of the enzyme extract and 0.5 ml of 0.01N catechol. The changes in the absorbance were recorded at 495 nm and 30 s interval for 3 min.

Phenyl alanine ammonia lyase (PAL) assay was conducted as per the method described by Ross and Sederoff (1992). The assay mixture containing 100 ml of enzyme, 500 ml of 50 mM Tris HCl (pH 8.8) and 600 ml of 1 mM L-phenylalanine was incubated for 60 min. The reaction was arrested by adding 2 N HCl. Later 1.5 ml of toluene was added, vortexed for 30 s, centrifuged (1000 rpm, 5 min) and toluene fraction containing trans-cinnamic acid was separated. The toluene phase was measured at 290 nm against the blank of toluene.

Ascorbic acid oxidase activity was analyzed spectrophotometrically by following procedure given by Drum et al. (1972).

#### Root damage assessment

Root damage assessment was done at 90 days after inoculation. All the roots collected from each plant were divided into two categories; dead and functional roots.

#### Root necrosis

The length of five selected roots was reduced to 10 cm and the roots were sliced lengthwise. Scoring was carried out at one half each of five roots for the per cent of root cortex showing necrosis. The maximum root necrosis per root half can be 20% giving a maximum root necrosis of 100% for the five halves together. Necrosis of the individual root was recorded. The sum is the total root necrosis of the sample (Carlier et al., 2002).

#### Root lesion index

Resistance rating for nematode screening under pot culture given by Sundararaju (1996) was followed (Table 2).

**Table 2.** Root lesion index: resistance rating for nematode screening under pot culture as given by Sundararaju (1996).

S/N	Number of lesions	Score
1	No infection	1
2	5-10 lesions	2
3	5-10 lesions with rooting	3
4	10-15 lesions	4
5	> 15 lesions with full rotting	5

### Corm damage assessment

Corm damage assessment was worked out as per the procedure suggested by Carlier et al. (2002) and expressed as corm lesion index in a 1 to 5 scale (Table 3).

### Host reaction

As per the terminology described by Bos and Parlevliet (1995), the host reaction was scored in banana which is described below:

- A susceptible plant allows nematode development and allows it to reproduce freely.
- A resistant plant suppresses the nematode development and reproduction
- A sensitive plant shows much injury, even when relatively lightly infected with nematode.
- A tolerant plant may suffer little injury, even when heavily infected with nematode.

## RESULTS

### Total phenols

Among the three mutants of Robusta, Ro Im V<sub>4</sub> 6-1-1 registered the highest total phenol content (108.80 mg/100 g) followed by Ro Im V<sub>4</sub> 6-2-1, whereas the susceptible check Robusta recorded 65.75 mg/100 g only. In Silk group, Si Im V<sub>4</sub> 10-5-3 maintained superiority (106.9 mg/100 g) than the susceptible check Rasthali (69.15 mg/100 g). The resistant check, Pisang Lilin ranked top among all the tested mutants (118.75 mg/100 g) while the tolerant check Anaikomban recorded the total phenol content of 103.4 mg/100 g.

### Tannin

The highest tannin in the leaves was recorded by Pisang Lilin (53.65 µg/g) which was the resistant check whereas the tolerant check Anaikomban recorded tannin content of 48.63 µg/g. Among the mutants, Si Im V<sub>4</sub> 10-5-3 and Ro Im V<sub>4</sub> 6-1-1 recorded the higher tannin contents. The susceptible checks recorded very low values for tannin content.

### Lignin

Lignin content in leaves at 90 DAI ranged from 0.185 to

**Table 3.** Corm damage assessment as per the procedure suggested by Carlier et al. (2002).

S/N	Lesion size and number	Score
1	No lesion	1
2	One small lesion	2
3	Several small lesion	3
4	One large lesion	4
5	Several large lesions	5

0.765 mg/100 g. The resistant check and the tolerant check registered very high values whereas the susceptible checks recorded very poor values. Among the mutants, Ro Im V<sub>4</sub> 6-1-1 recorded the highest value for lignin content followed by Si Im V<sub>4</sub> 10-5-3.

### Enzymes

Among the five *in vitro* derived banana mutants 2 belonging to Cavendish group and one belonging to Silk group recorded higher values than their respective susceptible checks for the enzymes peroxidase, polyphenol oxidase, phenylalanine ammonia lyase and ascorbic acid oxidase. In Cavendish group, the highest value was recorded by Ro Im V<sub>4</sub> 6-1-1 for peroxidase activity and in Silk group, the highest peroxidase activity was noticed in Si Im V<sub>4</sub> 10-5-3. For polyphenol oxidase activity, the data were statistically significant among the mutants and their respective checks of banana. The activity ranged from 41.30 to 115.70 units min<sup>-1</sup> g<sup>-1</sup> fresh weight. The mutants Ro Im V<sub>4</sub> 6-1-1 and Si Im V<sub>4</sub> 10-5-3 registered higher enzyme activity which was comparable with resistant check (Pisang Lilin). However the other mutants namely Ro Im V<sub>4</sub> 6-1-2 Si Im V<sub>4</sub> 6-2-5 recorded comparatively lower enzyme activity which was almost equivalent to susceptible checks. The phenyl alanine ammonia lyase activity differed significantly among the mutants and respective checks. Among the mutants of both Cavendish and Silk groups, the phenyl alanine ammonia lyase activity was the highest (303.38 units min<sup>-1</sup> g<sup>-1</sup>) in Si Im V<sub>4</sub> 10-5-3 which was on par with Ro Im V<sub>4</sub> 6-1-1 (300.81 units min<sup>-1</sup> g<sup>-1</sup>). Lower phenyl alanine ammonia lyase activity of 181.88 units min<sup>-1</sup> g<sup>-1</sup> and 186.12 units min<sup>-1</sup> g<sup>-1</sup> was recorded in Ro Im V<sub>4</sub> 6-1-2 and Si Im V<sub>4</sub> 6-2-5, respectively. Nematode infestation increased the activity of ascorbic acid oxidase in mutants in the order of Ro Im V<sub>4</sub> 6-1-1, Si Im V<sub>4</sub> 10-5-3, Ro Im V<sub>4</sub> 6-2-1, Si Im V<sub>4</sub> 6-2-5 and Ro Im V<sub>4</sub> 6-1-2. The susceptible checks of both Robusta and Rasthali registered lower ascorbic acid activity.

### Total number of roots and infected roots

The total number of roots at 90 DAI varied significantly among the mutants and their respective checks (Table 4).

**Table 4.** Root and corm parameters of tested mutants and respective checks at 90 DAI (days after inoculation).

Name	Genomic group	Total number of roots	Number of infested roots	% Root necrosis	Root lesion index	Corm lesion index	Host response
Ro Im V <sub>4</sub> 6-1-1 (Derived from Robusta)	Cavendish - AAA	68.4	6.99	15	2	1	R
Ro Im V <sub>4</sub> 6-1-2 (Derived from Robusta)	Cavendish - AAA	56.6	19.02	40	4	4	S
Ro Im V <sub>4</sub> 6-2-1 (Derived from Robusta)	Cavendish - AAA	67.8	12.67	20	3	3	MR
Robusta (Susceptible check)	Cavendish - AAA	52.4	26.13	85	5	5	HS
Si Im V <sub>4</sub> 10-5-3 (Derived from Rasthali)	Silk- AAB	72.0	6.75	15	2	1	R
Si Im V <sub>4</sub> 6-2-5 (Derived from Rasthali)	Silk- AAB	54.8	16.80	40	4	4	S
Rasthali (Susceptible check)	Silk- AAB	54.2	29.17	80	5	5	HS
Pisang Lilin - Resistant check	AA	70.6	7.47	10	1	1	R
Anaikomban - Tolerant check	AA	62.0	13.50	30	3	3	T
SED		0.544	0.915	-	-	-	-
CD 5%		1.23	2.07	-	-	-	-

R = Resistant; MR = moderately resistant; T = tolerant; S = susceptible; HS = highly susceptible.

The total number of roots ranged from 52.4 to 72.0. The highest total number of roots was noticed in Si Im V<sub>4</sub> 10-5-3 (72.0) which was significantly different from other mutants and it was followed by Pisang Lilin (70.6) which was a resistant check. The susceptible checks of both the groups recorded the lower values for total number of roots (54.8 in Silk and 52.4 in Cavendish). In the case of infested roots produced also statistically differed among the different mutants of banana. The number of infested roots ranged from 6.75 in Si Im V<sub>4</sub> 10-5-3 to 29.17 in Rasthali susceptible check. Among the mutants in

Cavendish group, the least number of infested roots was recorded in Ro Im V<sub>4</sub> 6-1-1 (6.99) while the highest number of infested roots was recorded in Ro Im V<sub>4</sub> 6-1-2 (19.02). Similarly in Silk group, between the two mutants, the least value was registered by Si Im V<sub>4</sub> 10-5-3, whereas the mutant Si Im V<sub>4</sub> 6-2-5 registered higher number of infested roots. In both the groups the susceptible checks relatively recoded higher values for num-

ber of infected roots.

### Root and corm damage assessment

The root and corm damage was assessed in terms of per cent root necrosis, root lesion index and corm lesion index (Table 4). The total root necrosis per cent varied from 10 to 85. In Cavendish group the least per cent (15%) was recorded by Ro Im V<sub>4</sub> 6-1-1 followed by Ro Im V<sub>4</sub> 6-2-1 (20%). In Silk group, the least per cent (15%) was registered by Si Im V<sub>4</sub> 10-5-3. In general, the susceptible checks of both groups comparatively recorded higher per cent of root necrosis. The root lesion index varied from 1 to 5 among the *in vitro* derived mutants of banana. The highest index (5) was recorded in susceptible checks while the lowest index (1) was recorded in Pisang Lilin which was a resistant check. Among the mutants of both the groups Ro Im V<sub>4</sub> 6-1-1 and Si Im V<sub>4</sub> 10-5-3 relatively recorded lower

index (1). Similarly corm lesion index also ranged from 1 to 5. The least index (1) was recorded by Ro Im V<sub>4</sub> 6-1-1 and Si Im V<sub>4</sub> 10-5-3 which was on par with resistant check Pisang Lilin. The susceptible checks of both the groups recorded maximum index (5).

Based on the per cent root necrosis, root lesion index and corm lesion index, the level of resistance was assessed among the *in vitro* derived banana mutants. The mutants Ro Im V<sub>4</sub> 6-1-1 and Si Im V<sub>4</sub> 10-5-3 were found to be resistant while the mutant Ro Im V<sub>4</sub> 6-2-1 was moderately resistant. The rest of the mutants namely Ro Im V<sub>4</sub> 6-1-2 and Si Im V<sub>4</sub> 6-2-5 were found to be susceptible to nematodes.

### DISCUSSION

The biochemical basis for resistance to nematode was studied in mutants. The mutants Ro Im V<sub>4</sub> 6-1-1, Ro Im V<sub>4</sub> 6-2-1 and Si Im V<sub>4</sub> 10-5-3 posses-

**Table 5.** Biochemical parameters in the leaves of tested mutants and respective checks at 90 DAI (days after inoculation).

Name	Genomic group	Total phenol (mg/100 g)	Tannin ( $\mu\text{g/g}$ )	Lignin (%)	Peroxidase ( $\text{min}^{-1} \text{g}^{-1}$ fresh wt)	Polyphenol oxidase ( $\text{min}^{-1} \text{g}^{-1}$ fresh wt)	Phenyl alanine ammonia lyase ( $\text{min}^{-1} \text{g}^{-1}$ fresh wt)	Ascorbic acid oxidase ( $\text{min}^{-1} \text{g}^{-1}$ fresh wt)
Ro Im V <sub>4</sub> 6-1-1 (Derived from Robusta)	Cavendish - AAA	108.80	48.57	0.751	48.65	110.11	300.81	52.64
Ro Im V <sub>4</sub> 6-1-2 (Derived from Robusta)	Cavendish - AAA	70.23	25.53	0.238	22.25	55.56	181.88	57.14
Ro Im V <sub>4</sub> 6-2-1 (Derived from Robusta)	Cavendish - AAA	100.28	39.87	0.695	43.82	90.28	260.72	53.64
Robusta (Susceptible check)	Cavendish - AAA	65.75	18.16	0.185	14.31	41.30	131.70	18.75
Si Im V <sub>4</sub> 10-5-3 (Derived from Rasthali)	Silk- AAB	106.90	46.76	0.743	48.64	107.65	303.38	57.32
Si Im V <sub>4</sub> 6-2-5 (Derived from Rasthali)	Silk- AAB	71.65	26.18	0.223	21.52	53.66	186.12	57.57
Rasthali (Susceptible check)	Silk- AAB	69.15	18.15	0.192	16.83	41.70	130.75	20.70
Pisang Lilin - Resistant check	AA	118.75	53.65	0.765	50.70	115.70	310.70	66.28
Anaikomban - Tolerant check	AA	103.40	40.06	0.705	44.80	91.15	265.70	50.64
SED		0.5615	0.5615	0.0071	0.6575	0.7071	2.2001	0.7122
CD 5%		1.2703	1.2703	0.0160	1.4872	1.5996	4.9772	1.6112

sed higher quantity of total phenol, tannin and lignin in the leaves (Table 5). The higher total phenol, tannin and lignin in the roots were attributed to polyphenol oxidase, phenylalanine ammonia lyase, ascorbic acid oxidase and peroxidase activities. Fogain (1996) and Valette et al. (1997) found higher amounts of phenolics in the resistant cultivar Yangambi km 5. Increased activity of peroxidase in tomato, phenylalanine ammonia lyase in brinjal was positively correlated with nematode resistance (Rajasekar et al., 1997; Sirohi and Dasgupta, 1993).

Several physiological processes in the host are stimulated due to the activation of certain enzymes. Enzymes like peroxidase, polyphenol oxidase, phenylalanine ammonia lyase and ascorbic acid oxidase were found to be involved in the plant defense mechanism (Abbastista and Matta, 1975). The mutants exhibited higher activities of peroxidase, polyphenol oxidase, phenylalanine ammonia lyase and ascorbic acid oxidase (Table 5). Enhanced peroxidase activity has been associated with resistant hybrids to


nematodes (Fogain and Gowen, 1996; Vallette et al., 1997). The PAL and ascorbic acid oxidase enzymes might have increased the chlorogenic acid and ascorbic acid content in the leaves, which act as toxic compounds against nematodes. Transcriptional activation of the genes encoding PAL has been observed within 5 min of elicitor treatment (Lam et al., 1986), confirming its role in defense mechanism. In susceptible mutants, the nematodes would have broken the defense barrier by producing certain offensive chemicals while in the case of resistant/moderately resistant mutants, the plant would have synthesized certain toxic compounds known as phytoalexins. Usually these phytoalexins are low molecular weight antimicrobial compounds that are synthesized and accumulated in the cells. It has been well established that phenylalanine ammonia lyase is the prime enzyme involved in plant defense mechanism which is involved in the phenyl propanoid pathway in plant system. Lignin and phenol are synthesized via phenyl propanoid pathway which impart resistance against nematode attack. The

role of phytoalexins and other toxic compounds like phenols and lignin (which are otherwise called phytoanticipins and are synthesized as a part of the normal plant development) in resistance mechanism have been reported by earlier workers (Reuveni et al., 1992; Sariah et al., 1999).

Good root development with healthy roots and corm favours resistance. In the present investigation, the resistant and moderately resistant mutants showed lower percentage of infected roots (Figure 1).

## Conclusion

The study vividly indicated the presence of resistance and tolerance in the *in vitro* derived mutants of banana. Further the role of different biochemical parameters and enzymes has also been well established in developing resistance against nematodes in banana. The resistant and moderately resistant mutants of banana could be further used in breeding programmes as well as



**Figure 1.** Effects of nematodes on infestation of *in vitro* derived mutants of banana.

be recognized as potential cultivars of commerce.

## REFERENCES

- Abbattista G, Matta A (1975). Production of some effects of ethylene in relation to Fusarium wilt of tomato. *Physiol. Plant Pathol.* 5: 27-35.
- Bos L, Parlevliet JE (1995). Concepts and terminology on plant/part relationships towards consensus in plant pathology and crop protection. *Ann. Rev. Phytopath.* 33: 69-102.
- Carlier J, De Waele D, Vincent E (2002). INIBAP Technical guidelines on Global evaluation of *Musa* germplasm for resistance to Fusarium wilt, *Mycosphaerella* leaf spot disease and nematodes. Eds. By Anne Veniza and Claudine Picq. Publ. by INIBAP, France.
- Chesson A (1978). The maceration of line flax under anaerobic condition. *J. Appl. Bacteriol.* 45: 219-230.
- Drumm H, Brning K, Mohr H (1972). Activities of ascorbic acid oxidase in relation to nematode resistance in potato. *Planta*, 106: 259-267.
- Fogain R (1996). Evaluation of *Musa* spp. for susceptibility to nematodes and study of resistance mechanisms. *Acta Hort.* 540: 215-224.
- Hartec EF (1955). Haematin compounds. In: *Modern Methods of plant analysis* Vol. 4., Peach K, Tracq M (Eds.) Springer-verlag, New York, pp. 197-245.
- Lam CJ, Corbin DR, Lawton MA, Sauer N, Wingate VPM (1986). Recognition and response in plants: Pathogen interaction, recognition in microbe plant symbiotic interaction, (B. Legtenberg, ed.), Springer Symbiotic Interaction, pp. 333-344.
- Mayer AM, Haul E, Shaul RB (1965). Assay of catechol oxidase, a critical comparison of method. *Phytochemical.* 5: 783-789.
- Rajasekar SP, Ganguly AK, Swain SC (1997). Quantitative changes in superoxide dismutase, catalase and peroxidase with reference to resistance in tomato to *Meloidogyne incognita*. *Indian J. Nematol.* 27: 79-85.
- Reuveni O, Israeli Y, Golubowicz S (1992). Factors influencing the occurrence of somoclonal variation in micropropagated bananas. *Acta Hort.* 336: 357-364.
- Robert EB (1971). Method for estimation of tannin in grain sorghum. *Agron. J.* 63: 511.
- Ross WW, Sederoff RR (1992). Phenylalanine ammonia lyase from loblolly pine; Purification of the enzyme and isolation of complementary DNA clones. *Plant Physiol.* 98: 380-386.
- Sariah M, Lim CL, Tariq SA (1999). Use of biochemical marker as a measure of resistance towards *Fusarium* wilt of banana. Proc. Intl. workshop on the banana *Fusarium* wilt disease, 18-20 Oct. 1999, Malaysia, pp. 243-249.
- Sirohi A, Dasgupta DR (1993). Mechanism of resistance in cowpea to root-knot nematode, *Meloidogyne incognita* race 1; I. early induction of phenylalanine ammonia lyase and chlorogenic acid. *Indian J. Nematol.* 23: 37-41.
- Spies JR (1955). *Methods in Enzymology*. Eds. S.P. Colowick and W.O. Kaplan. Academic Press, pp. 467-468.
- Sundararaju P (1996). Nematode pests of banana and their management. Souvenir, In: Conference on Challenges of banana production and utilization in First century held at Trichy on September, 24-25, pp. 17-19.
- Sundararaju P, Cannayane I (2002). Integrated nematode management on banana in India. Global conference on Banana and Plantain, organized by AIPUB, held at Bangalore, between 28-31 October, 2002, Souvenir: 73-76.
- Valette C, Nicole M, Sarah JL, Boisseau M, Boher B, Fargette M, Geiger JP (1997). Ultrastructure and cytochemistry of interactions between banana and the nematode *Radopholus similis*. *Fundam. Appl. Nematol.* 20: 65-77.