

Full Length Research Paper

## ***Polistes olivaceous* decreases biotic surface colonization**

Wings T. Y. Loo<sup>1</sup>, L. J. Jin<sup>1</sup>, Mary N. B. Cheung<sup>2</sup>, Yi-ding Dou<sup>2\*</sup> and J. P. Chen<sup>3</sup>

<sup>1</sup>Faculty of Dentistry, the University of Hong Kong, Hong Kong.

<sup>2</sup>Jinhua Dentistry, Chengdu, Sichuan, China.

<sup>3</sup>School of Chinese Medicine, University of Hong Kong, Hong Kong.

Accepted 10 September, 2009

The objective of this investigation was to evaluate the anti-bacterial efficacy of the honeycomb of *Polistes olivaceous* on oral biotic surface (biofilm) model by means of pH response, population of oral bacteria and enamel mineralization. Three copies of a three-organism-bacterial consortium was grown on hydroxyapatite (HA) surfaces in a continuous culture system and exposed to repeated solution pulses of sucrose solution every 12 h to construct a cariogenic biofilm on the HA discs in the flow cells. One flow cell was only pulsed with 500 µmol/ml of sucrose (S group). The second flow cell was pulsed with 500 µmol/ml sucrose and 2.5 mg/ml *P. olivaceous* extract (P group). The third flow cell was pulsed with 500 µmol/ml sucrose, 230 mg/L sodium fluoride and 0.2% chlorhexidine digluconate (C group). During the course of carbohydrate supplement, the pH of the S group dropped sharply compared with the others. The P group demonstrated pH recovery to baseline more easily than the S group ( $p < 0.05$ ). The C group demonstrated very little pH drop. The P group displayed a lower level of colonization than the S group, which was reflected by a lower cariogenic bacterial count and a less compact biofilm especially after the third pulse. *P. olivaceous* suppresses bacteria growth and accelerates pH recovery. *P. olivaceous* may have stabilizing effect against cariogenic shift on the oral biofilm, preventing tooth decay.

Comment [p1]: mineralization

Comment [p2]: Check it

Comment [p3]: especially

**Key words:** *Polistes olivaceous*, oral biotic surface (biofilm), hydroxyapatite, cariogenic bacteria, sucrose.

### INTRODUCTION

A biotic surface (biofilm) is defined as a community of bacteria intimately associated with each other within an exo-polymer matrix. This biological unit exhibits its own properties that are quite different from those shown by each individual species in planktonic form (Bortolaia and Sbordone, 2002). The micro-organisms which have been correlated with the most common oral pathologies reside in the biofilm (Bortolaia and Sbordone, 2002). The importance of the biofilm mode of growth for the expression of specific physiological characteristics has been established in a number of studies (Brown and Gilbert, 1993; Hardie, 1993; Wennstrom and Lindhe, 1985). It is reported that micro-organisms within biofilms tend to be more resistant to antimicrobial agents (up to 1000-fold), even

though these same bacteria are sensitive to the agents if grown under planktonic conditions (Brown and Gilbert, 1993; Gander, 1996). The reason for this is not clear, however, some speculations include the inactivation of antimicrobial agents or the failure of the antimicrobial agent to penetrate into the biofilm. In addition, it is also reported that biofilm increases the opportunity for gene transfer between bacteria (Xie et al., 2000).

Comment [p4]: microorganisms

Cariogenic bacteria, such as *Streptococcus mutans* etc, can colonize tooth surface and initiate plaque formation by synthesizing extracellular polysaccharides from sucrose. The polysaccharide matrix of biofilm forms a barrier that prevents the diffusion of acids produced by the bacteria. The acids accumulate *in situ* and decalcify minerals in the enamel and result in dental caries (Bradshaw et al., 1989).

Many researchers have emphasized the importance of controlling dental plaque to prevent dental caries and to maintain oral health (Kanchanakamol et al., 1993;

Comment [p5]: All "et all" should be Italic

\*Corresponding author. E-mail: wtyloo@126.com. Tel: 86-13709003409. Fax: 86-28-86677743.

Sauvetre et al., 1995; Sharma et al., 1994; Yankell et al., 1996). A wide variety of sources have been explored in the search for some effective anti-plaque agents (Marsh and Bradshaw, 1997; Southard et al., 1984; Wennstrom and Lindhe, 1985). In recent years, reports on the isolation and identification of natural plaque inhibiting substances have generated much interest in the research of caries prevention (Gander, 1996; Taniguchi and Kubo, 1993). A recent study has demonstrated antimicrobial activity against selected oral pathogens from natural sources and herbs (Yankell et al., 1993).

From the native American plant *Ceanothus americanus*, ceanothic acid and ceanothetric acid demonstrated growth-inhibitory effects against *S. mutans*, *A. viscosus* and *P. gingivalis* (Li et al., 1997). Bakuchiol, isolated from *P. corylifolia*, have antimicrobial activities against oral microorganisms *in vitro* (Katsura et al., 2001). Ethanolic extracts of propolis showed antimicrobial activity against three oral microorganisms (Koo et al., 2000; Xiao et al., 2006). Most of these studies focused on antimicrobial activities against batch cultivated cariogenic bacteria *in vitro* (Katsura et al., 2001; Koo et al., 2000; Kostiuchenkov and Farashchuk, 1991). Some natural herbs have been given as a dentifrice and were proven to show antimicrobial effects (Lee et al., 2004).

*Polistes olivaceous*, extracted from honeycomb of *De Geer*, is generally used in clinics of conventional Chinese Medicine as an anti-oxygenation agent for treating renal deficiency (Kostiuchenkov and Farashchuk, 1991). It has been reported to show antineoplastic effects in combination with other herbs used to treat parotid gland cancer, mastocarcinoma, gastric and liver cancer *in vivo* and *in vitro* in traumatology to treat microorganism infection (Zhang and Zhou, 2004). The application of the powdered form of *P. olivaceous* directly to carious teeth for relieving toothache has been well known among Chinese medical and dental practitioners.

Although *P. olivaceous* is widely used, only few studies have been carried out to investigate the effect of *P. olivaceous* has on dental biofilm bacterial colonization (Xiao et al., 2006; Xiao et al., 2007). The cariostatic efficacy of *P. olivaceous* was evaluated by studying the influence of *P. olivaceous* on pH response, the population of cariogenic organisms and enamel mineralization within a biofilm model invented by Hodgson (Hodgson et al., 2001).

## MATERIALS AND METHODS

### The preparation of *P. olivaceous*

*P. olivaceous* was provided by the School of Chinese Medicine, the University of Hong Kong and was tested by Société Générale de Surveillance (SGC) in Hong Kong to be free of micro-organisms. The solid *P. olivaceous* was dried and grounded into a powder which was then extracted with methanol (Lin et al., 2001). Prior to use, it was dissolved in a sucrose solution at a final concentration of 2.5 mg/ml (Huang et al., 2003) and filtered with 0.22 µm Cellulose

Acetate (Corning, NY, USA) for sterilization.

*Streptococcus sanguis* (*S. sanguis*) ATCC 10557, *Streptococcus salivarius* (*S. salivarius*) ATCC 25975 and *Streptococcus mutans* (*S. mutans*) ATCC 25175 were purchased from American Type Culture Collection (ATCC, Manassas, VA, USA).

They were cultured in our laboratory. The mixed bacterial culture was grown in a 250 ml vessel, operated at 37°C under a gas phase of 5% (v/v) CO<sub>2</sub> in nitrogen, at a dilution rate of 0.1 h<sup>-1</sup> for 20 h. The pH was maintained at 7.0 by the automated addition of 0.5 N NaOH.

Total aerobic (15% CO<sub>2</sub> in air atmosphere) and anaerobic counts were performed on supplemented Brain Heart Infusion (BHI) Agar (BioMerieux, Inc. Wilsonville, Oregon, USA) containing 5% (v/v) defibrinated horse blood (BHIS). *S. sanguis* and *S. salivarius* were counted on tryptone-yeast extract cysteine (TYC) agar (Scharlab S.L. Chemical Plant; Barcelona, Spain) and *S. mutans* on tryptone-yeast-cysteine-sucrose-bacitracin (TYCSB) agars (Microdiagnostik, Brisbane, Australia) (Wan et al., 2002; Van Palenstein Helderman et al., 1983). Identification was based on colony morphology and Gram staining reaction (Holt, 1984).

### Biofilm preparation

The growth biofilm medium contained 2.5 mg/ml hog gastric mucin Type III (Sigma-Aldrich, MO, USA), 2 mg/ml proteose peptone (Sigma-Aldrich, MO, USA), 1 mg/ml trypticase peptone (Sigma-Aldrich, MO, USA), 1 mg/ml yeast extraction (Sigma-Aldrich, MO, USA), 0.5 mg/ml glucose, 2.5 mg/ml KCl, 0.1 mg/ml cysteine-HCl (Sigma-Aldrich, MO, USA). The medium was filtered with 0.22 µm Cellulose Acetate (Corning, NY, USA) for sterilization and adjusted to pH 7.5 (Harvey, 2000).

### Inoculation of organism

When a stable population containing all three organisms had been established, three parallel connecting flow cells were inoculated simultaneously with a mixture of the planktonic phase from the continuous culture and fresh medium (in a 1:9 ratio). After inoculation, the consortium was allowed to establish for at least 7 days prior to the start of biofilm experiments. The constant liquid level was maintained by means of a weir system (20 ml working volume). The growth medium was supplemented with 5 µmol/ml phosphate buffer, pH 8.0. A total flow rate of 15 ml/h of combined inoculum or fresh medium was used to obtain a dilution rate of 0.80 h<sup>-1</sup>. The contents of the flow cells were re-circulated (60 ml/min) to achieve constant mixing. Liquid feeds to the flow cell, recirculation of contents and removal of waste were controlled by peristaltic pumps. The pH of the planktonic phase was monitored over the course of the experiments using electrodes; these were cleaned in sodium hypochlorite and washed with normal saline prior to use.

### *Polistes olivaceous* minimal and optimal concentration determination

The pilot studies to find out the minimal and optimal concentration of *P. olivaceous* had been conducted in the Hospital of Ningxia Medical University, Yinchuan, China, a total of 550 university students were randomly recruited in the test between 2007 and 2009. The ethics committees at Ningxia Medical University approved the research protocols, also informed consents were obtained from these subjects before the joint study. The minimal and optimal concentration of *P. olivaceous* to be used in the pulse experiment was determined by doses response tests from 0.5, 1, 1.5, 2, 2.5, 3, 3.5, 4 and 5 mg/ml in total 450 patients. Another 50 subjects used oral rinses of 0.9% normal saline to serve as a negative control and


Comment [p8]: Bold

Comment [p9]: Bold

Comment [p10]: concentration

Comment [p6]: Bold

Comment [p7]: Correct it



**Figure 1.** pH response to pulses of sucrose alone, sucrose with *Polistes olivaceus* and sucrose with sodium fluoride and chlorhexidine digluconate. ↓ : indicates one pulse to the flow cell; △: sucrose pulsed group (S); □: sucrose, sodium fluoride and chlorhexidine digluconate pulsed group (C); ○: Sucrose and *Polistes olivaceus* pulsed group (P).

50 used oral rinses of 0.2% chlorhexidine to serve as a positive control. Each dose group consisted of 25 males and 25 females; they were aged between 20 and 35 with the mean age 26.8. Each tested concentration of *P. olivaceus* was rinsed by 10 individuals for three days at three times per day. After three days, an oral swab from the enamel surface was used for a bacterial culture to determine the number of colony forming units (CFU). Observations from the culture results (data did not show in this study) determined that 2.5 mg/ml of *P. olivaceus* was the optimal and minimal concentration which was statistically different from the CFU obtained by the saline rinse ( $p < 0.05\%$ ). As the concentration increased from 2.5 mg/ml, the CFU counts approached those obtained by patients who used the chlorhexidine rinse, but concentrations above 2.5 mg/ml caused discomfort to the subjects. The oral bacteria were unresponsive to concentrations of *P. olivaceus* below 2.5 mg/ml and CFU counts were similar to patients who used the 0.9% normal saline. Thus 2.5 mg/ml serving as the optimal and minimal concentration of *P. olivaceus* was used for the pulse experiments.

resin. Representative sections were prepared using an LKB/Wallac 8801A Ultratome III Microtome (Diversified Equipment Company, Inc. Lorton, VA, USA) with a diamond knife. Thick sections (1  $\mu$ ) were stained with toluidine blue stain (Newcomer Supply, Middleton, WI, USA). Thin sections (800 Å) were stained with aqueous uranyl acetate, counterstained with Reynolds' lead citrate and examined by a Leica Stereoscan S440 Scanning Electron Microscope (Leica Microsystems GmbH, Wetzlar, Germany) in the Electron Microscope Unit, Queen Mary Hospital, the University of Hong Kong.

#### Statistical analysis

Statistical analysis of the mean colony formation units per ml ( $\log_{10}$  cfu/mL) on the agar plates was calculated by SPSS 15(USA) using one-way ANOVA test.

## RESULTS

### Pulse conditions

Three cariogenic biofilms on hydroxyapatite (HA) discs in the flow cells were created, each of them was pulsed every 12 h for 72 h with a sucrose solution (500  $\mu$ mol/ml in water) to produce an initial concentrations of 50  $\mu$ mol/ml in the bulk liquid phase.

The first flow cell was pulsed with sucrose alone (S group). The second flow cell was pulsed with sucrose, 230 mg/L sodium fluoride and 0.2% chlorhexidine digluconate (C group). The final flow cell was pulsed with 2.5 mg/ml *P. olivaceus* extract (initial concentration of 0.5 mg/ml) together with 500  $\mu$ mol/ml sucrose solution (P group).

### Scanning electron microscopy

The HA discs were fixed by modified Karnovsky's fixative (2.5% glutaraldehyde, 2% paraformaldehyde in 0.1 M sodium cacodylate buffer, pH 7.4). Samples were fixed at 4°C for 12 h. All equipments were then rinsed briefly in buffer, post-fixed in 1% buffered osmium tetroxide and dehydrated in graded ethanol. Specimens were then rinsed in propylene oxide then infiltrated and embedded in Araldite

### pH response to pulses

The flow cells resided at a steady-state of pH 6.0 during the initial phase of biofilm growth. After the first pulse, the S group presented with a pH drop from 6.0 to 4.3, which recovered to 6.0 before the second pulse. The pH recovered to 6.0 after the second pulse, but remained at 4.3 from the third pulse onwards. The P group demonstrated similar pH drops and recoveries as the S group. However, unlike the S group, the pH continued to recover after every pulse and did not remain at 4.3. Also, after around 30 h, there was a significant difference ( $p < 0.05$ ) between S and P group. The pH of S group fluctuated less, stayed around 4 and 5 and became less dependent to the pulse after 31 h while the pH for P group continued to demonstrate a pH drops and recoveries after 31 h. Meanwhile the pH value of the C group did not show any sharp pH drops and remained stable at about pH 6.2 (Figure 1).

Comment [p13]: Bold

Comment [p11]: Bold

Comment [p12]: Bold

**Table 1.** Bacterial count of *S. mutans*, *S. sanguis* and *S. salivarius* in each flow.

Bacteria	Log <sub>10</sub> CFU/disc		
	S group	C group	P group
<i>S. mutans</i>	8.20 ± 0.15*	6.10 ± 0.10	6.50 ± 0.11 *
<i>S. sanguis</i>	10.00 ± 0.18*	7.40 ± 0.12	9.10 ± 0.12
<i>S. salivarius</i>	9.10 ± 0.17*	8.40 ± 0.15	8.20 ± 0.09

Values are mean ± standard deviation.

\*Statistically significant ( $p < 0.05$ ).

S group: pulses of sucrose alone, P; group: pulses of sucrose and *Polistes olivaceus*; C; group: pulses of sucrose, sodium fluoride and chlorhexidine digluconate.

### Response of biofilm populations to pulses

The bacterial population was cultured after the biofilm experiment was performed for four days. The mean colony formation units (CFU) was calculated and expressed in terms of log<sub>10</sub> cfu/mL of organism in planktonic phase (Table 1).

A significant increase ( $p < 0.05$ ) in total biofilm population for all three species was observed between the biofilms of the S and the C group.

While between the S and the P group, the population of *S. mutans* was significantly inhibited ( $p < 0.05$ ).

### Biofilm microstructure of (Scanning Electron Microscope) SEM imaging

The HA discs were taken out from the flow cells after 72 h for observation under a SEM. In the C group, the circumference was rather clear without sucrose pulsed in (Figure 2a). While in the micro-radiography of the S group, the biofilm had much more extracellular polysaccharide, which created a fuzzy appearance indicating a sign of demineralization (Figure 2b). As compared to the S group, the P group showed less demineralization (Figure 2c).

### DISCUSSION

It is now clear that dental plaque plays an essential role in the pathogenesis of dental caries. When supplied with carbohydrate, there will be a shift from a symbiotic biofilm to a cariogenic biofilm, which contains a larger population of cariogenic bacteria (De Stoppelaar et al., 1970; Dennis et al., 1975; Minah et al., 1985; Staat et al., 1975). These effects have also been demonstrated in animal models (Beighton and Hayday, 1986).

To control dental plaque, a wide variety of sources have been screened. The use of natural products has been one of the most successful strategies for the discovery of new medicines (Harvey, 2000). Seventy eight percent of

new antibiotics and 61% of new antitumor drugs approved by the Food and Drug Administration, USA or comparable entities in other countries from 1983 to 1994 were natural products or derived from natural products (Cragg et al., 1997).


Extensive studies showed *P. olivaceus* have antimicrobial activity against oral microorganisms and can inhibit the activity of glucosyltransferase *in vitro* (Koo et al., 2000; Koo et al., 2002). In this context, we studied *P. olivaceus* influences on the cariogenic shift of oral biofilm.

The pH measurements made in the planktonic phase of flow cells showed pulse-like responses to sucrose challenge. With sucrose pulsed in, the pH dropped quickly. Acid was cleared over a 12 h period, with about four hours at minimum pH; this was compared with the C group which had no significant pH drop. Since the third pulse, the pH of the S group had been inhibited at a low level of pH 4-5, which was the critical pH of enamel decalcification (Ingram and Silverstone, 1981). While in the P group, though there was the same concentration of sucrose pulsed in, the pH could be recovered 6 - 7 h after every pulse, which might contributed to the effect of *P. olivaceus* extract.

Biofilm structure was also clearly affected by the sucrose pulsing, which produced a more compact biofilm structure compared with the C group. These findings confirmed previous work using a similar biofilm system (Bradshaw et al., 1989; Singleton et al., 1997). Increased density of biofilm structure had been implicated as a barrier to host immune responses (Anwar et al., 1992). Furthermore, extracellular polysaccharides produced, as a result of sucrose metabolism, might play a role in enhancing the cariogenic challenge. These polysaccharides influenced sugar and acid diffusions (Dibdin and Shellis, 1988; McNee et al., 1982), or facilitate fermentable substrate penetration to deeper regions of the biofilm, where the reduced fixed buffer effect from lower numbers of bacteria may allow a more pronounced pH drop at the plaque enamel interface (Dibdin and Shellis, 1988; Zero et al., 1986). While with the *P. olivaceus* pulsed together with sucrose, the biofilm of the P group was less compact than that of the S group even though both groups contained the same amount of sucrose.

The ecological plaque hypothesis for dental caries (Marsh and Bradshaw, 1997) states that generation of a low pH environment from sugar fermentation results in an ecological shift to a more cariogenic microflora. The results presented here, using the continuous culture system, confirm that the ecological responses were consistent with this hypothesis. With carbohydrate supplement in the flow cell, the population of cariogenic bacteria decreased significantly, together with more extracellular polysaccharides and more compact biofilm structure. While with the *P. olivaceus* pulsed in (P group), the growth of *S. mutans*, one of the primary cariogenic organisms, was inhibited significantly when compared with

**Comment [p14]:** You should put a, b, ... to separate significant means



**Figure 2.** The micro-radiograph taken from scanning electron microscope of the enamel sections of HA discs under a) pulses of sucrose, sodium fluoride and chlorhexidine digluconate, b) pulses of sucrose alone and c) pulses of sucrose and *Polistes olivaceous* (Magnification 5000x).

that of the S group. The microstructure of the biofilm was also less compact and pH recovery was accelerated. All these results indicated that cariogenic shift of plaque was inhibited to some extent by the pulsed medicine extract, *P. olivaceous*.

Current oral rinses contain chlorhexidine which may induce discomfort after long term use. Further studies may involve monitoring of the long term use of *P. olivaceous* to determine its suitability and long term safety profile.

In summary, *P. olivaceous* may help in the clearance of acid accumulation from carbohydrate metabolism. A lower level of colonization of the HA surface was observed in the *P. olivaceous* pulsed group and this was reflected in both the total viable count and biofilm imaging, which has less cariogenic bacteria and a less compact biofilm (Xiao et al., 2006, 2007). These revealed that *P. olivaceous* could suppress bacteria growth, as well as accelerating pH recovery. These events could reduce the pain due to acids and cytotoxins produced by oral bacteria and could

help patients recover from tooth decay (Xiao et al., 2006, 2007). From the present study, *P. olivaceus* may have inhibitory effects on the cariogenic shift of oral biofilm and it appears to be a promising source of new agents that may prevent dental caries. A phase I clinical trial would soon be conducted to investigate the efficacy of direct application of *P. olivaceus* in mouth rinse.

## ACKNOWLEDGEMENT

We were grateful to Professor J Wei for helping recruiting healthy subjects in Ningxia Medical University in the pilot study of the minimal and optimal concentration of *P. olivaceus*.

## REFERENCE

- Wan AKL, Seow WK, Walsh LJ, Bird PS (2002). Comparison of five selective media for the growth and enumeration of *Streptococcus mutans*. *Aust. Dental J.* 47: 21-26
- Anwar H, Strap JL, Costerton JW (1992). Establishment of aging biofilms: possible mechanism of bacterial resistance to antimicrobial therapy. *Antimicrob. Agents Chemother.* 36: 1347-1351.
- Beighton D, Hayday H (1986). The influence of diet on the growth of streptococcal bacteria on the molar teeth of monkeys (*Macaca fascicularis*). *Arch. Oral Biol.* 31: 449-454.
- Bortolaia C, Sbordone L (2002). Biofilms of the oral cavity. Formation, development and involvement in the onset of diseases related to bacterial plaque increase. *Minerva Stomatol.* 51: 187-192.
- Bradshaw DJ, McKee AS, Marsh PD (1989). Effects of carbohydrate pulses and pH on population shifts within oral microbial communities in vitro. *J. Dental Res.* 68: 1298-1302.
- Brown MR, Gilbert P (1993). Sensitivity of biofilms to antimicrobial agents. *J. Appl. Bacteriol.* 74(Suppl): 87S-97S.
- Cragg GM, Newman DJ, Snader KM (1997). Natural products in drug discovery and development. *J. Nat. Prod.* 60: 52-60.
- De Stoppelaar JD, Van Houte J, Backer DO (1970). The effect of carbohydrate restriction on the presence of *Streptococcus mutans*, *Streptococcus sanguis* and iodophilic polysaccharide-producing bacteria in human dental plaque. *Caries Res.* 4: 114-123.
- Dennis DA, Gawronski TH, Sudo SZ, Harris RS, Folke LE (1975). Variations in microbial and biochemical components of four-day plaque during a four-week controlled diet period. *J. Dental Res.* 54: 716-722.
- Dibdin GH, Shellis RP (1988). Physical and biochemical studies of *Streptococcus mutans* sediments suggest new factors linking the cariogenicity of plaque with its extracellular polysaccharide content. *J. Dental Res.* 67: 890-5.
- Gander S (1996). Bacterial biofilms: resistance to antimicrobial agents. *J. Antimicrob. Agents Chemother.* 37: 1047-1050.
- Hardie JM (1993). Oral microbiology: current concepts in the microbiology of dental caries and periodontal disease. *Br. Dental J.* 172: 271-278.
- Harvey A (2000). Strategies for discovering drugs from previously unexplored natural products. *Drug Discov. Today*, 5:294-300.
- Hodgson RJ, Lynch RJ, Watson GK, Labarber R, Treloar R, Allison C (2001). A continuous culture biofilm model of cariogenic responses. *J. Appl. Microbiol.* 90: 440-448.
- Holt JG (1984). *Bergey's Manual of Systematic Bacteriology*. Baltimore, Williams & Wilkins, pp. 141-100
- Huang ZW, Zhou XD, Li J, Liu T, Li H, Zhu B (2003). The Effects of Traditional Chinese Medicines on the Adherence of *Streptococcus mutans* to Salivary Acquired Pellicle in vitro. *J. Sichuan Univ. (Medical Science Edition)*, 34: 135-137.
- Ingram GS, Silverstone LM (1981). A chemical and histological study of artificial caries in human dental enamel in vitro. *Caries Res.* 15:393-398.
- Kanchanakamol U, Srisilapanan P, Umprivan R (1993). Dental plaque removal in adults using a newly developed, Concept 45 Degrees toothbrush. *Int. Dent. J.* 43: 116-120.
- Katsura H, Tsukiyama RI, Suzuki A, Kobayashi M (2001). In vitro antimicrobial activities of bakuchiol against oral microorganisms. *Antimicrob. Agents Chemother.* 45: 3009-3013.
- Koo H, Gomes BP, Rosalen PL, Ambrosano GM, Park YK, Cury JA (2000). In vitro antimicrobial activity of propolis and *Arnica montana* against oral pathogens. *Arch. Oral Biol.* 45: 141-148.
- Koo H, Rosalen PL, Cury JA, Park YK, Bowen WH (2002). Effects of compounds found in propolis on *Streptococcus mutans* growth and on glucosyltransferase activity. *Antimicrob. Agents Chemother.* 46: 1302-1309.
- Kostiuchenkov VN, Farashchuk NF (1991). Study of biological activity of *Polistes olivaceus* infusions during physical exercise. [Article in Russian] *Kosmicheskaiia Biologiia I Aviakosmicheskaiia Meditsina.* 25: 58-59.
- Lee SS, Zhang W, Li Y (2004). The antimicrobial potential of 14 natural herbal dentifrices: results of an in vitro diffusion method study. *J. Am. Dent. Assoc.* 135: 1133-1141.
- Li XC, Cai L, Wu CD, (1997). Antimicrobial compounds from *Ceanothus americanus* against oral pathogens. *Phytochemistry*, 46: 97-102.
- Lin MC, Lin JH, Wen KC (2001). Detection and Determination of Phenformin in Chinese Medicinal Capsules by GC-MS and HPLC. *J. Food Drug Anal.* 9: 139-144.
- Marsh PD, Bradshaw DJ (1997). Physiological approaches to the control of oral biofilms. *Adv. Dent. Res.* 11: 176-185.
- McNee SG, Geddes DA, Weetman DA, Sweeney D, Beeley JA (1982). Effect of extracellular polysaccharides on diffusion of NaF and [14C]-sucrose in human dental plaque and in sediments of the bacterium *Streptococcus sanguis* 804 (NCTC 10904). *Arch. Oral Biol.* 27: 981-986.
- Minah GE, Solomon ES, Chu K (1985). The association between dietary sucrose consumption and microbial population shifts at six oral sites in man. *Arch. Oral Biol.* 30: 397-401.
- Sauvetre E, Rozow A, de Meel H, Richebe A (1995). Comparison of the clinical effectiveness of a single and a triple-headed toothbrushes in a population of mentally retarded patients. *Bull Group Int. Rech. Sci. Stomatol. Odontol.* 38: 115-119.
- Sharma NC, Galustians J, McCool JJ, Rustogi KN, Volpe AR (1994). The clinical effects on plaque and gingivitis over three-month's use of four complex-design manual toothbrushes. *J. Clin. Dent.* 5: 114-118.
- Singleton S, Treloar R, Warren P, Watson GK, Hodgson R, Allison C (1997). Methods for microscopic characterization of oral biofilms: analysis of colonization, microstructure, and molecular transport phenomena. *Adv. Dent. Res.* 11: 133-49.
- Southard GL, Boulware RT, Walborn DR, Groznik WJ, Thorne EE, Yankell SL (1984). Sanguinarine, a new antiplaque agent: retention and plaque specificity. *J. Am. Dent. Assoc.* 108: 338-341.
- Staat RH, Gawronski TH, Cressey DE, Harris RS, Folke LE (1975). Effects of dietary sucrose levels on the quantity and microbial composition of human dental plaque. *J. Dent. Res.* 54: 872-80.
- Taniguchi M, Kubo I (1993). Ethnobotanical drug discovery based on medicine men's trials in the African savanna: screening of east African plants for antimicrobial activity II. *J. Natl. Proc.* 56: 1539-1546.
- Van Palenstein Helderma WH, Ijsseldijk M, Huis in 't Veld JH (1983). A selective medium for the two major subgroups of the bacterium *Streptococcus mutans* isolated from human dental plaque and saliva. *Arch. Oral Biol.* 28: 599-603.
- Wennstrom J, Lindhe J (1985). Some effects of a Sanguinarine-containing mouthrinse on developing plaque and gingivitis. *J. Clin. Periodontol.* 12: 867-872.
- Xie H, Cook GS, Costerton JW, Bruce G, Rose TM, Lamont RJ (2000). Intergeneric communication in dental plaque biofilms. *J. Bacteriol.* 182: 7067-7069.
- Xiao J, Liu Y, Zuo YL, Li JY, Ye L, Zhou XD (2006). Effects of *Nidus Vespae* extract and chemical fractions on the growth and acidogenicity of oral microorganisms. *Arch. Oral Biol.* 51(9): 804-813
- Xiao J, Zuo Y, Liu Y, Li J, Hao Y, Zhou X (2007). Effects of *Nidus Vespae* extract and chemical fractions on glucosyl-transferases,

Comment [p15]: concentration

- adherence and biofilm formation of *Streptococcus mutans*. Arch. Oral Biol. 52: 869-875.
- Yankell SL, Emling RC, Perez B (1996). A six-month clinical evaluation of the Dentrust toothbrush. J. Clin. Dent. 7: 106-109.
- Yankell SL, Edvardsen S, Braaten S, Emling RC (1993). Laboratory and clinical evaluations of the Jordan Exact toothbrush. J. Clin. Dent. 4: 67-70.
- Zero DT, van Houte J, Russo J (1986). The intra-oral effect on enamel demineralization of extracellular matrix material synthesized from sucrose by *Streptococcus mutans*. J. Dent. Res. 65: 918-923.
- Zhang CM, Zhou ZY (2004). Clinical Application of Complex Decoction on the Treatment of Malignant Tumor. [Article in Chinese] Hunan Zhongyiyao daobao. 5: 1-10.