

Full Length Research Paper

Effect of soaking of seeds in potassium silicate and uniconazole on germination and seedling growth of tomato cultivars, Seogeon and Seokwang

Iyyakkannu Sivanesan¹, Moon Sook Son², Chae Shin Lim³ and Byoung Ryong Jeong^{1,2,4*}

¹Institute of Agriculture and Life Science, Gyeongsang National University, Jinju 660-701, Korea.

²Department of Horticulture, Division of Applied Life Science (BK21 Program), Graduate School, Gyeongsang National University, Jinju 660-701, Korea.

³Agriculture Technology Education Center, Gyeongsangnam-do Agricultural Research and Extension Services, Jinju 660-985, Korea.

⁴Research Institute of Life Science, Gyeongsang National University, Jinju 660-701, Korea.

Accepted 10 June, 2011

Experiments were conducted to investigate the effects of soaking seeds in potassium silicate and uniconazole on seed germination and seedling growth of two tomato cultivars. Tomato (*Lycopersicon esculentum* Mill. 'Seogeon and Seokwang') seeds were put in a Petri dish filled with 15 ml of a solutions containing either 50 or 100 mg L⁻¹ potassium silicate or uniconazole and were placed in an environment controlled chamber (25°C, 80% RH, dark) for 12 or 24 h. After the soaking treatment, seeds were washed in distilled water and were dried in a growth chamber (25°C, 80% RH, and in the dark) for 4 h. Seeds were sown in 288-cell (11 cc) plug trays containing a Tosilee medium and trays were layed out in a randomized complete block design on beds in a glasshouse. A nutrient solution was supplied uniformly for all treatments once a day through a sub-irrigation system. Soaking seeds in potassium silicate or uniconazole solution reduced germination percentage in both cultivars when compared to the control. In both cultivars, soaking treatment of uniconazole significantly reduced length of stem, hypocotyls, internode, leaf area and dry weight of stem and root, as compared to the control and other treatments. Root length increased significantly in all treatments when compared with the control. Hypocotyl length and plant height of 'Seogeon' seedlings were suppressed in the 100 mg L⁻¹ potassium silicate treatment as compared to the control and water soaking. In contrast, height of 'Seokwang' seedlings increased by potassium silicate treatment. The chlorophyll fluorescence ratio (Fv/Fm) increased by low concentration of uniconazole treatment as compared to the control and other treatments. The growth of tomato seedlings was efficiently regulated by uniconazole 50 mg L⁻¹ (12 h soaking) treatment.

Key words: Chlorophyll fluorescence, plant growth retardants, plug plants, potassium silicate, seed treatment, silicon, uniconazole.

INTRODUCTION

As a result of the merits of reduction of labor in mass production of uniform transplants and specialization of transplant production, plug transplants have been used by many growers. However, one of the problems in plug transplant production is stretchiness of transplants

caused by high plant densities in plug trays. Many methods, including plant growth retardants (PGRs), withholding water or nutrients, temperature control, clipping shoots and mechanical stimulation (brushing), are used to control transplant height. Among many methods used for the control of plant height, the use of PGRs is most popular. If PGRs are used to control plant height, height can be influenced by concentration (LeCain et al., 1986; McDaniel, 1986; Ruter, 1992), time of application (Miranda and Carlson, 1980; Gilbertz, 1992), mode of

*Corresponding author. E-mail: brjeong@gnu.ac.kr. Tel: +82-55-772-1913. Fax: +82-55-757-7542.

Table 1. Treatments used in this experiment.

Treatment	Chemical	Concentration (mg L ⁻¹)	Soaking duration (h)
1	Control (unsoaked)	0	0
2	Distilled water	0	12
3		0	24
4		50	24
5	Potassium silicate	100	24
6		50	12
7	Uniconazole	50	24
8		100	12
9		100	24

application (Cathey, 1975), formulation (McDaniel, 1986; Ruter, 1992), and media composition (Barrett, 1982). In many cases, by the time growers apply PGR, stretching of the hypocotyl had already occurred and application to seedlings are ineffective (Pasian and Bennett, 2001). Seedling height can be controlled by applying the PGR directly to the seed (Pasian and Bennett, 2001; Pill and Gunter, 2001; Still and Pill, 2003). Treatment of tomato seeds with hypertonic solutions containing uniconazole (0.1, 1, or 10 mg L⁻¹) was of little practical value in protecting seedlings from freeze damage, although T₅₀ (days) and seedling height was reduced (Davis et al., 1990). Soaking pepper seeds in 1 to 100 mg L⁻¹ uniconazole for 24, 72 or 120 h significantly suppressed hypocotyls length and seedling growth (Shin and Jeong, 2002). In our previous study, the growth of tomato seedlings was efficiently regulated by 100 mg L⁻¹ uniconazole (one day soaking) treatment, but significantly affected the percent emergence.

Silicon (Si) is the second most abundant element both on the surface of the Earth's crust and in the soils. Although silicon has not been considered to be an essential element for higher plants (Epstein, 1999), yet its beneficial effects have been demonstrated for many plants, especially when they are subjected to biotic or abiotic stresses (Ma and Yamaji, 2006; Sivanesan et al., 2011). Application of silicon has been reported to enhance stress (Gunes et al., 2007; Romero-Aranda et al., 2006), and disease resistance (Dannon and Wydra, 2004) in tomato. Addition of Si either increased or decreased plant height in many plant species (Kim et al., 2002; Gong et al., 2003; Mattson and Leatherwood, 2010). In marigold, application of silicon decrease seedlings height (Sivanesan et al., 2010).

This study was conducted to evaluate the effect of seed soaking treatment with potassium silicate and uniconazole on seed germination and seedlings growth of two tomato cultivars.

MATERIALS AND METHODS

Tomato (*Lycopersicon esculentum* Mill. 'Seogeon' and 'Seokwang')

seeds were placed in a Petri dish (87 x 15 mm) filled with a 15 ml solution containing either 50 or 100 mgL⁻¹ potassium silicate or uniconazole (MB cell, Seoul, Korea) (Table 1), and were placed in an environment controlled chamber (25°C, 80% RH and in the dark) for 12 or 24 h. After soaking, seeds were washed in distilled water and dried in a growth chamber (25°C, 80% RH, in the dark) for 4 h. Seeds were sown in 288-cell (11 cc) plug trays containing a commercial germination medium (Tosilee medium; Shinan Grow, Jinju, Korea), and were placed on germination beds with a fogging system for three days in a glasshouse. After seedlings had emerged, trays were laid out in a randomized complete block design on beds in a glasshouse. Each treatment consisted of three replicates. Nutrient composition (Shin et al., 2009) was supplied uniformly for all treatments once a day through a sub-irrigation system. Plant height, stem diameter, hypocotyls length, number of leaves, leaf area, root length, and fresh and dry weights of shoot and root were recorded after 32 days of cultivation. Leaf area was determined with a leaf area meter (LI-3100 area meter, LI-COR. Inc., Lincoln, Nebraska, USA). Dry weight was measured after 72 h of drying at 60°C in a dry oven. Chlorophyll fluorescence parameters were measured using a chlorophyll fluorometer (PAM-2100, Walz, Germany). Seedlings were dark adapted for at least 30 min before the chlorophyll fluorescence measurements. Temperatures of the greenhouse were measured during the experimental period by digital thermometers (Thermo Recorder TR-71S, T&D Crop., Japan). Maximum, minimum and mean temperatures of the greenhouse during the culture period were 35.4, 12.6 and 24.4°C, respectively. Data collected were analyzed for statistical significance by the SAS (Statistical Analysis System, V. 6.12, Cary, NC, USA) program. The experimental results were submitted to an analysis of variance (ANOVA).

RESULTS AND DISCUSSION

Differences among the two cultivars in their responses were observed in germination and seedling growth. Soaking seeds in potassium silicate or uniconazole solution reduced germination percentage in both cultivars when compared with the control. However, high concentration (100 mg L⁻¹) of uniconazole and 24 h soaking period decreased germination percentage significantly (Figure 1). These results are in agreement with other report (Pasian and Bennett, 2001; Shin and Jeong, 2002; Still and Pill, 2003; Shin et al., 2009); they reported that seed germination was lower with an increase in PGRs concentration and soaking duration of seeds. This might




Figure 1. Effect of soaking seeds in potassium silicate and uniconazole on germination of tomato seeds recorded at 9 days after sowing in plug trays.

be due to inhibition of gibberellins biosynthesis. Gibberellins are known to stimulate germination of dormant as well as non-dormant seeds of several plant species (Bewley and Black, 1982). The growth retardants paclobutrazol, ancymidol and uniconazole are known to block the oxidation of ent-kaurene to ent-kaurenoic acid which is a step in the GA synthesis pathway of gibberellins (Pressman and Shaked, 1991).

Plant height, length of hypocotyl and inter node, number of leaves and leaf area were significantly affected by the treatments (Figure 2). In 'Seogeon', plant height and internode length were suppressed in the 100 mg L⁻¹ potassium silicate treatment, as compared to the control and distilled water soaking, while plant height was not reduced in 'Seokwang'. Application of Si either increased or decreased plant height in many plant species (Kim et al., 2002; Gong et al., 2003; Mattson and Leatherwood, 2010). Uniconazole treatment reduced plant height and internode length of the two tomato cultivars when compared with other treatments and control, however, increasing concentration of uniconazole or soaking duration did not affect the plant height significantly (Table 2). Therefore, soaking seeds in the 50 mg L⁻¹ uniconazole solution significantly reduced plant height with good germination percentage. The plant growth retardant uniconazole is known to affect the levels of endogenous

gibberellins. Gibberellins are known to induce elongation (Kurepin et al., 2006), while triazole PGRs reduces gibberellins levels and cause a decrease in shoot growth (Davis and Curry, 1991). Uniconazole reduce stem length in many plant species such as hibiscus (Wang and Gregg, 1991), chrysanthemum (Schuch, 1994), kalanchoe (Hwang et al., 2008, 2009) and pepper (Shin and Jeong, 2002). In both cultivars, soaking seeds in distilled water, potassium silicate or uniconazole solution reduced stem diameter and number of leaves when compared with the control. Leaf area was not reduced when the seeds were treated with potassium silicate, but significantly was reduced by uniconazole treatment when compared with the control (Table 2).

Root length was increased in all treatments when compared with the control. When seeds were treated with potassium silicate at 50 and 100 mg L⁻¹, there was increase in root length in 'Seogeon' and 'Seokwang', respectively (Table 3). The highest root length was observed in 'Seogeon' when seeds were soaked in distilled water for 12 h, while in 'Seokwang', at 50 mg L⁻¹ uniconazole, there was significant increase in the root length when compared with the control. Fresh and dry weights of shoot significantly decreased when seeds were soaked in 50 or 100 mg L⁻¹ uniconazole. Soaking seeds in potassium silicate solution increased the fresh



Figure 2. Effect of seed treatment with potassium silicate and uniconazole on growth of tomato seedlings measured at 32 days after sowing in plug trays. A, 'Seogeon'; B, 'Seokwang'

Table 2. Effect of seed treatment in potassium silicate and uniconazole on growth characteristics of tomato seedlings measured at 32 days after sowing in plug trays.

Cultivar (A)	Treatment (B)	Plant height (cm)	Stem diameter (mm)	Hypocotyl length (cm)	Internodal length (cm)	Number of leaf	Leaf area (cm ²)
Seogeon	1	27.2 ^{bc1}	2.60 ^{cd}	7.53 ^b	9.91 ^{bc}	4.2 ^{e-g}	38.0 ^{de}
	2	27.8 ^b	2.54 ^{cd}	8.12 ^a	10.40 ^{ab}	4.1 ^g	39.9 ^{b-d}
	3	28.1 ^{ab}	2.50 ^d	8.46 ^a	10.43 ^{ab}	3.9 ^g	35.8 ^e
	4	25.3 ^d	2.53 ^{cd}	6.59 ^c	8.76 ^e	3.9 ^g	36.2 ^{de}
	5	17.1 ^e	2.28 ^e	4.77 ^d	4.64 ^f	4.0 ^g	27.5 ^f
	6	11.1 ^{f-i}	2.10 ^f	3.42 ^e	2.13 ^g	4.0 ^g	20.2 ^g
	7	10.5 ^{hi}	2.28 ^e	3.31 ^e	2.20 ^g	4.1 ^g	19.3 ^g

Table 2. continues.

	8	10.6 ^{hi}	2.13 ^{ef}	3.39 ^e	1.99 ^g	4.0 ^g	18.4 ^g
	9	10.4 ⁱ	2.14 ^{ef}	3.36 ^e	1.96 ^g	4.0 ^g	18.9 ^g
Seokwang	1	25.1 ^d	2.88 ^a	6.96 ^c	8.92 ^{de}	5.1 ^a	43.9 ^a
	2	29.0 ^a	2.69 ^{bc}	7.97 ^{ab}	10.69 ^a	4.8 ^{a-c}	43.2 ^{ab}
	3	27.8 ^b	2.59 ^{cd}	7.92 ^{ab}	10.05 ^{a-c}	4.7 ^{b-d}	42.2 ^{a-c}
	4	24.9 ^d	2.68 ^{bc}	6.96 ^c	9.56 ^{cd}	4.7 ^{b-d}	39.1 ^{c-e}
	5	26.7 ^c	2.84 ^{ab}	8.44 ^a	10.23 ^{a-c}	4.9 ^{ab}	45.4 ^a
	6	12.0 ^f	2.26 ^{ef}	3.33 ^e	2.07 ^g	4.6 ^{b-d}	20.6 ^g
	7	10.9 ^{g-i}	2.28 ^e	3.28 ^e	1.99 ^g	4.4 ^{d-f}	19.6 ^g
	8	11.8 ^{fg}	2.21 ^{ef}	3.54 ^e	1.96 ^g	4.1 ^{fg}	19.2 ^g
	9	11.5 ^{f-h}	2.13 ^{ef}	3.30 ^e	2.00 ^g	4.5 ^{c-e}	18.3 ^g
A		***	***	**	***	***	***
B		***	***	***	***	***	***
A*B		***	***	***	***	*	***

¹Means separation within columns by Duncan's multiple range test ($P \leq 0.05$). NS, non-significant; *significant at $P = 0.05$; ** $P = 0.01$; *** $P = 0.001$, respectively.

Table 3. Effect of seed treatment in potassium silicate and uniconazole on root length, fresh and dry weights of tomato seedlings measured at 32 days after sowing in plug trays.

Cultivar (A)	Treatment (B)	Root length (cm)	Fresh weight (mg)		Dry weight (mg)	
			Shoot	Root	Shoot	Root
Seogeon	1	8.20 ^{c-f1}	2.42 ^{bc}	0.31 ^{b-e}	0.16 ^{bc}	0.025 ^{a-c}
	2	10.26 ^a	2.41 ^{bc}	0.30 ^{c-f}	0.15 ^{cd}	0.027 ^a
	3	9.42 ^{a-d}	2.18 ^d	0.27 ^{d-g}	0.14 ^{de}	0.017 ^{c-e}
	4	9.04 ^{a-d}	2.22 ^{cd}	0.29 ^{c-g}	0.14 ^{c-e}	0.017 ^{c-e}
	5	8.52 ^{b-f}	1.80 ^e	0.32 ^{b-d}	0.12 ^{ef}	0.018 ^{c-e}
	6	8.81 ^{b-e}	1.27 ^f	0.32 ^{b-d}	0.10 ^{f-h}	0.019 ^{c-e}
	7	9.52 ^{a-c}	1.38 ^f	0.37 ^{ab}	0.09 ^{gh}	0.020 ^{b-d}
	8	9.17 ^{a-d}	1.28 ^f	0.38 ^a	0.08 ^{gh}	0.025 ^{ab}
	9	9.67 ^{ab}	1.26 ^f	0.32 ^{b-d}	0.07 ^h	0.022 ^{a-c}
Seokwang	1	7.65 ^{ef}	2.56 ^{ab}	0.24 ^g	0.19 ^a	0.013 ^e
	2	8.65 ^{b-f}	2.52 ^{ab}	0.26 ^{e-g}	0.17 ^{bc}	0.016 ^{c-e}
	3	8.75 ^{b-e}	2.53 ^{ab}	0.32 ^{b-d}	0.18 ^{ab}	0.017 ^{c-e}
	4	8.93 ^{a-e}	2.28 ^{cd}	0.30 ^{c-f}	0.17 ^{bc}	0.018 ^{c-e}
	5	9.37 ^{a-d}	2.69 ^a	0.33 ^{a-c}	0.20 ^a	0.014 ^{de}
	6	9.46 ^{a-c}	1.27 ^f	0.24 ^g	0.09 ^{hg}	0.018 ^{c-e}
	7	8.34 ^{c-f}	1.30 ^f	0.32 ^{b-d}	0.12 ^{ef}	0.025 ^{ab}
	8	8.12 ^{d-f}	1.20 ^f	0.25 ^{fg}	0.08 ^{gh}	0.017 ^{c-e}
	9	8.48 ^{b-f}	1.23 ^f	0.29 ^{c-g}	0.10 ^{fg}	0.018 ^{c-e}
A		**	***	***	***	**
B		*	***	*	***	NS
A*B		*	***	**	***	**

¹Means separation within columns by Duncan's multiple range test ($P \leq 0.05$). NS, non significant and significant; * $P = 0.05$; ** $P = 0.01$; *** $P = 0.001$.

weight of shoot in 'Seokwang', while it was decreased in 'Seogeon' when compared with the control. In both

cultivars, fresh weight of root increased significantly when seeds were treated with uniconazole, however, root dry




Figure 3. Treatments induction changes in chlorophyll fluorescence (Fv/Fm) ratio of tomato seedlings.

weight was reduced in 'Seogeon' (Table 3).

Photosynthesis is an important parameter used to monitor plant response to abiotic stress. Chlorophyll fluorescence is a rapid, non-destructive method to detect changes in the photosynthetic systems induced by stress. Measurement of Fv/Fm ratio value varied between the two cultivars (Figure 3). In 'Seogeon', distilled water and the 24 h uniconazole (100 mg L^{-1}) soaking treatment decreased Fv/Fm ratio, while potassium silicate and uniconazole (50 mg L^{-1}) soaking treatment increased the Fv/Fm ratio when compared with the control. In 'Seokwang', all the treatments increased the Fv/Fm ratio, except 24 h soaking in the distilled water when compared with the control. Low concentration of potassium silicate increased the Fv/Fm ratio in both cultivars. High concentration of uniconazole and the 24 h soaking period decreased the Fv/Fm ratio in both cultivars which reflected stressful condition of seedlings, while low concentration of uniconazole increased the Fv/Fm ratio, indicating that the seedlings were normal.

In conclusion, a soaking treatment of uniconazole 50 mg L^{-1} significantly reduced seedlings height with good germination percentage and a mean value of Fv/Fm ratio was typical of a well functioning photosynthetic apparatus. Hypocotyl length and plant height of 'Seogeon' were suppressed in the 100 mg L^{-1} potassium

silicate treatment as compared to the control. These results would be helpful in allowing plant nurseries to control the stretching of tomato seedling.

ACKNOWLEDGEMENTS

This research was supported by the Technology Development Program for Agriculture and Forestry, Ministry for Food, Agriculture, Forestry and Fisheries, Republic of Korea (Project No 108102-5). Moon Sook Son was supported by a scholarship from the BK21 Program, the Ministry of Education, Science and Technology, Korea.

REFERENCES

- Barrett JE (1982). Chrysanthemum height control by ancymidol, pp333, and EL-500 dependent on medium composition. Hort. Sci. 17: 896-897.
- Bewley JD, Black M (1982) Physiology and biochemistry of seeds in relation to germination. II. Viability, dormancy and environment control. Springer-Verlag, Berlin.
- Cathey HM (1975). Comparative plant growth retarding activities of ancymidol with ACPC, phosfon, chlormequat, and SADH on ornamental plant species. Hort. Sci., 10: 204-216.
- Dannon EA, Wydra K (2004). Interaction between silicon amendment, bacterial wilt development and phenotype of *Ralstonia solanacearum* in tomato genotypes. Physiol. Mol. Plant Pathol. 64: 233-243.

- Davis TD, Curry EA (1991). Chemical regulation of vegetative growth. *Crit. Rev. Plant Sci.* 10: 151-188.
- Davis TD, Eils JE, Walser RH (1990). Emergence, growth, and freezing tolerance of tomato seedlings grown from uniconazole-treated seed. *Hort. Sci.* 25: 312-313.
- Epstein E (1999). Silicon. *Ann. Rev. Plant Physiol. Plant Mol. Biol.* 50: 641-664.
- Gilbert DA (1992). Chrysanthemum response to timing of paclobutrazol and uniconazole sprays. *Hort. Sci.*, 27: 322-323.
- Gong H, Chen K, Chen G, Wang S (2003). Effects of silicon on growth of wheat under drought. *J. Plant Nutr.* 26: 1055-1063.
- Gunes A, Inal A, Bagci EG, Pilbeam DJ (2007). Silicon-mediated changes of some physiological and enzymatic parameters symptomatic for oxidative stress in spinach and tomato grown in sodic-B toxic soil. *Plant Soil*, 290: 103-114.
- Hwang SJ, Lee MY, Sivanesan I, Jeong BR (2008). Growth control of kalanchoe cultivars Rako and Gold Strike by application of paclobutrazol and uniconazole as soaking treatment of cuttings. *Afr. J. Biotechnol.* 7: 4212-4218.
- Hwang SJ, Lee MY, Sivanesan I, Jeong BR (2009). Suppression of stretchiness in pot *Kalanchoe blossfeldiana* Poelln. 'Rako' by application of plant growth retardants as recycled subirrigational supply. *Propag. Orn. Plants*, 9: 26-34.
- Kim SG, Kim KW, Park EW, Choi DI (2002). Silicon induced cell wall fortification of rice leaves: A possible cellular mechanism of enhanced host resistance to blast. *Phytopathology*, 92: 1095-1103.
- Kurepin LV, Pharis RP, Richard P, Reeid DM, Chinnappa CC (2006). Involvement of gibberellins in the stem elongation of sun and shade ecotypes of *Stellaria longipes* that is induced by low light irradiance. *Plant Cell Environ.* 29: 1319-1328.
- LeCain DR, Schekel KA, Wample RL (1986). Growth-retarding effects of paclobutrazol on weeping fig. *Hort. Sci.*, 21: 1150-11520.
- Ma JF, Yamaji N (2006). Silicon uptake and accumulation in higher plants. *Trends Plant Sci.* 11: 392-397.
- Mattson NS, Leatherwood WR (2010). Potassium silicate drenches increased leaf silicon content and affect morphological traits of several floricultural crops grown in a pear-based substrate. *Hort. Sci.*, 45: 43-47.
- McDaniel GL (1986). Comparison of paclobutrazol, flurprimidol and tetcyclacis for controlling poinsettia height. *Hort. Sci.*, 21: 1161-1163.
- Miranda RM, Carlson WH (1980). Effect of timing and number of applications of chlormequat and ancymidol on the growth and flowering of seed geraniums. *J. Am. Soc. Hort. Sci.* 105: 273-277.
- Pasian CC, Bennett MA (2001). Paclobutrazol soaked marigold, geranium, and tomato seeds produce short seedlings. *Hort. Sci.*, 36: 721-723.
- Pill WG, Gunter JA (2001). Emergence and shoot growth of cosmos and marigold from paclobutrazol-treated seed. *J. Environ. Hort.* 19: 11-14.
- Pressman E, Shaked R (1991). Interactive effects of Gas, CKs and growth retardants on the germination of celery seeds. *Plant Growth Regul.* 10: 65-72.
- Romero-Aranda MR, Jurado O, Cuartero J (2006). Silicon alleviates the deleterious salt effect on tomato plant growth by improving plant water status. *J. Plant Physiol.* 163: 847-855.
- Ruter JM (1992). Growth and flowering response of butterfly-bush to paclobutrazol formulation and rate of application. *Hort. Sci.*, 27: 929.
- Schuch UK (1994). Response of chrysanthemum to uniconazole and daminozide applied as dip to cuttings or as foliar spray. *J. Plant Growth Regul.* 13: 115-121.
- Shin WG, Hwang SJ, Sivanesan I, Jeong BR (2009). Height suppression of tomato plug seedlings by an environment friendly seed treatment of plant growth retardants. *Afr. J. Biotechnol.* 8: 4100-4107.
- Shin WG, Jeong BR (2002). Seed treatment of growth retardants for height suppression of pepper plug seedlings. *J. Kor. Soc. Hort. Sci.* 43: 565-570.
- Sivanesan I, Son MS, Lee JP, Jeong BR (2010). Effects of Silicon on Growth of *Tagetes patula* L. 'Boy Orange' and 'Yellow Boy' Seedlings Cultured in an Environment Controlled Chamber. *Propag. Orn. Plants*, 10: 136-140.
- Sivanesan I, Song JY, Hwang SJ, Jeong BR (2011). Micropropagation of *Cotoneaster wilsonii* Nakai-a rare endemic ornamental plant. *Plant Cell Tissue Organ Cult.* 105: 55-63.
- Still JR, Pill WG (2003). Germination and emergence, and seedling growth of tomato and impatiens in response to seed treatment with paclobutrazol. *Hort. Sci.*, 38: 1201-1204.
- Wang YT, Gregg (1991). Modification of Hibiscus growth by treating unrooted cuttings and potted plants with uniconazole or paclobutrazol. *J. Plant Growth Regul.* 10: 47-51.