

Full Length Research Paper

Sorghum stem yield and soluble carbohydrates under different salinity levels

A. Almodares^{1*}, M. R. Hadi² and H. Ahmadpour¹

¹Department of Biology, Faculty of Science, University of Isfahan-Iran.

²Research Department of Potato Biotechnology, University of Isfahan-Iran.

Accepted 25 August, 2008

The aim of this study was to select the most suitable cultivar for salty land in this geographical area. Two sweet sorghum cultivars (Keller and Sofra) and one grain sorghum cultivar (Kimia) were grown in greenhouse benches under four salinity levels of 2, 4, 8 and 12 dSm⁻¹ to evaluate the effects of salinity on stem yield and soluble carbohydrate (sucrose, glucose and fructose). The results showed that in all cultivars as salinity increased, the amount of stem yield and soluble carbohydrate decreased. In all salt concentrations, Keller and Kimia had the highest and the lowest stem yield and sucrose, respectively. At the highest salt concentration (12 dSm⁻¹), Keller had the lowest stem yield reduction (less than 1%) and the highest sucrose content while Kimia had the highest stem yield reduction (more than 18%) and the lowest sucrose content. Therefore, Keller and Kimia can be considered as salt tolerance and salt sensitive cultivars, respectively. As salinity increased, the amount of glucose and fructose in Keller decreased while they increased in Sofra. Increasing glucose and fructose in Sofra is not an indication of its salt tolerance. At the physiological maturity stage, the plant has the highest stem yield and sucrose content while it has the lowest glucose and fructose content than flowering stage. Based on the results, Keller is recommended to be planted under soil salinity conditions and harvested at physiological maturity stage.

Key words: Sweet sorghum, grain sorghum, salinity, stem yield, sucrose, glucose, fructose.

INTRODUCTION

Soil salinity is one of the main problems for agriculture, especially in countries where irrigation is an essential aid to agriculture (Ahloowalia et al., 2004). Saline soils are estimated about 5 – 10% of the world's arable land (Szabolcs, 1994), and the area affected by salinity is increasing steadily (Ghassemi et al., 1995). There are vast areas in Iran with salinity-affected soils and as Kehl (2006) reported, moderate saline soils occupy approximately 25.5 million ha and strong saline soils cover about 8.5 million ha. Soil salinity reduces yield production of most crops (Munns et al., 2002). Therefore, there is a need to improve salinity tolerance of important crops. Sorghum [*Sorghum bicolor* (L.) Moench] is a potential crop for moderately saline areas (Almodares and Sharif, 2007) and shown to contain intraspecific variability for salinity (Igartua et al., 1995). However, Salinity reduced sorghum growth and biomass production (Ibrahim, 2004).

Nevertheless, the development of high-yielding salinity-tolerant sorghums is the best option to increase the productivity in such soils (Igartua et al. 1994). Krishnamurthy et al. (2007) reported that there are large genotypic variations for tolerance to salinity in sorghum. Ibrahim (2004) reported that in sorghum, total soluble sugar increased with increasing salinity level. Sucrose content of sorghum could be an indicator for its salt tolerance (Juan et al., 2005). In sorghum, the fructose level was always higher than that of the glucose in response to various salinity treatments (Gill et al., 2001). The present study was conducted to determine how salinity affects stem yield, sucrose and invert sugars (glucose and fructose) of two sweet and one grain sorghum cultivars.

MATERIALS AND METHODS

Experimental location, plant material and experimental design

This experiment was conducted in the Isfahan University green

*Corresponding author. E-mail: aalmodares@yahoo.com.

house benches (3 m long, 1.2 m length and 0.5 m depth) in 2006. Seed of two popular sweet sorghum and one grain sorghum [*Sorghum bicolor* (L.) Moench] cultivars were provided from Isfahan University Research Station in Iran. The experimental design was split-split plot in complete randomize block with three replications. Four levels of salinity (2, 4, 8 and 12 dSm⁻¹) were assigned to the main plots. Two sweet sorghum (Keller and Sofra) and grain sorghum (Kimia) cultivars were assigned to the subplots. Two harvesting dates (flowering and physiological maturity) were assigned to the sub-subplots. Each sub-subplot consisted of 3 rows 3 m long and 0.4 m apart. The inter-row space was 0.1 m.

Plant growth and salt treatment

Seeds of the above cultivars were planted in the benches. Salt treatments were started six weeks after germination following plant establishment (Almodares et al., 2008). The plants were irrigated with NaCl solution based on the experimental design once a week. Plants were grown under stress conditions until physiological maturity stage. Plants were harvested at flowering and physiological maturity.

Measurement of stem yield and carbohydrate content

Three plants samples were taken at flowering and physiological maturity. After the leaves and panicle were removed, stalk was weighed and immediately placed into 100°C oven. They were kept at that temperature for 1.5 h, and then dried at 70°C. After oven drying, the samples were ground and passed through 0.05 mm screen, mixed and stored for carbohydrates analysis. Sucrose and invert sugar were extracted with 80% ethanol from 100 mg of sample by AOAC method (1975). Sucrose content was determined according to Varma (1988) and invert sugar according Lane-Eynon method (1970). The amount of glucose was determined by Pearson (1970) method. Fructose content was determined by subtracting glucose from invert sugar (total of glucose and fructose).

Statistical analysis

Statistical analysis was performed using Statistical Analysis System (SAS) computer program. The means were compared according to Duncan multiple rang test.

RESULTS AND DISCUSSION

The effect of salinity on the stem yield and sucrose was significant at 1% level and on glucose and fructose at 5% level. As salt concentration increased, the stem yield decreased (Figure 1). It was 54.07 t h⁻¹ at 2 dSm⁻¹ and 50.65 t h⁻¹ at 12 dSm⁻¹. El-Sayed et al. (1994) reported that sorghum growth was significantly reduced from 50 to 150 mM NaCl. Also, Almodares and Sharif (2007) indicated the salinity of water has an adverse effect on sweet sorghum biomass. They reported that sweet sorghum cultivar SSV108 and Rio had the lowest and the highest biomass under water qualities of 2, 5, and 8 dS m⁻¹. In addition, Silva et al. (2003) showed that salt stress reduced root and shoot dry matter. Therefore, it seems that as Netondo et al. (2004) mentioned, sorghum grown in salt affected soils may suffer from drought stress, ion toxicity, and mineral deficiency leading to reduced growth and productivity. As salt concentration increased, the




Figure 1. Mean comparisons^{*} among salt concentrations for stem yield in sweet sorghum cultivars. Values within each column followed by the same letter are not significantly different at 5% level, using Duncan multiple rang test.




Figure 2. Mean comparisons^{*} among salt concentrations for percent carbohydrates in sweet sorghum cultivars. Values within each carbohydrate column followed by the same letter are not significantly different at 5% level, using Duncan multiple rang test.

amount of sucrose, glucose and fructose decreased (Figure 2). The amount of sucrose was 17.4% at 2 dSm⁻¹ and 12.8% at 12 dSm⁻¹. The amount of glucose and fructose was not significantly different at 2 dSm⁻¹ and 4 dSm⁻¹. However, they decreased significantly at 8 dSm⁻¹ and 12dSm⁻¹. Almodares et al. (2008) reported that in sweet sorghum, as salinity increased, the amount of sucrose and glucose in Sofra cultivar decreased. Also, Anjum (2008) reported that concentrations of sugars i.e. sucrose, glucose and fructose in the leaves of Cleopatra mandarin and both leaves and roots of Troyer citrange in seedling stage decreased with increase in salinity level in the irrigation water containing 0, 40 or 80 mM NaCl for 12 weeks. On the contrary, Serraj and Sinclair (2002) reported that salt-stressed sorghum plants additionally accumulate sugars but its accumulation can be due to a reduced utilization during salt stress period. However, reduction in carbohydrates concentrations has been related to tissue re-hydration during stress recovery by some authors (Lacerda et al., 2005). Therefore, the increases of carbohydrate content under salt stress have not been detected here under the experimental conditions used. It seems that carbohydrates reduction

Table 1. Interaction* between salt concentrations and cultivars for measured characteristics in sweet sorghum.

Salt concentration (dSm ⁻¹)	Cultivar	Stem yield (t ha ⁻¹)	Sucrose (%)	Glucose (%)	Fructose (%)
2	Keller	67.43 ^a	21.73 ^d	6.50 ^b	3.06 ^b
	Sofra	56.00 ^b	18.66 ^e	5.70 ^e	3.66 ^a
	Kimia	39.20 ^e	12.03 ⁱ	4.56 ^f	2.16 ^e
4	Keller	67.60 ^a	22.26 ^c	5.86 ^d	2.86 ^{bc}
	Sofra	56.70 ^b	18.00 ^f	5.80 ^{de}	3.73 ^a
	Kimia	36.26 ^f	11.30 ^j	4.30 ^g	1.96 ^e
8	Keller	67.60 ^a	22.43 ^b	6.60 ^c	2.66 ^{cd}
	Sofra	53.67 ^c	15.80 ^g	6.40 ^b	3.73 ^a
	Kimia	34.40 ^g	7.20 ^k	4.43 ^{fg}	1.96 ^e
12	Keller	66.90 ^a	22.90 ^a	5.73 ^{de}	2.46 ^d
	Sofra	51.33 ^d	13.33 ^h	6.83 ^a	3.53 ^a
	Kimia	32.23 ^h	2.46 ⁱ	2.73 ^h	1.43 ^f

* Values within each column followed by the same letter are not significantly different at $P < 0.05$.

could be due to the long time exposure of plants under salinity and or high concentration of salt.

Interaction between salt concentrations and cultivars were significant at 1% level for stem yield and sucrose; and at 5% level for glucose and fructose. Keller and Kimia had the highest and lowest stem yield at all salt concentrations (Table 1). At 12 dSm⁻¹, stem yield of Keller and Kimia was 66.90 t ha⁻¹ and 32.23 t ha⁻¹, respectively.

De lacerda et al. (2003) compared seedlings of two forage sorghum genotypes (salt tolerance and salt sensitive) under salinity of 0 and 100 mM NaCl. They reported that biomass of salt sensitive plant was significantly reduced as salt concentration increased. Also, Sunseri et al. (1998) compared four sweet sorghum cultivars under soil salinity. They reported that salt tolerant lines did not show significant differences in leaf dry weight and stem dry weight, with increasing level of soil salt stress. While, the salt sensitive lines showed a different behaviour; significantly reducing yield performances with increasing level of soil salt stress. Thereby, it seems that Keller is more salt tolerant than Sofra and Kimia. Sucrose content of Keller was the highest at the 12 dSm⁻¹. As the amount of salt concentration increased, the sucrose content of Keller significantly increased.

Silva et al. (2003) reported that in sorghum under salt stress, the salt-tolerant cultivar showed greater enhancement in soluble carbohydrate content. They suggested that the accumulation of soluble carbohydrates were significantly related to salt tolerance in relation to leaf osmotic adjustment and soluble carbohydrate contents of leaves were significantly correlated with the acclimatization to salt stress. Thus, these parameters may be used as physiological markers of salt tolerance in sorghum. Also as Juan et al. (2005) reported that in tomato, sucrose content of plant parts is an indicator of salt tolerance. Therefore, it seems that Keller is more tolerant than both Sofra and Kimia. On the other hand, as salinity

increased, the amount of invert sugar in Keller decreased while the amounts of glucose and fructose of Sofra increased. Sofra had the highest amount of glucose and fructose at 12 dSm⁻¹ (6.83 and 3.53 %, respectively). Balibera et al. (1997) reported that in tomato as salinity increased, plant accumulates more sucrose in its stalk, root and leaf whereas the amount of hexose such as glucose and fructose decreased in salt tolerant cultivars. Therefore, it seems that increasing glucose and fructose in Sofra is not an indication of its salt tolerance but rather as salt sensitive cultivars.

Also, invert sugar reduction in Keller could be due to hydrolysis inhibition of sucrose or as Barreto et al. (1995) showed that reduction of hexose in sorghum could be due to plant energetic cost for active exclusion of Na⁺ as mechanism salt tolerance. The effect of harvesting stage on stem yield and carbohydrates were significant at 1% level (data not shown) and their interactions were significant at 1% for sucrose and at 5% for stem yield, glucose and fructose (Table 2). Stem yield and sucrose content of all cultivars were significantly higher at physiological maturity than flowering. In all salt concentrations Keller had the highest stem yield and sucrose at both flowering and physiological maturity. While, Kimia had the lowest stem yield and sucrose at both of the above stages. Almodares et al. (1994) reported that in sweet sorghum, biomass and carbohydrate content was higher at the physiological maturity than flowering. Tarpley and Vietor (2007) reported that amount of sucrose in sorghum culm is the highest at ripening stage (physiological maturity) and sucrose can be radially transferred to the intracellular compartment of mature ripening sorghum internode without being hydrolysed which is in agreement with our results. However, Tew and Cobill (2006) reported that in sorghum, mean sugar yields of M81E, Theis and Topper were not significantly different from each other at the each harvest date in July, August or September. Therefore, it seems that at the physiological

Table 2. Interaction between harvesting stage and cultivars for measured characteristics in sweet sorghum.

Harvesting stage	Cultivar	Stem yield (t ha ⁻¹)	Sucrose (%)	Glucose (%)	Fructose (%)
Flowering	Keller	65.63 ^b	17.70 ^c	6.10 ^b	3.76 ^a
	Sofra	51.33 ^d	11.73 ^d	7.53 ^a	3.96 ^a
	Kimia	33.80 ^f	7.10 ^f	5.13 ^d	2.73 ^c
Physiological maturity	Keller	69.10 ^a	26.93 ^a	5.93 ^c	1.73 ^d
	Sofra	57.46 ^c	21.16 ^b	4.83 ^e	3.36 ^b
	Kimia	37.20 ^e	9.40 ^e	2.83 ^f	1.03 ^e

* Values within each column followed by the same letter are not significantly different at P<0.05.

maturity stage the plant has the highest stem yield and sucrose content. In all cultivars, glucose and fructose at physiological maturity were lower than at flowering.

Lingle (1987) reported that in sorghum at physiological maturity stage, sucrose content increased while invert sugar decreased. The author indicated that at this stage, invert sugars (glucose and fructose) were converted to sucrose, which is in agreement with our results. Therefore, it seems that low sucrose content at flowering could be due high invertase activities which invert sucrose to glucose and fructose. In contrast, at physiological maturity due to low invertase activity, sucrose does not invert to glucose and fructose. Based on the above results, it seems that Keller is better adapted to salinity than others and it is recommended to plant Keller and harvest at physiological maturity in salty land.

ACKNOWLEDGMENTS

The authors wish to thank Dr. A. Mostajeran of the Department of Biology, University of Isfahan for his valuable cooperation in this research.

REFERENCES

- Ahloowalia BS, Meluzynski M, Nichterlein K (2004). Global impact of mutation-derived varieties. *Euphytica* 135: 187-204.
- Almodares A, Sharif ME (2007). Effects of irrigation water qualities on biomass and sugar contents of sugar beet and sweet sorghum cultivars. *J. Environ. Biol.* 28(2): 213-218.
- Almodares A, Hadi MR, Dosti B (2008). The effects of salt stress on growth parameters and carbohydrates contents in sweet sorghum. *Res. J. Environ. Sci.* 2(4): 298-304.
- Almodares A, Sephai A, Dallotojary H, Ghvami R (1994). Effects of phenological stages on biomass and carbohydrate contents of sweet sorghum cultivars. *Ann. Plant Physiol.* 8: 42-48.
- Anjum MA (2008). Effect of NaCl concentrations in irrigation water on growth and polyamine metabolism in two citrus rootstocks with different levels of salinity tolerance. *Acta. Physiol. Plant.* 30(1): 43-52.
- Association of Official Agricultural Chemist (AOAC) (1975). Official methods of analysis of the AOAC. 12th ed. Washington DC.
- Balibera ME, Rus-Alvarez AM, Bolarin MC, Perez-Alfocea F (1997). Fast changes in soluble carbohydrates and proline contents in tomato seedlings in response to ionic and non-ionic iso-osmotic stresses. *J. Plant Physiol. (Germany)* 151(2): 221-226.
- Barreto LP, Bezerra Neto E, Azevedo Neto AD, Tabosa JN, Santos PM, Mala JLT (1995). Seleção de materiais genéticos de sorgo forrageiro (*Sorghum bicolor*) com tolerância ao estresse salino. In: Congresso Brasileiro De Ciencia Do Solo, 25., Viçosa., Resumos expandidos. Viçosa, Sbcsof, 4v. il. p. 1.351-1.353.
- De Lacerda CF, Cambraia J, Oliva MA, Ruiz HA, Prisco JT (2003). Solute accumulation and distribution during shoot and leaf development in two sorghum genotypes under salt stress. *Environ. Exp. Bot.* 49(1): 107-120.
- El-Sayed HM, El-Haddad, O'Leary JW (1994). Effect of salinity and K/Na ratio of irrigation water on growth and solute content of *Atriplex amnicola* and *Sorghum bicolor*. *Irrig. Sci.* 14(3): 127-133.
- Ghassemi F, Jakerman AJ, Nix HA (1995). Salinisation of land and water resources: human causes, extent, management and case studies. CAB International, Wallingford, UK, p. 526.
- Gill PK, Sharma AD, Singh P, Bhullar SS (2001). Effect of various abiotic stresses on the growth, soluble sugars and water relations of sorghum seedlings grown in light and darkness. *Bulg. J. Plant Physiol.* 27(1-2): 72-84.
- Ibrahim AH (2004). Efficacy of exogenous glycine betaine application on sorghum plants grown under salinity stress. *Acta. Bot. Hungarica.* 43(3-4): 307-318.
- Igartua E, Gracia MP, Lasa JM (1994). Characterization and genetic control of germination, emergence responses of grain sorghum to salinity. *Euphytica* 76(3): 185-193.
- Igartua E, Gracia MP, Lasa JM (1995). Field responses of grain sorghum to a salinity gradient. *Field Crops Res.* 42: 15-25.
- Juan M, Rivero RM, Romero L, Ruiz JM (2005). Evaluation of some nutritional and biochemical indicators in selecting salt-resistant tomato cultivars. *Environ. Exp. Bot.* 54(3): 193-201.
- Kehl M (2006). Saline soils of Iran with examples from the alluvial plain in of Korbal, Zagros Mountains, Proceedings of the International Conference Soil and Desertification – Integrated Research for the Sustainable Management of Soils in Dry lands. 5-6 May, Hamburg, Germany.
- Krishnamurthy LR, Serraj CT, Hash AJ, Dakheel Reddy BVS (2007). Screening sorghum genotypes for salinity tolerant biomass production. *Euphytica* 156(1-2): 15-24.
- Lacerda CF, Cambraia J, Oliva MA, Ruiz HA (2005). Changes in growth and in solute concentrations in sorghum leaves and roots during salt stress recovery. *Environ. Exp. Bot.* 54: 69-76.
- Lane -Eynon (1970). *Encyclopedia of Industrial Chemical Analysis.* 10: 562-563 (New York).
- Lingle SE (1987). Sucrose metabolism in the primary culm of sweet sorghum during development. *Crop Sci.* 27: 1214-1219.
- Munns R, Husain S, Rivelli AR, James RA, Condon AG, Lindsay MP, Lagudah ES, Schachtman DP, Hare RA (2002). Avenues for increasing salt tolerance of crops, and the role of physiologically based selection traits. *Plant Soil* 247: 93-105.
- Netondo GW, Onyango JC, Beck E (2004). Sorghum and salinity. I. Response of growth, water relations, and ion accumulation to NaCl salinity. *Crop Sci.* 44: 707-710.
- Pearson D (1970). *The chemical analysis of food* (6th edition). 29: 132 (Now York).
- Serraj R, Sinclair TR (2002). Osmolyte accumulation: can it really help increase crop under drought conditions? *Plant Cell Environ.* 25: 333-341.

- Silva JV, Lacerda CF, de Azevedo Neto AD, Costa PHA, Prisco JT, Enéas Filho J, Gomes Filho E (2003). Growth and osmoregulation in two sorghum genotypes under salt stress. *Revista Ciência Agron.* 34(2): 125-131.
- Sunseri F, Palazzo D, Montemurro N, Montemurro F (1998). Salinity Tolerance in Sweet Sorghum (*Sorghum bicolor* L. Moench): Field Performance under Salt Stress *Ital. J. Agron.* 2(2): 111-116.
- Szabolcs I (1994). Soils and salinization. In: Pessarakali M (ed) *Handbook of plant and crop stress*. Marcel Dekker, New York, pp. 3-11.
- Tarpley L, Vietor DM (2007). Compartmentation of sucrose during radial transfer in mature sorghum culm. *BioMed. Central Plant Biol.* 7(33): 1-10.
- Tew TL, Cobill RM (2006). Evaluation of sweet sorghum as a complementary bioenergy crop to sugarcane in Louisiana. *J. Am. Soc. Sugar Cane Tech.* 26(1): 57-58.
- Varma NC (1988). *System of technical control for cane sugar factories in India*. The sugar Technologist's Associations. India.