

## Short Communication

# DNA cleavage agents from *Schisandra propinqua* var. *sinensis*

Ye-gao Chen<sup>1,2\*</sup>, Li-na Hai<sup>3</sup>, Ying Liu<sup>1</sup>, and Xin-rong Liao<sup>3</sup>

<sup>1</sup>Department of Chemistry, Yunnan Normal University, Kunming 650092, China.

<sup>2</sup>Engineering Research Center of Sustainable Development and Utilization of Biomass Energy, Ministry of Education, Yunnan Normal University, Kunming 650092, China.

<sup>3</sup>Department of Traditional Chinese Herbs, Yunnan Institute of Traditional Chinese Medicine, Kunming 650011, China.

Accepted 10 July, 2009

**Constituents of *Schisandra propinqua* (Wall.) Bail var. *sinensis* Oliv were investigated and 8 compounds were isolated from the stems and their structures were elucidated as (+)-catechin (1), (-)-gallicocatechin (2), p-(2'-hydroxyethyl)-phenol- $\beta$ -D-glucopyranoside (3), daucosterol (4), manwuweizic acid (5), cerotic acid 1-monoglyceride (6), octocosoic acid (7) and  $\beta$ -sitosterol (8) respectively. Based on MS and NMR techniques, (+)-Catechin, (-)-gallicocatechin, p-(2'-hydroxyethyl)-phenol- $\beta$ -D-glucopyranoside, manwuweizic acid, cerotic acid 1-monoglyceride and octocosoic acid were isolated from the plant for the first time. (+)-Catechin and (-)-gallicocatechin mediated DNA cleavage with good efficiency at 1  $\mu$ g/ml. (-)-Gallicocatechin show DNA strand-scission activity for the first time.**

**Key words:** Chemical composition, *Schisandra propinqua* var. *sinensis*, DNA cleaving activity, (-)-gallicocatechin, (+)-catechin.

## INTRODUCTION

DNA strand breakage process is involved in various biological stages such as inflammation, mutagenesis, carcinogenesis, or aging (Mibu et al., 2003; Chen et al., 2006). As a consequence of the clinical utility of DNA cleavage agents such as bleomycin, considerable effort has been made to identify and characterize naturally occurring molecules capable of mediating DNA strand scission, as such species may serve as lead structures for the development of novel anti-tumor drugs (Ma et al., 2004; Seo et al., 2003; Fukuhara et al., 1998). *Schisandra propinqua* (Wall.) Bail var. *sinensis* Oliv (Schisandraceae) is a plant used in folk medicine to promote blood circulation and to treat fracture, chronic gastritis, rheumatoid arthritis and irregular menstruation (Yunnan provincial crude drugs company, 1993). Previously, 3 lignans, isoschizandrolic acid,  $\beta$ -sitosterol and stearic acid were isolated from the plant roots and stems (Liu et al., 1998). In the course of our search for plant derived DNA cleavage agents, we isolated 2 DNA strand-nicking principles

from the plant stems of *S. propinqua* var. *sinensis* and the results are reported herein.

## MATERIALS AND METHODS

### Materials

The stems of *S. propinqua* var. *sinensis* was collected from E-Shan county of Yunnan province, China in November, 2001 and identified by Dr. Yong-Fa Wang, a botanist of Yunnan Institute of Traditional Chinese Medicine, where a voucher specimen (No.0102035) is deposited. Silica gel (200-300 mesh) was used for column chromatography and silica gel GF<sub>254</sub> for TLC (Qingdao Marine Chemical Co., China). Solvents were of the Indus-trial purity and distilled prior to use.


### Methods

MS were determined on an API Qstar Pulsa LC/TOF mass spectrometer and NMR spectra were measured on a Bruker DRX-500 spectrometer.

### Extraction and isolation

The dried powered stems of *S. propinqua* var. *sinensis* (9.0 kg)

\*Corresponding author. E-mail: ygchen48@gmail.com. Tel: +86-871-5516063. Fax: +86-871-5516061.



**Figure 1.** Compounds from *Schisandra propinqua* var. *sinesis*.

were extracted with 95% EtOH exhaustively at room temperature. The EtOH extract was evaporated *in vacuo* to yield a dark brown residue (840 g). H<sub>2</sub>O (2.5 l) was added to the residue and the resulting solution was extracted with chloroform, EtOAc and n-BuOH, successively. The n-BuOH extract (85 g) was applied to a silica gel column, eluting with chloroform containing increasing amounts of methanol to offer 8 fractions (A-I). Fraction B, F and G were subjected to repeated column chromatography to yield 1 g of (+)-catechin, 500 mg of (-)-gallocatechin, 9 mg of p-(2'-hydroxyethyl)-phenol-β-D-glucopyranoside and 100 mg of daucosterol (Figure 1). The EtOAc fraction (60 g) was applied to a silica gel column, eluting with petroleum ether containing increasing amounts of EtOAc. The fractions obtained from petroleum ether- EtOAc (4:1) elution were combined and subjected to repeated column chromatography to yield 12 mg of manwuweizic acid and 200 mg of cerotic acid 1-monoglyceride. The chloroform fraction (64 g) was applied to a

silica gel column, eluting with petroleum ether containing increasing amounts of EtOAc to obtained 500 mg of octocosoic acid and 700 mg of β-sitosterol.

### Identification of compounds

(+)-Catechin, white amorphous powder; <sup>1</sup>H NMR (CD<sub>3</sub>COCD<sub>3</sub>) δ 6.90 (1H, d, J= 1.8), 6.79 (1H, d, J = 8.1), 6.71 (1H, dd, J = 1.8, 8.1), 6.02 (1H, d, J = 2.3), 5.87 (1H, d, J = 2.3), 4.54 (1H, d, J = 7.9), 3.99 (1H, m), 2.92 (1H, dd, J = 5.5, 16.2), 2.50 (1H, dd, J = 8.2, 16.2); <sup>13</sup>C NMR (CD<sub>3</sub>COCD<sub>3</sub>) δ 157.0, 156.4, 156.2, 144.9, 131.4, 119.3, 114.9, 114.5, 99.8, 95.4, 94.6, 82.0, 67.6, 28.1; EIMS m/z (rel. int.): 290 (M<sup>+</sup>, 90), 167 (15), 152 (77), 139 (100), 123 (78), 110 (23), 97 (1), 85 (2), 69 (14).

(-)-Gallocatechin, white amorphous powder; <sup>1</sup>H NMR (CD<sub>3</sub>COCD<sub>3</sub>) δ 8.15 (1H, s, OH), 7.98 (1H, s, OH), 7.82 (2H, s, OH), 7.25 (H, s, OH), 6.45 (2H, s), 6.01 (H, d, J = 2.3), 5.87 (1H, d, J = 2.3), 4.51 (1H, d, J = 7.5), 3.90 (1H, m), 2.86 (1H, dd, J = 5.4, 16.1), 2.50 (1H, dd, J = 8.2, 16.1); <sup>13</sup>C NMR (CD<sub>3</sub>COCD<sub>3</sub>) δ 156.8, 156.4, 156.0, 145.4, 132.4, 130.7, 106.4, 99.8, 95.3, 94.6, 81.9, 67.5, 27.6; EIMS m/z (rel. int.): 306 (M<sup>+</sup>, 10), 168 (26), 139 (100), 126 (24), 110 (13), 98 (16), 83 (15), 69 (23), 55 (29).

p-(2'-Hydroxyethyl)-phenol-β-D-glucopyranoside, white amorphous powder; <sup>1</sup>H NMR (CD<sub>3</sub>OD) δ 7.17 (2H, dd, J= 6.7, 2.1 Hz), 7.04 (2H, dd, J= 6.7, 2.1 Hz), 4.87 (1H, d, J= 7.4 Hz), 4.67 (2H, t, J= 7.0 Hz), 3.88 (1H, dd, J= 12.1, 2.1 Hz), 3.68 (1H, dd, J= 12.1, 5.5 Hz), 3.41 (4H, m), 3.21 (2H, t, J= 7.0 Hz); <sup>13</sup>C NMR (CD<sub>3</sub>OD) δ 77.5 (C-1), 33.8 (C-2), 131.7 (C-1'), 130.7 (C-2', 6'), 118.0 (C-3', 5'), 158.3 (C-4'), 102.4 (C-1''), 78.1 (C-2''), 78.0 (C-3''), 74.9 (C-4''), 71.4 (C-5''), 62.5 (C-6''); EIMS m/z 167 (70), 149 (17), 120 (100), 107 (21), 91 (29), 77 (16).

Daucosterol, white amorphous powder, mp 296-298°C; FABMS m/z 577 [M+H]<sup>+</sup>; Identified by mixed melting point, co-TLC and comparison of <sup>1</sup>H and <sup>13</sup>C NMR spectrum with that of authentic sample.

Manwuweizic acid (5), white amorphous powder; <sup>1</sup>H NMR (CD<sub>3</sub>OD) δ 5.96 (1H, t, J= 7.6 Hz, H-24), 4.92, 4.71 (each 1H, s, H-28), 1.87 (3H, d, J= 1.0 Hz, H-27), 1.78 (3H, s, H-29), 0.98 (3H, s, H-19), 0.97 (3H, s, H-30), 0.95 (3H, d, J= 6.4 Hz, H-21), 0.79 (3H, s, H-18); <sup>13</sup>C NMR (CD<sub>3</sub>OD) δ 25.3 (C-1), 22.8 (C-2), 178.1 (C-3), 148.7 (C-4), 48.3 (C-5), 32.1 (C-6), 30.5 (C-7), 130.8 (C-8), 140.5 (C-9), 41.5 (C-10), 37.0 (C-11), 33.9 (C-12), 45.6 (C-13), 52.0 (C-14), 32.4 (C-15), 29.0 (C-16), 51.6 (C-17), 16.5 (C-18), 21.0 (C-19), 37.6 (C-20), 19.1 (C-21), 27.5 (C-22), 27.0 (C-23), 144.1 (C-24), 128.5 (C-25), 171.7 (C-26), 21.0 (C-27), 114.4 (C-28), 23.4 (C-29), 25.6 (C-30); EIMS m/z 470 (M<sup>+</sup>, 46), 455 (14), 397 (100), 261 (7), 235 (9), 161 (15), 149 (16), 119 (20), 107 (23), 95 (37), 69 (21).

Cerotic acid 1-monoglyceride, white amorphous powder; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 4.16 (2H, m, H-1), 3.91 (1H, m, H-2), 3.68 (1H, dd, J= 11.1, 5.0 Hz, H-3), 3.58(1H, dd, J= 11.1, 4.5 Hz, H-3), 2.32 (2H, t, J= 7.5 Hz, H-2'), 1.61(2H, m, H-3'), 0.87 (3H, t, J= 6.7 Hz, H-26'); EIMS m/z 470 (M<sup>+</sup>, 2), 452 (M<sup>+</sup>-H<sub>2</sub>O, 1), 442 (2), 424 (5), 411 (15), 396 (16), 368 (25), 351 (33), 340 (16), 323 (21), 294 (6), 134 (56), 112 (45), 98 (71), 57 (100).

Octocosoic acid, white amorphous powder; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 2.39 (2H, t, J= 7.5 Hz, H-2), 1.67 (2H, m, H-3), 1.30 (48H, br s, H-4~ H-27), 0.93(3H, t, J= 7.0Hz, H-28); EIMS m/z 424 (28, M<sup>+</sup>), 396 (100), 368 (94), 354 (14), 297 (4), 241 (3), 185 (6), 129 (30), 73 (39), 57 (48).

β-Sitosterol (8), white needles, mp 136 - 137°C; EIMS m/z 414 (M<sup>+</sup>); Identified by mixed melting point, co-TLC and comparison of <sup>1</sup>H and <sup>13</sup>C NMR spectrum spectrum with that of authentic sample.

### DNA cleavage assay

The DNA strand-scission assay was performed based on a modified

Hecht procedure (Deng et al., 2000; Huang et al., 1996). Compounds were dissolved in DMSO-MeOH (1:1); 1  $\mu$ l of each sample was added to a 13  $\mu$ l reaction mixture (total volume) containing 600 ng of PBR 322 DNA and 20  $\mu$ M  $\text{CuCl}_2$  in 10 mM Tris-HCl, pH 8.0. The reactions were incubated at 37°C for 1 h, terminated by addition of 2  $\mu$ l of 0.125% bromophenol blue in 30% glycerol and applied to a 1% agarose gel containing 0.7  $\mu$ g/ml ethidium bromide. The gel was run in 89 mM Tris containing 8.9 mM boric acid and 2.0 mM  $\text{Na}_2\text{-EDTA}$  at 110 V for 1 h, then visualized by UV irradiation. Agarose gels were quantified for percent DNA cleavage utilizing 1D Image Analysis software, Windows Extra Package, Version 3.5.

## RESULTS AND DISCUSSION

The stems of *S. propinqua* var. *sinensis* was extracted with 95% EtOH and then fractionated into chloroform, ethyl acetate and n-butanol soluble fractions successively. The n-butanol fraction was subjected to repeated column chromatography on silica gel to yield (+)-catechin, (-)-gallocatechin, p-(2'-hydroxyethyl)-phenol- $\beta$ -D-glucopyranoside and daucosterol identified by spectroscopic analysis (NMR and MS), literature comparison and comparison with authentic samples (Nonaka et al., 1977, 1981; Zhang et al., 1995; Tanahashi et al., 1997). The ethyl acetate fraction was subjected to repeated column chromatography on silica gel to yield manwuweizic acid and cerotic acid 1-monoglyceride and the chloroform fraction obtained octocosoic acid and  $\beta$ -sitosterol by similar chromatography (Liu et al., 1988; Chen et al., 2001).

(+)-Catechin, (-)-gallocatechin and p-(2'-hydroxyethyl)-phenol- $\beta$ -D-glucopyranoside and manwuweizic acid, cerotic acid 1-monoglyceride and octocosoic acid were isolated from the plant for the first time.

In view of the complete DNA cleaving activity of n-butanol fraction at 100  $\mu$ g/ml in the presence of 20  $\mu$ M  $\text{Cu}^{2+}$  in initial bioassay test, (+)-catechin, (-)-gallocatechin and p-(2'-hydroxyethyl)-phenol- $\beta$ -D-glucopyranoside were tested for their ability to relax PBR 322 plasmid DNA, a supercoiled, covalently closed, circular DNA in the presence of  $\text{Cu}^{2+}$ , using a cell free DNA cleavage assay (Deng et al., 2000). (+)-Catechin, showed 52% relaxation of supercoiled DNA (Form I) to nicked DNA (Form II) at 1  $\mu$ g/ml in the presence of 20  $\mu$ M  $\text{Cu}^{2+}$ , while (-)-gallocatechin was found to relax 39% of form I DNA to form II DNA at 1  $\mu$ g/ml in the presence of 20  $\mu$ M  $\text{Cu}^{2+}$ . p-(2'-Hydroxyethyl)-phenol- $\beta$ -D-glucopyranoside showed no activity. (-)-Gallocatechin was reported to show DNA strand-scission activity for the first time.

The results of this study provided some understanding on the chemical constituents of *S. propinqua* var. *sinensis* and its use in folk medicine to treat fracture, chronic gastritis and rheumatoid arthritis.

## ACKNOWLEDGEMENTS

This investigation was supported by a grant (No. 2005DFA30670) for international collaborative research by Ministry of Science and Technology, China, and a program for New Century Excellent Talents in University (NCET-05-0824).

## REFERENCES

- Chen YG, Qin GW, Cao L, Xie YY (2001). Triterpenoid acids with cytotoxic effect on rat luteal cells and human decidual cells in vitro from *Schisandra propinqua*. *Fitoterapia*, 72: 434-436.
- Chen YG, Song XP, Hai LN, Lv YP, A F, Halaweish F, Liao XR (2006). Compounds with DNA cleaving activity from *Kadsura ananosma*. *Pharmazie*. 61: 891-892.
- Deng JZ, Newman DJ, Hecht SM (2000). Use of Compare analysis to discover functional analogues of bleomycin. *J. Nat. Prod.* 63: 1269-1272.
- Fukuhara K, Miyata N (1998). Resveratrol as a new type of DNA-cleaving agent. *Bioorg. Med. Chem. Lett.* 8: 3187-3192.
- Huang L, Wall ME, Wani MC, Navarro H, Santisuk T, Reutrakul V, Seo EK, Farnsworth NR, Kinghorn AD (1996). New compounds with DNA strand-scission activity from the combined leaf and stems of *Uvaria hamiltoni*. *J. Nat. Prod.* 59: 290-292.
- Liu JS, Huang MF, Tao Y (1988). Anwuweizic acid and manwuweizic acid, the putative active principle of *Schisandra propinqua*. *Can. J. Chem.* 66: 414-415.
- Liu JS, Ma YT, Huang MF (1998). Chemical constituents from the stem and roots of *Schisandra propinqua* (Wall.) Bail var. *sinensis* Oliv. *Acta Chem. Sin.* 46: 345-348. (in Chinese)
- Ma J, Hecht SM (2004). Javaniside, a novel DNA cleavage agent from *Alangium javanicum* having an unusual oxindole skeleton. *Chem. Commun.* pp. 1190-1191.
- Mibu N, Yukawa M, Kashige N, Iwase Y, Goto Y, Miake F, Yamaguchi T, Ito S, Sumoto K (2003). Synthesis and DNA strand breakage activity of some 1, 4-diazepines. *Chem. Pharm. Bull.* 51: 27-31.
- Nonaka G, Minami M, Nishioka I (1977). Studies on rhubarb (rhei rhizoma). III. Stilbene glycosides. *Chem. Pharm. Bull.* 25: 2300-2305.
- Nonaka G, Nishioka I, Nagasawa T, Oura H (1981). Tannins and related compounds. I. Rhubarb. *Chem. Pharm. Bull.* 29: 2862-2870.
- Seo EK, Lee D, Shin YG, Chai HB, Navarro HA, Kardono LBS, Rahman I, Cordell GA, Farnsworth NR, Pezzuto JM, Kinghorn AD, Wani MC, Wall ME (2003). Bioactive prenylated flavonoids from the stem bark of *Artocarpus kemando*. *Arch. Pharm. Res.* 26: 124-127.
- Tanahashi T, Takenaka Y, Akimoto M, Okuda A, Kusunoki Y, Suekawa C, Nagakawa N (1997). Six secoiridoid glucosides from *Jasminum polyanthum*. *Chem. Pharm. Bull.* 45: 367-372.
- Yunnan Provincial Crude Drugs Company (1993). Name Lists of Chinese Herbal Medicine Resources in Yunnan. Science Press, Beijing, p. 151. (in Chinese)
- Zhang WJ, Liu YQ, Li XC, Yang CR (1995). Chemical constituents of ecological tea from Yunnan. *Acta Botanica Yunnanica* 17: 204-208. (in Chinese).