

Full Length Research Paper

***In vitro mecA* gene transfer among *Staphylococcus aureus* in Malaysian clinical isolates**

Negar Shafiei Sabet^{1*}, Geetha Subramaniam¹, Parasakthi Navaratnam² and Shamala Devi Sekaran¹

¹Department of Medical Microbiology, Faculty of Medicine, University of Malaya, 50603, Kuala Lumpur, Malaysia.

²School of Medicine and Health Sciences, Monash University Malaysia, No. 20 & 22, Jalan PJS 11/5, Bandar Sunway, 46150, Petaling Jaya, Malaysia.

Accepted 19 September, 2011

A simple experiment was performed to observe *in vitro* transfer of *mecA* gene from methicillin-resistant *Staphylococcus aureus*, Malaysian clinical isolates, to methicillin-sensitive one. PCR-based method was used in combination with the selective antibiotic screening method to study the transfer of resistance among clinical isolates carrying different genetic markers. After performing the mixed liquid culture experiments, two types of the transconjugants were obtained. Transconjugants were analyzed by PCR for the presence of *mecA* and *ccr* recombinase genes. The *mecA*, *ccrA3* and *ccrB2* genes were detected in donor strains as well as in both types of transconjugants. Transfer of *mecA*, *ccrA3* and *ccrB2* genes from donor to recipient strains that were present in close proximity during *in vitro* experiment may suggest possibility of horizontal gene transfer (HGT) among *S. aureus* clinical strains. Bacterial co-infections *in vivo* where the bacteria could be in close proximity are likely to be a possible point of horizontal gene transfer (HGT) with regard to certain antibiotic resistant genes.

Key words: *MecA*, horizontal gene transfer, *SCCmec*, *ccr* genes, methicillin-resistant *Staphylococcus aureus* (MRSA).

INTRODUCTION

Staphylococcus aureus has been recognized as a bacterium that can cause a broad spectrum of diseases and is a major cause of hospital-acquired infections worldwide. Methicillin-resistant *S. aureus* (MRSA) often referred to as the hospital 'superbug' hence clinical isolates, are often resistance to a wide variety of antibiotics. Horizontal gene transfer (HGT) among bacteria and their mobile genetic elements (MGEs) is a primitive mode of transfer and spread of antibiotic resistance in clinically important pathogens. The rapid acquisition of antibiotic resistance in *S. aureus* in the early 1960s was mediated by plasmid transfer. However, horizontal gene transfer (HGT) events have played a prominent role in the rapid acquisition of antibiotic resistance in *S. aureus* in recent years (Brody et al., 2008). The cellular mechanism by which *S. aureus* is capable of withstanding the

inhibitory effects of all β -lactam antibiotics, including the semi-synthetic ones such as methicillin, is based on acquisition of the *mecA* gene (Jansen et al., 2006).

Transfer of *mecA* is mediated by the staphylococcal cassette chromosome *mec* (*SCCmec*) (Wang and Archer, 2010). Although the origins of the *mecA* gene (the principle component of methicillin resistance in staphylococci) are unknown, a *mecA* homologue (88% similarity) is ubiquitous in the antibiotic-susceptible *Staphylococcus sciuri*, and may be a possible evolutionary precursor of the *mecA* gene of the MRSA strains. Furthermore, *S. aureus mecA* was identical to that identified in a *Staphylococcus epidermidis* isolate from the same individual, thus suggesting that MRSA strain has arisen *in vivo* by horizontal transfer of *mecA* between two staphylococcal species (Brody et al., 2008; Wielders et al., 2001; Deurenberg et al., 2007). It has been hypothesized that all MRSA clones have a common ancestor, while another theory suggests that *SCCmec* was introduced several times into different *S. aureus* lineages (Juhas et al., 2009).

*Corresponding author. E-mail: negar@um.edu.my. Tel: +603-79676672. Fax: +603-79676672.

Table 1. Antibiotic resistance and sensitivity profiles of the donors, recipients and transconjugants that were obtained by disc diffusion method.

| Strain profile | Antibiotic | | | | | |
|------------------|------------|-----|-----|-----|-----|------|
| | Ox | Amp | Gen | Ery | Tet | Amik |
| Recipients | S | R | R | R | S | S |
| Donors | R | R | S | R | R | R |
| Transconjugant A | R | R | R | R | R | R |
| Transconjugant B | R | R | R | R | S | S |

Ox, Oxacillin; Amp, ampicillin; Gen, gentamicin; Ery, erythromycin; Tet, tetracycline; Amik, amikacin; R, resistant; S, sensitive.

MecA is carried on a genetic element, *SCCmec*, which inserts precisely into the *S. aureus* chromosome at *orfX* (Hiramatsu et al., 2001). Genes encoding cassette chromosome recombinases (*ccr*) are located within the *SCCmec* elements. For movement, *SCCmec* carries specific genes (*ccr*), which encode recombinases of the invertase/resolvase family. Cassette chromosome recombinase (*ccr*) genes (*ccrC* or the pair of *ccrA* and *ccrB*) encode recombinases mediating integration and excision of *SCCmec* into and from the chromosome. Four different homologous pairs of *ccr AB* genes and one *ccr C* gene have been reported (Hanssen and Sollid, 2006). These genes are designated as, *ccrA1*, *ccrA2*, *ccrA3*, *ccrA4*, *ccrB1*, *ccrB2*, *ccrB3* and *ccrB4* (Ito et al., 2004). Integration of a *SCCmec* element into the chromosome converts drug-sensitive *S. aureus* into hospital pathogen methicillin-resistant *S. aureus* (MRSA), which is resistant to practically all β -lactam antibiotics (Hiramatsu et al., 2001)

The mechanism(s) responsible for *mecA* transfer is still obscure. The assumption is that the *ccr* and *mec* genes were brought together in coagulase-negative staphylococci (CoNS) from an unknown source where deletion in the *mec* regulatory genes occurred before the genes were transferred into *S. aureus* to generate MRSA (Hanssen et al., 2004). Studies with human staphylococcal strains indicate that *S. epidermidis* is a reservoir of antibiotic resistance genes that can be transferred to *S. aureus* under *in vitro* and *in vivo* conditions (Weilders et al., 2002). This study was therefore performed to observe *in vitro* resistance transfer of *mecA* gene from MRSA clinical isolate to methicillin-sensitive one that carries different phenotypic markers. Simple experiment was conducted using PCR-based method in combination with the selective antibiotic screening test for determining direction of resistance transfer.

MATERIALS AND METHODS

Bacterial strains

Clinical isolates were obtained from clinical samples processed at the diagnostic microbiology laboratory at University Malaya Medical Centre (UMMC, Kuala Lumpur, Malaysia). For strain identification,

S. aureus isolates were grown on Colombia Blood agar plates (Oxoid, Basingstoke, UK) at 37°C for 24 h. The *S. aureus* isolates used in this study were identified primarily as *S. aureus* by colony morphology, Gram stain, catalase reaction and coagulase production. Identification of each isolate was confirmed using API staph. System (bioMérieux, Marcy-l'Étoile, France).

Determining antibiotic resistance profile of bacterial strains

Resistance to various antibiotics was determined using the method recommended by the Clinical and Laboratory Standards Institute (CLSI, 2006). Disk diffusion was performed on Mueller–Hinton agar using Kirby–Bauer method (Oxoid, Basingstoke, UK). Antibiotics; oxacillin, ampicillin, gentamicin, erythromycin, tetracycline and amikacin, were used for determining the sensitivity of the bacterial strains.

Criteria for choosing donors and recipients and their antibiotic profiles

In order to differentiate the donor from recipient, the strain carrying different genetic marker was selected. Donor strains that are resistant to methicillin (oxacillin) were selected based on the presence of *mecA* gene (*mecA*⁺) and recipient strains that are sensitive to methicillin were selected based on absence of *mecA* gene (*mecA*⁻). After studying antibiotic resistant profile of the isolates, the donor strain with the following genetic marker was chosen depending on the availability of the clinical isolates in our laboratory (Table 1). The donor that was chosen was sensitive to gentamicin and resistant to tetracycline and oxacillin, while recipient that was chosen was resistant to gentamicin and sensitive to tetracycline and oxacillin.

Transfer of antibiotic resistance in mixed liquid cultures

Bacterial cultures were grown for 12 h in Luria-Bertani (LB) broth. Equal volumes (200 ml) of the donor and recipient strains were mixed in a flask. The cultures were supplemented with 200 ml of LB broth and incubated at 37°C without shaking. Aliquots of 50 ml each were withdrawn after 6 h and placed on two different antibiotics - tetracycline (30 μ g/ml) and gentamicin (10 μ g/ml) containing agar plates respectively after appropriate dilution. After 24 h, colonies resistant from each plate were picked up as probable transconjugants. The method used was a modification of the experiment performed by Khan et al. (2000). The choice of these antibiotics was purposefully made so that the resistant colonies might represent the transconjugants when the transfer of oxacillin resistance from resistant to sensitive strains or the transfer of gentamicin resistance from sensitive to resistant strains had taken




Figure 1. Representative pictures of antibiotic resistance and sensitivity profiles of the donors, A; recipients, B; transconjugant A, C; and transconjugant B, D; that was obtained by disc diffusion method. Ox: Oxacillin, Amp: Ampicillin, Gen: Gentamicin, Ery: Erythromycin, Tet: tetracycline, Amik: Amikacin.

place. In either case, the transconjugants would be resistant to oxacillin and gentamicin. Colonies from these plates were placed on oxacillin containing agar and blood agar plates.

Detection and amplification of *mecA* and *ccrAB* genes by PCR

DNA extraction was performed following the method described by Unal et al. (1992). The *mecA* primer that could produce 214 bp PCR product was used to screen donors, recipients and transconjugants for the presence of *mecA* gene (Sabet et al., 2006). The donors, recipients and transconjugants were subjected to the PCR for the presence of genes as well. Six primers pairs were used to screen the presence of *ccrAB* genes according to the work by Hanssen et al. (2004). The mastermix gave a final concentration of; 10 mM Tris-HCl (pH 8.3), 50 mM KCl, 1.4 mM MgCl₂, 0.09 mM of each dNTP and 0.4 μM of each of the primers. The 45 μl mastermix were mixed with 5 μl template. The following

PCR amplification conditions for *ccrA1*, *ccrB1*, *ccrA3* and *ccrB3* primer sets were used; 1 min at 95°C, 30 cycles with 30 s at 95°C, 10 s at 57°C, 60 s at 72°C and the final extension was 4 min at 72°C. For *ccrA2* and *ccrB2* sets of primers, the following conditions were used; 1 min at 94°C, 30 cycles with: 30 s at 94°C 30 s at 62°C, 60 s at 72°C and the final extension was 4 min at 72°C. 10 μl of PCR-amplified reaction mixture was resolved by electrophoresis on agarose gel. The sizes of amplification product were estimated by comparison with a 100-bp molecular size standard ladder (New England BioLabS, UK).

DNA sequencing of *mecA* and *ccrAB* genes

Polymerase chain reaction products for both *mecA* and *ccrAB* genes were purified using the PCR Purification Kit (Qiagen, Hilden, Germany). The PCR DNA sequencing was carried out using an automated DNA sequencer (an ABI Prism 377 DNA sequencer;




Figure 2. An ethidium bromide-stained gel demonstrating PCR amplification of *mecA* gene in donors, recipients and transconjugants. Lanes 1 and 2, donors; lanes 3 and 4, transconjugants A; lanes 5 and 6, transconjugants B; lanes 7 to 10, recipients; lane 11, negative control; lane 12, 100 bp DNA marker.

Perkin- Elmer ABI, Wellesley, MA). Nucleotide sequences were analyzed and compared by the use of the BLAST computer program (National Center for Biotechnology Information <http://www.ncbi.nlm.nih.gov>).

RESULTS

Strain Identification and antibiotic resistance profile of bacterial isolates

The Gram stain characteristics of the all bacterial strains were Gram positive cocci (GPC) in clusters. All the strains were catalase and tube coagulase positive. Using API Staph. system for the identification, all stains were identified as *S. aureus*. Antibiotic resistance profiles of the donor strains (MRSA) were resistant to oxacillin, ampicillin, erythromycin, tetracycline and amikacin, but sensitive to gentamicin (Table 1). The recipient strains (MSSA) were resistant to ampicillin, gentamicin and erythromycin, but sensitive to oxacillin, tetracycline and amikacin (Table 1).

Antibiotic resistance profile of transconjugants

After performing the mixed liquid cultures experiments and plating on tetracycline and gentamicin-containing agar plates, the growth observed indicated that these

colonies are transconjugants. Two types of the transconjugants were obtained (Table 1). The antibiotic resistance profile of the transconjugant A is shown in Figure 1C. The transconjugant A was resistant to oxacillin, ampicillin, gentamicin, erythromycin, tetracycline and amikacin. The transconjugant B was resistant to oxacillin, ampicillin, gentamicin and erythromycin, but sensitive to tetracycline and amikacin (Figure 1D).

Detection of *mecA* and *ccr AB* genes at gene level

Prior to mixing, the bacterial lysates from donor and recipient strains were subjected to PCR analysis. The *mecA* gene was detected in donor strain but not amplified in recipient strains (Figure 2). To determine whether the transfer of the *mecA* gene had taken place or not, the transconjugants were analyzed by PCR for the presence of *mecA* gene.

The PCR amplified products from donors, recipients and transconjugant A and transconjugant B are shown in Figure 2. As shown in the Figure, the *mecA* gene was present in both transconjugant A and B and donor strains. The donors, recipients and transconjugants were subjected to the PCR assay for the presence of *ccrAB* genes. The PCR results show the amplification of *ccrA3* and *ccrB2* genes from donors, transconjugant A and transconjugant B. Amplification of *ccrA3* gene and *ccrB2*




Figure 3. An ethidium bromide-stained gel demonstrating PCR amplification of *ccrA3* gene in donors, recipients and transconjugants. Lane 1, donor; lane 2, transconjugant A; lane 3, transconjugant B; lanes 4,5, and 6, recipients; lane 7, negative control; lane 8, positive control (MRSA ATCC 43300); lane 9, 100bp DNA Marker.




Figure 4. An ethidium bromide-stained gel demonstrating PCR amplification of *ccrB2* gene in donors, recipients and transconjugants. Lane 1, 100 bp DNA Marker; lane 2, donor; lane 3, transconjugant A; lane 4, transconjugant B; lanes 5 to 7, recipients; lane 8, negative control.

genes produced PCR product of 873 and 1128 bp, respectively (Figures 3 and 4). Among the six primers used for screening *ccrAB* genes in the recipient strains, no amplification were observed among recipient strains.

DNA sequencing of *mecA* and *ccrAB* genes

The *mecA* gene showed 98% homology with *S. aureus*, *mecA* gene for PBP2' (penicillin binding protein 2') (Gene Bank accession number; X52593). The *ccrA3* gene showed 100% homology with *S. aureus* subsp. *aureus* strain ATCC 33592 CcrA3 (Gene Bank accession number; DQ460218). The *ccrB2* gene showed 100% homology with *S. epidermidis* strain ATCC 12228 CcrB (Gene Bank accession number; AE015929).

DISCUSSION

The resistance mechanism by which the *mecA* can be transferred has been studied extensively because of the importance of resistant strains in the clinical setting. This study was performed to observe possibility of *in vitro* transfer of *mecA* gene from local MRSA to MSSA clinical isolate that do not carry *mecA* gene, using cost effective and simple approach. Many studies support that horizontal transfer play an important role in the dissemination of the *mecA* gene in the *S. aureus* population and specially transfer *mecA* gene more frequently to MSSA (Weilders et al., 2002). Also, it was suggested that different staphylococcal species acquire SCC*mec* locally by horizontal gene transfer (Hanssen et al., 2004). The study by Khan et al. (2000) showed the usefulness of PCR and gene-specific probing along with the conven-

tional selective antibiotic screening methods to study the transfer of drug resistance. This resistance transfer was observed between selective isolates that were resistant to the same antibiotic but carry different genetic markers (Khan et al., 2000). The method presented in their study uses a simple preliminary unique approach to analyze the transconjugants.

We have used the same approach in this study to observe and study the transfer of resistant genes with unique focus to study transfer of *mecA* gene *in vitro*. The mixing of donor and recipient strains in liquid culture was chosen as it allows transfer of antibiotic resistance between the strains that are in close proximity *in vitro*. The results obtained showed the use of gentamicin and tetracycline as selective antibiotics was helpful in observing transfer of oxacillin resistance to transconjugants. The transconjugant A is resistant to all the antibiotics, therefore transconjugant A with the phenotype resistant to oxacillin, tetracycline and gentamicin having a new phenotype, resembles neither the recipient nor the donor. The direction of the transfer of oxacillin also could not be determined in the transconjugant A. This could be the donor strain that has received gentamicin resistance from recipient strain or this could be recipient strain that has received the oxacillin and tetracycline resistant from donor strain. However, the transconjugant B with the phenotype of resistance to oxacillin, gentamicin and sensitive to tetracycline has the phenotype resembling the recipient, while it is resistant to oxacillin, as well as a new phenotype which is different from the donor but resistant to oxacillin.

Furthermore, studies have found novel SCC*mec* types or SCC elements without *mecA*, which could be a reservoir for antibiotic resistance islands in *S. aureus*. In this study, *ccr* recombinases genes was not detected in methicillin-sensitive strains (recipient strains) therefore SCC*mec* were absent in these recipient strains. For integration into and excision of SCC element from the chromosome, genes encoding cassette chromosome recombinases (*ccr*) are involved. Detection of *ccrA3* and *ccB2* genes in donor, transconjugant A and B and their absence originally from the recipient strains showed that the *mecA* gene transfer have taken place along with *ccr* recombinases genes. In this study, we focused on using very simple and cost effective experiment to observe *in vitro* resistance transfer in particular *mecA* gene between clinical *S. aureus* isolates.

In conclusion, *ccr* recombinases genes detection and simple conventional selective antibiotic screening methods was useful in observing *mecA* gene *in vitro* transfer among clinical isolates. This observation can further support theory of horizontal gene transfer among same species of clinically important strains of *S. aureus*.

The observation in this study further highlights the crucial role of horizontal gene transfer in the evolution of bacteria and dissemination of variable genes, including antibiotic resistance and virulence genes.

REFERENCES

- Brody T, Yavatkar AS, Lin Y, Ross J, Kuzin A, Kundu M, Fann Y, Odenwald WF, (2008). Horizontal gene transfers link a human MRSA pathogen to contagious bovine mastitis bacteria. PLoS ONE 3(8): e3074.
- Clinical and Laboratory Standards Institute (2006). Performance standards for antimicrobial susceptibility testing; sixteenth informational supplement, 16th ed. M100-S16. Wayne, Pa.
- Deurenberg RH, Vink C, Kalenic S, Friedrich AW, Bruggeman CA, Stobberingh EE (2007). The molecular evolution of methicillin-resistant *Staphylococcus aureus*. Clin. Microbiol. Infect. 13: 222-235
- Hanssen AM, Kjeldsen G, Sollid JUE (2004). Local variants of staphylococcal cassette chromosome *mec* in sporadic methicillin-resistant *Staphylococcus aureus* and methicillin-resistant coagulase-negative staphylococci: Evidence of horizontal gene transfer? Antimicrob. Agents Chemother. 48: 285-296.
- Hanssen AM, Sollid JUE (2006). SCC*mec* in staphylococci: genes on the move. FEMS Immunol. Med. Microbiol. 46: 8-20.
- Hiramatsu K, Cui L, Kuroda M, Ito T (2001). The emergence and evolution of methicillin-resistant *Staphylococcus aureus*. Trends Microbiol. 9: 486-493.
- Ito T, Ma XX, Takeuchi F, Okuma k, Yuzawa H, Hiramatsu K (2004). Novel type V staphylococcal cassette chromosome *mec* driven by a novel cassette chromosome recombinase, *ccrC*. Antimicrob. Agents. Chemother. 48: 2637-2651.
- Jansen WTM, Beitsma MM, Koeman CJ, Wamel WJB, Verhoef J, Fluit AC, (2006). Novel mobile variants of staphylococcal cassette chromosome *mec* in *Staphylococcus aureus*. Antimicrob. Agents Chemother. 50: 2072-2078.
- Juhas M, Roelof van der Meer J, Gaillard M, Harding RM, Hood DW, Crook DW (2009). Genomic islands: tools of bacterial horizontal gene transfer and evolution. FEMS. Microbiol. Rev. 33: 376-393.
- Khan SA, Nawas MS, Khan AA, Cerniglia CE (2000). Transfer of erythromycin resistance from poultry to human clinical strains of *Staphylococcus aureus*. J. Clin. Microbiol. 38:1832-1838.
- Sabet NS, Subramaniam G, Navartnam P, Sekaran SD (2006). Simultaneous species identification and detection of methicillin resistance in staphylococci using triplex real-time PCR assay. Diag. Microbiol. Infect. Dis. 56: 13-18.
- Unal S, Hoskins J, Flokowsch JE, Wu CYE, Preston DA, Skatrud PL (1992). Detection of methicillin-resistant staphylococci by using the polymerase chain reaction. J. Clin. Microbiol. 30: 1685-1691.
- Wang L, Archer GL (2010). Roles of *CcrA* and *CcrB* in excision and integration of Staphylococcal Cassette Chromosome *mec*, a *Staphylococcus aureus* genomic island. J. Bacteriol. 192(12): 3204-3212.
- Wielders CLC, Vriens MR, Brisse S, Graaf-Miltenburg LAD, Troelstra A, Fleer A, Schmitz FJ, Verhoef J, Fluit AC (2001). Evidence for *in-vivo* transfer of *mecA* DNA between strains of *Staphylococcus aureus*. Lancet, 357: 1674-1675.
- Wielders CLC, Fluit AC, Brisse S, Verhoef J, Schmitz FJ (2002). *mecA* gene is widely disseminated in *Staphylococcus aureus* population. J. Clin. Microbiol. 40: 3970-3975.