

Full Length Research Paper

Bulked segregant analysis to detect quantitative trait loci (QTL) related to heat tolerance at grain filling rate in wheat using simple sequence repeat (SSR) markers

Mohamed Najeb Barakat^{1,2*}, Abdullah Abdulaziz Al-Doss¹, Adel Ahmed Elshafei¹ and Khaled Ahmed Moustafa¹

¹Plant Genetic Manipulation and Genomic Breeding Group, Plant Production Department, College of Food and Agricultural Sciences, King Saud University, Riyadh, Saudi Arabia.

²Biotechnology Lab., Crop Science Department, Faculty of Agriculture, University of Alexandria, Egypt.

Accepted 25 May, 2012

The grain-filling rate (GFR) plays an important role in determining grain yield. An F₂ population of wheat was developed from a cross between the 2 wheat cultivars, Ksu106 (heat-tolerant) and Yecora Rojo (heat-sensitive). The parents and 205 F₂ plants were planted on the 20th of January during the winter season of 2009 to evaluate heat tolerance during the grain-filling period. The sowing date in the present investigation represents the heat stress conditions in Saudi Arabia. Bulked – segregant analyses (BSA) was used in conjunction with simple sequence repeats (SSR) analysis to find markers linked to genes of heat tolerance. Composite interval mapping was used for mapping quantitative trait loci (QTL). The results reveal that 12 SSR markers: *Wmc24*, *Wmc168*, *Wmc326*, *Xgwm30*, *Xgwm456*, *Wmc25*, *Wmc44*, *Wmc94*, *Wmc161*, *Wmc273*, *Wmc327* and *Xgwm566* were linked to GFR by QTLs analysis of the F₂ population. The results show that regression analysis for the relationship between the 12 markers and the phenotypes of F₂ individuals were highly significant. The results demonstrate that SSR markers combined with bulked segregant analysis could be used to identify molecular markers linked to the grain filling rate as an indicator for heat tolerance in wheat.

Key words: Grain filling rate, QTL analysis, SSR marker, wheat.

INTRODUCTION

Due to the global change in climatic conditions that results from increasing atmospheric concentrations of carbon dioxide and other greenhouse gases, high temperature stress have become one of the major factors exerting serious influence on wheat production (Maestri et al., 2002). The heat stress especially occurring during flowering stage in wheat growth area results in a

reduction in both individual kernel weight and kernel number (Plaut et al., 2004; Hays et al., 2007). So the research on the genetic mechanism of heat tolerance is getting more and more important for the utilization of heat tolerant genes and the development of new wheat varieties with heat tolerance. Grain filling is a crucial and dynamic process of wheat growth. Its duration and rate determine the individual grain size, grain weight and as a result, the economic yield of the crop (Li and Pan, 2005). Grain filling duration is much influenced by temperature, particularly in the presence of stress (Wiegand and Cuellar, 1981; Knott and Gebeyehou, 1987), while grain-filling rate (GFR) appears to be largely under genetic control (Van Sanford, 1985).

Moreover, since wheat is harvested at the beginning of

*Corresponding author. E-mail: mnrbarakat@yahoo.com.

Abbreviations: BSA, Bulked segregant analysis; CIM, composite interval mapping; GFR, grain filling rate; QTL, quantitative trait loci; SSR, simple sequence repeats.

summer and some other crops are planted in succession in wheat growth area, the selection of genotypes with high GFR appears to be a successful strategy for increasing overall grain yield. The physiological mechanisms by which crops regulate GFR have been widely explored (Egli et al., 1989), but little is known until now about the genetics and loci affecting this complex trait. Availability of molecular maps helps to dissect complex quantitative traits into component loci and study their relative effects on a specific trait in a segregating population by quantitative trait loci (QTL) analysis (Roder et al., 1998). In recent years, with the development of molecular marker technique and widespread application, there were reports on genetic research of heat tolerance in wheat by utilizing molecular marker (Yang et al., 2002; Mason et al., 2010). Previous research on the genetic mechanism of heat tolerance in wheat was focused on the grain filling duration (Yang et al., 2002) and the heat susceptibility index (Mason et al., 2010) under high temperature as a measure of heat tolerance.

Also, QTLs for GFR under drought stress have been reported (Kirigwi et al., 2007; Wang et al., 2009). Recently, identification of new microsatellite markers linked to the grain filling rate as an indicator for heat tolerance genes in an F_2 wheat population have been reported (Barakat et al., 2011). However, in the present investigation, composite interval mapping (CIM) has been used for mapping QTLs which is considered statistically more powerful compared to single-point analysis (Lander and Botstein, 1989; Liu, 1998). In the present study, we present the mapping of 12 QTLs associated with heat tolerance using pooled bulks from the F_2 population of Ksu106 (heat tolerant) / Yecora Rojo (heat-sensitive) cross. This strategy is expected to contribute towards a deeper understanding of the genetic mechanism of wheat heat tolerance as well as the cloning of heat tolerance QTLs and marker assisted selection breeding.

MATERIALS AND METHODS

Population development

The wheat genotypes used in this study were Ksu106; advanced lines (F_8) selected from the wheat breeding program at the Plant Production Department, College of Food and Agricultural Sciences, King Saud University, Saudi Arabia, and Yecora Rojo; the adapted cultivar in Saudi Arabia. The pedigree of Ksu106 is Barouk /R1474-75-3-53-3-3 and it is heat tolerant (Al-Doss et al., 2009). The Yecora Rojo is a USA cultivar and the elite recommended cultivar adapted to Saudi Arabia environment since 1981. Yecora Rojo is a 2-gene dwarf high yielding cultivar but is very sensitive to any change in the environmental factors such as temperature, especially during the critical period of grain filling towards the end of the growing season. The increase in temperature during this period causes the plant to speed up its life cycle and shorten the time for grain filling and, thus, the kernel weight is reduced and consequently so is grain yield (Gandourah, 1989). Two wheat genotypes that had contrasting response to heat stress were crossed to generate F_1 seeds during the winter season 2008 at the

college of food and agricultural sciences, experimental research station, at Dierab near Riyadh, King Saud University. F_1 seeds population derived from the cross (Ksu106 × Yecora Rojo) were obtained. The F_2 seeds were obtained by selfing in the summer season in 2008 under green house conditions.

Evaluation of heat tolerance

205 F_2 plants and the parents were planted on 20th January in the winter season 2009 at the experimental research station, King Saud University, to evaluate the heat tolerance during the grain-filling period. The sowing date in the present investigation represents the heat stress conditions in Saudi Arabia. The cultural practices were carried out according to the recommended practices followed in Riyadh region. The agronomic traits such as grain filling duration, grain yield per plant and GFR were determined. GFR is the rate at which assimilates are transported from the source to the sink. It was estimated as the ratio between grain yield per plant and grain filling duration.

DNA extraction

Frozen young leaves (500 mg) of 205 F_2 plants and their parents were individually ground to a powder in a mortar with liquid nitrogen. The DNA extraction was done using the CTAB method (Sagahi-Marouf et al., 1984).

PCR amplification

A set of 100 microsatellite primers developed by several investigators (Roder et al., 1998; Gupta et al., 2002) were used in this study. The PCR reaction mixture consisted of 20 to 50 ng genomic DNA, 1 × PCR buffer, 2.0 mM $MgCl_2$, 100 μ M of each dNTP, 0.1 μ M primer and 1U *Taq* polymerase in a 25 μ l volume. The PCR cycle included an initial denaturation at 94°C for 3 min followed by 35 cycles of denaturation at 94°C for 1 min; annealing at 50, 55 or 60°C (depending on the individual microsatellite primer) for 1 min; and extension at 72°C for 2 min followed by a 17 min final extension at 72°C. The amplification products have been electrophoresed in 2 to 3% agarose gels.

Bulked segregant analysis

Bulked – segregant analyses (BSA) was used in conjunction with simple sequence repeats (SSR) analysis (Michelmore et al., 1991) to find markers linked to genes of heat tolerance. Tolerant and sensitive bulks were prepared from F_2 individuals by pooling aliquots, containing equivalent amounts of total DNA, approximately, 50 ng/ μ l from each of ten sensitive and ten tolerant F_2 plants selected, based on phenotypic assessments. SSR primers were, then, screened on the parents and the two bulk DNA samples, from which some primer combinations revealed bands that were polymorphic, not only among parental genotypes, but also between the pair of the bulk DNA. Based on the evaluations of DNA bulks, individual F_2 plants were analyzed with co-segregating primers to confirm SSR markers linkage to the grain filling rate as an indicator for heat tolerance genes.

Data and linkage analysis

The computer program Map Manager QTX Version 0.22 (Meer et al., 2002) was used to perform composite interval mapping (CIM)

(Zeng, 1994) to evaluate marker intervals putatively associated with trait phenotypes. Linkage was detected when a log of the likelihood ratio (LOD) threshold of 3.0 and maximum distance was 50 cm. The Kosambi's mapping function was used. Genetic loci with the most significant effect for each QTL were assembled into multiple regression models, using PROC REG of SAS version 9.1 software packages (SAS Institute, Cary, NC 2007), to determine the total amount of the phenotypic variation explained (Nelson, 1997).

RESULTS

SSR markers analysis

Out of the 100 SSR markers used in this study, only 25 primer pairs generated polymorphisms between the parents. Each of these markers was used to screen DNA bulks of the 10 tolerant and the 10 sensitive F_2 plants. Twenty SSR markers (*Wmc24*, *Wmc25*, *Wmc44*, *Wmc47*, *Wmc48*, *Wmc52*, *Wmc94*, *Wmc149*, *Wmc154*, *Wmc161*, *Wmc168*, *Wmc215*, *Wmc219*, *Wmc261*, *Wmc273*, *Wmc326*, *Wmc327*, *Xgwm30*, *Xgwm456*, and *Xgwm56*) only amplified polymorphic bands (Table 1). The amplification profiles of the twenty primer pairs were characterized by the F_2 progeny and their parents. The SSR primers *Wmc24*, *Wmc47*, *Wmc48*, *Wmc52*, *Wmc149*, *Wmc154*, *Wmc168*, *Wmc219*, *Wmc261*, *Wmc326*, *Xgwm30* and *Xgwm456* generated one polymorphic fragment at 152, 160, 130, 170, 230, 180, 310, 180, 120, 225, 160 and 140 bp, respectively which was present only in the tolerant bulk and Ksu106 (tolerant parent) and were missing in the sensitive bulk and Yecora Rojo (sensitive parent) (Figure 1).

The SSR primers *Wmc25*, *Wmc44*, *Wmc94*, *Wmc161*, *Wmc215*, *Wmc273*, *Wmc327*, and *Xgwm566* generated one polymorphic fragment at 200, 260, 120, 200, 207, 180, 200 and 140 bp, respectively, which was present only in the tolerant bulk and Ksu106 (tolerant parent) and another polymorphic fragments at 220, 290, 105, 190, 195, 170, 180, and 150 bp, respectively which were present only in the sensitive bulk and Yecora Rojo (sensitive parent) (Figure 1). These co-dominant microsatellite markers were able to identify the heterozygotes plants. A typical amplification pattern generated by *Wmc25* and *Wmc327* is shown in Figure 1. The *Wmc25* of the tolerant parent was smaller than the sensitive parent. This locus was inherited in a Mendelian co-dominant manner. There was clear co-segregation between the amplification of the smaller *Wmc25* and the F_2 plants showing tolerant phenotypes. In the homozygous sensitive F_2 plants, only the large *Wmc25* was amplified.

In a proportion of tolerant F_2 plants, both the larger and the smaller were amplified; these plants were presumably heterozygous. In contrast, the *Wmc327* of the tolerant parent was larger than the sensitive parent. This locus was also inherited in a Mendelian co-dominant manner. Again there was clear co-segregation between the

amplification of the larger *Wmc327* and the F_2 plants showing tolerant phenotypes. In the homozygous tolerant F_2 plants, only the large *Wmc327* was amplified. In a proportion of tolerant F_2 plants, both the larger and the smaller alleles were amplified, so these plants were presumably heterozygous. The co-dominant microsatellite markers *Wmc25*, *Wmc44*, *Wmc94*, *Wmc161*, *Wmc215*, *Wmc273*, *Wmc327*, and *Xgwm566* were able to identify the heterozygotes, and would serve as an important tool to rapidly transfer the heat tolerance genes into other wheat cultivars.

To check for potential co-segregation of DNA fragments and heat tolerant phenotypes, multiple regression analysis was carried out in order to confirm an association between the twenty SSR markers and the grain filling rate as an indicator for heat tolerance genes in all 205 F_2 progenies. The results show that the regression analysis for the relationship between the dominant markers (*Wmc47*, *Wmc48*, *Wmc52*, *Wmc149*, *Wmc154*, *Wmc219* and *Wmc261*) and the phenotypes of F_2 individuals were non-significant. However, the dominant markers *Wmc24*, *Wmc168*, *Wmc326*, *Xgwm30* and *Xgwm456* were significantly ($P < 0.05$) associated with the grain filling rate and explained 31, 27, 22, 29, and 33% of the variation. Also, the co-dominant microsatellite markers *Wmc25*, *Wmc44*, *Wmc94*, *Wmc161*, *Wmc273*, *Wmc327*, and *Xgwm566* were significantly ($P < 0.05$) associated with the grain filling rate and explained 45, 30, 39, 43, 48, 32 and 64% of the variation, respectively (Table 2). This indicates that the 12 markers were associated with the grain filling rate as an indicator for heat tolerance genes. The R^2 values suggested that *Wmc24*-linked QTL, *Wmc168*-linked QTL, *Wmc326*-linked QTL, *Xgwm30*-linked QTL, *Xgwm456*-linked QTL, *Wmc25*-linked QTL, *Wmc44*-linked QTL, *Wmc94*-linked QTL, *Wmc161*-linked QTL, *Wmc273*-linked QTL, *Wmc327*-linked QTL, and *Xgwm566*-linked QTL account 0.31, 0.27, 0.22, 0.29, 0.33, 0.45, 0.30, 0.39, 0.43, 0.48, 0.32 and 0.64 of the total phenotypic variation, respectively, in heat tolerance in the F_2 population.

QTL analysis

The linkage relationship between the 12 SSR markers (*Wmc24*, *Wmc168*, *Wmc326*, *Xgwm30*, *Xgwm456*, *Wmc25*, *Wmc44*, *Wmc94*, *Wmc161*, *Wmc273*, *Wmc327*, and *Xgwm566*) and the grain filling rate as an indicator for heat tolerance genes were estimated using the F_2 population, deriving from the cross, Ksu106 × Yecora Rojo. The genetic distance between the 12 SSR markers and heat tolerance genes were determined as 9.9, 9.0, 9.0, 7.6, 8.9, 7.8, 9.7, 8.4, 7.3, 7.0, 8.6, and 5.7cM, respectively, with LOD scores of (29.8, 31.7, 31.2, 34.0, 29.9, 56.6, 52.2, 56.0, 59.4, 60.5, 53.5, and 66.4, respectively) (Table 2). Therefore, these SSR markers

Table 1. Characteristics of the twenty SSR markers generated polymorphism between the parents.

Marker	Primer sequences	Annealing temperature (°C)	Product size (bp)
<i>Wmc24-2A</i>	GTGAGCAATTTTGATTATACTG TACCCTGATGCTGTAATATGTG	51	152
<i>Wmc25-2AS,2BS,2DS</i>	TCTGGCCAGGATCAATATTACT TAAGATACATAGATCCAACACC	51	166
<i>Wmc44-1B</i>	GGTCTTCTGGGCTTTGATCCTG TGTTGCTAGGGACCCGTAGTGG	61	232 to242
<i>Wmc47-4B,5A</i>	GAAACAGGGTTAACCATGCCAA ATGGTGCTGCCAACACATACA7A	61	127 to141
<i>Wmc48-4A,4B,4D</i>	GAGGGTTCTGAAATGTTTTGCC ACGTGCTAGGGAGGTATCTTGC	61	139 to 190
<i>Wmc52-1B,4D</i>	TCCAATCAATCAGGGAGGAGT AGAACGCATCAAGGCATGAAGTA	61	192
<i>Wmc94-7D</i>	TTCTAAAATGTTTGAAACGCTC GCATTTTCGATATGTTGAAGTAA	51	107
<i>Wmc149-3B,2A,2B,5B</i>	ACAGACTTGGTTGGTGCCGAGC ATGGGCGGGGGTGTAGAGTTTG	61	180 to230
<i>Wmc154-2B</i>	ATGCTCGTCAGTGTCATGTTTG AAACGGAACCTACCTCACTCTT	61	147 to 149
<i>Wmc161-4A,5D</i>	ACCTTCTTTGGGATGGAAGTAA GTACTIONCACTTGTAACGCA	61	200
<i>Wmc168-7A</i>	AACACAAAAGATCCAACGACAC CAGTATAGAAGGATTTTGAGAG	51	319
<i>Wmc215-5D,3A,5A</i>	CATGCATGGTTGCAAGCAAAAG CATCCCGGTGCAACATCTGAAA	61	207 to 211
<i>Wmc219-4A</i>	TGCTAGTTTGTCTATCCGGGCGA CAATCCCGTTCTACAAGTTCCA	61	204
<i>Wmc261-2A,1D,2B,3B,7B</i>	GATGTGCATGTGAATCTCAAAAGT AAAAGAGGGTCACAGAATAACCTAAA	61	110
<i>Wmc273-6A,7B,7D</i>	AGTTATGTATTCTCTCGAGCCTG GGTAACCACTAGAGTATGTCCTT	51	182-235
<i>Wmc326-3B,3A,5B</i>	GGAGCATCGCAGGACAGA GGACGAGGACGCCTGAAT	61	186
<i>Wmc327-5A</i>	TGCGGTACAGGCAAGGCT TAGAACGCCCTCGTCGGA	61	183
<i>Xgwm30-2D</i>	ATCTTAGCATAGAAGGGAGTGGG TTCTGCACCCTGGGTGAT	60	156
<i>Xgwm456-1D</i>	TCTGAACATTACACAACCCTGA TGCTCTCTCTGAACCTGAAGC	55	138-165
<i>Xgwm566-3B</i>	TCTGTCTACCCATGGGATTTG CTGGCTTCGAGGTAAGCAAC	60	122-131

were linked to the quantitative trait loci (QTL) for the grain filling rate as an indicator for heat tolerance genes. All of

the QTLs had a positive additive effect indicating contribution of alleles increasing GFR by the tolerant




Figure 1. Selective genotyping of F₂ progeny of Ksu 106 X Yecora Rojo wheat cultivars with the *Wmc326-3B, 34A, 5B*; *Wmc25-2AS, 2BS,2DS* and *Wmc327-5A* markers for heat tolerance. Lane M: Molecular weight followed by P₁ and P₂ parents Ksu106 and Yecora Rojo, respectively. T=F₂ Tolerant plant; S=F₂ Sensitive plant; TS = F₂ Heterozygous plant. Arrow points to polymorphic bands of the SSR markers.

Table 2. Genetic characteristics of QTL related to the grain filling rate as indicator for heat tolerance genes in the 205 F₂ plants population of Ksu106 X Yecora Rojo.

Marker	Chromosome arm	Type of marker	QTL (cm)	LOD	R ² (%)	P value	Additive effect
<i>Wmc24</i>	1AS	Dominant	9.9	29.8	31	0.05	122.48
<i>Wmc25</i>	2AS,2BS,2DS	Co-dominant	7.8	56.6	45	0.01	135.89
<i>Wmc44</i>	1BL	Co-dominant	9.7	52.2	30.00	0.05	103.99
<i>Wmc47</i>	4B,5A	Dominant	10.9	29.4	15.00	0.21	52.64
<i>Wmc48</i>	4A,4BS,4DS	Dominant	44.9	2	-	0.16	-
<i>Wmc52</i>	1B,4DS	Dominant	32.9	6.9	8.00	0.29	68.71
<i>Wmc94</i>	7DL	Co-dominant	8.4	56	39	0.05	121.61
<i>Wmc149</i>	3B,2A,2B,5BS	Dominant	20.7	14.5	8.00	0.22	77.98
<i>Wmc154</i>	2Bs	Dominant	26.8	10.1	8.00	0.55	22.9
<i>Wmc161</i>	4A,5DL	Co-dominant	7.3	59.4	43	0.05	131.06
<i>Wmc168</i>	7AS	Dominant	9.0	31.7	27	0.05	87.76
<i>Wmc215</i>	5D,3A,5AL	Co-dominant	16.4	35.4	28.00	0.37	99.24
<i>Wmc219</i>	4AL	Dominant	16.3	20.8	16.00	0.38	23.85
<i>Wmc261</i>	2AL,1D,2B,3B,7B	Dominant	17.2	19.8	17.00	0.70	23.39
<i>Wmc273</i>	7AL,7B,7D	Co-dominant	7.0	60.5	48.00	0.01	148.23
<i>Wmc326</i>	3B,3A,5B	Dominant	9.0	31.2	22	0.05	141.2
<i>Wmc327</i>	5AL	Co-dominant	8.6	53.5	32	0.05	112.95
<i>Xgwm30</i>	2D	Dominant	7.6	34.0	29	0.01	117.54
<i>Xgwm456</i>	1D	Dominant	8.9	29.9	33	0.05	61.11
<i>Xgwm566</i>	3B	Co-dominant	5.7	66.4	64	0.05	158.6

parent 'Ksu106' (Table 2). Positive additive effect of the QTL on chromosomes 1A, 2A, 3A, 4A, 5A, 7A, 1B, 2B, 3B, 5B, 7B, 1D, 2D, 5D and 7D indicates contribution of QTL alleles in these loci from the tolerant parent, 'Ksu106'. In all of the detected QTLs, additive effects were positive. This indicates the relative importance of additive gene effects in controlling GFR as an indicator for heat tolerance in F_2 population. The present study indicated that SSR markers, combined with bulked segregant analysis, could be used to identify molecular markers linked to the grain filling rate as an indicator for heat tolerance genes in wheat and suggested that marker-assisted selection with microsatellite primers might be useful for developing improved cultivars.

DISCUSSION

Breeding for heat tolerance is still in its infancy and warrants more attention than it has been given in the past. It is unfortunate that the literature contains relatively little information on breeding for heat tolerance in different crop species. However, despite all the complexity of heat tolerance and difficulties encountered during transfer of tolerance, some heat-tolerant inbred lines and hybrid cultivars with commercial acceptability have been developed and released, at least in a few crop species such as tomato (Scott et al., 1986, 1995). Composite interval mapping (CIM) has been used in the present investigation for mapping QTLs associated with heat tolerance using pooled bulks from the F_2 population of Ksu106 (heat tolerant)/Yecora Rojo (heat-sensitive) cross. This method combines interval mapping with linear regression and includes additional genetic markers in the statistical model in addition to an adjacent pair of linked markers for interval mapping (Jansen, 1993; Jansen and Stam, 1994; Zeng, 1994). The main advantage of CIM is that it is more precise and effective at mapping QTLs compared to single-point analysis and interval mapping.

Microsatellites are highly popular genetic markers because of their co-dominant inheritance, high abundance, enormous extent of allelic diversity, and the ease of assessing SSR size variation by PCR with pairs of flanking primers. The reproducibility of microsatellites is such that they can be used efficiently by different research laboratories to produce consistent data (Saghai-Marouf et al., 1984). In the present study, the markers *Wmc24*, *Wmc168*, *Wmc326*, *Xgwm30*, *Xgwm456*, *Wmc25*, *Wmc44*, *Wmc94*, *Wmc161*, *Wmc273*, *Wmc327*, and *Xgwm566* were assigned to chromosomes 1A, 7A, (3B, 3A, 5B), 2D, 1D, (2A, 2B, 2D), 1B, 7D, (4A, 5D), (7A, 7B, 7D), 5A, and 3B, respectively following several investigators (Roder et al., 1998; Somers et al., 2004; Golabadi et al., 2011). Homoeologous groups of chromosomes 2, 3, 5 and 7 of wheat contain a number of genes that are important for tolerance to abiotic stress (Dubcovsky et al., 1995; Somers et al., 2004; Golabadi et

al., 2011).

Some markers tightly linked to genes were found by using bulked segregant analysis (BSA) (Xu et al., 1995; Mackay and Caligari, 2000; Altinkuet and Gozukirmizi, 2003; Podlich et al., 2004; Govindaraj et al., 2005; Zhang et al., 2009; Barakat et al., 2010; Milad et al., 2011). BSA was firstly reported by Michelmore et al. (1991) to identify random amplified polymorphic DNA (RAPD) markers tightly linked to genes for resistance to lettuce downy mildew. Using a method inspired by BSA, we were able to identify the aforementioned 12 SSR markers associated with the grain filling rate in wheat under heat-stress. These markers should be useful for marker-assisted selection. The present results support the idea that BSA can provide fast detection of molecular markers linked to genes of interest.

The quantitative and molecular characterization of heat tolerance in hexaploid wheat has previously been investigated (Yang et al., 2002). They reported that two markers, *Xgwm11* and *Xgwm293*, were linked to the grain filling duration (GFD) by quantitative trait loci (QTL) analysis of an F_2 population. Recently, a bulked segregant analysis to detect QTLs related to heat tolerance in rice (*Oryza sativa* L.) using SSR markers has been reported (Zhang et al., 2009). In this study, multiple regressions of GFR under heat stress on the 5 dominant SSR markers and 7 co-dominant SSR markers were significant. The R^2 values suggested that *Wmc24*-linked QTL, *Wmc168*-linked QTL, *Wmc326*-linked QTL, *Xgwm30*-linked QTL, *Xgwm456*-linked QTL, *Wmc25*-linked QTL, *Wmc44*-linked QTL, *Wmc94*-linked QTL, *Wmc161*-linked QTL, *Wmc273*-linked QTL, *Wmc327*-linked QTL, and *Xgwm566*-linked QTL account 31, 27, 22, 29, 33, 45, 30, 39, 43, 48, 32, and 64% of the total phenotypic variation, respectively, in heat tolerance in the F_2 population.

Conclusion

The aforementioned results, although preliminary, because of the low number of SSR primers and SSR markers associated with the grain filling rate in wheat under heat-stress, are nevertheless encouraging. Moreover, it is suggested that using more SSR primers to construct a linkage map from $F_{4,5}$ families in future studies in our breeding program will help to detect more closely linked markers associated with the grain filling rate in wheat under heat-stress.

Acknowledgements

The investigators acknowledges the Center of Excellence in Biotechnology Research (grant CEBR2 - 08), King Saud University and Ministry of Higher Education, Saudi Arabia for this generous financial support.

REFERENCES

- Al-Doss AA, Moustafa KA, Ahmed EI, Elshafei AA, Barakat MN (2009). Assessment of genetic diversity in Saudi wheat genotypes under heat stress using molecular markers and agronomic traits. *Int. J. Plant Breed. Global Sci. Books* 3:103-110.
- Altinkuet A, Gozukirmizi N (2003). Search for microsatellite markers associated with water-stress tolerance in wheat through bulked segregant analysis. *Mol. Biotechnol.* 23:97-106.
- Barakat MN, El-Shafei AA, Al-Doss AA (2010). Molecular mapping of QTLs for resistance to northern corn leaf blight in maize. *J. Food Agric. Environ.* 8:547-552.
- Barakat MN, Al-Doss AA, Elshafei AA, Moustafa KA (2011). Identification of new microsatellite marker linked to the grain filling rate as indicator for heat tolerance genes in F₂ wheat population. *Australian J. Crop Sci.* 5(2):104-110.
- Dubcovsky J, Luo MC, Dvorák J (1995). Linkage relationships among stress-induced genes in wheat. *Theor. Appl. Genet.* 91:795-801.
- Egli DE, Ramseur EL, Yu ZW, Sullivan CH (1989). Source-sink alterations affect the number of cells in soybean cotyledons. *Crop Sci.* 29:732-735.
- Gandourah MO (1989). Grain yield and its stability of some selected wheat varieties in Saudi Arabia. *J. King Saud Univ.* 1:75-85.
- Golabadi M, Arzani A, Tabatabaei BES, Maibody SM, Mohammadi SA (2011). Identification of microsatellite markers linked with yield components under drought stress at terminal growth stages in durum wheat. *Euphytica* 177:207-221.
- Govindaraj P, Arumugachamy S, Maheswaran M (2005). Bulk segregant analysis to detect main effect QTL associated with grain quality parameters in Basmati 370/ASD16 cross in rice (*Oryza sativa* L.) using SSR markers. *Euphytica* 144:61-68.
- Gupta PK, Balyan HS, Edwards KJ (2002). Genetic mapping of 66 new microsatellite (SSR) loci in bread wheat. *Theory Appl. Genet.* 105:413-422.
- Hays DB, Do JH, Mason RE, Morgan G, Finlayson SA (2007). Heat stress induced ethylene production in developing wheat grains induces kernel abortion and increased maturation in a susceptible cultivar. *Plant Sci.* 172:1113-1123.
- Jansen R, Stam P (1994). High resolution of quantitative traits into multiple loci via interval mapping. *Genetics* 136:1447-1455.
- Jansen R (1993) Interval mapping of multiple quantitative trait loci. *Genetics* 135:205-211.
- Kirigwi FM, Ginkel MV, Gill GB, Paulsen GM, Fritz AK (2007). Markers associated with a QTL for grain yield in wheat under drought. *Mol. Breed.* 20:401-413.
- Knott DR, Gebeyehou G (1987). Relationship between the lengths of the vegetative and grain filling periods and the agronomic characters in three durum wheat crosses. *Crop Sci.* 27:857-860.
- Lander E, Botstein D (1989). Mapping Mendelian factors underlying quantitative traits using RFLP linkage maps. *Genetics* 121:185-199.
- Li XJ, Pan ZD (2005). A study on the grain filling characteristic of different weight wheat. *Rev China Agric. Sci. Tech.* 7:26-30.
- Liu BH (1998). *Statistical Genomics*. CRC Press, Boca Raton, FL. pp. 387-416.
- Mackay IJ, Caligari PDS (2000). Efficiencies in F₂ and backcross generations for bulked segregant analysis using dominant markers. *Crop Sci.* 40:626-630.
- Maestri E, Klueva N, Perrotta C, Gulli M, Nguyen HT, Marmioli N (2002). Molecular genetics of heat tolerance and heat shock proteins in cereals. *Plant Mol. Biol.* 48:667-681.
- Mason RE, Pacheco A, Hays DB, Mondal S, Jampala B, Beecher FW Ibrahim AM (2010). QTL associated with heat susceptibility index in wheat (*Triticum aestivum* L.) under short-term reproductive stage heat stress. *Euphytica* DOI 10.1007/s10681-010-0151-x
- Meer J, Robert HC, Kenneth FM (2002). Map manager version 0.22. <http://manager.roswellpark.org/mmQTX.html>
- Michelmore RW, Paran I, Kesseli RV (1991). Identification of markers linked to disease-resistance genes by bulked segregant analysis: A rapid method to detect markers in specific genomic regions by using segregating populations. *Proc. Natl. Acad. Sci. USA* 88:9828-9852.
- Milad SI, Wahba LE, Barakat MN (2011). Identification of RAPD and ISSR markers associated with flag leaf senescence under water-stressed conditions in wheat (*Triticum aestivum* L.). *Aust. J. Crop Sci.* 5(3):334-340.
- Nelson JC (1997). QGENE: Software for marker-based genomic analysis and breeding. *Mol. Breed.* 3:239-245.
- Plaut Z, Butow BJ, Blumenthal CS, Wrigley CW (2004). Transport of dry matter into developing wheat kernels and its contribution to grain yield under post-anthesis water deficit and elevated temperature. *Field Crops Res.* 86:185-198.
- Podlich DW, Winkler CR, Cooper M (2004). Mapping as you go: An effective approach for marker-assisted selection of complex traits. *Crop Sci.* 44:1560-1571.
- Röder MS, Korzun V, Gill BS, Ganal MW (1998). The physical mapping of microsatellite markers in wheat. *Genome* 41:278-283.
- Saghai-Marouf MA, Soliman KM, Jorgensen RA, Allard RW (1984). Ribosomal DNA spacer-length polymorphism in barley: Mendelian inheritance, chromosomal location, and population dynamics. *Proc. Natl. Acad. Sci.* 81:8014-8018.
- SAS (2007). SAS/STAT. Guide for personal computers. Version 9 eds. SAS end. SAS Institute, Cary, N.C, U.S.A.
- Scott JW, Volin RB, Bryan HH, Olson SM (1986). Use of hybrids to develop heat tolerant tomato cultivars. *Proc. Fla. State Hortic. Soc.* 99:311-315.
- Scott JW, Olson SM, Howe TK, Stoffella PJ, Bartz JA, Brya HH (1995). 'Equinox' heat-tolerant hybrid tomato. *Hort. Sci.* 30:647-648.
- Somers DJ, Isaac P, Edwards K (2004). A high-density microsatellite consensus map for bread wheat (*Triticum aestivum* L.). *Theor. Appl. Genet.* 109:1105-1114.
- Xu BY, Shen ZT, Chen Y (1995). Molecular mapping for quantitative trait loci controlling rice yield. *Acta Genetica Sinica* 22:46-52.
- Van Sanford DA (1985). Variation in kernel growth characters among soft red winter wheats. *Crop Sci.* 25:626-630.
- Wang RX, Hai L, Zhang XY, You GX, Yan CS, Xiao SH (2009). QTL mapping for grain filling rate and yield-related traits in RILs of the Chinese winter wheat population Heshangmai 3 Yu8679. *Theor. Appl. Genet.* 118:313-325.
- Wiegand CL, Cuellar JA (1981). Duration of grain filling and kernel weight as affected by temperature. *Crop Sci.* 21:95-101.
- Yang J, Sears RG, Gill BS, Paulsen GM (2002). Quantitative and molecular characterization of heat tolerance in hexaploid wheat. *Euphytica* 126:275-282.
- Zeng ZB (1994). Precision mapping of quantitative trait loci. *Genetics* 136:1457-1468.
- Zhang G, Chen L, Xiang G, Xiang Y, Chen X, Zhang S (2009). Bulk segregant analysis to detect QTL related to heat tolerance in rice (*Oryza sativa* L.) using SSR markers. *Agric. Sci. China* 8:482-487.