

Full Length Research Paper

Overexpression and characterization of a thermophilic and hemolytic phospholipase of *Vibrio vulnificus* cloned in *Escherichia coli* (phospholipase of *V. vulnificus*)

Young-Sun Lim, Tianshi Wang, Wan-Seok Oh and Jung-Wan Kim*

Division of Life Sciences, University of Incheon, Incheon, 406-772, Korea.

Accepted 8 June, 2012

A phospholipase (PLase) gene of *Vibrio vulnificus* was cloned in *Escherichia coli* and the properties of the gene product were investigated. The PLase structural gene was composed of 1,251 bp, encoding 417 amino acids for a protein with a predicted molecular mass of 47,187 Da including a putative signal sequence. The predicted protein sequence was 87 and 82% identical to those of hemolysins from *Vibrio* spp. and that of lecithinase from *V. cholerae*, respectively. A lipid binding motif, GDSL, conserved among various PLases and lipases was also observed. Over-expression of PLase caused inclusion body formation in *E. coli*, but not that of the PLase subclone without the signal sequence (45 kDa). Purified PLase exhibited hemolytic activity on red blood cells and hydrolyzed phosphatidylcholine, phosphatidylethanolamine, phosphatidylinositol, and soya-lecithin mainly to fatty acid and 1,2-diacylglycerol, indicating that it was a PLase with unique catalytic activity. PLase from *V. vulnificus* had temperature and pH optimum at 45°C and 7.0 in 50 mM Tris-HCl buffer, respectively, but was quite active at temperatures up to 55°C and in a broad range of pH 5 to 10. The activity of the enzyme was enhanced by divalent cations such as Ca²⁺, Co²⁺, Mg²⁺, and Mn²⁺, but not by ethylenediaminetetraacetic acid (EDTA).

Key words: Phospholipase, *Vibrio vulnificus*, hemolytic, *VpIA*.

INTRODUCTION

Vibrio vulnificus is a Gram negative marine bacterium that is pathogenic to both human and marine animals. The bacterium has become a great concern during summer time in Korea due to the increase of serious wound infections and fatal septicemia with high mortalities (>50%; Blake et al., 1979). The life cycle of *V. vulnificus* between marine environment and human hosts implies that the pathogen has delicate regulatory systems sensing and responding to environmental changes and that there is a close inter-relationship among the systems that affect the survival of the pathogen. Virulence of the bacterium is likely to be controlled by such complicate

and delicate networks of various regulatory systems.

V. vulnificus produces various virulent factors such as polysaccharide capsules (Simpson et al., 1987), cytolytic hemolysins (Chang et al., 1997; Lee et al., 2004), elastolytic protease (Kothary and Kreger, 1987), siderophore (Simpson and Oliver, 1983), collagenase (Smith and Merkel, 1982), and phospholipase (PLase; Testa et al., 1984) etc. Among them, PLases are known to be associated with virulence in many pathogens (Titball, 1993; Mengaud et al., 1991; Portnoy et al., 1981; Gustafson and Tagesson, 1990). Due to their catalytic capability, PLases may also be involved in scavenging phosphate when it became a factor limiting bacterial growth. PLase C of *Pseudomonas aeruginosa* has well been characterized for its role in phosphate scavenging and pathogenesis (Gray et al., 1982). Many members of the family Vibrionaceae produce and secrete PLases,

*Corresponding author. E-mail: kjw5864@incheon.ac.kr. Tel: +82-32-835-8244, +82-10-9286-8641. Fax: +82-32-835-0763.

some of which have been shown to be hemolysins (Testa et al., 1984; McCarter and Silverman, 1987; Pal et al., 1997; Shinoda et al., 1991) and others glycerophospholipid-cholesterol acyltransferases (Shaw et al., 1994; Thornton et al., 1988). PLases from other family also have been reported to be involved in the induction of apoptosis (Kirschnek and Gulbins, 2006) or in the regulation of flagella (Schmiel et al., 2000), indicating that their expression might be under the control of multiple regulatory systems.

In previous studies, the PLase activity in the lysate of *V. vulnificus* was shown to be dependent on phosphate concentration, temperature, pH, osmotic pressure, and also subjected to catabolite repression by glucose (Oh et al., 2007). Therefore, PLase that is controlled by multiple environment factors would make a good system to investigate inter-relationship among various regulatory systems. As the first step toward such research goal, a gene encoding PLase of *V. vulnificus* was cloned based on its capability of forming hemolytic zone on a blood agar plate and over-expressed in *E. coli*. The physico-chemical properties of the purified enzyme were also characterized in this study.

MATERIALS AND METHODS

Bacteria, plasmids and media

V. vulnificus ATCC29307 was cultured either in Luria-Bertani broth (LB; 0.5% yeast extract, 1% bacto tryptone, Difco, USA) containing 2.5% NaCl or heart infusion broth (beef heart infusion 1.0%, tryptose 1.0%, Difco, USA) containing 2.5% NaCl at 37°C with shaking at 200 rpm. *E. coli* MC1061 [*hsdR*, *merB*, *araD139Δ(araABC-leu)7679*, *ΔlacX79*, *galU*, *galK*, *rpsL*, *thi*] and pUC119 was used for cloning the PLase gene; *E. coli* BL21 (DE) (*clts857*, *ind1*, *sam7*, *nin5*, *lacUV5-T7 gene1*) and pET22b(+) for over-expression of the clone. The *E. coli* strains were cultured in LB broth and ampicillin (100 µg/ml) was added when it was necessary.

Cloning of the PLase gene

The PLase gene of *V. vulnificus* ATCC29307 was cloned in *E. coli* MC1061 by shotgun cloning. The *V. vulnificus* genomic DNA was digested partially with either *Sau3AI* or *HindIII* and the DNA fragments in the range of 2.0 to 9.4 kb were isolated from agarose gel by electro-blotting onto DEAE membrane. Isolated DNA fragments were ligated to pUC119 at the *BamHI* or *HindIII* site to construct a sub-genomic DNA library and the resulting library was transformed into *E. coli* MC1061. Putative PLase clones were isolated based on their hemolytic activity on sheep blood agar plates. Nucleotide sequence analysis of the insert on the positive clone (pHPLC3) was determined by the dideoxy chain termination sequencing method using an AB1377 PRISM DNA sequencer (Perkin-Elmer, USA). Nucleotide sequence was analyzed using Lasergene (DNASTAR Inc., USA) and putative signal sequence was predicted using SignalP 3.0 Server (Technical University of Denmark).

Over-expression of the PLase gene

The PLase structural gene (*VplA*) cloned on pHPLC3 was

subcloned onto pET22b(+) for over-expression in *E. coli*. The structural gene was amplified by polymerase chain reaction (PCR) using two primers, VPLF2 (5'-GAAGATAGTCATATGAAGAAGATA-3') and VPLR1 (5'-GTTGACCTCGAGATTAAGCGTTG-3'), carrying a *NdeI* and a *XhoI* site (underlined), respectively. In order to remove the signal sequence from the gene another primer set, VPLF3 (5'-AAACATATGGATGAGCCAGCTCTCTCTC C-3') and VPL4 (5'-AAAGTCGACGCTAAAAATTA AAGCGTTGC-3'), carrying an *NdeI* and a *SalI* site (underlined), respectively, was used. The 1.25 kb long PCR product was digested with *NdeI* and *XhoI* (or *SalI*) and ligated to the corresponding sites on pET22b(+) to place the *VplA* gene under the control of the T7 promoter and to tag the 3' end of the gene with the sequence encoding 6xHis. Putative subclones were screened for their capability of forming clear zone around the colony by producing hemolysin and/or PLase activities upon induction with 0.5 mM Isopropyl-β-D-thiogalactopyranoside (IPTG) on a blood agar plate and a lecithin agar plate containing 1% lecithin, respectively. The subclone of PLase with or without the signal sequence was designated as pVPL10 and pVPL11, respectively.

Purification of recombinant PLase

PLase was over-expressed in *E. coli* BL21 (DE3) carrying one of the subclones, pVPL10 or pVPL11 from the T7 promoter by inducing with 0.5 mM IPTG during early exponential growth phase in LB broth containing 100 µg/ml of ampicillin. However, PLase with the signal sequence from pVPL10 was found in the insoluble fraction as inclusion body and the enzyme was refolded and purified according to Wingfield et al. (1995). The cell extract in the His-Tag binding buffer containing 8 M urea was subjected to chromatography using a Ni-NTA column (Qiagen, Germany) after sonication in the presence of 1 mM phenylmethylsulfonyl fluoride (PMSF) and filtration of the supernatant through 3M paper (Whatman, USA). The enzyme bound to the resin was renatured by sequential wash and incubation with renaturation buffers [100 mM NaCl, 20% glycerol, 20 mM Tris-HCl (pH 7.4)] containing 6, 4, 2, 1, and 0 M urea. Finally, the enzyme was washed with 20 mM Tris-HCl (pH 8.0) and eluted with 20 mM Tris-HCl (pH 8.0) containing 100 mM imidazole. PLase without the signal sequence from pVPL11 was purified from the supernatant of the cell lysate using regular buffers for Ni-NTA chromatography. Protein concentration was determined by the Bradford method (Bradford, 1976) and the purified enzyme was analyzed by sodium dodecyl sulphate-polyacrylamide gel electrophoresis (SDS-PAGE) using 10% polyacrylamide gel.

Enzyme assay

PLase activity was determined using an artificial substrate, *p*-nitrophenylphosphorylcholine (pNPPC; Sigma Co., USA; Kurioka and Matsuda, 1976). Enzyme solution (200 ng/100 µL) was mixed with pre-warmed 50 mM pNPPC (60 µL) dissolved in dH₂O and 2 M Tris-HCl (pH 7.0) containing 1 mM ZnCl₂ (140 µL). The reaction mixture was incubated at 37°C for 10 min or until yellow color was developed. The reaction was stopped by boiling for 3 min and mixing with 800 µL of ethanol and 200 µL of 2 M Tris-HCl (pH 7.0). Optical density of the supernatant was measured at 420 nm. One PLase unit was defined as the amount of enzyme that released 1 mmole of *p*-nitrophenol at 37°C for 1 h (Kuroshima and Hayano, 1982). Moreover, hemolytic activity of PLase was analyzed using sheep red blood cells according to Lee et al. (2002a) or on sheep blood agar plates for the capability of the clone forming clear zone. Lecithinase activity of PLase was examined using agar plates containing 1% lecithin based on the formation of clear zone around colonies (Sibinovic et al., 1971).




Figure 1. Agar plate assay of hemolytic (A) and lecithinase (B) activities of the *VplA* clones. Panel A represents a blood agar plate inoculated with *V. vulnificus* 29307 (1), *E. coli* BL21 (2), *E. coli* MC1061 harboring pHPLC3 (3), and *E. coli* BL21 harboring pVPL10 (4 to 7). Panel B represents a lecithin (egg yolk) agar plate inoculated with *E. coli* BL21 (1) and *E. coli* BL21 harboring pVPL10 (2 to 4).

Thin layer chromatography (TLC)

In other to assay phospholipid hydrolyzing activity of the cloned PLase, TLC was carried out as follows. Purified PLase was mixed with various phospholipids (0.5%; Sigma Co., USA.) homogenized in 50 mM Tris-maleic acid-NaOH buffer (pH 7.0) and incubated at 37°C for 1 h. The reaction was terminated by adding a mixture of methanol and chloroform (1: 2). The reaction mixture was spotted on a silica gel plate (Whatman, England) and separated using a mixture of n-hexane: diethyl ether: acetic acid (80: 20: 4). The plate was developed by dipping into the mixture of 5% H₂SO₄ and methanol.

RESULTS AND DISCUSSION

Screening of the sub-genomic DNA libraries for the putative PLase gene

From the two sub-genomic DNA libraries of *V. vulnificus*, several putative clones were isolated based on their capability of hemolyzing red blood cells. Five hemolytic clones from the *Sau3AI* sub-genomic DNA library (pSPLC1-5) had inserts of different lengths but with a common open reading frame (ORF) for the zinc metalloprotease gene of *V. vulnificus* (Chuang et al., 1997). The hemolytic zone of the clones on the blood agar plate was hazy and small. Three hemolytic clones from the *HindIII* sub-genomic DNA library (pHPLC1-3) carried an identical insert of 3.5 kb long (Supplementary data). *E. coli* MC1061 carrying pHPLC3 formed a clearer hemolytic zone than *V. vulnificus* (Figure 1a) and the metalloprotease clones (data not shown) on blood agar plates. They also showed small clear zone around the colonies on a lecithin agar plate (Figure 1b). The coding

region for the hemolytic and pNPPC hydrolyzing activities was defined to the 2.5 kb *HindIII* fragment by deletion mapping (data not shown). Therefore, nucleotide sequence analysis of the DNA fragment was carried out to determine the primary structure of the PLase gene.

Primary structure of the PLase gene

Nucleotide sequence analysis of the insert revealed an open reading frame (ORF) of 417 amino acid residues for a protein with a predicted molecular mass of 47,187 Da (Figure 2). The ORF was preceded by a putative ribosome binding site and potential promoter sequences at the 5' upstream region and followed by an inverted repeat for transcriptional termination at the 3' downstream. The gene was designated as *VplA*. The first 19 amino acids of the ORF with a stretch of hydrophobic residues were typical of a signal sequence for secretion, correlating well with the secreted nature of PLase in *V. vulnificus*.

A homology search of the GenBank/EMBL database revealed that the ORF shared homology with various PLases and hemolysins. The ORF was 99% identical to the PLase encoded by a gene (Gene ID 1181413, VV2_1483) on chromosome II of *V. vulnificus* CMCP6 strain. It also shared 87% identity with hemolysin of *Vibrio harveyi* (Zhang et al., 2001), 83% with thermolabile hemolysin of *Vibrio parahaemolyticus* (Taniguchi et al., 1986), 82% with lecithinase of *V. cholerae* (Fiore et al., 1997), and 69% with hemolysin of *Vibrio campbellii* (Luis and Hedreyda, 2006), and 64% with PLase A of *Vibrio mimicus* (Kang et al., 1998; Lee et al., 2002b) at the amino




Figure 2. DNA and predicted amino acid sequences of the *VplA* gene and its flanking regions. The sequences of putative promoters are boxed and ribosome binding sites (rbs) are shown in a black box. A transcription terminator-like sequence was underlined. Putative signal sequence was also underlined with a cleavage site indicated by an arrow. The conserved blocks for lipolytic enzymes are shown in shaded boxes and the residues constituting an active site were marked with asterisks.

acid sequence level. The ORF carried conserved sequences (Block I-V, Figure 3) frequently observed among various lipases/acetyltransferases in their active sites (Thornton et al., 1988; Brumlik and Buckley, 1996). The catalytic signature for lipolytic enzymes, GDSL, was observed in Block I and the residues constituting the

active site are all conserved except one in Block V, in which a D residue was substituted with a G at residue 386 (Figure 3). The significance of the substitution is yet to be elucidated. Based on the results of the comparative analysis, the cloned DNA fragment was likely to carry a putative hemolytic PLase gene (*VplA*). Southern blot analysis confirmed that the insert was originated from the chromosomal DNA of *V. vulnificus* (data not shown).

Expression of the *VplA* gene in *E. coli*

The *VplA* gene cloned on pHPLC3 was subcloned onto an expression vector, pET22b(+), since it expressed the PLase activity poorly in *E. coli* when pNPPC was used as the substrate (Supplementary data). The structural *VplA* gene amplified by PCR was placed under the control of T7 promoter and fused to six histidine residues at the carboxyl terminus on the expression vector. The resulting recombinant DNA was designated as pVPL10 and *E. coli* carrying the subclone formed even clearer and larger zones on both blood and lecithin agar plates than those carrying pHPLC3 (Figure 1). However, purification of *VplA* from the supernatant of the *E. coli* cell lysate using a Ni-NTA column was not successful, since the protein induced by the addition of IPTG was precipitated forming inclusion body (Figure 4a, lanes 3 and 4). Therefore, in order to recover active *VplA* from the insoluble fraction of the cell lysate, the proteins in the precipitate was solubilized in the presence of 8 M urea (Figure 4a, lane 5) and mixed with Ni-NTA agarose before being packed into a column. Then, the enzyme was refolded before eluted by sequential removal of urea as described above. Renatured *VplA* was present in the soluble fraction and purified to an apparent homogeneity (Figure 4a, lane 6).

Inclusion body formation of PLases that were overexpressed in heterologous systems has been reported previously for PLase C from *Bacillus cereus* (Tan et al., 1997), YplA from *Yersinia enterocolitica* (Hatic et al., 2002), and human pancreatic PLase (Han et al., 1997). However, when *VplA* was expressed without the signal sequence, the mature protein (ca. 45 kDa) did not form inclusion body and was purified from the soluble fraction of the cell lysate in one step using the Ni-NTA column (Figure 4b). The two PLases expressed from pVPL10 and pVPL11 showed identical enzymatic properties.

Substrate specificity of *VplA*

Substrate specificity of purified *VplA* was analyzed using 0.5% (w/v) phosphatidylcholine, phosphatidylethanolamine, phosphatidylinositol, phosphatidylserine, and soya-lecithin homogenized in 50 mM Tris-maleic acid-NaOH buffer (pH 7.0) at 37°C for 1 h. TLC analysis of apolar lipids in the reaction mixtures indicated that the

| | Accession Number | Block I | Block II | Block III | Block IV | Block V |
|--------------------------------|------------------|-----------------------------|----------------------------|------------------|----------------------|----------|
| <i>Vibrio vulnificus</i> | vv2_1483 | IVAF <u>GD</u> SLSDTG (36) | LPLYNWAV <u>GG</u> AA (34) | FTLEFGLNDFM (22) | AGAKNIMLMTLPD (101) | FWGVTHPT |
| <i>Aeromonas hydrophila</i> | P10480 | IVM <u>FGD</u> SLSDTG (37) | LTIANEA <u>EGG</u> PT (37) | VILWVGANDYL (26) | NGAKQIILLFNLPD (128) | FWDQVHPT |
| <i>Vibrio parahaemolyticus</i> | Q99289 | VVAL <u>GDS</u> SLSDTG (36) | LPLYNWAV <u>GG</u> AA (34) | FTLEFGLNDFM (22) | AGAKNFMLMTLPD (101) | FWNVTHPT |
| <i>Xenorhabdus uminescens</i> | P40601 | LYV <u>FGD</u> SLSDGG (34) | KGGTNYA <u>AGG</u> AT (31) | YVHWIGGNDVD (29) | AGAGLVIVPTVPD (155) | FADDFHPT |
| <i>Pseudomonas putida</i> | P40604 | MIV <u>FGD</u> SLSDAG (48) | PDGNWAV <u>GG</u> YT (45) | YYLTGGGNDFL (26) | GGARYIMVWLLPD (94) | FNDLVHPT |
| <i>Arabidopsis thaliana</i> | P40602 | VFF <u>FGD</u> SVFDTG (62) | LTGVSFAS <u>GG</u> AG (55) | AIVVGGSNLI (35) | YGARRIGVIGTTP (101) | FWDGVHPT |
| <i>Brassica napus</i> | P40603 | VFF <u>FGD</u> SIFDTG (68) | LTGVSFAS <u>GG</u> AG (45) | AIVVAGSNLI (35) | YGARRIGVIGTTP (101) | FWDGLHPS |
| <i>Escherichia coli</i> | P29679 | LLIL <u>GDS</u> LS-AG (27) | ISGDT <u>SOQ</u> LAR (12) | VLVELGGNDGL (36) | YGRRYNEAFSAIY (26) | QDDGIHPN |
| <i>Vibrio mimicus</i> | Q07792 | LLVL <u>GDS</u> LS-AG (30) | ISGDTTGN <u>L</u> AR (12) | VLIELGANDGL (36) | YCKRYSDFYDIY (26) | MDDGLHEK |

Figure 3. Comparison of conserved blocks among various lipolytic enzymes harboring the GDSL motif. Amino acid residues that are conserved in at least six of the enzymes are presented in shade and those constituting the catalytic sites are underlined. Numbers in parentheses represent the spacing between the conserved blocks (Brumlik and Buckley, 1996).




Figure 4. SDS-PAGE analysis of the purified VpIA enzymes from *E. coli* BL21 harboring pVPL10 (A) or pVPL11 (B). Proteins were subjected to 10% SDS-polyacrylamide gel and visualized by staining Coomassie-blue. A) Lane 1 was loaded with standard markers; lane 2, *E. coli* cell lysate without induction; lane 3, *E. coli* cell lysate that had been induced by 0.5 mM IPTG; lane 4, supernatant of the induced *E. coli* cell lysate; lane 5, denatured VpIA using 8 M urea; lane 6, purified VpIA using a Ni-NTA column. B) Lane 1 was loaded with purified VpIA without the signal sequence; lane 2, standard markers.




Figure 5. Substrate specificity of VpIA analyzed by TLC. VpIA was incubated with 0.5% of each substrate homogenized in 50 mM Tris-HCl (pH 7.0) at 37°C for 30 min and the reaction mixture was subjected to TLC. PC represents phosphatidylcholine; PE, phosphatidylethanolamine; PI, phosphatidylinositol; PS, phosphatidylserine; SO, soya-lecithin. Lanes B were spotted with reactions carried out in the absence of VpIA; lanes E, those carried out in the presence of the enzyme.

substrates were hydrolyzed mainly to free fatty acids at various degrees, suggesting that VpIA might be a PLase A (Figure 5; Flieger et al., 2000). PLase A has been known to hydrolyze phospholipids into lysophosphatidylcholine and fatty acid (Dessen, 2000). However, significant amounts of 1,2-diacylglycerol were also released, implying that it also had PLase C activity (Flieger et al., 2000). VpIA preferentially hydrolyzed various phospholipids in the order of phosphatidylcholine > soya-lecithin > phosphatidylinositol phosphatidylethanolamine > phosphatidylserine. VpIA could also hydrolyze pNPPC, an artificial substrate that was specific for PLase C (Figure 6) (Kuroshima and Hayano, 1982). The results, therefore, indicated that VpIA seemed to be a unique lecithin-dependent PLase A with PLase C activity.

However, VpIA did not share significant homology with any PLase C that has been documented. PLase A has been reported from various pathogenic bacteria such as *Mycobacterium tuberculosis* (Parker et al., 2007),

Legionella pneumophila (Broich et al., 2006) and *P. aeruginosa* (Kirschnek and Gulbins, 2006) as a potent virulence factor. Recently, Koo et al. (2007) reported that a hemolytic PLase A of *V. vulnificus* could have a crucial role in the pathogenesis of the bacterial infection using a cytolysin-negative mutant strain, CVD707. However, it is not clear yet whether the PLase A enzyme they reported is a protein identical to VpIA, since they did not identify the gene responsible for the activity.

Enzymatic properties of VpIA

Temperature and pH optima of purified VpIA were investigated using 50 mM pNPPC as the substrate. VpIA had the highest activity at 45°C with a broad temperature range for activity between 25 and 80°C in 50 mM Tris-HCl buffer (pH 7.0; Figure 6a). It is interesting to note that PLase from *V. vulnificus*, a mesophile, is a thermophilic




Figure 6. Effects of various factors on the activity of VpIA. (A) Effect of temperature on the enzyme activity was analyzed using 50 mM pNPPC in 50 mM Tris-HCl (pH 7.0) for 30 min. (B) Effect of pH on the activity was analyzed using 50 mM pNPPC in 50 mM sodium acetate for pH 5.0 and 6.0; 50 mM Tris-HCl for pH 7.0-9.0; 50 mM CHES for pH 10.0. The reactions were carried out at 45°C for 30 min. (C) Effect of various divalent cations and EDTA (5 mM) on the activity was analyzed using 50 mM pNPPC in 50 mM Tris-HCl (pH 7.0) at 45°C for 30 min.

enzyme. The enzyme showed enzyme activity between pH 5 to 10, being most active at neutral pH in 50 mM Tris-HCl buffer (Figure 6b) at 45°C. These results indicated that VpIA has quite stable structure for maintaining the activity under various conditions. To our knowledge, PLase A that was active at broad high temperature and pH ranges has not been reported from *Vibrio* spp. The effect of various metal ions on the activity of VpIA was also tested in 50 mM Tris-HCl buffer (pH 7.0) at 45°C. The activity of the enzyme was enhanced by divalent cations such as Ca²⁺, Co²⁺, Mg²⁺, and Mn²⁺; inhibited severely by Cu²⁺ and Fe²⁺; but not affected significantly by either Zn²⁺ or EDTA (Figure 6c).

Previously, extracellular PLase A₂ and Lyso-PLase of *V. vulnificus* E4125 have been characterized by Testa et al. (1984). However, the physico-chemical properties of PLase A₂ were different from those of VpIA in that it was larger in molecular mass (≥80 kDa), heat-labile, and active at pH 5.0 to 5.5. The enzyme was not affected by Ca²⁺, either. Lyso-PLase was heat-resistant with molecular mass of larger than 80 kDa. Both PLases reported by the group did not seem to be identical to VpIA. Moreover, the activity of hemolytic and membranous PLase A reported by Koo et al. (2007) was dependent on Ca²⁺ and exhibited the highest activity at 10 mM. From all these results, *V. vulnificus* was likely to

possess multiple or strain specific PLases with various activities and functions. Further analysis constructing the *VplA* mutant might prove the existence of multiple PLases in the pathogenic bacterium.

ACKNOWLEDGEMENT

This study was supported by the grant from the University of Incheon.

REFERENCES

- Blake PA, Merson MH, Weaver RE, Hollis DG, Heublein PC (1979). Disease caused by a marine *Vibrio*. Clinical characteristics and epidemiology. N. Engl. J. Med. 300:1-5.
- Bradford M (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. Anal. Biochem. 72:248-254.
- Broich M, Rydzewski K, MaNealy TL, Marre R, Flieger A (2006). The Global Regulatory Proteins LetA and RpoS Control Phospholipase A, Lysophospholipase A, Acyltransferase, and Other Hydrolytic Activities of *Legionella pneumophila* JR32. J. Bacteriol. 188:1218-1226.
- Brumlik MJ, Buckley JT (1996). Identification of the Catalytic Triad of the Lipase/ Acyltransferase from *Aeromonas hydrophila*. J. Bacteriol. 178:2060-2064.
- Chang TM, Chuang YC, Su JH, Chang MC (1997). Cloning and sequence analysis of a novel hemolysin gene (*vly*) from *Vibrio vulnificus*. Appl. Environ. Microbiol. 63:3851-3857.
- Chuang YC, Chang TM, Chang MC (1997). Cloning and characterization of the gene (*empV*) encoding extracellular metalloprotease from *Vibrio vulnificus*. Gene 189:163-168.
- Dessen A (2000). Structure and mechanism of human cytosolic phospholipase A(2). Biochim. Biophys. Acta. 1488:40-47.
- Fiore AE, Michalski JM, Russell RG, Sears CL, Kaper JB (1997). Cloning, Characterization, and Chromosomal Mapping of a Phospholipase (Lecithinase) Produced by *Vibrio cholerae*. Infect. Immun. 65:3112-3117.
- Flieger A, Gong S, Faigle M, Deeg M, Bartmann P, and Neumeister B (2000). Novel Phospholipase A Activity Secreted by *Legionella* Species. J. Bacteriol. 182:1321-1327.
- Gray GL, Berka RM, Vasil ML (1982). Phospholipase C Regulatory Mutation of *Pseudomonas aeruginosa* That Results in Constitutive Synthesis of Several Phosphate-Repressible Proteins. J. Bacteriol. 150:1221-1226.
- Gustafson C, Tagesson C (1990). Phospholipase C from *Clostridium perfringens* stimulates phospholipase A₂-mediated arachidonic acid release in cultured intestinal epithelium cells (INT 407). Scand. J. Gastroenterol. 29:243-247.
- Han SK, Lee BI, Cho W (1997). Bacterial expression and characterization of human pancreatic phospholipase. Biochim. Biophys. Acta. 1346:185-192.
- Hatic II SO, Picking WL, Young BM, Young GM, Picking WD (2002). Purification and Characterization of Two Active Derivatives of Recombinant YpIA, a Secreted Phospholipase from *Yersinia enterocolitica*. Biochem. Biophys. Res. Comm. 292:463-467.
- Kang JH, Lee JH, Park JH, Huh SH, Kong IS (1998). Cloning and identification of a phospholipase gene from *Vibrio mimicus*. Biochim. Biophys. Acta. 1394:85-89.
- Kirschnek S, Gulbins E (2006). Phospholipase A₂ Functions in *Pseudomonas aeruginosa*-Induced Apoptosis. Infect. Immun. 74:850-860.
- Koo B, Lee J, Kim S, Yoon H, Kim K, Kwon K, Kim H, Park J, Park B (2007). Phospholipase A as a potent virulence factor of *Vibrio vulnificus*. Intl. J. Mol. Med. 20:913-918.
- Kothary MH, Kregger AS (1987). Purification and characterization of an elastolytic protease of *Vibrio vulnificus*. J. Gen. Microbiol. 133: 1783-1791.
- Kurioka S, Matsuda M (1976). Phospholipase C assay using *p*-nitrophenylphosphorylcholine together with sorbitol and its application to studying the metal and detergent requirement of the enzyme. Anal. Biochem. 75:281-289.
- Kuroshima T, Hayano K (1982). Phospholipase C activity in soil. Soil Sci. Plant Nutr. 28:535-542.
- Lee JH, Ahn SH, Kim SH, Choi YH, Park KJ, Kong IS (2002a). Characterization of *Vibrio mimicus* phospholipase A (PhIA) and cytotoxicity on fish cell. Biochem. Biophys. Res. Comm. 298:269-276.
- Lee SE, Ryu PY, Kim SY, Kim YR, Koh JT, Kim OJ, Chung SS, Choy HE, Rhee JH (2004). Production of *Vibrio vulnificus* hemolysin *in vivo* and its pathogenic significance. Biochem. Biophys. Res. Comm. 324:86-91.
- Lee SH, Kim S, Park SC, Kim MJ (2002b). Cytotoxic Activities of *Leptospira interrogans* hemolysin SphH as a Pore-Forming Protein on Mammalian Cells. Infect. Immun. 70:315-322.
- Luis BBS, Hedreyda CT (2006). Analysis of a gene (*vch*) encoding hemolysin isolated and sequenced from *Vibrio campbellii*. J. Gen. Appl. Microbiol. 52:303-313.
- McCarter LL, Silverman M (1987). Phosphate Regulation of Gene Expression in *Vibrio parahaemolyticus*. J. Bacteriol. 169:3441-3449.
- Mengaud J, Braun-Brenton C, Cossart P (1991). Identification of phosphatidylinositol specific phospholipase C activity in *Listeria monocytogenes*: novel type of virulence factor? Mol. Microbiol. 5:367-372.
- Oh W, Im Y, Yeon K, Yoon Y, Kim J (2007). Phosphate and Carbon Source Regulation of Alkaline Phosphatase and Phospholipase in *Vibrio vulnificus*. J. Microbiol. 45:311-317.
- Pal S, Guhathakurta B, Sasmal D, Mallick R, Datta A (1997). Purification and characterization of a hemolysin with phospholipase C activity from *Vibrio cholerae* O139. FEMS. Microbiol. Lett. 147:115-120.
- Parker SK, Curtin KM, Vasil ML (2007). Purification and Characterization of Mycobacterial Phospholipase A: an Activity Associated with Mycobacterial Curinase. J. Bacteriol. 189:4153-4160.
- Portnoy DA, Moseley SL, Falkow S (1981). Characterization of plasmids and plasmid-associated determinants of *Yersinia enterocolitica* pathogenesis. Infect. Immun. 31:775-792.
- Schmiel D, Young GM, Miller VL (2000). The *Yersinia enterocolitica* Phospholipase Gene *ypIA* Is Part of the Flagellar Regulon. J. Bacteriol. 182:2314-2320.
- Smith GC, Merkel JR (1982). Collagenase activity of *Vibrio vulnificus*: potential contribution to its invasiveness. Infect. Immun. 35:1155-1156.
- Shaw JF, Chang RC, Chuang KH, Yen YT, Wang FG (1994). Nucleotide sequence of novel arylesterase gene from *Vibrio mimicus* and characterization of the enzyme expressed in *E. coli*. Biochem. J. 298: 675-680.
- Shinoda S, Matsuoaka H, Tsuchie T, Miyoshi SI, Yamamoto S, Taniguchi H, Mizuguchi Y (1991). Purification and characterization of a lecithin-dependent haemolysin from *Escherichia coli* transformed by a *Vibrio parahaemolyticus* gene. J. Gen. Microbiol. 137:2705-2711.
- Sibinovic KH, Brown FA, Pettigrew KD, Vought RL (1971). Lecithin-agar assay for lecithinase antibodies in serum. Appl. Microbiol. 21:98-103.
- Simpson LM, Oliver JD (1983). Siderophore production by *Vibrio vulnificus*. Infect. Immun. 41:644-649.
- Simpson LM, White VK, Zane SF, Oliver JD (1987). Correlation between virulence and colony morphology in *Vibrio vulnificus*. Infect. Immun. 55:269-272.
- Tan CA, Hehir MJ, Roberts MF (1997). Cloning, overexpression, refolding, and purification of the nonspecific phospholipase C from *Bacillus cereus*. Protein Expr. Purif. 10:365-372.
- Taniguchi H, Hirano H, Kobomura S, Higashi K, Mizuguchi Y. (1986). Comparison of the nucleotide sequences of the genes for the thermostable direct hemolysin and thermolabile hemolysin from *Vibrio parahaemolyticus*. Microb. Pathog. 1:425-432.
- Testa J, Daniel LW, Kregger AS. (1984). Extracellular phospholipase A₂ and lysophospho-lipase produced by *Vibrio vulnificus*. Infect. Immun. 45:458-463.
- Thornton J, Howard SP, Buckley JT (1988). Molecular cloning of a phospholipid-cholesterol acyltransferase from *Aeromonas hydrophila*. sequence homologies with lecithin-cholesterol acyltransferases and

- other lipases. *Biochim. Biophys. Acta.* 959:153-159.
- Titball RW (1993). Bacterial Phospholipases C. *Microbiol. Rev.* 57:347-366.
- Wingfield PT, Coligan JE, Dunn BM, Ploegh HL, Spicher DW (1995). Current protocols in protein science, pp. 6.2.1.-6.2.15, Wiley and Sons Inc., Hoboken.
- Zhang XH, Meaden PG, Austin B (2001). Duplication of Hemolysin Genes in a Virulent Isolate of *Vibrio harveyi*. *Appl. Environ. Microbiol.* 67:3161-3167.