

*Full Length Research Paper*

# Immunopurification and characterization of a rape (*Brassica napus* L.) seedling lipase

Belguith H.<sup>1\*</sup>, Fattouch S.<sup>2</sup>, Jridi T.<sup>1</sup> and Ben Hamida J.<sup>1</sup>

<sup>1</sup>Department of Biology, Faculty of Science, Bizerte, Tunisia.

<sup>2</sup>Biological Engineering Laboratory, INSAT, Tunis, Tunisia.

Accepted 30 March, 2009

Lipase or triacylglycerol acylhydrolase (E.C.3.1.1.3) was purified to homogeneity from rapeseed-germinated cotyledons (*Brassica napus* L.). The purification scheme involved homogenization, centrifugation, ultracentrifugation and affinity chromatography using polyclonal antibodies raised against porcine pancreatic lipase. The purified rapeseed lipase was homogenous and did not contain contaminating proteins detectable by SDS-PAGE and HPLC analysis. The specific activity of the purified preparation was increased about 1950 times, with an overall yield of 35%. The rapeseed lipase was found to be a cytosoluble, glycosylated and heat-labile serine-hydrolase. It was monomeric with a molecular mass of 38 kDa and a pH of 6.6. The purification method used in the present work is rapid, simple, and yields highly purified lipase. It may therefore be applicable in the purification of other uncharacterized plant lipases.

**Key words:** *Brassica napus* L., immuno-affinity, lipase, purification, triacylglycerol acyl hydrolase.

## INTRODUCTION

Interest in lipases (triacyl glycerol acylhydrolases: E.C.3.1.1.3) from micro-organisms, animals and plant sources has markedly increased in the last decade owing to their novel and multifold applications in industry, oleo-chemistry, organic synthesis, detergent formulation, nutrition and medicine. Lipases are unique in catalyzing the hydrolysis of fats into fatty acids and glycerols at the water-lipid interface and reversing the reaction in non-aqueous media. The comparison of the properties of these proteins provided valuable information on their evolutionary relationships. In addition, their structure and their function analysis may help us to understand the fundamental mechanisms in this family (Kanaya et al., 1998). Recently, the alignment of animal, bacterial and fungal lipase sequences suggested the presence of sequence homologies including a significant conserved region, Gly-X-Ser-X-Gly as the catalytic moiety (Kanaya et al., 1998; Saxena et al., 2003; Osterlund et al., 1996). This consensus sequence, found in the substrate binding site, contains a serine residue suspected to be essential

for binding to lipid substrates.

Lipases from a large number of bacterial (Muraoka et al., 1982; Gotz, 1991; Tyski et al., 1983; Vicente et al., 1990; Aires-Barros et al., 1991), yeast (Veeraragavan and Gibbs, 1989), fungal (Suzuki et al., 1986; Kundu et al., 1987; Sugihara et al., 1988; Isobe and Nokihara, 1991; Lang et al., 1991), and animal sources (Kanaya et al., 1998; Osterlund et al., 1996; Cambillau and Bourne, 1991; Cheng, 1985; Gargouri, 1989; Gubernator, et al., 1991; Jensen et al., 1982; Verger, 1984; Carriere, 1994) have been purified to homogeneity, but not from plants at our knowledge. Compared to bacterial, fungal and animal lipases, little is known about plant ones. This lack of information about plant lipases could be due to their complicated and laborious purification (WeselaKe and Jain, 1992). The difficulties involved in carrying out research on plant lipases, particularly from oil seeds, have been essentially attributed to their low abundance, their instability and the loss of activity using the traditional purification strategies. These procedures were generally based on clarification, precipitation, differential and density gradient centrifugations associated to a series of sequential combinations of different chromatographic steps giving up a low final yield (Ben and Mazliak, 1985). Many variant of methods have been applied to purify plant lip-

\*Corresponding author. E-mail: Hatem.Belguith@fsb.mnu.tn. Tel: 216 71 872 600. Fax: 216 71 885 480.

ases without reaching the homogeneity of preparations (Huang, 1993).

To overcome these difficulties, we have tried to apply a new technique essentially based on an immunopurification step to prepare the rapeseed lipase. Antibodies immobilized onto support matrices have been already used to immunopurify a wide variety of biological compounds such as bacterial proteins (Sjoberg and Holmgren, 1973), enzymes (Melchers and Messer, 1970), hormones (Akanuma et al., 1970; Murphy et al., 1973), viral proteins (Diacio et al., 1986), and virus particles (Kenyon et al., 1973).

We report here the immunopurification and characterization of a rapeseed lipase by affinity chromatography using an immunosorbent prepared by covalently coupling polyclonal antibodies raised against the commercial porcine pancreatic lipase (anti-PPL) to activated CNBr-sepharose 4B. In a previous work, we found that these anti-PPL antibodies cross-react with the rapeseed lipase (Belguith et al., 2001).

## MATERIALS AND METHODS

### Plant material

Rapeseed (*Brassica napus* L.), cultivated locally, were soaked for 24 h, then, surface sterilized with 5% (v/v)  $\text{CaCl}_2\text{O}_2$  in distilled water for 5 min. Sterilized seeds were allowed to germinate on filter paper moistened with distilled water in darkness at 26°C.

### Homogenization and fractionation

The rape (*B. napus* L.) seedling lipase was prepared as described in a previous work (Belguith et al., 1999). Briefly, cotyledons (20 to 30 g) from 3-days old seedlings were excised from the hypocotyls and homogenized using a pestle and a mortar in grinding buffer containing 0.15 M Tris, pH 7.5, 0.6 M Sucrose, 1 mM DTT, 1 mM Benzamidine and 10 mM KCl at a ratio of 1:5 (w/v). The resulting homogenate was filtered through a Miracloth layer and fractionated by centrifugation at 10,000 g for 30 min into a yellow fatty layer (removed carefully with a spatula), a supernatant solution (S10) and a pellet. The S10 fraction was centrifuged at 100,000 g for 3 h. The resulting supernatant (S100) was stored at -20°C. All the homogenization and centrifugation steps were performed at 4°C.

### Lipase activity assay

#### Colorimetric method

Lipase activity was determined by measuring free fatty acids produced by triacylglycerol (TAGs) hydrolysis using a colorimetric method (Duncombe, 1962). Experiments were carried out in a Teflon screw-top glass test tube, in a total volume of 1 ml. The reaction mixture contained 50 mM triolein emulsified in 5% (w/v) arabic gum, 1 M Tris-HCl pH 7.5 (Lin and Haung, 1983). Reactions were started by addition of 50  $\mu\text{l}$  enzymatic solution and allowed to proceed for 20 min in a shaking bath at 30°C. Appropriate controls were included and reactions were stopped with 5 ml of cold chloroform. Fatty acids released were converted to copper soap using 0.1% (w/v) sodium diethyl dithio-carbamate. The copper complex was subsequently estimated spectrophotometrically at 440 nm.

### Radioactive method

In radiometric assays, lipase activity was measured using TAGs containing radiolabelled acyl chains (243.092 cpm/ $\mu\text{l}$ ) as a substrate (Beisson et al., 1999).

Twenty microliter of the rapeseed lipase preparation was incubated at pH 7.5 in the presence of 10  $\mu\text{l}$  of radiolabeled triolein (22Ci mM) (Perkin-Elmer), 1% sodium taurodeoxycholate (NaTDC) and 7.6  $\mu\text{l}$  of 4 M  $\text{CaCl}_2$ , in a final volume of 200  $\mu\text{l}$ . After each 15 min, 50  $\mu\text{l}$  of the reaction mixture were added to 1 ml of the stopping buffer. Then, a mixture of methanol/chloroform/heptane (21/18/15; v/v/v) was added to extract the radiolabeled free fatty acids. After a centrifugation at 13,000 g for 2 min, 200  $\mu\text{l}$  of the aqueous upper phase was taken and mixed with 8 ml of scintillation liquid (Hionic FluorTM Packard BioScience B.V.). The radioactivity of the tritium resulting of the hydrolysis of radiolabeled TAGs was counted on a Beckman LS 1801 apparatus.

### Fluorimetric method

Fluorescent TAGs were extracted from the seed kernels of *Parinari glaberrimum*. Fifth mg of the crude lipidic extract was dissolved in 1 ml of diethylether containing 0.01% (w/v) butylhydroxytoluene (BHT) as antioxidant. The TAGs were isolated by preparative TLC under an argon atmosphere and the purity was checked by TLC. Purified TAGs were stored in an ethanol solution in the presence of 0.01% (w/v) BHT (stock solution), and stored in the dark at -20°C under an argon atmosphere.

After evaporating ethanol under a nitrogen stream, 3 mg of purified TAGs was placed in a 0.5-ml polypropylene microtube, 100  $\mu\text{l}$  of the following buffer was added: 50 mM Tris-HCl (pH 8) containing 3% gum arabic, 4 mM NaTDC, 100 mM NaCl, 6 mM  $\text{CaCl}_2$ , 0.001% (w/v) BHT. The microtube was kept closed under argon and the mixture was sonicated for 30 s in a sonicating bath (35 kHz and 30 W) (Beisson et al., 1999).

The incubation buffer (990  $\mu\text{l}$ ) and the TAG stock solution (10  $\mu\text{l}$ ) were added consecutively to a quartz cuvette of 1.5 ml (optic path-length 1 cm) containing a magnetic stirrer 8 mm in length. The final TAGs concentration was 18  $\mu\text{g}$  /ml. The above mixture is slightly turbid and requires continuous stirring to ensure homogeneity. The cuvette was kept under nitrogen using a teflon cap. After gently shaking the cuvette, it was left to equilibrate at 25°C.

Lipase solution was injected and the fluorescence due to the free fatty acids released was read at regular intervals under continuous stirring in a spectrofluorimeter (SFM 25 from Kontron). Excitation was at 324 nm and emission at 420 nm (Beisson et al., 1999).

The following standard incubation buffer (pH 8) was used: 4 mM NaTDC, 100 mM NaCl, 6 mM  $\text{CaCl}_2$ , 0.001% (w/v) BHT.

### Electrophoresis

SDS-PAGE (Laemmli, 1970) was carried out on a discontinuous 6% stacking and 12% resolving polyacrylamid gel with Tris-Glycine running buffer and staining with Coomassie Brilliant Blue R-250 or silver nitrate. Standards (Sigma, St. Louis, MO, USA): Alpha-lactalbumine (14.4 kDa), trypsin inhibitor (20.1 kDa), carbonic anhydrase (29 kDa), egg albumine (45 kDa), bovine albumin serum (66 kDa), and phosphorylase (97 kDa). The native-PAGE was done in the same gel and buffer conditions without SDS. Total proteins were quantified assayed according to the Bradford method (Bradford, 1976) using bovine serum albumin (Sigma) as standard.

### HPLC analysis

Reversed-phase HPLC (C8 column Ultra sphere octyl 5  $\mu\text{m}$ ; 25 x 0.46 cm), using a linear gradient of acetonitrile in 0.1% (v/v)

aqueous trifluoroacetic acid, was used to analyze the active immunopurified fraction.

### Polyclonal antibody production

Five-hundred  $\mu\text{l}$  (0, 5 mg protein) of porcine pancreatic lipase type VI-S (Sigma, St. Louis, MO) mixed with an equal volume of complete Freund adjuvant was injected into rabbits. Two further injections were given fortnight intervals with the same amount of the immunogen emulsified in incomplete Freund adjuvant. Two weeks after the booster injection, sera samples were collected in order to evaluate the immune response of the rabbits, using ring test and ELISA as described later. Another injection was carried out after 5 weeks and the animals were bled two weeks after that last injection.

### Direct-binding ELISA test

Enzyme-linked Immunosorbent assay was performed using the method of Kang et al (1988). For all ELISA procedures the following buffers were used. Coating buffer: PBS. Wash medium: PBS-Tween-20 (0.5 g . L<sup>-1</sup>). Saturating buffer: PBS-Tween-20 containing bovine serum albumin (BSA; 5 %). Substrate solution: O-phenylene-diamine (0,4 g.L<sup>-1</sup>) (Sigma, St. Louis, MO) dissolved in 0.05 M sodium phosphate/citrate buffer, pH 5, containing hydrogen peroxide. Stop solution: 3 M chloridric acid.

### Immunoprecipitation

To immunoprecipitate lipase IRS-lipase and HPL, 20  $\mu\text{l}$  of lipase was added to 20  $\mu\text{l}$  of anti- EC-A lipase immune serum at a 1:500 dilution. The total mixture was incubated at 37°C for 1 h, immunoprecipitate was centrifuged at 10,000 x g for 10 min and residual lipase activity remaining in the supernatant was determined by the fluorimetric and radiolabeled method.

### Immunoaffinity chromatography column

A CNBr-Sepharose 4B (Pharmacia Biotech AB) pre-activated gel was used to prepare the immunoaffinity column. 20 mg of the purified IgG anti-PPL was dissolved in the coupling buffer 0.1 M NaHCO<sub>3</sub> pH 8.3 containing 0.5 M NaCl.

The anti-PPL antibodies were first coupled to the CNBr-Sepharose 4B pre-activated gel according to the manufacturers' instructions and all free antibodies were removed by washing, and BSA (3%) was used to block unreacted gel groups. To control the efficiency of the adsorbance of antibodies to the gel, we measured the absorbance at 280 nm of the eluted fraction. Proteins of the S100 fraction (10 ml) concentrated by acetone precipitation were taken up in binding buffer (0.1 M NaHCO<sub>3</sub>, pH 8.2), applied to the immunoaffinity column equilibrated with the same buffer and allowed to react overnight at 4°C with the immuno-adsorbent to optimize the fixation of the rapeseed lipase by the antibodies Anti-PPL. The column was washed with binding buffer (4 x 5 mL), to remove unbound proteins.

To elute bound proteins, the column was washed with glycine buffer 0.25 M pH 2.2, and in order to make the pH close to the neutrality, each eluted volume was supplemented with 50  $\mu\text{l}$  of Tris-HCl 1 M (pH 8). The eluted fractions were submitted to a lipase activity test and a protein quantification test (measurement of the absorbance at 280 nm).

The supernatant S100 was obtained by ultracentrifugation at 100,000 g/3h and 10 ml subjected to chromatography on CNBr activated sepharose 4B coupled with the immune globulin G fraction of antiserum raised against PPL. Protein bound to the immunoaffinity matrix was eluted by adsorption with glycine buffer 0.25 M pH 2.2.

## RESULTS

### Enzyme purification

Rapeseed lipase activity was found in the soluble fraction of 3-day old seedlings. After ultra-centrifugation at 100,000 x g approximately 70% of rapeseed lipase activity was remained in the supernatant S100 (Belguith et al., 1999). In the present work we used an immunoaffinity chromatography to purify rapeseed lipase from the soluble fraction S100. The S100 proteins concentrate was subjected to immunoaffinity chromatography on immobilized polyclonal antibodies against PPL, which had been demonstrated in a previous work to present a cross reactivity with the rapeseed lipase (Belguith et al., 2001).

Figure 1 shows the elution profile of rapeseed lipase with 0.2 M glycine buffer, pH 2.2, from the affinity chromatography on CNBr-activated sepharose-4B column. Protein quantification and lipase activity test were performed for each fraction, only in the fractions 4, 5, 6 and 7 a lipase activity was detected. The overall purification protocol is summarized in the Table 1. The most striking aspect of the procedure is that the immunopurification step yielded relatively little protein, but with a specific activity of 91.83 nkat.mg<sup>-1</sup>. The enzyme preparation eluted from the immunoaffinity column which corresponded to a yield of about 35% and a 1953-fold enrichment of lipase activity of the original fraction (Table 1), was homogeneous as proved by SDS-PAGE analysis. The polypeptide eluted from the immunoaffinity column migrated as a single peptide protein species upon SDS-PAGE at approximately 39 kDa (Figure 2). This result suggests that the IRSL is a monomeric protein with a molecular weight about (39 kDa) similar to the *Euphorbia characias* lipase (Moulin et al., 1994).

In order to evaluate the purity of the IRSL, the active immunopurified fraction was analyzed by reverse phase HPLC. A single and symmetric peak was resolved at about 50% acetonitrile and eluted at 19.72 min (Figure 3). The obtained HPLC profile confirms that IRSL was homogeneous. Our results demonstrate the efficiency of the immunoaffinity chromatography to purify plant lipases. The immunoaffinity column allowed us to purify the rapeseed lipase with a high purification factor.


This type of strategy may be useful in the purification of low abundance plant proteins such as lipases.

### Biochemical characteristics of cytosolic rapeseed lipase

In a first step, different tests were used to evaluate and to confirm that the immunopurified rapeseed lipase is a true triacylglycerol acylhydrolase (E.C.3.1.1.3) of germination. In a second step we have characterized the IRSL.

### Lipase activity tests

In order to confirm that the IRSL is a true triacylglycerol



**Figure 1.** Elution profile from affinity chromatography on CNBr-activated sepharose-4B column. Enzyme preparation (S100) was applied onto the column with stepwise elution with 0.2 M glycine buffer, pH 2.2. Protein quantification and lipase activity test were performed for each fraction.

**Table 1.** Summary of the rapeseed lipase immunopurification.

Preparation step	Total proteins (mg)	Total activity (nkat)	Specific activity (nkat.mg <sup>-1</sup> )	Purification factor	Yield (%)
Crude extract	264	12.525	0.047	1	100
Supernatant S100	88.71	8.35	0.094	2	66.6
Immunoaffinity Column elute	0.048	4.408	91.83	1953.82	35.19


acylhydrolase (E.C.3.1.1.3) of germination, we tested the lipase activity using two different methods.

### Results obtained using radiolabeled TAGs

Twenty microliters of the IRSL were incubated at pH 7.5 in presence of 10  $\mu$ l radiolabeled acyl fatty chains

(243.092 cpm/ $\mu$ l), 1 % NaTDC, 7.6  $\mu$ l 4 M of CaCl<sub>2</sub>, in a final volume of 200  $\mu$ l. We quantify the cpm resulting of the hydrolysis of radiolabeled TAG. Figure 4 clearly shows that the IRS-lipase presents a high activity level with a specific activity of about 94 nkat.mg<sup>-1</sup>.

We observe a linear kinetic for 15 min, after that it reaches a plateau. No activity was observed in the heat-treated IRSL for 5 min at 90°C used as a control. These



**Figure 2.** SDS-PAGE analysis of the immunopurified rapeseed lipase on a 12% gel, stained with silver nitrate. Lane 1, 12  $\mu$ g of IRS-lipase; lane 2, molecular mass markers.

results also suggest that the IRS-lipase is a heat-labile enzyme.

### Result obtained using fluorimetric method

In this method we used a *Parinari glaberrimum* TAGs as a substrate. The fluorescence was recorded versus time 10 min for 180 min. A significant level of lipase activity was detected using 20  $\mu$ l of IRSL. Figure 5 shows that the increase of relative fluorescence was linear with time during 120 min of incubation, with a specific activity of about 92 nkat.mg<sup>-1</sup>, after that it reaches a plateau. These results are identical of the obtained ones using the radiolabeled method.

### Esterase activity test

This test was performed to investigate if the IRSL presents or not an esterase activity such as some described partially purified plant lipases. Figure 6 shows the result of the separation by native-PAGE of the IRSL and the crude 3-days germinating cotyledons extract, submitted after that to a specific revelation using the  $\alpha$ -naphthol acetate as a substrate. The obtained electrophoregramme revealed three esterase isoforms in the crude extract (lanes 2 and 3) as found in the sunflower germinating crude extract (Ben, 1986) and in the excised

sunflower cotyledons crude extract. No esterase activity was detected in the IRSL fraction (lane 1). This result indicates that the IRS-lipase did not exhibit any esterase activity. Thus, this immunopurified enzyme can be considered as a true lipase.

### Some structure and molecular properties of IRS-lipase

#### The rapeseed lipase is a glycoprotein

Generally plant proteins are glycosylated and characterized by the presence of a glycane structure related to the asparagines residues. 20  $\mu$ l of the IRSL was separated by SDS-PAGE, electrotransferred in a nitrocellulose membrane. After that we incubated it firstly with the biotine conjugated hydrazide and secondly with alkaline phosphatase conjugated avidine. A positive reaction was obtained showing that the rapeseed lipase is a glycolprotein (data not shown).

### Effect of antibodies anti-*E. characias* lipase on rapeseed lipase activity


In order to study the effect of the polyclonal anti-ECL antibodies (made by the laboratory of LLE-CNRS Marseille-France) on rapeseed lipase activity, we pre-incubated 20  $\mu$ l of the IRSL with an equal volume of these antibodies. After 1 h of pre-incubation at 37°C, a centrifugation at 10,000 x g for 2 min was performed and the supernatant (S10) was recuperated to measure the residual lipase activity using two different methods (radiolabeled and fluorimetric). Two control tests were carried out using pre-incubated IRS-lipase with a pre-immune rabbit serum and a heat-treated IRS-lipase for 5 min at 90°C. In these cases, no change in the lipase activity amount was observed, but a very low residual lipase activity was found in the recuperated supernatant (S10) with a 97 % inhibition rate (Figure 7). These results show an evident immunochemical cross-reactivity between ECL and rapeseed lipase, suggesting that these antibodies recognized the native IRSL and are bound to some residues located in the catalytic site or are related to it.

### Evidence of a cross-reactivity with ECL


To confirm the last results, we assessed the immunoblotting technique, using the anti-EC-lipase polyclonal immune serum at a 1: 500 dilution (Belguith et al., 2001). The immunoblot presented in Figure 8 demonstrates clearly that the anti-ECL antibodies cross-react with IRS-lipase. This result suggests an antigenic relationship between the two lipases.

Moulin et al. (1994) had demonstrated that the EC-






**Figure 4.** Lipolytic activity on radio-labeled triolein substrate. Test of the IRS-lipase activity: 20  $\mu$ l of native and heat treated ISR-lipase were incubated at pH 7.5 and 37°C in presence of 10  $\mu$ l of radiolabeled triolein.



**Figure 5.** Kinetics of hydrolysis of naturally fluorescent TAGs from *Parinari glaberrimum* upon incubation with 20  $\mu$ l of IRS-lipase incubated in 1 ml of the standard reaction medium (pH 8), containing 16  $\mu$ g of fluorescent TAGs, 50 mM Tris, 100 mM NaTDC, 6 mM CaCl<sub>2</sub>, and 0.001% BTH.

residue in the catalytic site of IRS-lipase and that it is a serine triacylglycerol acyl hydrolase as the major of lipases.




**Figure 6.** Electrophoregramme of esterases specific revelation. 6 to 12  $\mu$ g of proteins were separated by native PAGE. The gel was submitted after that to an esterase specific revelation, using  $\alpha$ -naphthol acetate as a substrate. Lane 1, 12  $\mu$ g of ISR-lipase; lanes 2 and 3: 6 and 12  $\mu$ g of the rapeseed crude extract (S10).

### Mass spectrometry analysis

Mass spectrometry is considered as an indispensable technique to analyze and characterize the primary structure of proteins. This technique allows us to evaluate the purity and to get the exact molecular weight of proteins. The MALDI-ToF specter of the rapeseed lipase shows a single peak with a MW of 38,0387 (Figure 10). This MW is very close to that of EC-lipase described by Moulin et al. (1994).

### DISCUSSION

We have performed a rapid, high yielding, inexpensive and reproducible procedure for rapeseed lipase purification; it seems to be efficient to purify low abundant plant lipases since the overall obtained yield in our case is about 35.19% with a 1950-fold purification factor. The purity of the immunopurified lipase was firstly tested by SDS-PAGE, a single band was stained in the polyacrylamide gel with a MW of about 39 kDa, secondly by HPLC where a single and symmetric peak was obtained. Finally, using mass spectrometry analysis, we determined that the rapeseed lipase is a monomeric protein having an exact MW of 38.0387 kDa, molecular weight identical to that of EC-lipase. Our results confirm that the immunopurification is one of the most selective and powerful affinity techniques for protein purification leading to purification of 1000- to 10 000-fold in a single step (Harlow and Lane, 1988). Very few examples of immunopurification of lipase are found in the literature.




**Figure 7.** Immunoprecipitation of IRS-lipase and HPL by antibodies anti-EC-A lipase. Residual lipase activity was measured using fluorimetric method (a) and the radiolabeled method (b).

This method is recommended to be used as a first step on purification procedures leading to high recovery yields.

It can be considered as a highly selective purification technique that requires less number of steps leading to






**Figure 8.** Immunoblot analysis of the IRS-lipase separated by SDS-PAGE, showing a cross reactivity with *Euphorbia characias* lipase. The western blot was probed with an anti-EC-lipase polyclonal immune serum at a 1:500 dilution. Lane 1, 15 µg of the IRS-lipase; 2, Protein markers.

an efficient separation and high recovery purification. Immunopurification techniques can be used in industries, because they are rapid, efficient, inexpensive, powerful affinity techniques for protein purification and amenable to large-scale operations.

We also demonstrate that the rapeseed lipase is glycosylated and it was inhibited by THL. It is worth noting that the total inactivation of the rapeseed lipase by THL supports the existence of an activated serine in the catalytic site and that is a true serine triacyl glycerol acylhydrolase.

This result suggests a structure similarities and a chemical relationship between lipases from different organisms. This suggestion is consistent with the pre-



**Figure 9.** Effect of the addition of tertahydrolipstain (THL) on the IRS-lipase activity. 5 µl of THL was added in the cuvette assay. Lipase activity was measured using the fluorimetric method.

sence in the primary rapeseed lipase amino acid sequence of the consensus motif Gly-X-Ser-X-Gly

Nearly, all lipases that undergo interfacial activation are somehow akin to a molecular structure covering the catalytic site. Using anti-EC lipase antibodies, we demonstrate an evident immunochemical cross reactivity between EC-lipase and IRS-lipase, reflecting a considerable degree of structure homology between these enzymes. Furthermore, these antibodies were able to total inhibit rapeseed lipase activity suggesting that they bind to some residue located in the catalytic site.

The different methods used to test lipase activity confirm that the immunopurified rapeseed lipase is a true lipase of germination and not an esterase. A short sequence of the IRS-lipase was obtained (D, S) INGGXATLPQX.

## ACKNOWLEDGMENTS

The authors gratefully acknowledge financial support from the Tunisian Ministry of Higher Education, Scientific Research and Technology and the University of 7 November at Carthage.




Figure 10. The Maldi-Tof spectrum of the rapeseed lipase.

#### REFERENCES

Aires-Barros MR, Cabral JMS (1991). Purification and kinetic studies of lipases using reversed micellar systems. *in*: Alberghina L, Schmid RD and Verger R (Eds.), *Lipases: Structure, Mechanism and Genetic Engineering*. GBF Monographs, Braunschweig, Germany 16: 407-416.

Akanuma Y, Kuzuya T, Hayashi M, Ide T, Kuzuya N (1970). Immuno-

logical reactivity of insulin to sepharose coupled with insulin-antibody: its used for extraction of insulin from serum. *Biochem. Biophys. Res. Com.* 38: 947-953.

Beisson F, Ferte N, Nari J, Noat G, Arondel V, Verger R (1999). Use of naturally fluorescent triacylglycerols from *Parinari glaberrimum* to detect low lipase activities from *Arabidopsis thaliana* seedlings. *J. Lip. Res.* 40: 2313-2321.

Belguith H, Jridi T, Ben Hamida J (1999). Subcellular repartition and

- characterization of lipase during rapeseeds (*Brassica napus* L.) germination. *Ann. de l'I.N.R.A.T.* 72: 197-211.
- Belguith H, Khodjet El, Kill H, Fattouch S, Jridi T, Ben Hamida J (2001). Contribution of blotting techniques to the study of rapeseeds (*Brassica napus* L.) lipases. *Electrophoresis*, 22: 18-22.
- Ben Hamida J (1986). Etude des lipides, des lipases et d'un inhibiteur des lipases, chez le tournesol (*Helianthus annuus* L.) Thèse de Doctorat d'Etat Es-Sciences Naturelles. FST.
- Ben Hamida J, Mazliak P (1985). Les lipases des graines oleagineuses. *Ann. Biol. T XXIV, Fasc. 3*: 201-232.
- Bradford MM (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principal of dye binding. *J. Biol. Chem.* 252: 248-254.
- Cambillau C, Bourne Y (1991). Crystallographic studies of the pancreatic lipase/colipase system. *in: Alberghina L, Schmid RD and Verger R (Eds.), Lipases: Structure, Mechanism and Genetic Engineering*. GBF Monographs, Braunschweig, Germany. 16: 31-34.
- Carriere F, Thirstrup K, Hjorth S, Boel E, Verger R, Thim L (1994). Structure-function relationships in naturally occurring mutants of pancreatic lipase. *Protein Eng.* 7: 563-569.
- Cheng CF, Bensadoun A, Bersot T, Hsu JST, Melford KH (1985). Purification and characterization of human lipoprotein lipase and hepatic triglyceride lipase. *J. Biol. Chem.* 260: 10720-10726.
- Diacio R (1986). Hill, H. and Durand, D. P. Purification of soybean mosaic virus by affinity chromatography using monoclonal antibodies. *J. Biol. Chem.* 261: 345-351.
- Duncombe WC (1962). The colorimetric determination of long chain fatty acids in the 0,05-0,5 mole range. *Biochem. J.* 83: p. 6.
- Gargouri Y, Moreau H, Verger R (1989). Gastric lipases: biochemical and physiological studies. *Biochem. Biophys. Acta* 1006: 255-271.
- Gotz F (1991). Staphylococcal lipases and phospholipases. *in: Alberghina L, Schmid RD and Verger R (Eds.), Lipases: Structure, Mechanism and Genetic Engineering*. GBF Monographs, Braunschweig, Germany, 16: 285-292.
- Gubernator K, Müller K, Winkler F (1991). The Structure of human pancreatic lipase suggests a locally inverted, trypsin-like mechanism. *in: Alberghina L, Schmid RD and Verger R (Eds.), Lipases: Structure, Mechanism and Genetic Engineering*. GBF Monographs, Braunschweig, Germany. 16: 9-16.
- Hadvary P, Sidler W, Meister W, Velter W, Wolf H (1991). Inhibition of pancreatic lipase in vitro by the covalent inhibitor tetrahydropyridin. *J. Bio. Chem.* 266: 2021-2027.
- Harlow E, Lane D (1988). *Antibodies* Cold Spring Harbour Publications, USA.
- Huang AHC (1993). Lipases. *in: Lipid Metabolism in plants*. Moore TS, (Eds). CRC Press Inc., Boca Raton. pp. 473-502.
- Isobe K, Nokihara K (1991). Physicochemical properties of mono- and diacylglycerol lipase from *Penicillium camembertii*. *in: Alberghina L, Schmid, R.D. and Verger, R. (Eds.), Lipases: Structure, Mechanism and Genetic Engineering*. GBF Monographs, Braunschweig, Germany. 16: 345-348.
- Jensen GL, Dagg B, Bensadoun A (1982). Triacylglycerol lipase, monoacylglycerol lipase and phospholipase activities of highly purified rat hepatic lipase. *Biochem. Biophys. Acta.* 710: 464-470.
- Kanaya S, Koyanagi T, Kanaya E (1998). An esterase from *Escherichia coli* with a sequence similarity to hormone-sensitive lipase. *Biochem. J.* 322: 75-80.
- Kang AS, Abott SJ, Harris N (1988). *Manipulating secondary metabolism in culture*, Cambridge University Press, Cambridge.
- Kenyon AJ, Gander JE, Lopez C, Good RA (1973). Isolation of Aleutian mink disease virus by affinity chromatography. *Sci.* 179: 187-189.
- Kundu M, Basu J, Guchhait M, Chakrabarti P (1987). Isolation and characterization of an extracellular lipase from the conidia of *Neurospora crassa*. *J. Gen. Microbiol.* 133: 149-153.
- Laemmli UK (1970). Clivage of structural proteins during the assembly of the head of bacteriophage T4. *Nat.* 227: 680-685.
- Lang S, Katsiwela E, Kleppe F, Wagner F (1991). Ustilago maydis lipolytic enzymes: Characterization and partial purification. *in: Alberghina L, Schmid RD and Verger R (Eds.), Lipases: Structure, Mechanism and Genetic Engineering*. GBF Monographs, Braunschweig, Germany. 16: 361-364.
- Lin YH, Haung ACH (1983). Lipase in lipid bodies of cotyledons of rape and mustard seedlings. *Arch. Biochem. Biophys.* 225: 360-369.
- Melchers F, Messer W (1970). The activation of mutant  $\alpha$ -galactosidase by specific antibodies. *Euro. J. Biochem.* 17: 267-272.
- Moulin A, Tiessere M, Bernard C, Pieroni G (1994). Lipases of the Euphorbaceae family: Purification of a lipase from *Euphorbia characias* latex and structure-function relationships with the B chain of ricin. *Biochem. J.* 301: 1132-1133.
- Muraoka T, Ando T, Okuda H (1982). Purification and properties of a novel lipase from *Staphylococcus aureus* 226. *J. Biochem.* 92: 1933-1939.
- Murphy RF, Buchanan KD, Elmore DT (1973). Isolation of glucagon-like immunoreactivity of gut by affinity chromatography on anti-glucagon antibodies coupled to sepharose 4B. *J. Biol. Chem.* 248: 118-127.
- Osterlund T, Danielsson B, Degerman E, Contreras JA, Edgren G, Davis RC, Schotz MC, Holm C (1996). Domain-structure analysis of recombinant rat hormone-sensitive lipase. *Biochem. J.* 319: 411-420.
- Saxena RK, Sheoran A, Giri B, Davidson W (2003). Purification strategies for microbial lipases. *J. Microbiol. Methods*, 52: 1-18.
- Sjoberg BM, Holmgren A (1973). Purification of thioredoxin from *Escherichia coli* and bacteriophage immunosorbent affinity chromatography. *Bioch. Biophys. Acta.* 315: 176-180.
- Sugihara A, Shimada Y, Tominaga Y (1988). Purification and characterization of *Aspergillus niger* lipase. *Agric. Biol. Chem.* 52: 1591-1592.
- Suzuki M, Yamamoto H, Mizugaki M (1986). Purification and general properties of a metal-insensitive lipase from *Rhizopus japonicus* NR 400. *J. Biol. Chem.* 261: 1207-1213.
- Tyski S, Hryniewicz W, Jeljaszewicz J (1983). Purification and some properties of the coccal extracellular lipase. *Biochem. Biophys. Acta.* 749: 312-317.
- Veeraragavan K, Gibbs BF (1989). Detection and partial purification of two lipases from *Candida rugosa*. *Biotechnol. Lett.* 11: 345-348.
- Verger R (1984). Pancreatic Lipases. In B. Borgström and H.L. Brockman (Eds.). *Lipases* Science Publishers, Amsterdam.
- Vicente MLC, Aires-Barros MR, Cabral JMS (1990). Purification of *Chromobacterium viscosum* lipases using reversed micelles. *Biotechnol. Techniques*, 4: 137-142.
- WeselaKe R, Jain JC (1992). Strategies in the purification of plant proteins. *Physiol. Plant.* 84: 301-309.