Full Length Research Paper

Monitoring of uncultured *Dunaliella* sp. in an Egyptian solar saltern field based on RuBisCO-encoding gene *cbbL*

Hosam Easa Elsaied

Genetics and Biotechnology Research Group, Department of Aquaculture, National Institute of Oceanography and Fisheries, NIOF, Cairo, Egypt.

Accepted 8 August, 2013

Culture-independent molecular approach was used to explore and evaluate the diversity of *Dunaliella* species living at the salt field Malahat El-Max Alexandria, Egypt. Bulk genomic DNA was extracted directly from the collected salt water samples. Specific PCR primers and methodology were designed to amplify the gene *cbbL*, which encodes the large subunit of the enzyme ribulose-1,5-bisphosphate carboxylase/oxygenase (RuBisCO, EC 4.1.1.39) of only *Dunaliella* species, from the extracted microbial metagenome. The 700 bp-PCR amplicons were cloned and *cbbL* clone library was constructed and analyzed by sequencing. Rarefaction curve was saturated at sequence analyses of 23 clones, obtaining 19 phylotypes of *Dunaliella cbbL*, representing the total composition of *Dunaliella* in the collected sample. All recorded phylotypes had the known deduced amino acid *cbbL* motive sequence and catalytic sites. Fingerprint sequence, characterizing *Dunaliella cbbL*, was recorded. The *cbbL* phylotypes were grouped into two distinct phylogenetic clusters. The cluster 1, consisting of 18 current *cbbL* phylotypes was rooted with a cluster containing *cbbL*s of *Dunaliella salina*, *Dunaliella bioculata*, *Dunaliella primolecta* and *Dunaliella tertiolecta*. The single phylotype, uncultured *Dunaliella* EIMax.3, forming cluster 2, showed a unique phylogenetic lineage in the evolution of *Dunaliella cbbL*. This study introduced the first functional gene markers for exploring *Dunaliella* species in salt waters without culture.

Key words: Uncultured *Dunaliella*, RuBisCO *cbbL*, solar saltern water, diversity.

INTRODUCTION

The genus *Dunaliella*, belonging to Chlorophyceae, has been represented by 27 known species, of which 23 are from salt water. Species of *Dunaliella* occur in freshwater, euryhaline habitats of all continents, oceans including the Dead Sea and even the salt lakes of the Antarctic (Tafreshi and Shariati, 2009). Thus, *Dunaliella* is a global genus of algae, thriving in habitats with wide ranges of salinity, pH, light intensity and temperature. The characterization of economically important algae in solar salterns has become a key focus for many research

groups all over the world (Chen and Jiang, 2009). *Dunaliella* species that are living in these fields, have an economic importance due to its highly production of anti-oxidants (Oren, 2010). In Egypt, there are several coastal solar salterns, where *Dunaliella* is living, mainly on and near to the Mediterranean coast. Studies on *Dunaliella* in Egyptian aquatic habitats have been focused on characterization, environmental stresses and biotechnological applications of isolated species, including *Dunaliella bardawil* (Adam, 1997), *Dunaliella salina* (Abd El-Baky

Figure 1. A map showing the sites of sampling marked by black circles.

et al., 2004; Dardir and Abd El-Tawab, 2005; Shafik, 2008), *Dunaliella parva* (Said, 2009) and *Dunaliella tertiolecta* (Tammam et al., 2011). Molecular phylogeny techniques have been applied to the taxonomic study of *Dunaliella* from 1999 onwards. These studies have encompassed the 18S rRNA genes and the internal transcribed spacer regions, ITS and have been based on gene sequence comparisons as well as on restriction fragment length polymorphism studies (Gomez and Gonzalez, 2001; Assunção et al., 2012). Little correlation was found between the molecular data and the traditional morphological-physiological attributes to delineate species within the genus *Dunaliella* (González et al., 1999, 2001).

On the basis of 18S rRNA gene sequencing, Olmos et al. (2000) could differentiate between *D. salina*, *D. parva* and *D. bardawil* as species containing one, two and three introns, respectively, within the 18S rRNA gene. Also, Hejazi et al. (2010) reported a new 18S rRNA gene arrangement in term of intron localization and nucleotide sequence in a *Dunaliella* isolated from Iranian salt lakes. However, molecular phylogenetic studies of *Dunaliella* in Egyptian solar salterns have not yet been done; remaining the diversity of *Dunaliella* species in these fields are poorly described. The gene, *cbbL* that encodes the large subunit of the enzyme RuBisCO, the key of Calvin cycle has been used as an efficient molecular tool for studying the functional diversity of several phytoplankton (Tabita, 1999; Yoon et al., 2001; Ghosh and Love, 2011). The RuBisCO large subunit is distributed in almost all phytoplankton, through five different forms: form IA, IB, IC, ID and II (Watson and Tabita, 1997; John et al., 2007). The halophylic *Dunaliella* species, the pink micro-algae, are differentiated from

other species in the class Chlorophyceae by harboring the RuBisCO form ID (Lin and Carpenter, 1997). This study developed a molecular approach for first monitoring of the diversity of *Dunaliella* based on PCR amplification, cloning and sequencing of the RuBisCO gene *cbbL* in metagenomic DNA extracted from the Saltern field, Malahat El-Max, Alexandria, Egypt. Statistical analysis was applied to determine the total composition of *Dunaliella cbbL* phylotypes and the phylogenetic tree was constructed showing the phylogenetic distribution of the recovered *Dunaliella cbbL* phylotypes.

MATERIALS AND METHODS

Sampling

Pink-color *Dunaliella*-containing surface waters, 5 L was collected from five stations, 1 L from each station, covering mostly the total area of the saltern field, Malahat El-Max, Alexandria, Egypt (Figure 1). Each liter of the collected water was filtered on a single cylindrical 0.2 µm filter membrane unit (type Sterivex-GS, Millipore Corp., USA). In order to remove the trapped pink carotenoid pigments, which may link with the extracted DNA and hinder PCR amplification, the filters were washed with 10 ml of 80% ethanol in sterile SET buffer (20% sucrose, 50 mM EDTA, 50 mM Tris-HCl, pH 7.6) several times. The inlet and outlet of the filters were capped, and the filters were stored at -30°C until processed.

Genomic DNA extraction and amplification of the *Dunaliella* RuBisCO gene *cbbL*

Bulk microbial DNA, metagenome, was extracted essentially within the Sterivex-GS filters housing according to the method of Somerville et al. (1989), with modifications. Cell lysis and proteolysis were carried out within the filter housing using 5M guanidine

Figure 2. Efficiency of the PCR primers ID 537F and ID 1212R for amplification of RuBisCO form ID from genomes of known *Dunaliella* species and other phytoplankton. 1: amplification from *Dunaliella bardawil* strain TS-1; 2: *Dunaliella bioculata* CCAP 19/4; 3: *Dunaliella parva* IAM C-527; 4: *Dunaliella salina* CCAP 19/3; 5: *Chlamydomonas reinhardtii*; 6: *Synechococcus* sp. strain WH7803; 7: *Rhodospirillum rubrum*; 8: *Chlorobium tepidum*. M: 100 bp-DNA ladder marker.

thiocyanate in SET buffer at 70°C for 20 min with shake. Crude, which contained high-molecular-weight nucleic acids, was then drawn off the filter and purified using high pure PCR template preparation kit, Roche, Catalog no. 11796828001, Mannheim, Germany. The purified metagenomic DNAs, extracted from all samples, were combined and run on 0.9% agarose gel electrophoresis followed by staining with ethidium bromide and UV visualization. A new PCR primer set, Duna-537f (5'-RTW AAA CCT AAA TTA GGT YTA TCW GGT-3') and Duna-1212r (5'-AAT AGT ACC ACC ACC RAA TTG-3') were designed from multiple alignments of *cbbL* sequences of *Dunaliella* species, incorporated to date in the DNA databases, including *D. salina* strain CCAP 19/18 (acc. no. AY531529, GQ250046), *D. acidophila* (HQ142901), *D. bioculata* (AB127991), *D. primolecta* strain TS-3 (AB127992), *D. tertiolecta* (AY882012), *D. bardawil* strain TS-1 (AB127990) and *D. parva* strain UTEX 1983 (AJ001877). The primers corresponded to positions 511 to 537 (forward primer) and 1191 to 1212 (reverse primer) of the *cbbL* gene of *Synechococcus* sp. (Shinozaki et al., 1983). PCR reaction mixture (50 µl), contained 10x EX buffer II (Mg²⁺ plus), 0.2 µM primer, 400 µM dNTP each, 2.25U Takara EX-Taq Polymerase (Takara, Japan) and 5 to 30 ng DNA template. The PCR mixture composition was performed with an initial denaturation step of 3 min at 95°C. The reaction was continued with 30 cycles of 1 min at 95°C, 2 min at 46°C, and 3 min at 72°C, with a final extension of 10 min at 72°C.

PCR products had 3'-A overhangs to facilitate TA-cloning into TOP10 *Escherichia coli* using a TOPO XL PCR-cloning kit according to the manufacturer's instructions (Catalog no. K4750-20, Invitrogen Life Technologies, Carlsbad, CA, USA). Only cells containing XL-TOPO vector with the insert were competent to grow with kanamycin. Colonies of these cells were screened directly by sequencing using vector primers T7 and an ABI 3730xl 96-capillary DNA analyzer (Applied Biosystems, Foster City, CA, USA).

Sequence and bioinformatics analyses of *cbbL* genes

Sequencing results were introduced to FASTA (<http://fasta.ddbj.nig.ac.jp/top-e.html>), to determine their similarity to

known *Dunaliella cbbL* sequences, deposited in DNA database. Nucleotide sequences for *cbbL* open reading frames (ORFs) were submitted to Transeq, (<http://www.ebi.ac.uk/emboss/transeq/>), to obtain the inferred amino-acid sequences. The correct *cbbL* ORFs were identified from the presence of diagnostic *cbbL* CO₂ fixation motifs. Grouping of sequences into phylotypes was done based on genetic distances using the Mothur software package V.1.7.2 (Schloss et al., 2009), where *cbbL* sequences, having 100% deduced amino acid identities, were grouped into a single phylotype. The expected total composition of *Dunaliella cbbL* was determined by rarefaction analyses in the Mothur software package. The phylogenetic analysis was done based on deduced amino acid and nucleotide sequences, and included the current *cbbL* sequences and their *Dunaliella cbbL* homologues, with significant homologies, *e*-value < 0.001, from the DNA database bank. The process was performed by applying the neighbor-joining algorithm and drawing the trees using the MEGA 3.1 software (<http://www.megasoftware.net/>). The branching patterns of the constructed phylogenetic tree were confirmed by reconstruction of the phylogenies using two other methods of analysis, namely maximum parsimony and maximum-likelihood, contained within the Phylip package.

The sequences of recovered *Dunaliella cbbL* were deposited in DNA data base bank under accession numbers from AB781569 to AB781587.

RESULTS AND DISCUSSION

Efficiency of primers for PCR amplification of *Dunaliella cbbL*

This work aimed to cover the molecular diversity of almost all *Dunaliella* species in the collected saline water sample. In order to achieve this target, several considerations were involved in the design of an efficient primer set for amplification of *Dunaliella cbbL*. The primers must be designed from *cbbL* region, which encodes CO₂ catalytic site of *Dunaliella* RuBisCO form ID. Hence, the codon region of the *cbbL* catalytic site amino acid group T(V) KPKLG (Newman and Gutteridge, 1993) represented specific sequence for the design of the primer Duna-537f. The primer Duna-1212r was designed from the codons that encode the catalytic amino acids QFGGGTI, characterizing the *Dunaliella* RuBisCO ID *cbbL*. These primer sequences were differed from other *cbbL* sequences, located in other autotrophic algae, in several nucleotides, sufficient to make each of the primers binds only with *Dunaliella cbbL*. This was clear in the positive PCR amplifications of *Dunaliella cbbL*, while negative amplifications were noticed for *cbbL* from other phytoplankton species (Figure 2). These primer sets provided *cbbL* product size, 700 bp, longer than that produced by Paul et al. (2000). This gave an extended view to screen the diversity of *Dunaliella cbbL*.

In order to reduce the possibility of bias in amplification of *Dunaliella cbbL*, the PCR was tested using a range from 26 to 30 cycles, and then the amplicons were combined for cloning (Suzuki and Giovannoni, 1996). This was clear in the rarefaction curves, for the recovered *Dunaliella cbbL*, which showed saturation after analysis of 23 clones, yielding 19 phylotypes (Supplementary

		K	K		G	X	D	F	X	K	
Uncultured <i>Dunaliella</i> ElMax-1	GTAA	AAA	CCT	AAA	TTAGGTTTATCTGGT	27	TGGATTAGACTTTACCAAA	84			
Uncultured <i>Dunaliella</i> ElMax-2					C.....	27		84			
Uncultured <i>Dunaliella</i> ElMax-3					T.....	27	G..T.....T..C..T..G	84			
Uncultured <i>Dunaliella</i> ElMax-4						27	T..A.....C..T..C..A	84			
Uncultured <i>Dunaliella</i> ElMax-5						27		G..G	84		
Uncultured <i>Dunaliella</i> ElMax-6						27		C..A	84		
Uncultured <i>Dunaliella</i> ElMax-7					C.....	27			84		
Uncultured <i>Dunaliella</i> ElMax-8					T.....	27		G..G	84		
Uncultured <i>Dunaliella</i> ElMax-9						27			84		
Uncultured <i>Dunaliella</i> ElMax-10					C.....	27			84		
Uncultured <i>Dunaliella</i> ElMax-11					T.....	27		C...	84		
Uncultured <i>Dunaliella</i> ElMax-12					C.....	27		G...	84		
Uncultured <i>Dunaliella</i> ElMax-13						27		G.G	84		
Uncultured <i>Dunaliella</i> ElMax-14						27		CA.A	84		
Uncultured <i>Dunaliella</i> ElMax-15					T.....	27			84		
Uncultured <i>Dunaliella</i> ElMax-16						27			84		
Uncultured <i>Dunaliella</i> ElMax-17						27			84		
Uncultured <i>Dunaliella</i> ElMax-18						27		T..	84		
Uncultured <i>Dunaliella</i> ElMax-19						27		A..	84		
<i>Dunaliella bioculata</i>	A.T....	A.....			A... 534			G..G	591		
<i>Dunaliella primolecta</i>					534				591		
<i>Dunaliella tertiolecta</i>					547				603		
<i>Dunaliella bardawil</i>					531			C..A	588		
<i>Dunaliella salina</i>					504	...T.....		G..G	561		
<i>Dunaliella parva</i>					516			C..T...	573		
<i>Chlamydomonas reinhardtii</i>	..C...A.T...	C.C...	C.T.....	546	C.T.....	A	603				
<i>Synechococcus</i> sp.	G..G..G.....	C..GAGC..C	522	C.....G.....	C..G	579					
<i>Rhodospirillum rubrum</i>					C.TCCC 519	...--C.....		T....	573		

Figure 3. Multiple alignments, within the amplification size, 700 bp, of recorded uncultured *Dunaliella* ElMax.1 to 19, and *cbbL*s from other known phytoplankton recorded in database. The nucleotide alignments are only in the sites of RuBisCO *cbbL* motive sequence, marked by shading, and catalytic amino acids, bold face. Alignments in the box represent the finger print sequence of *Dunaliella cbbL*. Species used in the alignments had accession numbers listed in Figure 4.

Figure 1). This result may reflect the efficiency of the designed primers and PCR methodology in covering the total composition of *Dunaliella* in the studied water sample.

The recovered *cbbL* phylotypes were belonged to uncultured *Dunaliella* species

All recovered *Dunaliella cbbL* phylotypes were from metagenomic DNAs and, consequently, the source *Dunaliella* species were uncultured and had no common synonyms. Hence, nomenclatures were given to these phylotypes, whereby each phylotype was the descriptor of uncultured *Dunaliella* followed by clone name, which included sampling site name and clone number, for example, uncultured *Dunaliella* ElMax.1 and so on. *cbbL* phylotypes have been recovered from uncultured autotrophic microbial communities, which have no obvious species identifications (Guo et al., 2013).

The recovered phylotypes had the known *cbbL* deduced amino acid catalytic sites, beside specific sequence characterizing *Dunaliella cbbL*

Alignments of 702 bp-*cbbL* phylotypes, which yielded 234 deduced amino acids, were done with those recovered from known *Dunaliella* species (Figure 3). The positions of catalytic sites were corresponded to the *cbbL* of *Synechococcus* sp. (Newman and Gutteridge, 1993). All the current *Dunaliella cbbL* phylotypes possessed the characteristic RuBisCO motif sequence, GXDFXKXDE. Each of the amino acids GDFKE in the motif sequence was represented by two codons, for example, the amino acid G was represented by the codon "GGT" in the phylotype uncultured *Dunaliella* ElMax.3 and the codon "GGA" in other recovered phylotypes (Figure 3). The recovered *cbbL* phylotypes contained the known *cbbL* catalytic deduced amino acid sequences. The nucleotide positions from 714 to 741, of *cbbL* of *Synechococcus* sp. were found to contain the finger print sequence, 5'-

	K	S	G	G	G
Uncultured <i>Dunaliella</i> ElMax-1	AAA 483	TCTGGTGGT 624	GGTGGT 693		
Uncultured <i>Dunaliella</i> ElMax-2	... 483	 624	 693		
Uncultured <i>Dunaliella</i> ElMax-3	... 483	 624	 693		
Uncultured <i>Dunaliella</i> ElMax-4	... 483	 624	 693		
Uncultured <i>Dunaliella</i> ElMax-5	... 483	 624	 693		
Uncultured <i>Dunaliella</i> ElMax-6	... 483	 624	 693		
Uncultured <i>Dunaliella</i> ElMax-7	... 483	 624	 693		
Uncultured <i>Dunaliella</i> ElMax-8	... 483	 624	 693		
Uncultured <i>Dunaliella</i> ElMax-9	... 483	 624	 693		
Uncultured <i>Dunaliella</i> ElMax-10	... 483	 624	 693		
Uncultured <i>Dunaliella</i> ElMax-11	... 483	 624	 693		
Uncultured <i>Dunaliella</i> ElMax-12	... 483	 624	 693		
Uncultured <i>Dunaliella</i> ElMax-13	... 483	 624	 693		
Uncultured <i>Dunaliella</i> ElMax-14	... 483	 624	 693		
Uncultured <i>Dunaliella</i> ElMax-15	... 483	 624	 693		
Uncultured <i>Dunaliella</i> ElMax-16	... 483	 624	 693		
Uncultured <i>Dunaliella</i> ElMax-17	... 483	 624	 693		
Uncultured <i>Dunaliella</i> ElMax-18	... 483	 624	 693		
Uncultured <i>Dunaliella</i> ElMax-19	... 483	 624	 693		
<i>D. bioculata</i>	... 990	 1131	 1200		
<i>D. primolecta</i>	... 990	 1131	 1200		
<i>D. tertiolecta</i>	... 1002	 1143	 1212		
<i>D. bardawil</i>	... 987	 1128	 1197		
<i>D. salina</i>	... 960	 1101	 1170		
<i>D. parva</i>	... 972	 1111	-----		
<i>C. reinhardtii</i>	... 1002	..A..C... 1143	 1212		
<i>Synechococcus</i> sp.	..G 978	..C..... 1119	 1189		
<i>R. rubrum</i>	... 980	AG.....C 1110	..C..C 1182		

Figure 3. Contd.

GCTGAAGGAATGCTTCAACGTGCACAA-3', of the *Dunaliella cbbL* (Figure 3). Consequently, the amino acids AGLQQ (Figure 3), encoded by this finger print nucleotide sequence, were considered as *Dunaliella* RuBisCO key amino acids, which differentiate it from those of in other phytoplankton. *cbbL* finger print sequences have been recorded for several autotrophic bacterial endosymbionts and used as nucleotide probes for *in situ* localization of the corresponding *cbbL* gene in the host tissues (Elsaied et al., 2002, 2006; Kimura et al., 2003).

On the other hand, red-like RuBisCO *cbbL* marker sequence has been used as a TaqMan probe for detection and quantification of soil autotrophic bacteria (Selesi et al., 2007). Hence, this current *Dunaliella cbbL* finger print sequence may be used as a biomarker for detection of uncultured *Dunaliella* by probing and hybridization techniques.

The recovered *Dunaliella cbbL* phylotypes were phylogenetically unique

Phylogenetic tree of *cbbL* was constructed in order to demonstrate the classification of the recovered *Dunaliella cbbL* (Figure 4). The nineteen *cbbL* phylotypes were distributed in two unique phylogenetic clusters. The cluster 1 harbored 18 *cbbL* phylotypes, which showed an average value of 95%_{ND} (nucleotide identity), that is, 97%_{aa} (amino acid identity) with each others. On the other hand, the *cbbL* phylotypes, constituting cluster 1, were distinct from other known *Dunaliella* species located in the same *Dunaliella* phylogenetic branch by showing homologies of 89 to 90%_{ND} (92 to 93%_{aa}) (Figure 4). The uncultured *Dunaliella* ElMax.3, forming the cluster 2, had a unique *cbbL* phylogenetic lineage. It showed average identity of 87%_{ND} (89%_{aa}) with other current *Dunaliella*

Figure 4. A tree shows the phylogenetic relationship between the current uncultured *Dunaliella* clones, EIMax.1 to 19, marked with bold face, and other *cbbL* sequences from *Dunaliella* species available in database, beside *cbbL* sequences from other phytoplankton, as out groups. The consensus tree was constructed by neighbour-joining analysis and confirmed by maximum-parsimony and maximum-likelihood methods. Bootstrap values of >50% are indicated at the nodes. The bar represents 0.1 changes per amino acid or nucleotide.

	K	S	G	G	G
D-1	AAA 483	TCTGGTGGT 624	GGTGGT 693		
D-2	... 483	 624	 693		
D-3	... 483	 624	 693		
D-4	... 483	 624	 693		
D-5	... 483	 624	 693		
D-6	... 483	 624	 693		
D-7	... 483	 624	 693		
D-8	... 483	 624	 693		
D-9	... 483	 624	 693		
D-10	... 483	 624	 693		
D-11	... 483	 624	 693		
D-12	... 483	 624	 693		
D-13	... 483	 624	 693		
D-14	... 483	 624	 693		
D-15	... 483	 624	 693		
D-16	... 483	 624	 693		
D-17	... 483	 624	 693		
D-18	... 483	 624	 693		
D-19	... 483	 624	 693		
<i>D. bioculata</i>	... 990	 1131	 1200		
<i>D. primolecta</i>	... 990	 1131	 1200		
<i>D. tertiolecta</i>	... 1002	 1143	 1212		
<i>D. bardawil</i>	... 987	 1128	 1197		
<i>D. salina</i>	... 960	 1101	 1170		
<i>D. parva</i>	... 972	 1111	-----		
<i>C. reinhardtii</i>	... 1002	..A..C... 1143	 1212		
<i>Synechococcus</i> sp.	..G 978	..C..... 1119	 1189		
<i>R. rubrum</i>	... 980	AG.....C 1110	..C..C 1182		

Figure 4. Contd.

cbbL phylotypes and those of known *Dunaliella* species.

These results may indicate that dynamically studied solar saltern field may select for different and diverse uncultured *Dunaliella* lineages, which added to *Dunaliella* phylogeny. In conclusion, this study could monitor uncultured *Dunaliella* based on specific *cbbL*-PCR approach. This strategy will help in finding diverse *Dunaliella* species yet to be discovered and open a new window to study the phylogeny of uncultured *Dunaliella* in aquatic environments.

ACKNOWLEDGEMENTS

We wish to express our deep gratitude to Dr. Akihiko Maruyama, National Institute of Advanced Industrial Science and Technology, AIST, Tsukuba Science city, Japan, for providing the genomes of *Dunaliella* species. This research was supported by the Grant in-Aid for Scientific Research, National Institute of Oceanography, Egypt.

REFERENCES

- Abd El-Baky H, El Baz F, El-Baroty G (2004). Production of Antioxidant by the Green Alga *Dunaliella salina*. Int. J. Agric. Biol. 6: 49–57.
- Adam MS (1997). Metabolic response of the halotolerant green alga *Dunaliella bardawil* to nitrogen: Phosphorus ratios in batch culture. Folia Microbiol. 42: 357–360.
- Assunção P, Jaén-Molina R, Caujapé-Castells J, de la Jara A, Carmona L, Freijanes K, Mendoza H (2012). Molecular taxonomy of *Dunaliella* (Chlorophyceae), with a special focus on *D. salina*: ITS2 sequences revisited with an extensive geographical sampling. Aquat. Biosyst. 8: 2.
- Chen H, Jiang JG (2009). Osmotic responses of *Dunaliella* to the changes of salinity. J. Cell Physiol. 219: 251–258.
- Dardir AA, Abd EL-Tawab A (2005). Usage lake quaroun water to cultivation of *Dunaliella Salina* as biotechnology in Fayoum, Egypt. Egypt J. Aquat. Res. 31: 341–351.
- Elsaied H, Kaneko R, Naganuma T (2006). Molecular Characterization of a deep-Sea methanotrophic mussel symbiont that carries a RuBisCO gene. Mar. Biotechnol. 8: 511–520.
- Elsaied H, Kimura H, Naganuma T (2002). Molecular characterization and endosymbiotic localization of the gene encoding ribulose-1,5-bisphosphate carboxylase/oxygenase (RuBisCO) form II in the deep-sea vestimentiferan trophosome. Microbiology 148: 1947–1957.
- Ghosh S, Love N (2011). Application of *rbcl* based molecular diversity analysis to algae in wastewater treatment plants. Bioresour. Technol.

- 102: 3619–3622.
- Gomez PI, Gonzalez MA (2001). Genetic polymorphism in eight Chilean strains of the carotenogenic microalga *Dunaliella salina* Teodoresco (Chlorophyta). *Biol. Res.* 34: 23–30.
- González MA, Coleman AW, Gómez PI, Montoya R (2001). Phylogenetic relationship among various strains of *Dunaliella* (Chlorophyceae) based on nuclear ITS rDNA sequences. *J. Phycol.* 37: 604–611.
- González MA, Gómez PI, Montoya R (1999). Comparison of PCR-RFLP analysis of the ITS region with morphological criteria of various strains of *Dunaliella*. *J. Appl. Phycol.* 10: 573–580.
- Guo X, Yin H, Cong J, Dai Z, Liang Y, Liu X (2013). RubisCO gene clusters found in a metagenome microarray from acid mine drainage. *Appl. Environ. Microb.* 79: 2019–2026.
- Hejazi MA, Barzegari A, Gharajeh NH, Hejazi MS (2010). Introduction of a novel 18S rDNA gene arrangement along with distinct ITS region in the saline water microalga *Dunaliella*. *Saline Systems* 6:4.
- John DE, Wang ZA, Liu X, Byrne RH, Corredor JE, López JM, Cabrera A, Bronk DA, Tabita FR, Paul JH (2007). Phytoplankton carbon fixation gene (RuBisCO) transcripts and air-sea CO₂ flux in the Mississippi River plume. *ISME J.* 1: 517–531.
- Kimura H, Sato M, Sasayama Y, Naganuma T (2003). Molecular characterization and *in situ* localization of endosymbiotic 16S Ribosomal RNA and RuBisCO genes in the pogonophoran tissue. *Mar. Biotechnol.* 5: 261–269.
- Lin S, Carpenter, E (1997). Rubisco of *Dunaliella tertiolecta* is redistributed between the pyrenoid and the stroma as a light /shade response. *Mar. Biol.* 127: 521–529.
- Newman J, Gutteridge S (1993). The X-ray structure of *Synechococcus* PCC6301 ribulose-bisphosphate carboxylase/oxygenase-activated quaternary complex at 2.2-Å resolution. *J. Biol. Chem.* 268: 25876–25886.
- Olmos J, Paniagua J, Contreras R (2000). Molecular identification of *Dunaliella* sp. utilizing the 18S rDNA gene. *Lett. Appl. Microbiol.* 30: 80–84.
- Oren A (2010). Industrial and environmental applications of halophilic microorganisms. *Environ. Technol.* 31: 825–834.
- Paul JH, Alfrieder A, Wawrik B (2000). Micro- and macrodiversity in rbcL sequences in ambient phytoplankton populations from the southeastern Gulf of Mexico. *Mar. Ecol. Prog.* 198: 9–18.
- Said HA (2009). Changes in levels of cellular constituents of *Dunaliella parva* associated with inorganic phosphate depletion. *Middle East J. Sci. Res.* 4: 94–99.
- Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB *et al.* (2009). Introducing Mothur: open source, platform-independent, community-supported software for describing and comparing microbial communities. *Appl. Environ. Microb.* 75: 7537–7541.
- Selesi D, Pattis I, Schmid M, Kandeler E, Hartmann, A (2007). Quantification of bacterial RubisCO genes in soils by cbbL targeted real-time PCR. *J. Microbiol. Methods* 69: 497–503.
- Shafik M (2008). Phytoremediation of some heavy metals by *Dunaliella salina*. *Glob. J. Environ. Res.* 2: 01–11.
- Shinozaki K, Yamada C, Takahata N, Sugiura M (1983). Molecular cloning and sequence analysis of the cyanobacterial gene for the large subunit of ribulose-1,5-bisphosphate carboxylase/oxygenase. *Proc. Natl. Acad. Sci. USA* 80: 4050–4054.
- Somerville C, Knight IT, Straube WL, Colwell R (1989). Simple, rapid method for direct isolation of nucleic acids from aquatic environments. *Appl. Environ. Microb.* 55: 548–554.
- Suzuki M, Giovannoni S (1996). Bias caused by template annealing in the amplification of mixtures of 16S rRNA genes by PCR. *Appl. Environ. Microb.* 62: 625–630.
- Tabita F (1999). Microbial ribulose 1,5-bisphosphate carboxylase/oxygenase: a different perspective. *Photosyn. Res.* 60: 1–28.
- Tafreshi A, Shariati M (2009). *Dunaliella* biotechnology: methods and applications. *J. Appl. Microb.* 107: 14–35.
- Tammam A, Fakhry E, El-Sheekh M (2011). Effect of salt stress on antioxidant system and the metabolism of the reactive oxygen species in *Dunaliella salina* and *Dunaliella tertiolecta*. *Afr. J. Biotechnol.* 10: 3795–3808.
- Watson G, Tabita F (1997). Microbial ribulose 1,5-bisphosphate carboxylase/oxygenase: a molecule for phylogenetic and enzyme-logical investigation. *FEMS Microbiol. Lett.* 146: 13–22.
- Yoon HS, Lee J, Boo S, Bhattacharya D (2001). Phylogeny of *Alariaceae*, *Laminariaceae*, and *Lessoniaceae* (Phaeophyceae) based on plastid-encoded RuBisCO spacer and nuclear-Encoded ITS sequence comparisons. *Mol. Phylo. Evol.* 21: 231–243.

Supplementary Figure 1. Rarefaction curves for the expected number of *Dunaliella cbbL* gene phylotypes.