

## Review

# Synthetic seeds: A review in agriculture and forestry

M. Chandrasekhara Reddy<sup>1</sup>, K. Sri Rama Murthy<sup>1\*</sup> and T. Pullaiah<sup>2</sup>

<sup>1</sup>Department of Botany and Biotechnology, School of Conservation Biology and Plant Biotechnology, Montessori Mahila Kalasala, Vijayawada - 520 010 Andhra Pradesh, India.

<sup>2</sup>Department of Botany, Sri Krishnadevaraya University, Anantapur - 515 055, Andhra Pradesh, India.

Accepted 15 June, 2012

**Production of synthetic seeds has unraveled new vistas in *in vitro* plant propagation technology, because it offers many useful advantages on a commercial scale for the propagation of a variety of crop plants. These tools provide methods for production of synthetic seeds for conversion into plantlets under *in vitro* and *in vivo* conditions. This technology is useful for multiplying and conserving the elite agricultural and endangered medicinal plant species, which are difficult to regenerate through conventional methods and natural seeds. The synthetic seed technology was developed in different economically important plant species such as vegetable crops, forage legumes, industrially important crops, cereals, spices, plantation crops, fruit crops, ornamental plants, orchids, medicinal plants and wood yielding forest trees etc. All these aspects are presented in this review.**

**Key words:** Synthetic seeds, *in vitro*, *in vivo* plant propagation.

## INTRODUCTION

Nowadays, artificial seed technology is one of the most important tools to breeders and scientists of plant tissue culture. It has offered powerful advantages for large scale mass propagation of elite plant species. In general, synthetic seeds are defined as artificially encapsulated somatic embryos, shoot tips, axillary buds or any other meristematic tissue, used for sowing as a seeds and possess the ability to convert into whole plant under *in vitro* and *in vivo* conditions and keep its potential also after storage (Capuano et al., 1998). The somatic embryo can be encapsulated, handled and used like a natural seed was first suggested by Murashige (1977) and efforts to engineer them into synthetic seed have been ongoing ever since Kitto and Janick (1982), Gray (1987).

Bapat et al. (1987) proposed the encapsulation of shoot tip in *Morus indica*; this application has made the concept of synthetic seed set free from its bonds to somatic embryos and broaden the technology to the encapsulation of various *in vitro* derived propagules. An implementation of artificial seed technology to somatic embryogenesis or the regeneration of embryos is based

on the vegetative tissues as an efficient technique that allows for mass propagation in a large scale production of selected genotype (Ara et al., 2000). The aim and scope for switching towards artificial seed technology was for the fact that the cost-effective mass propagation of elite plant genotypes will be promoted. There would also be a channel for new transgenic plants produced through biotechnological techniques to be transferred directly to the greenhouse or field.

The artificial seed technology has been applied to a number of plant species belonging to angiosperms. Present review aimed to give a brief description of methodology involved in synseed preparation, types of synthetic seeds, species in which this technique has been developed successfully.

## SYNTHETIC SEEDS PREPARATION

### Induction of somatic embryogenesis

One prerequisite for the application of synthetic seed technology in micropropagation is the production of high quality vigorous somatic embryos. Induction of somatic embryogenesis requires a change in the fate of a

\*Corresponding author. E-mail: [drksrmurthy@yahoo.com](mailto:drksrmurthy@yahoo.com).

vegetative (somatic) cell. In most cases, an inductive treatment is required to initiate cell division and establish a new polarity in the somatic cell. The inductive treatment is most commonly given by phytohormones. Among the plant growth regulators, auxins are known to be essential for the induction of somatic embryogenesis. In some plant species, 2,4-dichlorophenoxyacetic acid (2,4-D) is the most commonly used auxin. Other auxins may be required for certain species (Ammirato, 1983). Embryogenic cells are unique: superficially they resemble meristematic cells, though they generally are smaller, more isodiametric in shape, have larger, more densely staining nuclei and nucleoli, and have a denser cytoplasm (Williams and Maheswaran, 1986; Carman, 1990). In the carrot model described by Komamine et al. (1992), competent single cells formed embryogenic cell clusters (State 1) in the presence of auxin. These single cells are considered as predetermined for embryogenesis. During this phase, the cell clusters gained the ability to develop into embryos when auxin was removed from the medium, leading to the development of state 1 cell clusters (Nomura and Komamine, 1985; Komamine et al., 1992).

Once the induction of an embryogenic state is complete, the mechanisms of pattern formation that lead to the zygotic embryo are common passing through globular, heart-shaped and torpedo-shaped stages. Somatic embryo induction is usually promoted by auxins (Williams and Maheswaran, 1986). In some plant species, a combination of 2,4-D or  $\alpha$ -naphthalene acetic acid (NAA) with cytokinin was reported to be essential for the induction of somatic embryogenesis (Kao and Michayluk, 1981; Gingeas and Lineberger, 1989; Schuller et al., 1989). The process can be enhanced with the application of osmotic stress, manipulation of medium nutrients, reducing humidity etc.

### Suspension culture

The suspension culture is mainly used for large scale production of viable materials (somatic embryos) to produce synthetic seeds. In suspension, proembryogenic cell clusters form and can be separated from the single cells and the larger clumps of callus by sequentially sieving through nylon membranes of 500 and 224  $\mu$ m pore size. Generally the differentiation of somatic embryos in solid or liquid medium is highly asynchronous. Since synchronous embryo maturation is extremely important with regard to the artificial seed technology, several approaches have been developed to achieve it. This is achieved through selecting cells (Fujimura and Komamine, 1979) or pre-embryonic cell clusters of certain size (Altman et al., 1990) and manipulation of light and temperature (McKersie and Bowley, 1993), temporary starvation (Lee et al., 2001). Use of growth regulators to physiologically synchronize development

has proved most effective (Dobrev et al., 2002).

### Somatic embryo development

Even though somatic embryogenesis has been reported in several crop species, the quality of somatic embryos with regard to their conversion into plantlets has been very poor. This is because somatic embryos are normally incomplete in their development. Unlike seed embryos the somatic embryos actually do not go through the final phase of embryogenesis called 'embryo maturation'. For the maturation of somatic embryos however, transfer to media containing a low concentration or devoid of 2,4-D was essential (Ammirato, 1983; Cheema 1989; Van der Valk et al., 1989). The final stage of maturation is achieved by transferring the embryos to a medium containing abscisic acid (ABA). ABA prevents germination and promotes normal development of embryos by suppression of secondary embryogenesis and is reported to promote embryo maturation in several species (Ammirato, 1983). Senaratna et al. (1989, 1990) were able to confer desiccation tolerance of alfalfa somatic embryos by treating them with ABA. Buchheim et al. (1989) observed that the conversion of soybean somatic embryos was increased from 50 to 96% when matured in the presence of 10% sucrose. Other physical treatments including cold, heat, osmotic or nutrient stress can elicit a similar response, presumably because they stimulate the endogenous synthesis of ABA (McKersie et al., 1990).

### Encapsulation

Somatic embryogenesis is the only clonal propagation system economically viable. However somatic embryos would require mechanical strength for planting. It would be desirable to convert them into encapsulated units (synthetic seeds). Basic requirements for the encapsulation to form synthetic seeds are mentioned below.

### Explants used for encapsulation

Ever since synthetic seed technique was developed, the somatic embryos were largely used because they possess the radical and plumule that are able to develop into root and shoot in one step (Kitto and Janick, 1982, 1985 a, b; Kim and Janick, 1987, 1989, 1990; Janick et al., 1989; Redenbaugh et al., 1984; Redenbaugh et al., 1991b; Gray et al., 1991; Redenbaugh, 1993; McKersie and Bowley, 1993). Later on vegetative propagules e.g. shoot tips in *M. indica* (Bapat et al., 1987), axillary buds in *Camellia sinensis* (Mondel et al., 2002), calli in *Allium sativum* (Kim and Park, 2002), bulblets in *A. sativum* (Bekheet, 2006), cell aggregates derived from Horse-radish hairy roots (Repunte et al., 1995) and protocorm

like bodies in *Geodorum densiflorum* (Datta et al., 1999) were also used. In addition to the other *in vitro* derived meristematic tissues like microtubers, rhizomes and corms can also been used (Bapat and Minal, 2005).

### **Encapsulating agents**

Eight chemical compounds were tested for the production of synthetic seed coats, 'Polyox', water soluble resin was the most suitable agent for the encapsulation of somatic embryos (Kitto and Janick 1982, 1985c). However, Redenbaugh et al. (1984, 1986 and 1987) proposed that sodium alginate was the most suitable for the encapsulation of somatic embryos in few species such as alfalfa, celery, cauliflower and carrot. Sodium alginate was the most accepted hydro-gel and frequently used as a matrix for synthetic seeds, because of its low toxicity, low cost, quick gellation and bio compatibility characteristics (Saiprasad, 2001). In the previous studies several gelling agents, such as polyox, polyco 2133, agar, agarose, alginate, carboxy methylcellulose, carrageenan, guar gum, gelrite, tragacanth gum, sodium pectate ethylcellulose and nitrocellulose, polyacrylamide were tested for synthetic seed production (Ara et al., 2000; Saiprasad, 2001; Lambardi et al., 2006).

### **Synthetic endosperm**

It is believed that the encapsulated synthetic seeds should contain nutrients and plant growth regulators to serve as synthetic endosperm to the encapsulated propagules which results in increase in the efficiency of viability and germination of synthetic seeds. The quality of artificial seeds depends on the temporal, qualitative, quantitative supply of growth regulators and nutrients along with an optimal physical environment (Senaratna, 1992). Murashige and Skoog medium (MS) without hormones and MS + 6-benzyladenine (BA, 4.4  $\mu$ M) were used as artificial endosperm in *Morus* species (Pattnaik et al., 1995; Pattnaik and Chand, 2000). Refouvelet et al. (1998) used  $\frac{1}{2}$  MS + BA (5 mg/l) + NAA (0.01 mg/l) for encapsulation of axillary buds of *Syringa vulgaris*. Mariani (1992) reported that gibberellic acid ( $GA_3$ ) and sucrose showed negative effect on synthetic seed germination in eggplant. Many investigators reported that the addition of fungicides (Antonietta et al., 2007b), pesticides, fertilizers, microorganisms (Rhizobia), mycorrhiza fungus (Tan et al., 1998), marine cyanobacterial extracts (Wake et al., 1992) bactericides and activated charcoal (Ganapathi et al., 1992) to the encapsulation solution will protect the encapsulated propagules from micro-organisms, to reduce the release of toxic compounds and to enhance the germination capacity of seed.

### **Encapsulation procedure**

The most appropriate method for the preparation of

synthetic seeds was the hydro-gel encapsulation method developed by Redenbaugh et al. (1987). In this method, sodium alginate of different concentrations (2 to 5%) was prepared by mixing with calcium free liquid MS medium and then the explants were mixed with the solution. Explants were sucked with a pipette along with the sodium alginate solution and dropped into the calcium chloride solution, where the ion exchange reaction occurs and sodium ions were replaced by calcium ions forming calcium alginate beads. The whole process must be done under aseptic conditions. The size of the capsule depends on the inner diameter of the pipette nozzle. The shape and texture of the beads depends on the concentration of the sodium alginate, calcium chloride solutions and duration of complexion. Molle et al. (1993) suggested the use of a dual nozzle pipette in this embryos flow through the inner pipette and the alginate solution passes through the outer pipette. As a result, the embryos are positioned in the center of the capsule for better protection.

### **Desiccation**

Normally desiccation means to preserve by removing the moisture. Kitto and Janick (1982) for the first time encapsulated carrot somatic embryos followed by their desiccation. Desiccation tolerance is a characteristic of somatic embryos that must be induced and therefore requires a pretreatment with ABA or stress to elicit the desired response (Helal, 2011). Fujii et al. (1993) achieved the high conversion frequencies of the somatic embryos of celery by adding ABA and mannitol to the maturation medium. The survival of coated celery somatic embryos could be improved from 35 to 86% as reported by Kim and Janick (1987) and Janick et al. (1989). The vigor of the seedlings from dried somatic embryos was greater than those from embryos which had not been dried, but remained substantially lower than those from true seeds (Senaratna et al. 1990)

### **Germination and field planting**

The synthetic seeds have the possibility of being an alternative planting material meant for forestry sector in the future, especially for the highly demanded species (Asmah et al., 2011). Artificial seeds would allow direct planting of plant propagules into the greenhouse or field, thus bypassing many of the intermediate steps. Fujii et al. (1989) found that maturation of alfalfa somatic embryos with ABA showed high soil conversion of 48 to 64%. Addition of fungicide to the alginate beads prevents the contamination and increased the survival of mulberry buds when sown in soil (Bapat and Rao, 1990). Encapsulated somatic embryos (artificial seeds) and naked (uncoated) somatic embryos of alfalfa (*Medicago sativa*) were planted directly into the field to demonstrate

the feasibility of using artificial seeds for direct sowing (Fujii et al., 1992) and reported successful field planting of alfalfa artificial seeds derived from embryoids encapsulated in calcium alginate with 23% plant conversion. Preliminary experiments on calcium-sodium alginate encapsulation of somatic embryos from anther culture were performed, in order to evaluate the effect of this technique on the recovery of plantlets, and the applicability of the synthetic seed technology to *Citrus reticulata* (Antonietta et al., 1999). However sowing on soil mix medium did not result in satisfactory conversion. The behavior of sugarcane plants cv. CP-5243 derived from artificial seed compared with traditional and isolated bud methods was studied by Nieves et al. (2003). They reported that plants from artificial seed were taller and had a smaller diameter at eight months, but these differences disappeared at 12 months. With respect to sugar analysis and yield, no differences in all parameters evaluated were found between artificial seed-derived plants and plants derived from the two other methods. Successful field planting and conversion of artificial seeds were reported by several authors (Datta et al., 1999; Mandal et al., 2000; Anand and Bansal, 2002; Martin, 2003; Ikhtaq et al., 2010). The effective germination rate in *Acacia* hybrid (73.3 to 100%) and in *Pseudostellaria heterophylla* (80%) were reported by Asmah et al. (2011) and Ma et al. (2011).

### Types of synthetic seeds

According to the available literature, two types of synthetic seeds were developed, that is, desiccated and hydrated synthetic seeds (Bhojwani and Razdan, 2006). The desiccated synthetic seeds were first introduced from somatic embryos either naked or encapsulated in polyox followed by their desiccation (Kitto and Janick, 1982, 1985a, b).

Desiccation was achieved either slowly over a period of one or two weeks sequentially using chambers of decreasing relative humidity or rapidly by leaving the Petri dishes overnight on the bench of laminar airflow chamber (Ara et al., 2000). The hydrated synthetic seed technology was first produced by encapsulating hydrated somatic embryos of *M. sativa* (Redenbaugh et al., 1984). These hydrated synthetic seeds are used to produce plant species that their somatic embryos are recalcitrant and sensitive to desiccation. Hydrated artificial seeds are normally prepared by encapsulating the somatic embryos or other propagules in a hydro gel capsules. Several methods have been examined to produce hydrated artificial seeds of which calcium alginate encapsulation has been mostly used (Redenbaugh et al., 1993).

### Synthetic seeds

Synthetic seeds in plant propagation were successfully

studied in number of the plant species (Table 1). Plant propagation using artificial or synthetic seeds derived from somatic embryos or other vegetative propagules opens up new vistas in agriculture and forestry. Here, according to the convenience and importance of the plant species, artificial seed technology was developed and categorized into different groups.

### Vegetable crops

The production of synthetic seeds was by the encapsulation of multiple carrot somatic embryos (Kitto and Janick, 1982). In *Daucus carota*, production of desiccated synthetic seeds, hydrated synthetic seeds by using somatic embryos were reported (Kitto and Janick., 1982, 1985 a, b; Janick et al., 1989; Liu et al., 1992; Janick et al., 1993; Timbert et al., 1995; Timbert et al., 1996; Sakamoto et al., 1992 and Latif et al., 2007). 100% germination of encapsulated axillary buds by adding 0.5 mg/l NAA and 1.0% activated charcoal and advanced synthetic seed production systems by using somatic embryos in *Ipomoea batatas* were reported (Jeon et al., 1986; Cantliffe, 1993, Onishi et al., 1992, 1994). Encapsulation of celery and cauliflower somatic embryos and their conversion into plantlets were studied (Redenbaugh et al., 1986; Onishi et al., 1992). Propagation of *Solanum melongena* through encapsulation of various explants (somatic embryos, nodal segments with buds) and effect of carbon source on encapsulated propagules were described (Rao and Singh, 1991; Huda and Bari, 2007; Huda et al., 2007; Huda et al., 2009) and reported that sucrose showed better performance when nodal segment with axillary bud was used as explant, but when somatic embryos were used, sucrose + sorbitol (1:1) was found to be efficient. Encapsulation of different explants (somatic embryos, plantlets, cell aggregates from hairy roots), conservation of root regeneration potential of cell aggregates in coated capsules even after stored at 25°C up to 60 days and plant regeneration from them were observed respectively in *Armoracia rusticana* (Shigeta and Sato, 1994; Nakashimada et al., 1995; Repunte et al., 1995).

Encapsulation of different explants (somatic embryos, nodal segments, shoot tips and cell suspension cultures) and estimation of yield and canopy of field cultivated plants derived from synthetic seeds of *Solanum tuberosum* (Sarkar and Naik, 1998a; Fiegert et al., 2000; Nyende et al., 2003; Schafer-menuhr et al., 2003; Nyende et al., 2005 and Sarkar and Naik, 1998b) were investigated. Phonkajornyod et al. (2004) reported the dry synthetic seed production and desiccation tolerance induction in somatic embryos of *Capsicum annum*. Encapsulation of nodal segments and shoot tips of *Manihot esculenta* (Cassava) germplasm was reported (Danso and Ford-iioyd 2003; Cid et al., 2009). Malek (2009) studied the propagation of *Trichosanthes*

**Table 1.** List of plants and type of explants used for synthetic seeds.

Name of the species	Type of explant	Reference
<i>Acacia mangia</i> Willd x <i>A. auriculaformis</i> Cunn. ex Benth	Shoots and axillary buds	Asmah et al., 2011
<i>Acca sellowiana</i> (O.Berg) Burret	Pregerminated somatic embryos	Inocente et al., 2007
<i>Actinidia deliciosa</i> Liang & Ferguson	Shoot buds	Piccioni and Standardi, 1995
	Nodal	Gardi et al., 1999
	<i>In vitro</i> derived buds	Adriani et al., 2000
	Mirocuttings	Romay et al., 2002
<i>Adhatoda vasica</i> Nees	Shoot buds	Anand and Bansal, 2002
<i>Agave vera-cruz</i> Mill.	Shoot tips and somatic embryos	Tejavathi et al., 2006
<i>Allium sativum</i> L.	Calli	Kim and Park, 2002
	Bulblets	Bekheet, 2006
<i>Ananas comosus</i> L. Merr.	Micro shoots	Soneji et al., 2002
<i>Anthurium andreaum</i> Lind.	Embryogenic calli	Nhut et al., 2004
	Organogenic calli	Onishi et al., 1992
<i>Apium graveolens</i> Dulce	Somatic embryos	Redenbaugh et al., 1986
<i>Arachis hypogaea</i> L.	Somatic embryos	Padmaja et al., 1995
<i>Armoracia rusticana</i> , Gaertn., Mey & Scherb	Hairy roots	Uozumi et al., 1992
	Somatic embryos	Shigeta and Sato, 1994
	Shoots derived from hairy roots	Nakashimada et al., 1995
	Cell aggregates of hairy roots	Repunte et al., 1995
<i>Arnebia euchroma</i> (Royle) Johnston	Cotyledonary stage embryos	Manjkholia et al., 2005
<i>Asparagus cooperi</i> Baker	Somatic embryos	Ghosh and Sen, 1994.
<i>Asparagus officinalis</i> L.	Somatic embryos	Mamiya and Sakamoto, 2001
<i>Avena sativa</i> L.	Somatic embryos	Redenbaugh et al., 1987
<i>Bacopa monnieri</i> L.	Nodal microcuttings	Ramesh et al., 2009
<i>Begoniex hiemalis</i> Forch	Somatic embryos	Awal et al., 2007
<i>Beta vulgaris</i> L.	Somatic embryos	Tsai and Saunders, 1999
<i>Betula pendula</i> Roth.	Shoot buds	Piccioni and Standardi, 1995
<i>Betula platyphylla</i> Sukaczev var. japonica	Axillary buds	Kinoshita and Saito, 1990
<i>Brassica campestris</i> L.	Apical shoot buds	Arya et al., 1998
<i>Brassica oleracea</i> L. var. <i>botrytis</i>	Somatic embryos	Redenbaugh et al., 1986
<i>Camellia sinensis</i> (L.) O.Kuntze	Nodal explants	Mondal et al., 2002
<i>Camellia japonica</i> L.	Somatic embryos	Janeiro et al., 1997
	Shoot tips, nodal segments	Ballester et al., 1997
<i>Cannabis sativa</i> L.	Axillary buds	Lata et al., 2009
	Nodal segments	Lata et al., 2011
<i>Capsicum annuum</i> L.	Somatic embryo	Phonkajornyod et al., 2004
<i>Carica papaya</i> L.	Somatic embryos	Castillo et al., 1998
<i>Catharanthus roseus</i> (L.) G.Don	Somatic embryos	Maqsood et al., 2012

Table 1. Contd.

<i>Cedrela fissilis</i> Vellozo	Shoot tips	Nunes et al., 2003
<i>Cedrela odorata</i> L.	Shoot tips, axillary buds Shoot tips	Maruyama et al., 1997a Maruyama et al., 1997b
<i>Celogyne odoratissima</i> var. <i>angustifolia</i> Lindl.	Protocorms	Kamalakkannan et al., 1999
<i>Chlorophytum borivianum</i> Sant. et Fernand.	Shoot buds	Dave et al., 2004
<i>Cineraria maritima</i> L.	Microshoots	Srivastava et al., 2009
<i>Cinnamomum zeylanicum</i> (Linn.) Cor.	Shoot buds/Somatic embryos	Sajina et al., 1997
<i>Citrus nobilis</i> Lour. × <i>C. deliciosa</i> Tenora	Somatic embryos	Singh et al., 2007
<i>Citrus reticulata</i> Blanco.	Somatic embryos	Kitto and Janick, 1980 Kitto and Janick, 1985c Germana et al., 1999 Antonietta et al., 1999 Antonietta, et al., 2007b
<i>Cleopatra tangerine</i> L.	Embryos	Nieves et al., 1998
<i>Coelogyne breviscapa</i> Lindl	Protocorm like bodies	Mohanraj et al., 2009
<i>Coffea arabica</i> L.	Shoot buds	Nassar, 2003
<i>Coptis chinensis</i> Franch	Somatic embryos	Ke et al., 1995
<i>Coriandrum sativum</i> L.	Somatic embryos Somatic embryos	Chen et al., 1991; 1995 Stephen and Jayabalan, 2000
<i>Corymbia torelliana</i> (F. Muell.) Hill & Johnson. 2x <i>C. citriodora</i> (Hook.) Hill and Johnson	Shoot tips and Nodes	Hung and Trueman, 2011
<i>Crataegus oxyacantha</i> L.	Nodal segments Shoot buds	Piccioni and Standardi, 1995 Piccioni and Standardi, 1995
<i>Cremastra appendiculata</i> (D. Don) Makino	Protocorm	Zhang et al., 2009
<i>Cucumis sativus</i> L.	Somatic embryos	Tabassum et al., 2010
<i>Curcuma longa</i> L.	Adventitious buds	Sajina et al., 1997
<i>Cyclamen persicum</i> Mill.	Somatic embryos	Winkelmamm et al., 2004
<i>Cymbidium giganteum</i> Wall ex Lindl.	Protocorm Somatic embryos	Corrie and Tandon, 1993 Nhut et al., 2005
<i>Dalbergia sissoo</i> Roxb.	Somatic embryos Nodal segments	Singh and Chand, 2010 Chand and Singh, 2004
<i>Daucus carota</i> L.	Somatic embryos Somatic embryos Somatic embryos Somatic embryos Somatic embryos Asexual embryos Somatic embryos Somatic embryos Somatic embryos Somatic embryos Somatic embryos Somatic embryos	Kitto and Janick, 1985a Timbert et al., 1996 Wake et al., 1992 Latif et al., 2007 Sakamoto et al., 1992 Kitto and Janick, 1985b Liu et al., 1992 Janick et al., 1989 Timbert et al., 1995 Kamada et al., 1989 Kitto and Janick, 1982 Onishi et al., 1994

Table 1. Contd.

<i>Dendranthema x grandiflora</i> (RAMAT)	Nodal segments	Pinker and Abdel-rahman, 2005
<i>Dendrobium huoshanense</i> Tang ex Cheng.	Protocorm like bodies	Saiprasad and Polisetty, 2003
<i>Dendrobium densiflorum</i> Lindl.	Protocorm like bodies	Vij et al., 2001
<i>Dendrobium wardianum</i> Warner	Protocorm like bodies	Sharma et al., 1992
<i>Dendrocalamus strictus</i> L.	Somatic embryos	Mukunthakumar and Mathur, 1992
<i>Dianthus caryophyllus</i> L.	Shoot tips	Halmagyi and Deliu, 2007
<i>Dioscorea bulbifera</i> L.	Shoot tips	Narula et al., 2007
<i>Dioscorea</i> spp	Nodal segments	Hasan and Takagi, 1995
<i>Olea europaea</i> L. Shift	Nodal explants	Gardi et al., 1999
<i>Elaeis guineensis</i> Jacq.	Somatic embryos	Mariani et al., 2008
<i>Elektaria cardamom</i> Maton	Shoot tips	Ganapathi et al., 1994
<i>Eleusine coracana</i> Gaertn	Somatic embryos	George and Eapen, 1995
<i>Eucalyptus citriodora</i> Hook	Somatic embryos	Muralidharan and Mascarenhos, 1995
<i>Eustoma grandiflorum</i> (Raf.) Shinners	Somatic embryos	Ruffoni et al., 1994
<i>Flickingeria nodosa</i> (Dalz.) Seidenf.	Protocorm like bodies	Nagananda et al., 2011
<i>Genista monosperma</i> Lam.	Somatic embryos	Ruffoni et al., 1994
<i>Geodorum densiflorum</i> (Lam) Schltr.	Protocorm like bodies	Datta, et al., 1999
<i>Gerbera jamesonii</i> Bolus ex Hook.f	Microshoots and somatic embryos	Taha et al., 2009a
<i>Gladiolus</i> cultivars	Somatic embryos	Ramakrishnappa, 1998
<i>Goiabeira serrana</i>	Somatic embryos	Guerra et al., 2001
<i>Guazuma crinita</i> Mart	Shoot tips	Maruyama et al., 1997b
	Shoot tips, axillary buds	Maruyama et al., 1997a
<i>Gypsophila paniculata</i> L.	Shoot tips	Rady and Hanafy, 2004
<i>Helianthus annuus</i> L.	Shoot tips	Katouzi et al., 2011
<i>Hibiscus moscheutos</i> Welw. ex Hiern.	Nodal segments	Preece and West, 2006
<i>Hopea parviflora</i> Bedd.	Excised embryos	Sunilkumar et al., 2000
<i>Hordeum vulgare</i> L.	Microspore derived embryos	Datta and Potrykus, 1989
<i>Hyoscyamus muticus</i> L.	Somatic embryos	Pandey and Chand, 2005
<i>Ipomoea aquatica</i> Forsk	Axillary buds	Shaohu et al., 1994
<i>Ipomoea batatas</i> L.	Somatic embryos	Chee and Cantliffe, 1992.
	Somatic embryos	Cantliffe, 1993
	Somatic embryos from true seeds	Jeon et al., 1986
	Somatic embryos	Cantliffe et al., 1987
<i>Ipsea malabárica</i> (Reichb. F.) J. D. Hook.	Bulbs	Martin, 2003
<i>Jacaranda mimosaefolia</i> D. Don	Shoot tips or axillary buds	Maruyama et al., 1997a
	Shoot tips	Maruyama et al., 1997b
Kiwifruit CV Hayward, Tomuri	Nodal	Gardi et al., 1999
<i>Laeliocattleya okarchee</i> Addison (Cattleya)	Protocorm like bodies	Saiprasad and Polisetty, 2003
<i>Lilium longiflorum</i> Thunb.	Bulblets	Standardi et al., 1995
<i>Eustoma grandiflorum</i> (Raf.) Shinn.	Somatic embryos	Ruffoni et al., 1993
<i>Lycopersicon esculentum</i> Mill.	Seeds	Garrett et al., 1991
<i>Malus pumila</i> Mill. M 26 apple root stock	Apical buds	Micheli et al., 2002
	Nodal micro cuttings	Gardi et al., 1999

Table 1. Contd.

	Micropropagated buds	Piccioni, 1997
	Buds	Standardi and Piccioni, 1997
	Shoot tips	Sicurani et al., 2001
	Machine produced cuttings	Brischia et al., 2002
	Apical and axillary buds	Capuano et al., 1998
	Nodal explants	Gardi et al., 1999
	Micropropagated buds	Piccioni and Standardi, 1995
<i>Mangifera indica</i> L.	Somatic embryos	Ara et al., 1999
<i>Manihot esculenta</i> L.	Axillary buds	Cid et al., 2009
	Nodal segments and shoot tips	Danso and Ford-Lloyd, 2003
<i>Medicago sativa</i> L.		McKersie et al., 1989
		Fujii, et al., 1992.
		Redenbaugh et al., 1986
		Mckersie et al., 1990
	Somatic embryos	McKersie and Brown, 1996
		Redenbaugh et al., 1984,
		Redenbaugh and Walker 1990
		McKersie and Bowley, 1993
		Senaratna et al., 1990
<i>Mentha arvensis</i> L.	Axillary buds	Ahuja et al., 1989
<i>Morus alba</i> L.	Axillary buds	Pattnaik and Chand, 2000
	Axillary buds	Bapat, 1993
	Axillary buds	Machii, 1992
<i>Morus australis</i> Poir.	Axillary buds	Pattnaik and Chand, 2000
<i>Morus bombycis</i> Koidz	Axillary buds	Pattnaik and Chand, 2000
<i>Morus cathyana</i> Hemsl	Axillary buds	Pattnaik and Chand, 2000
<i>Morus indica</i> L.	Shoot buds	Bapat et al., 1987
	Axillary buds	Bapat and Rao, 1990
	Axillary buds	Kavyashree et al., 2006
	Apical buds	Kavyashree et al., 2004
<i>Morus latifolia</i> Poir.	Axillary buds	Pattnaik and Chand, 2000
<i>Morus nigra</i> L.	Axillary buds	Pattnaik and Chand, 2000
<i>Morus alba</i> L.	Axillary buds	Pattnaik et al., 1995
<i>Musa balbisiana</i> 'Kluai Hin' (BBB group)	Microshoots	Kanchanapoom and Promsom, 2012
		Ganapathi et al., 2002
		Ganapathi et al., 1992
		Hassanein et al., 2005
<i>Musa paradisiaca</i> L.	Shoot tips	Hassanein et al., 2011
		Matsumoto et al., 1995
		Priya and Arumugam, 2003
		Rao et al., 1993



Table 1. Contd.

<i>Musa paradisica</i> L. cv. <i>rasthali</i>	Somatic embryos	Srinivas., 2002
<i>Musa paradisica</i> L. cv. <i>grand naine</i>	Microshoots	Suprasanna et al., 2000
<i>Nerium oleander</i> L.	Axillary buds	Ganapathi et al., 2001.
<i>Nothafagus alpina</i> (Poepp. & Endl.) Oerst.	Somatic embryos	Sandoval-yugar et al., 2009
<i>Ocimum americanum</i> L.	Axillary buds	Ozden et al., 2008
<i>Ocimum basilicum</i> L.	Axillary buds	Cartes et al., 2009
<i>Ocimum gratissimum</i> L.	Axillary buds	Mandal et al., 2000
<i>Ocimum sanctum</i> L.	Axillary buds	Mandal et al., 2000
<i>Olea europaea</i> L. cv. <i>moraiolo</i>	Microcuttings	Mandal et al., 2000
<i>Olea europaea</i> L.	Nodal segments	Micheli et al., 2007
<i>Olive</i> cv. Camino, Moraiolo, <i>Ascolana tenera</i> and <i>Dolce agogia</i>	Nodal segments	Gardi et al., 1999
<i>Olea europaea</i> L.	<i>In vitro</i> derived shoots	Gardi et al., 1999
<i>Oncidium</i> 'Gower ramsay'	Protocorm like bodies	Ikhtiaq et al., 2010
<i>Oryza sativa</i> L.	Protoplast cultures	Saiprasad and Polisetty, 2003
	Androgenic proembryos	Giri and Reddy, 1994
	Somatic embryos	Roy and Mandal., 2008
	Somatic embryos	Suprasanna et al., 1996
	Somatic embryos	Arunkumar et al., 2005
<i>Oryza sativa</i> L. cv. <i>basmati</i> 370	Somatic embryos	Suprasanna et al., 2002
<i>Parkia speciosa</i> Hassk.	Embryos	Ummi et al., 2011
<i>Paulownia elongata</i> Hu.	Direct somatic embryos	Ipekci and Gozukirmizi, 2003
<i>Pelargonium domesticum</i> Bailey	Somatic embryos	Marsolais et al., 1991
<i>Pelargonium x hortorum</i> (zonal geranium)	Somatic embryos	Marsolais et al., 1991
<i>Pelargonium hortorum</i> Bailey.	Somatic embryos	Gill et al., 1994
<i>Phaius tankervilleae</i> (Banks) Blume	Protocorm like bodies	Malemngaba et al., 1996
<i>Phoenix dactylifera</i> L.	Somatic embryos	Bekheet et al., 2002 ; 2005
<i>Photinia fraseri</i> Dress	Axillary buds	Ozden et al., 2008
<i>Phyllanthus amarus</i> L.	Somatic embryos	Singh et al., 2006a
<i>Picea abies</i> L.	Somatic embryos	Gupta et al., 1987
<i>Picrorhiza kurroa</i> Royle. ex. Benth.	Micro shoots	Mishra et al., 2011
Pineapple <i>Ananas comosus</i> L.	Micro shoots	Gangopadhyay et al., 2005
<i>Piper hispidinervum</i> C.DC.	Pre germinated seeds	Pereira et al., 2008
<i>Piper nigrum</i> L.	Somatic embryos	Fair and Gupta, 2007
	Shoot buds	Sajina et al., 1997
<i>Pistachio vera</i> L.	Embryoids	Onay et al., 1996
<i>Plantago asiatica</i> L.	Shoot tips	Makowczynska and Andrzejewska-golec, 2006
<i>Pogonatherum paniceum</i> (Lam.) Hack	Shoot buds	Wang et al., 2007
<i>Pogostemon cablin</i> Benth.	Nodal segments	Swamy et al., 2009
<i>Populus tremuloides</i> L. x <i>P. tremula</i> Michx.	Shoot tips	Tsvetkov et al., 2006
Hybrid aspen	Shoot tips	Tsvetkov et al., 2006
<i>Pseudostellaria heterophylla</i> (Miquel) Pax.	Micro-tubers	Ma et al., 2011
<i>Psidium guajava</i> L.	Somatic embryos	Akhtar et al., 1997
	Somatic embryos	Rai and Jaiswal, 2008
	Shoot tips	Rai et al., 2008a

Table 1. Contd.

<i>Punica granatum</i> L.	Nodes	Rai et al., 2008b
<i>Pyrus communis</i> L.	Somatic embryos	Rai et al., 2008c
	Nodal segments	Naik and Chand, 2006
	Shoot tips	Ahmad et al., 2007
<i>Rauvolfia serpentina</i> (L.) Benth. ex Kurz.	Shoot tips	Ray and Bhattacharya, 2008
	Microshoots	Faisal et al., 2012
	Callus	Reddy et al., 2005
<i>Rauvolfia tetraphylla</i> L.	Microcuttings	Faisal et al., 2006
	Microshoots	Alatar and Faisal, 2012.
<i>Rhodiola kirilowii</i> (Regel) Maxim	Axillary buds and callus	Zych et al., 2005
<i>Rosa hybrida</i> Kings Ransom	Somatic embryos	Jayasree and Devi, 1997
<i>Rubus idaeus</i> L.	Shoot buds	Piccioni and Standardi, 1995
	Nodal	Gardi et al., 1999
	Nodal	Gardi et al., 1999
	Microcuttings	Romay et al., 2003
<i>Ruta graveolens</i> L.	Roots	Vdovitchenko and Kuzovkina, 2011
<i>Saccharum officinaum</i> L	Somatic embryos	Naik and Chikkagouda, 1997
		Sandeep, 2008
		Boonpeng et al., 2003
<i>Saintpaulia ionantha</i> Wendl.	Somatic embryos	Taha et al., 2009b
	Micro shoots	Daud et al., 2008
<i>Salvia officinalis</i> L.	Shoot tips	Grzegorzcyk and Wysokinska, 2011
<i>Santalum album</i> L.	Somatic embryos	Bapat, 1993
		Bapat and Rao, 1992
		Bapat and Rao, 1988
<i>Scutellaria baicalensis</i> Georgi	Roots	Vdovitchenko and Kuzovkina, 2011
Siberian ginseng	Somatic embryos	Choi and Jeong, 2002
<i>Simmondsia chinensis</i> L.	Shoot apical and axillary buds	Hassan, 2003
<i>Solanum melongena</i> L.	Somatic embryos	Rao and Singh, 1991
	Nodal segment with bud	Huda et al, 2007
	Somatic embryos	Huda and Bari, 2007
	Axillary buds	Huda et al., 2009
	Somatic embryos	Mariani, 1992
<i>Solanum nigrum</i> L.	Shoot tips	Verma et al., 2010
<i>Solanum tuberosum</i> L.	Nodal segments	Sarkar and Naik, 1998b

Table 1. contd.

	Shoot tips	Nyende et al., 2005
	Shoot tips	Patel et al., 2000
	Shoot tips	Nyende et al., 2005
	Nodal segments	Sarkar and Naik, 1998a
	Shoot tips	Nyende et al., 2003
	Shoot tips	Aggrey et al., 2003
<i>Solanum tuberosum</i> L. cv. <i>clarissa</i>	Cell suspension culture	Schafer-menuhr et al., 2003
<i>Solanum tuberosum</i> L. cv. <i>tomensa</i>	Somatic embryos	Fiegert et al., 2000
	Protocorm like bodies	Singh, 1991
<i>Spathoglottis plicata</i> BL.	Seeds and protocorms	Khor et al., 1998
	Seeds	Tan et al., 1998
	Protocorms	Singh, 1991
<i>Stevia rebaudiana</i> Bertoni	Shoot tips	Andlib et al., 2011
	Shoot tips, axillary buds	Ali et al., 2012
<i>Syringa vulgaris</i> L.	Axillary buds	Refouvelet et al., 1998
	Axillary buds	Ozden et al., 2008
<i>Theobroma cacao</i> L.	Embryos	Sudhakara et al., 2000
<i>Trichosanthes dioica</i> Roxb.	Shoot tips	Malek, 2009
	Shoots	Faisal and Anis, 2007
<i>Tylophora indica</i> (Burm.f.) Merrill	Somatic embryos	Chandrasekhar et al., 2006
	Somatic embryos	Devendra et al., 2011
<i>Valeriana wallichii</i> DC.	Apical and axial shoot buds	Mathur et al., 1989
<i>Vanda coerulea</i> Griff.	Protocorm like bodies	Sarmah et al., 2010
<i>Vanilla wightiana</i> Lindl.	Shoot buds	Sajina et al., 1997
<i>Vitex negundo</i> L.	Nodal segments	Ahmad and Anis, 2010
<i>Vitis vinifera</i> L.		Nirala et al., 2010
	Somatic embryos	Das et al., 2006
<i>Withania somnifera</i> (L.) Dunal	Shoot tips	Singh et al., 2006b
<i>Zea mays</i> L. var <i>saccharata</i>	Somatic embryos	Thobunluepop et al., 2009
<i>Zingiber officinale</i> Rosc.	Shoot buds	Sharma et al., 1994
	Micro shoots	Sundararaj et al., 2010

*dioica* through encapsulated shoot tips. Umami et al. (2011) investigated the effect of different storage intervals on encapsulated embryo and germination of *Parkia speciosa*. The encapsulated *P. speciosa* zygotic embryo (without storage) showed that germination occurred after seven days of culture on the germination medium. The synthetic seed stored at 4°C remained viable and

germination is initiated on day 14 after culture.

### Forage legumes

Among the forage legumes somatic embryogenesis and the development of synthetic seed, technology was

extensively studied in *Medicago sativa* (alfalfa). Encapsulation in hydrogel remains to be the most studied method of artificial seed production in alfalfa (Redenbaugh et al., 1986; Redenbaugh and Walker, 1990). Fujii et al. (1998) found improvement of plantlet growth and vigor through artificial seeds. Maturation and green house planting of alfalfa artificial seeds were studied by Fujii et al. (1987, 1989 and 1992) and observed that embryo maturation with ABA (5  $\mu$ M) was essential for high soil conversion (50 to 64%). Mckersie et al. (1989) developed artificial seeds of hybrid alfalfa. Induction of desiccation tolerance in *M. sativa* somatic embryos by ABA treatment and production of synthetic seeds were studied by Senaratna et al. (1989, 1990). They found that somatic embryos treated with ABA showed about 60% survival and conversion into plantlets when placed on moist filter paper or sown directly onto sterile soil. However, efficient coating and encapsulation methods for desiccated embryos of *M. sativa* are yet to be developed (Redenbaugh et al., 1991a, b). Somatic embryos of alfalfa (*M. sativa*) desiccated to 10 to 15% could be stored at room temperature for one year without a decline in their germinability (Mc Kersie and Bowley, 1993).

### Industrially important crops

Synthetic seed technology started from the mid 1980's in the industrially important crops such as mulberry, sandalwood, sugarcane etc. Mulberry is one of the most important crops which play an important role in silk industry, because its leaves serve as chief source for feeding silkworms (Yu et al., 2008). Bapat et al. (1987) and Bapat and Rao (1990) reported the successful *in vivo* growth of encapsulated shoot tip of *Morus indica* by the addition of fungicide to the alginate beads without contamination. Several reports have been published on axillary buds as encapsulation propagules in *Morus* spp. such as *Morus alba* (Machii, 1992), three years old mature mulberry trees of three indigenous and two Japanese varieties (Pattnaik et al., 1995) and in six mulberry species *M. alba*, *Morus australis*, *Morus bombycis*, *Morus cathyana*, *Morus latifolia* and *Morus nigra* (Pattnaik and Chand, 2000). Kavyashree et al. (2004, 2006) reported that the synthetic seeds were successfully produced by encapsulating non-embryogenic propagules like apical and axillary buds excised from aseptic shoots of mulberry (*M. indica* cv. S54). Sandalwood is a commercially valuable forest tree of India. It plays an important role in many industries such as pharmaceutical, cosmetics, soap and perfume industries etc. Plantlet regeneration from encapsulated somatic embryos of *Santalum album* was reported by Bapat and Rao (1988, 92) and Bapat (1993). *Saccharum officinarum* is well known as an extremely important commercially cultivated crop in India, which plays an

important role in sugar and fermentation industries, but, normally these crops are more susceptible to many fungal and bacterial diseases and propagate by using culm cuttings of mature stems which are very expensive. So Naik and Chikkagouda (1997) found that mass propagate sugarcane *in vitro*, sugarcane plantlets were regenerated from encapsulated somatic embryos derived from callus of sugarcane lines GSBT - 1. A model system for synchronous somatic embryo production combined with the formation of synthetic seeds was studied by Boonpeng et al. (2003) and found that 60% of germination successfully on Murashige and Skoong medium. Nieves et al. (2003) reported that plants *in vivo* of *Saccharum officinarum* from artificial seeds were taller and had a smaller diameter at eight months, but these differences disappeared at 12 months.

### Medicinal plants

Naturally most of the important medicinal plants are rare, endangered and endemic category. It is due to the low fruit and seed formation, poor germination capacity of seeds and due to the many other environmental conditions such as habitat modification, urbanization, climatic change and pollution etc. So, it is important to propagate and conserve these plant species. The production of synthetic seeds by encapsulating somatic embryos and vegetative propagules is rapidly becoming an applied technique with potential for mass propagation of medicinal plant species. Propagation of *Valeriana wallichii* using encapsulated apical and axial shoot buds with plantlet conversion under both *in vitro* (98%) and *in vivo* (64%) was reported by Mathur et al. (1989). Hasan and Takagi (1995) used nodal segments for encapsulation in *Dioscorea* spp. Alginate encapsulations of axillary buds of *Ocimum sanctum*, *Ocimum basilicum*, *Ocimum americanum*, *Ocimum gratissimum* and their regrowth even after 60 days of storage at 4°C were reported by Mandal et al. (2000). Anand and Bansal (2002) found that maximum conversion of the encapsulated shoot buds of *Adhatoda vasica* was on Gamborg's medium containing 4.65  $\mu$ M KN and 50 mg/l phloroglucinol. Direct somatic embryogenesis and synthetic seed production from *Paulownia elongata* was studied and their germination frequencies was reported as 53% before storage at 4°C and 43.2 and 32.4%, respectively after 30 and 60 days of storage by Ipekci and Gozukirmizi (2003). Dave et al. (2004) studied the *in vitro* propagation of rare Indian medicinal herb *Chlorophytum borivillianum* via encapsulated shoot buds and found that 80% of sprouting at 28 $\pm$ 2°C after three weeks and 90% of sprouting even after seven days of storage at 4°C. Pandey and Chand (2005) reported the efficient plant regeneration (60%) from encapsulated somatic embryos of *Hyoscyamus muticus*.

Micropropagation of *Rhodiola kirilowii* using the encapsulated axillary buds and callus was reported by

Zych et al. (2005). They found that 100% plantlet conversion even after six weeks of storage at 4°C. Studies on different plant species using various explants, cotyledonary embryos in *Arnebia euchroma* (Manjkhola et al., 2005), somatic embryos in *Tylophora indica* (Chandrasekhar et al., 2006), shoot tips in *Phyllanthus amarus* (Singh et al., 2006a), microcuttings in *Rauvolfia tetraphylla* (Faisal et al., 2006), shoot tips in *Withania somnifera* (Singh et al., 2006b), shoot tips in *Plantago asiatica* (Makowczynska and Andrzejewska-golec, 2006), shoot tips in *Dioscorea bulbifera* (Narula et al., 2007) and shoot tips in *Rauvolfia serpentina* (Ray and Bhattacharya 2008) has been reported. Ramesh et al. (2009) studied the effect of fungicide (bavistin) on conversion from encapsulated nodal microcuttings of micropropagated *Bacopa monnieri* and reported that 3.0 mg/l bavistin showed no reduction in plant conversion and generated maximum number of shoots even after storage up to 45 days at 18°C. The conversion to plantlet from alginate encapsulated axillary buds of *Cannabis sativa* was studied by Lata et al. (2009) and found that maximum conversion under *in vitro* was on MS medium supplemented with thidiazuron (TDZ, 0.5 µM) and plant preservative mixture (PPM, 0.07%) whereas *in vivo* 1:1 potting mix fertilone with coco natural growth medium, moistened with full strength MS + 3% sucrose + 0.5% PPM was observed. Cartes et al. (2009) encapsulated somatic embryos and zygotic embryos from mature seeds of rauli-beech (*Nothofagus alpina*) in different artificial endosperms in order to generate a cover and fulfill the function of nourishment, protection of the embryos facilitating their later germination. Srivastava et al. (2009) found the genetic stability of plants derived from encapsulated microshoots following six months of storage in *Cineraria maritima*.

Plant regeneration was achieved from encapsulated nodal segments in *Vitex negundo*, shoot tips in *Solanum nigrum* and microshoots in *Picrorhiza kurroa* (Ahmad and Anis, 2010; Verma et al., 2010; Mishra et al., 2011). Lata et al. (2011) prepared synthetic seeds of *C. sativa* by using the *in vitro* derived axillary buds and studied the genetic stability of synthetic seeds during *in vitro* multiplication and storage for six months at different growth conditions using inter simple sequence repeat (ISSR) DNA fingerprinting which showed homogeneity in the regrown clones and mother plant. Ma et al. (2011) produced a protocol for encapsulation and germination of microtubers of *P. heterophylla*. Andlib et al. (2011) reported that the synthetic seeds produced by the encapsulation of shoot tips are an alternative source for quick regeneration of *Stevia rebaudiana*. Direct somatic embryogenesis and synthetic seed production in *Tylophora indica* was reported by Devendra et al. (2011).

## Cereals

The application of synthetic seed technology to the

cereals started from the year 1989. Most of investigations were carried out to increase their yield and vigor. Artificial seeds are playing a major role in increasing the genetically transformed plant material and haploid plant production. Datta and Potrykus (1989) reported synthetic seeds derived from embryos of *Hordeum vulgare*. After this, Giri and Reddy (1994) reported alginate encapsulation of *Oryza sativa*. George and Eapen (1995) reported the encapsulation of somatic embryos in *Eleusine coracana*. Suprasanna et al. (1996) showed that the encapsulation of somatic embryos and conversion into plantlets of *Oryza sativa*. Suprasanna et al. (2002) studied the viability of encapsulated embryos derived from five year old long term culture of *Oryza sativa* cv. *basmati* 370. Arunkumar et al. (2005) reported the addition of protectants, bavistin and streptomycin as constituents of synthetic endosperm and found that there was no negative effect on germination and conversion. They also studied the conversion of synthetic seeds into seedlings in hybrid rice and reported that the application of self-breaking gel beads technology increased the germination (52%) and conversion (47%) of synthetic seeds. Roy and Mandal (2008) reported the development of synthetic seeds involving androgenic and pro-embryos in elite *Oryza sativa*. Model systems for synchronous somatic embryo production combined with encapsulation to form synthetic seeds were studied in *Zea mays* var. *saccharata* (Thobunluepop et al., 2009).

## Spices and plantation crops

Chen et al. (1991) reported 82% germination capacity of artificial seeds and survival rate of 83% in *Coriandrum sativum*. Stephen and Jayabalan (2000) produced artificial seeds in *Coriandrum sativum* by using somatic embryos derived from hypocotyls explants. Production of disease free encapsulated shoot buds of *Zingiber officinale* and their conversion into plantlets were reported by Sharma et al. (1994). High frequency plant regeneration from *Allium sativum* encapsulated calli and bulblets were reported, respectively by Kim and Park (2002) and Bekheet (2006). *In vitro* plant regeneration from encapsulated somatic embryos of *Piper nigrum* was reported by Nair and Gupta (2007). Sundararaj et al. (2010) showed the microshoots encapsulation of *Zingiber officinale*. Maximum conversion of the encapsulated shoot tips of *Elettaria cardamomum* into plantlets was on White's medium (Ganapathi et al., 1994). Cold storage of shoot cultures and alginate encapsulation of shoot tips of *Camellia japonica*, *Citrus reticulata* and propagation of tea (*C. sinensis*) by shoot proliferation of alginate-encapsulated nodal explants stored at 4°C were reported by Janeiro et al. (1997), Ballester et al. (1997) and Mondal et al. (2002). Induction of somatic embryogenesis, production of synthetic seeds and 70% of germination in *Elaeis guineensis* was studied by Mariani et al. (2008).

## Fruit crops

In most of the commercial fruit crops, the seed propagation has not been successful because of heterogeneity of seeds; minute seed size and presence of reduced endosperm, low germination rate and in some crops have desiccation sensitive and recalcitrant seeds which cannot be stored for longer time (Rai et al., 2009). Recently many of the crops available are seedless varieties. Propagation of *Musa paradisiaca* (Ganapathi et al., 1992; Matsumoto et al., 1995; Hassanein et al., 2005.) and *Musa paradisiaca* cv. *grand naine* (Sandoval-Yugar et al., 2009) was carried out through encapsulated shoot tips. In banana cv. *rasthali* (*Musa* spp. AAB group), plantlet regeneration was from alginate encapsulated somatic embryos (Ganapathi et al., 2001). Rooting induction and plantlet regeneration from *Malus pumila* var. M 26 apple rootstock synthetic seeds prepared by using apical and axillary micropropagated buds were reported (Standardi and Piccioni, 1997; Piccioni, 1997; Capuano et al., 1998; Sicurani et al., 2001; Brischia et al., 2002). Attempts for saving labor by using mechanical tools in the production of adventitious shoot tips suitable for encapsulation were tried by Sicurani et al. (2001). Micheli et al. (2002) stated that the presence of second layer of alginate (double encapsulation) and of a thin external coating layer on the alginate (encapsulation coating) did not show any detrimental effects on viability, sprouting and regrowth of the encapsulated microcuttings in M 26 apple rootstock. Effect of encapsulation on *Citrus reticulata* somatic embryo and their plantlet conversion, prospective of the encapsulation technology in the nursery activity of *Citrus* were studied by different authors (Antonietta et al., 1999; Antonietta et al., 2007 a, b). As compared to non-encapsulated or encapsulated with a growth regulator free artificial endosperm, somatic embryos encapsulated with an artificial endosperm containing GA<sub>3</sub> has greater ability to plantlet conversion (Antonietta et al., 1999).

The addition of thiophosphate – methyl (fungicide) in the artificial endosperm resulted in high levels of sprouting with prominent frequencies of root development and conversion into plantlets. Under *in vitro* and *ex vitro* conditions, Antonietta et al. (2007b) and Singh et al. (2007) studied the effect of storage conditions on percentage of germination of encapsulated and non encapsulated somatic embryos of kinnow mandarin (*Citrus nobilis* x *C. deliciosa* Tenora). Encapsulation of different explants were reported which includes: somatic embryos in *Carica papaya* and *Mangifera indica* (Castillo et al., 1998; Ara et al., 1999), micro shoots in *Ananas comosus* (Soneji et al., 2002; Gangopadhyay et al., 2005), nodal segments in *Punica granatum* (Naik and Chand 2006), shoots tips in *Pyrus communis* (Ahmad et al., 2007), shoot tips in *Psidium guajava* (Rai et al., 2008a) and somatic embryos in *Vitis vinifera* (Nirala et al., 2010).

## Forest trees

Mukunthakumar and Mathur (1992) observed that the additional coating of mineral oil on artificial seeds of *Dendrocalamus strictus* produced by encapsulating somatic embryos showed 56% plantlet conversion frequency at *in vivo* condition. The possibility for encapsulation of shoot tips and axillary buds in the production of artificial seeds in *Cedrela odorata*, *Guazuma crinita* and *Jacaranda mimosaeifolia* were studied by Maruyama et al. (1997a, b). Hassan (2003) produced encapsulated shoot apical and axillary buds of *Simmondsia chinensis*. *In vitro* conservation of *Cedrela fissilis* via encapsulation of shoot tips, cotyledonary and epicotyl nodal segments were reported by Nunes et al. (2003). Plant regeneration from alginate encapsulated nodal segments and somatic embryos of *Dalbergia sissoo*, respectively was described by Chand and Singh (2004) and Singh and Chand (2010). Inocente et al. (2007) reported that the pre-germinated somatic embryos encapsulated in a sodium alginate with BA (0.5 μM) and GA<sub>3</sub> (1 μM) developed radicals of *Acca sellowiana*. Hung and Trueman (2011) studied the alginate encapsulation of shoot tips and nodes for short term storage of *Corymbia torelliana* x *Corymbia citriodora*. They found that the seeds pretreated with indole 3 butyric acid (IBA) stored about six weeks effectively at 25°C (50 to 84% of germination) than that of 4°C (0 to 4% of germination). Highest frequencies of plantlet formation (46 to 80%) under *in vivo* conditions when pre converted shoot tip derived synthetic seeds were transferred on to an organic compost substrate and their 100% survival in nursery was described. Asmah et al. (2011) developed a protocol for encapsulation of *Acacia* hybrid *in vitro* derived shoots and axillary buds and found that the germination rate was within 73.3 to 100% in the duration of six to 20 days.

## Ornamental plants and orchids

In ornamental plants and orchids, the synthetic seeds have very much commercial importance, because of their minute seed size and presence of reduced endosperm in seeds (Lambardi et al., 2006). Ruffoni et al. (1994) produced synthetic seeds of somatic embryos in two ornamental species (*Eustoma grandiflorum* and *Genista monosperma*). Piccioni and Standardi (1995) produced synthetic seeds of shoot tips in *Betula pendula* (birch). Standardi et al. (1995) produced synthetic seeds of bulbs in *Lilium longiflorum*. A new protocol was presented for producing synthetic seeds of axillary buds in *Syringa vulgaris* (Refouvelet et al., 1998). Nhut et al. (2004) studied the propagation of *Anthurium andreanum* by the encapsulation of embryogenic calli. Rady and Hanafy (2004) reported the synthetic seed for encapsulation and regrowth of *in vitro* derived *Gypsophila paniculata* shoot tips. In various ornamental plant species using different

explants such as: shoot tips, microshoots and axillary nodes of hybrid aspen (Tsvetkov et al., 2006), *Saintpaulia ionantha* (Daud et al., 2008), *Nerium oleander*, *Photinia fraseri* and *Syringa vulgaris* (Ozden et al., 2008). Katouzi et al. (2011) encapsulated shoot tips of *Helianthus annuus* by adding salicylic acid for cold preservation.

In orchids the most widely used explants for the preparations of synthetic seeds are the seeds, protocorms and protocorm like bodies (Murthy et al., 2006). Khor et al. (1998), developed two-coat system for encapsulation of *Spathoglottis plicata* seeds and protocorms. The seeds and protocorms could withstand the encapsulation treatment with high viability 64 and 40%, respectively. Datta et al. (1999) produced an encapsulation of protocorm like bodies in *Geodorum densiflorum* an endangered orchid. Martin (2003) produced synthetic seeds in *Ipsea malabarica* by using *in vitro* formed bulbs.

In the recent years, the researchers are mostly concentrating on the enhancement of protocorm like bodies (PLBs) to make synthetic seed system commercial for the propagation of orchids. Zhang et al. (2009) showed a protocorm suspension culture of *Cremastra appendiculata* by liquid suspension culture system and formation synthetic seeds. Production and storage of synthetic seeds of *Coelogyne breviscapa* and *Vanda coerulea* using PLBs were reported, respectively by Mohanraj et al. (2009) and Sarmah et al. (2010). Nagananda et al. (2011) reported the regeneration of encapsulated protocorm like bodies of *Flickingeria nodosa*.

## THE GENETIC STABILITY OF SYNTHETIC SEEDS

Synthetic seeds have been widely used for micro-propagation of many plant species. The molecular studies to determine genetic stability of synthetic seeds derived plantlets were started from the last decade, but no modifications were revealed at the biochemical and/or molecular levels. The potential advantage of synthetic seeds for genetically identical to natural plants was supported by many reports (Nyende et al., 2003). The genetic stability of plantlets derived from encapsulated *Ananas comosus* micro shoots was proved by random amplified polymorphic DNA (RAPD) and ISSR techniques (Gangopadhyay et al., 2005). Bekheet (2006) reported that in *Allium sativum* both plantlets derived from encapsulated bulblets as well as normally *in vitro* were genetically similar to those that the *in vivo*. RAPD analysis showed the genetic stability of *in vitro* plantlet derived from encapsulated shoot tips of *Dioscorea bulbifera* (Narula et al., 2007). Srivastava et al (2009) reported that *Cineraria maritana*, analysis of the RAPD profiles revealed an average similarity coefficient of 0.944, they confirmed the molecular stability of plants derived from encapsulated microshoots followed by six months of storage. The genetic stability between mother

plants and somatic embryo derived synthetic seeds showed resemblance in *Cucumis sativus* and proved by using RAPD markers (Tabassum et al., 2010). In *Picrorhiza kurrooa*, the genetic stability of plants derived from encapsulated microshoots following three months of storage was proved by using cluster analysis of RAPD profile (Mishra et al., 2011). Lata et al. (2011) reported genetic stability of synthetic seed derived plants of *Cannabis sativa* studied by using ISSR- DAN fingerprinting and gas chromatography (GC) analysis of six major cannabinoids and showed homogeneity in the regrown clones and the mother plant.

## CONCLUSION

Synthetic seeds technique is a rapid tool of plant regeneration because of its wide use in conservation and delivery of tissue cultured plants. Protocols of encapsulation were already optimized for various plant species, but the commercial scale production of synthetic seeds was restricted to few species only due to several major problems, such as: asynchronous development of somatic embryos, improper maturation of somatic embryos, poor conversion rate of somatic embryos, lack of dormancy, and limited production of viable mature somatic embryos. Such investigations need a lot of efforts to perfect this technology and to make it available on a commercial scale.

In other cases where synthetic seeds were developed by encapsulating vegetative propagules, *in vitro* raised plantlets were used as the source of explants. So, optimized micropropagation systems would be required prior to synthetic seed development. In some plant species (trees) conversion of synthetic seeds into plantlets is another major problem for commercial application. Better understanding about manipulations in the composition of synthetic endosperm, explant size, media composition, change in the formulation of medium and type of medium, optimization of growth regulators and addition of other additives to the synthetic endosperm are required to enhance the germination frequency of encapsulated propagules. However further detailed research is needed mainly for improvement in conversion frequency of synthetic seeds and subsequent plantlet growth in soil.

## ACKNOWLEDGEMENTS

The receipt of financial assistance from the Council of Scientific and Industrial Research (CSIR) and University Grants Commission (UGC), New Delhi, are gratefully acknowledged.

## Abbreviations

**MS**, Murashige and Skoog medium; **BA**, 6- benzyl-

adenine;  $\mu\text{M}$ , micro moles;  $\text{mg/l}$ , milligrams per liter; **NAA**,  $\alpha$ - naphthalene acetic acid; **GA<sub>3</sub>**, gibberellic acid; **ABA**, abscisic acid; **Synseed**, synthetic seed; **KN**, Kinetin; **TDZ**, thidiazuron; **PPM**, plant preservative mixture; **PLBs**, protocorm like bodies; **ISSR**, inter simple sequence repeat; **RAPD**, random amplified polymorphic DNA; **IBA**, indole 3 butyric acid.

## REFERENCES

- Adriani M, Piccioni E, Standard A (2000). Effect of different treatments on the conversion of 'Hayward' kiwifruit synthetic seeds to whole plants following encapsulation of *in vitro*-derived buds. *NZJ Crop. Horticult. Sci.* 28:59-67.
- Aggrey NS, Siegfied G, Wagner MK, Micheal GJ (2003). Production storability and regeneration of shoot tips of potato (*Solanum tuberosum* L.) encapsulated in sodium alginate hollow beads. *In vitro Cell Dev. Biol. Plant.* 39:540 - 544.
- Ahmad AN, Ali EAM, Rizkalla AA (2007). Synthetic seeds of pear (*Pyrus communis* L.). Rootstock storage *in vitro*. *Aust. J. Basic Appl. Sci.* 1: 262- 270.
- Ahmad N, Anis M (2010). Direct plant regeneration from encapsulated nodal segments of *Vitex negundo*. *Biologia Plantarum.* 54:748-752.
- Ahuja PS, Mathur J, Lai N, Mathur A, Kukreja AK (1989). Towards developing artificial seeds by shoot bud encapsulation. In: Kukreja AK, Mthur A, Ahuja PS and Thakur RS (eds), *Tissue culture and Biotechnology of medicinal and aromatic plants*. Lucknow. India. CIMAP pp. 20-78.
- Akhtar N (1997). Studies on induction of somatic embryogenesis and production of artificial seeds for micropropagation, of a tropical fruit tree guava (*Psidium guajava* L.). Ph.D. dissertation, Banaras Hindu University, Varanasi.
- Alatar A, Faisal M (2012). Encapsulation of *Rauvolfia tetraphylla* microshoots as artificial seeds and evaluation of genetic fidelity using RAPD and ISSR markers *J Med. Plants Res.* 6:1367-1374.
- Ali A, Gull I, Majid A, Saleem A, Naz S, Naveed NH (2012). *In vitro* conservation and production of vigorous and desiccated tolerant synthetic seeds in *Stevia rebaudiana*. *J. Med. Plants Res.* 6:1327-1333.
- Altman A, Nadel BL, Falash Z, Levin N (1990). Somatic embryogenesis in Celery: induction control and changes in polyamines and proteins. In Nijkamp HJJ, Vander Plas LHW and Van Aartrijk J (eds), *progress in plant cellular and molecular biology*. Kluwer, Dordrecht pp. 454-459.
- Ammirato PV (1983). Embryogenesis. In: *Handbook of Plant Cell Culture*, (Eds.): Evans DA, Sharp WR, Ammirato PV and Yamada Y. Macmillan, New York. 1:82-123.
- Anand Y, Bansal YK (2002). Synthetic seeds: A novel approach of *in vitro* plantlet formation in vasaka (*Adhatoda vasica* Nees). *Plant Biotechnol.* 19:159-162.
- Andlib A, Verma RN, Batra A (2011). Synthetic seeds an alternative source for quick regeneration of a zero calorie herb - *Stevia rebaudiana* Bertoni. *J. Pharm. Res.* 4:2007-2009.
- Antonietta GM, Ahmad HI, Maurizio M, Alvaro S (2007a). Preliminary research on conversion of encapsulated somatic embryos of *Citrus reticulata* Blanco, cv. Mandarin Tardivo di Ciaculli. *Plant Cell Tiss. Org. Cult.* 88:117-120.
- Antonietta GM, Micheli M, Pulcni L, Standardi A (2007b). Perspectives of the encapsulation technology in the nursery activity of *Citrus*. *Caryologia* 60:192-195.
- Antonietta GM, Emanuele P, Alvaro S (1999). Effects of encapsulation on *Citrus reticulata* Blanco. somatic embryo conversion. *Plant Cell Tiss. Org. Cult.* 55:235-237.
- Ara H, Jaiswal U, Jaiswal VS (1999). Germination and plantlet regeneration from encapsulated somatic embryos of mango (*Mangifera indica* L.). *Plant Cell Rep.* 19:166-170.
- Ara H, Jaiswal U, Jaiswal VS (2000). Synthetic seed: prospects and limitations. *Curr. Sci.* 78:1438-1444.
- Arunkumar MB, Vakeswaran V, Krishnasamy V (2005). Enhancement of synthetic seed conversion to seedlings in hybrid rice. *Plant Cell Tiss. Org. Cult.* 81:97-100.
- Arya KR, Beg MU, Kukreja AK (1998). Microcutting and propagation of endosulphan tolerant genotypes of mustard (*Brassica campestris*) through apical shoot bud encapsulation. *Indian J. Exp. Biol.* 36:1162-1164.
- Asmah HN, Hasnida HN, Zaimah NAN, Noraliza A, Salmi NN (2011). Synthetic seed technology for encapsulation and regrowth of *in vitro*-derived *Acacia* hybrid shoot and axillary buds. *Afr. J. Biotechnol.* 10: 7820-7824.
- Awal A, Taha RM, Hasbullah NA (2007). *In vitro* formation of synthetic seeds of *Begoniax hiemalis* Foch. *Indian J. Environ. Sci.* 2:189-192.
- Ballester A, Janeiro LV, Vieitez AM (1997). Cold storage of shoot cultures and alginate encapsulation of shoot tips of *Camellia japonica* L. and *Camellia reticulata* Lindley. *Sci. Hort.* 71:67-78.
- Bapat VA (1993). Studies on synthetic seeds of sandalwood and mulberry. In K. Redenbaugh (ed), *synthetic seeds; application of synthetic seeds to crop improvement*. CRC Press, Boca Raton pp. 381-384.
- Bapat VA, Mahatre M, Rao PS (1987). Propagation of *Morus indica* L. (Mulberry) by encapsulated shoot buds. *Plant Cell Rep.* 6:393-395.
- Bapat VA, Rao PS (1988). Sandalwood plantlets from synthetic seeds. *Plant Cell Rep.* 7:434-436.
- Bapat VA, Rao PS (1990). *In vivo* growth of encapsulated axillary buds of mulberry (*Morus indica* L.) *Plant Cell Tiss. Org. Cult.* 20:69-70.
- Bapat VA, Rao PS (1992). Plant regeneration from encapsulated and non encapsulated desiccated somatic embryos of a forest tree: Sandalwood (*Santalum album* L.). *J. Plant Biochem. Biotechnol.* 1:109-113.
- Bapat V, Minal M (2005). Bioencapsulation of somatic embryos in woody plants. In SM Jain and PK Gupta (eds) *Protocol for somatic embryogenesis in woody plants forestry sciences*. Springer Netherlands 77:539-552.
- Bekheet SA (2006). A synthetic seed method through encapsulation of *in vitro* proliferated bulblets of garlic (*Allium sativum* L.). *Arab J. Biotech.* 9:415-426.
- Bekheet SA, Taha HS, El-Bahr MK (2005). Preservation of date palm cultures using encapsulated somatic embryos. *Arab J. Biotech.* 8:319-328.
- Bekheet SA, Taha HS, Saker MM, Moursya HA (2002). A synthetic seed system of date palm through somatic embryos encapsulation. *Anal. Agric. Sci. Cairo.* 47:325-337.
- Bhojwani SS, Razdan MK (2006). *Plant Tissue Culture: Theory and Practice*, Elsevier Science, Netherland pp. 125-166.
- Boonpeng S, Vearasilp S, Apavatjrit P, Suriyong S, Pawelzir E (2003). Possibility of sugarcane synthetic seed production. *Deutscher Tropentag*, October 8-10.
- Brischia R, Piccioni E, Standardi A (2002). Micropropagation and synthetic seed in M 26 apple rootstock (11): A new protocol for production of encapsulated differentiating propagules. *Plant Cell Tiss. Org. Cult.* 68:137-141.
- Buchheim JA, Colburn SM, Ranch JP (1989). Maturation of soybean somatic embryos and the transition to plantlet growth. *Plant Physiol.* 89: 768-775.
- Cantliffe DJ (1993). Advanced propagation system for biomass species: A model system based on Sweet potato. *Biomass Bioenergy* 5:63-69.
- Cantliffe DJ, Liu JR, Schultheis JR (1987). Development of artificial seeds of sweet potato for clonal propagation through somatic embryogenesis. In: Smith WH and Frank JR (eds.), *Methane from biomass a systems approach*. Elsevier Applied Publ. Ltd. New York pp. 183-195.
- Capuano C, Piccioni E, Standardi A (1998). Effect of different treatments on the conversion of M.26 apple rootstock synthetic seeds obtained from encapsulated apical and axillary micropropagated buds. *J. Hort. Sci.* 73:299-305.
- Carman JG (1990). Embryogenic cells in plant tissue cultures: occurrence and behaviour. *In Vitro Cell. Dev. Biol.* 26:746-753.
- Cartes PR, Castellanos HB, Rios DL, Saez KC, Spiercolli SH, Sanchez YMO (2009). Encapsulation of somatic embryos and zygotic embryos for obtaining artificial seeds of rauli-beech (*Nothofagus alpina* (Poepp. & Endl.) Oerst. Chilean J. Aric. Res. 69:112-118.
- Castillo B, Smith MAL, Yadav UL (1998). Plant regeneration from



- encapsulated somatic embryos of *Carica papaya* L. Plant Cell Rep. 17: 172-176.
- Chand S, Singh AK (2004). Plant regeneration from encapsulated nodal segments of *Dalbergia sissoo* Roxb. a timber yielding leguminous tree species. J. Plant Physiol. 161:237-243.
- Chandrasekhar T, Hussain MT, Gopal RG, Rao SJV (2006). Somatic embryogenesis of *Tylophora indica* (Burm.f.) Merrill., an important medicinal plant. Int. J. Appl. Sci. Eng. 4: 33-40.
- Chee RP, Cantlieff DJ (1992). Improved production procedures for a somatic embryos of sweet potato for a synthetic seed system. Hort Sci. 27:1314-1315.
- Cheema GS (1989). Somatic embryogenesis and plant regeneration in *Carica papaya* L. tissue culture derived from root explants. Plant Cell Rep. 6:348-351.
- Chen RR, Zhang JT, Li BP, Guo SS, Hao JP, Zhou XM (1991). Studies on the production of artificial seeds of Coriander. Chin. J. Biotechnol. 7: 127-134.
- Chen RR, Zhang JT, Ping LB, Guo SS, Hao JP (1995). Somatic embryogenesis and synthetic seeds in coriander (*Coriandrum sativum*). In: Bajaeg YPS (ed), Biotechnology in Agriculture and Forestry. Somatic embryogenesis and Synthetic seeds 11. Berlin. Springer. Verlag. 31:334-342.
- Choi YE, Jeong JH (2002). Dormancy induction of somatic embryos of Siberian ginseng by high sucrose concentrations enhances the conservation of hydrated artificial seeds and dehydration resistance. Plant cell Rep. 20:1112-1116.
- Cid LPB, Cruz ARR, Carvalho LJCB (2009). Encapsulation of Cassava nodal segments for germplasm storage. Embrapa Recursos Genéticos e Biotecnologia, Brasília, DF. pp. 1-10.
- Corrie S, Tandon P (1993). Propagation of *Cymbidium giganteum* well through high frequency conversion of encapsulated protocorms under *in vivo* and *in vitro* conditions. Indian J. Exp Biol. 31:61-64.
- Danso KE, Ford-Lloyd BV (2003). Encapsulation of nodal cuttings and shoot tips for storage and exchange of Cassava germplasm. Plant Cell Rep. 21:718-725.
- Das DK, Nirala NR, Reddy MK, Sopory SK, Upadhyaya KC (2006) Encapsulated somatic embryos of grape (*Vitis vinifera* L.): An efficient way for storage and propagation of pathogen-free plant material. Vitis 45:179-184
- Datta KB, Kanjilal B, Sarker D (1999). Artificial seed technology. Development of a protocol in *Geodorum densiflorum* (Lam.) Schltr. An endangered orchid. Curr. Sci. 76:1142-1145.
- Datta SK, Potrykus I (1989). Artificial seeds in barley: encapsulation of microspore derived embryos. Theor. Appl. Genet. 77:820-824.
- Daud N, Taha RM, Hasbullah NA (2008). Artificial seed production from encapsulated microshoots of *Saintpaulia ionantha* Wendl. (African violet). J. Appl. Sci. 8: 4662-4667.
- Dave A, Joshi N, Purohit SD (2004). *In vitro* propagation of *Chlorophytum borivilianum* using encapsulated shoot buds. Eur. J. Hort Sci. 69:37-42.
- Devendra BN, Srinivas N, Naik GR (2011). Direct somatic embryogenesis and synthetic seed production from *Tylophora indica* (Burm.f.) Merrill. An endangered, medicinally important plant. Int. J. Bot. 7:216-222.
- Dobrev P, Motyka V, Gaudinová A, Malbeck J, Trávníčková A, Kamínek M, Valková R (2002). Transient accumulation of *cis*- and *trans*-zeatin type cytokinins and its relation to cytokinin oxidase activity during cell cycle of synchronized tobacco -2 cells. Plant Physiol. Biochem. 40: 33-337.
- Fair RR, Gupta SD (2007). *In vitro* plant regeneration from encapsulated somatic embryos of Black pepper (*Piper nigrum* L.) J. Plant Sci. 2: 283-292.
- Faisal M, Anis M (2007). Regeneration of plants from alginate encapsulated shoots of *Tylophora indica* (Burm. f) Merrill. An endangered medicinal plant. J. Hort. Sci. Biotechnol. 82:351-354.
- Faisal M, Ahmad N, Anis M (2006). *In vitro* plant regeneration from alginate encapsulated microcuttings of *Rauvolfia tetraphylla* L. American-urasian J. Agric. Environ. Sci. 1:1-6.
- Faisal M, Alatar AA, Ahmad N, Anis M, Hegazy AK (2012). Assessment of genetic fidelity in *Rauvolfia serpentina* plantlets Grown from Synthetic (encapsulated) seeds following *in vitro* storage at 4° C. Molecules 17: 5050-5061.
- Fiegert AK, Mix-wagner G, Vorlop KD (2000). Regeneration of *Solanum tuberosum* L. Tomensa induction of somatic embryogenesis in liquid culture for the production of artificial seeds. Landbauforsch. Volk 50: 199-202.
- Fujii JA, Slade D, Redenbaugh K (1989). Maturation and greenhouse planting of alfalfa artificial seeds. *In Vitro* Cell. Dev. Biol. Plant 25: 1179-1182.
- Fujii JA, Slade D, Redenbaugh K (1998). Improving plantlet growth and vigour from alfalfa artificial seeds. *In vitro* Cell Dev. Biol. Plant 24: 70A.
- Fujii JA, Slade D, Aguirre Rascon J, Redenbaugh K (1992). Field planting of alfalfa artificial seeds. *In Vitro* Cell. Dev. Biol. Plant. 28:73-80.
- Fujii JA, Slade D, Redenbaugh K, Walker KA (1987). Artificial seeds for plant propagation. Trends Biotech. 5:335-339.
- Fujii JA, Slade D, Redenbaugh K (1993). Planting artificial seeds and somatic embryos In: Redenbaugh K (eds), synseeds: Application of synthetic seeds to crop improvement. CRC press, Boca Raton, FL. pp. 183-202.
- Fujimura T, Komamine A (1979). Synchronization of somatic embryogenesis in a carrot cell suspension culture. Plant Physiol. 64: 162-164.
- Ganapathi TR, Bapat VA, Rao PS (1994). *In vitro* development of encapsulated shoot tips of cardamom. Biotech. Techniques 8:239-244.
- Ganapathi TR, Srinivas L, Suprasanna P, Bapat VA (2001). Regeneration of plants from alginate-encapsulated somatic embryos of banana cv. Rasthali (*Musa* spp. AAB group). *In vitro* Cell Dev. Biol. Plant 37:178-181.
- Ganapathi TR, Suprasanna P, Kulkarni VM, Bapat VA, Rao PS (2002). Strategies for *in vitro* propagation and synthetic seeds in banana. BARC News letter. Founders day special issue: pp. 168-176.
- Ganapathi TR, Suprasanna P, Bapat VA, Rao PS (1992). Propagation of banana through encapsulated shoot tips. Plant Cell Rep. 11:571-575.
- Gangopadhyay G, Bandyopadhyay T, Poddar R, Gangopadhyay SB, Mukherjee KK (2005). Encapsulation of pineapple microshoots in alginate beads for temporary storage. Curr. Sci. 88:972-976.
- Gardi T, Piccioni E, Standardi A (1999). Effect of bead nutrient composition on regrowth of stored *in vitro*-derived encapsulated microcuttings of different woody species. J. Microencapsulation 16: 13-25.
- Garrett RE, Mehlchou NE, Smith NE, Redenbaugh KM (1991). Gel encapsulation of tomato seeds. Appl. Eng. Agric. 7:25-51.
- George L, Eapen S (1995). Encapsulation of somatic embryos in finger millet. Indian J. Exp. Biol. 33:291-293.
- Germana MA, Piccioni E, Standardi A (1999). Effect of encapsulation on *Citrus reticulata* Blanco somatic embryo conversion. Plant Cell Tiss. Org. Cult. 55:235-237.
- Ghosh B, Sen S (1994). Plant regeneration from alginate encapsulated somatic embryos of *Asparagus cooperi* Baker. Plant Cell Rep. 13: 381-385.
- Gill R, Senaratna T, Saxena PK (1994). Thidiazuron induced somatic embryogenesis enhances viability of hydrogel encapsulated somatic embryos of *Geranium*. J. Plant Physiol. 143:726-729.
- Gingear VM, Lineberger RD (1989). Asexual embryogenesis and plant regeneration in *Quercus*. Plant Cell Tiss. Org. Cult. 17:191-203.
- Giri CC, Reddy GM (1994). Alginate encapsulation technique for indica rice protoplast culture and plant regeneration. Curr. Sci. 67:542-545.
- Gray DJ (1987). Synthetic seed technology for the mass cloning of crop plants: problems and prospects. Horti. Sci. 22:795-814.
- Gray DJ, Purohit A, Trigiano RN, Triglanoph DRN (1991). Somatic embryogenesis and development of synthetic seed technology. Crit. Rev. Plant Sci. 10:33-61.
- Grzegorzczuk I, Wysokińska H (2011). A Protocol for synthetic seeds from *Salvia officinalis* L. shoot tips. Acta Biologica Cracoviensia Series Botanica 53:80-85.
- Guerra MP, Vesco LD, Ducroquet JPHJ, Nodari RO, Dosreis M (2001). Somatic embryogenesis in *Goiabeira serrana*: Genotype response, auxinic shock and synthetic seeds. Revista Brasileira de Fisiologia Veg. 13:117-128.
- Gupta PK, Shaw D, Durjan DJ (1987). Micropropagation, somatic

- embryogenesis and encapsulation In Donja JM and Durjan DJ (eds), Cell and tissue culture in forestry. Dordrecht. The Netherlands: Martinus Nijhoff Publ. pp. 101-108.
- Halmagyi A, Deliu C (2007). Cryopreservation of carnation (*Dianthus caryophyllus* L.). Shoot tips by encapsulation-vitrification. *Sci. Hortic.* 113: 300-306.
- Hasan SMZ, Takagi H (1995). Alginate coated nodal segments of Yam (*Dioscorea* spp) for germplasm exchange and distribution. *Plant Genet Resour. Newslett.* 103: 32- 35.
- Hassan NS (2003). *In vitro* propagation of Jojoba (*Simmondsia chinensis* L.) through alginate-encapsulated shoot apical and axillary buds. *Int. J. Agric. Biol.* 5:513-516.
- Hassanein AM, Ibrahiem IA, Galal AA, Salem JMM (2005). Micropropagation factors essential for mass propagation of Banana. *J. Plant Biotechnol.* 7:175-181.
- Hassanein AM, Ibraheim IA, Galal AA, Salem JMM (2011). Conversion of Banana Synseed influenced by the bead type and seed coat. *Am. J. Plant Sci.* 2:467-475.
- Helal NAS (2011). The green revolution via synthetic (artificial) seeds: A review. *Res. J. Agric. Biol. Sci.* 7:464-477.
- Huda A, Bari MA (2007). Production of synthetic seed by encapsulating asexual embryo in eggplant (*Solanum melongena* L.). *Int. J. Agric. Res.* 2:832-837.
- Huda A, Bari MA, Rahman M (2009). Asexual propagation of eggplant (*Solanum melongena* L.) through encapsulated axillary buds. *Plant Tissue Cult. Biotech.* 19:263-288.
- Huda AKMN, Rahman M, Bari MA (2007). Effect of carbon source in alginate bead on synthetic seed germination in eggplant (*Solanum melongena* L.). *J. Plant Sci.* 2:538-544.
- Hung CD, Trueman SJ (2011). Alginate encapsulation of shoot tips and nodal segments for short-term storage and distribution of the eucalypt *Corymbia torelliana* x *C. citriodora*. *Acta Physiol. Plant* 34:117-128.
- Ikhtaq M, Hafiz IA, Micheli M, Ahmad T, Abbasi NA, Standardi A (2010). *In vitro* storage of synthetic seeds: effect of different storage conditions and intervals on their conversion ability. *Afr. J. Biotechnol.* 9:5712-5721.
- Inocente GCC, Vesco LLD, Steinmacher D, Torres AC, Guerra MP (2007). Improvements in somatic embryogenesis protocol in Feijoa (*Acca sellowiana* (Berg) Burret): induction, conversion and synthetic seeds. *Sci. Hortic.* 111:228-234.
- Ipekci Z, Gozukirmizi N (2003). Direct somatic embryogenesis and synthetic seed production from *Paulownia elongata*. *Plant Cell Rep.* 22: 16-24.
- Janeiro LV, Ballester A, Vieitez AM (1997). *In vitro* response of encapsulated somatic embryos of *Camellia*. *Plant Cell Tiss. Org. Cult.* 51:119-125.
- Janick J, Kitto SL, Hwankim Y (1989). Production of synthetic seeds by desiccation and encapsulation. *In Vitro Cell. Deve. Biol. Plant.* 25: 1167 -1172.
- Janick Y, Kim YH, Kitto S, SarangaY (1993). Desiccated synthetic seeds. In: Redenbaugh K (ed), *Synthetic seeds: Application of synthetic seeds to crop improvement*. CRC press. Boca Raton. FL. pp. 11-33.
- Jayasree N, Devi BP (1997). Production of synthetic seeds and plant regeneration in *Rosa hybrida* L cv Kings ranson. *Indian J. Exp. Biol.* 35: 310-312.
- Jeon JH, Liu JR, Yang SG, Lee HS, Joung H, Han MH (1986). Development of a modal system for artificial seed production, 1: Encapsulation of somatic embryos by alginic acid. *Kor. J. Plant Tiss. Cult.* 13:119-128.
- Kamada H, Kobayashi K, Kiyosue T, Harada H (1989). Stress induced somatic embryogenesis in carrot and its application to synthetic seed production. *In vitro Cell. Dev. Biol. Plant* 25:1163-1166.
- Kamalakaran R, Narmatha BV, Jeyekudi L (1999). Plantlet regeneration from encapsulated protocorms of the endemic orchid *Celogyne odoratissima* var. *angustifolia* Lindl. *J. Phytol. Res.* 12:21-23.
- Kanchanapoom K, Promsorn N (2012). Micropropagation and *in vitro* germplasm conservation of endangered *Musa balbisiana* 'Kluai Hin' (BBB group). *Afr. J. Biotechnol.* 11:6464-6469.
- Kao AN, Michayluk MR (1981). Embryoid formation in alfalfa cell suspension cultures from different plants. *In Vitro*, 17: 645-648.
- Katouzi SSS, Majd A, Fallahian F, Bernard F (2011). Encapsulation of shoot tips in alginate beads containing salicylic acid for cold preservation and plant regeneration in sunflower (*Helianthus annuus* L.). *Aust. J. Crop Sci.* 5:1469-1474
- Kavyashree R, Gayatri MC, Revanasiddaiah MH (2004). Regeneration of apical buds of mulberry variety S54 (*Morus indica* L.). *Seriocologia* 44: 83-89.
- Kavyashree R, Gayatri MC, Revanasiddaiah MH (2006). Propagation of mulberry S54 by synthetic seeds of axillary buds. *Plant Cell Tiss. Org. Cult.* 84: 245-249.
- Ke S, Gui Y, Skirvin RM (1995). Somatic embryogenesis and synthetic seeds in *Coptis chinensis* In: Bajaj YPS (ed), *Biotechnology in Agriculture and Forestry somatic embryogenesis and synthetic seeds II*. Berlin. Springer Verlag. 31:323-333.
- Khor E, Ng WF, Loh CS (1998). Two-coat system for encapsulation of *Spathoglottis plicata* (Orchidaceae) seeds and protocorms. *Biotechnol. Bioeng.* 59:635-639.
- Kim MA, Park JK (2002). High frequency plant regeneration of garlic (*Allium sativum* L.) calli immobilized in calcium alginate gel. *Biotechnol. Bioprocess Eng.* 7:206-211.
- Kim YH, Janick J (1987). Production of synthetic seeds of celery. *Hort. Sci.* 22: 1065.
- Kim YH, Janick J (1989). ABA and polyox encapsulation of high humidity increase survival of desiccated somatic embryos of celery. *Hort. Sci.* 27: 674-676.
- Kim YH, Janick J (1990). Synthetic seed technology improving desiccation tolerance of somatic embryos of celery. *Acta Hortic.* 280: 23-28.
- Kinoshita I, Saito A (1990). Propagation of Japanese White Birch by encapsulated axillary buds. *J. Jpn. For. Soc.* 72:166-170.
- Kitto SK, Janick J (1982). Polyox as an artificial seed coat for asexual embryos. *Hort. Sci.* 17: 488-490.
- Kitto SL, Janick J (1980). Water soluble resins as artificial seed coats. *Hort. Sci.* 15:439.
- Kitto SL, Janick J (1985a). Production of synthetic seeds by encapsulating asexual plant embryos of carrot. *J. American Soc. Hort. Sci.* 110:277-282.
- Kitto SL, Janick J (1985b). Hardening treatments increase survival of synthetically-coated asexual embryos of carrot. *J. Am. Soc. Hort. Sci.* 110: 283-286.
- Kitto SL, Janick J (1985c). A citrus embryo assay to screen water soluble resins as synthetic seed coats. *Hort. Sci.* 20:98-102.
- Komamine A, Kawahara R, Matsumoto M, Sunabori S, Toya T, Fujiwara A, Tsukihara M, Smith J, Ito M, Fukuda H, Nomura K, Fujimura T (1992). Mechanisms of somatic embryogenesis in cell cultures: physiology, biochemistry, and molecular biology. *In Vitro Cell. Dev. Biol.* 28:11-14.
- Lambardi M, Benelli C, Ozudogru EA (2006). Synthetic seed technology in ornamental plants. In Teixeira da silva JA (ed), *Floriculture, ornamental and plant biotechnology*. Global Science books. UK. 2:347-354.
- Lata H, Chandra S, Natascha T, Khan IA, ElSohly MA (2011). Molecular analysis of genetic fidelity in *Cannabis sativa* L. plants grown from synthetic (encapsulated) seeds following *in vitro* storage. *Biotechnol. Lett.* 33:2503-2508.
- Lata H, Chandra S, Khan IA, Elsohly, MA (2009). Propagation through alginate encapsulation of axillary buds of *Cannabis sativa* L. -an important medicinal plant. *Physiol. Mol. Boil. Plant* 15: 79-86.
- Latif Z, Nasir IA, Riazuddin S (2007). Indigenous production of synthetic seeds in *Daucus carota*. *Pak. J. Bot.* 39: 849-855.
- Lee EK, Cho DY, Soh WY (2001). Enhanced production and germination of somatic embryos by temporary starvation in tissue cultures of *Daucus carota*. *Plant Cell Rep.* 20:408-415.
- Liu JR, Jeon J, Yang SG, Lee HS, Song NH, Jeong WJ (1992). Dry type of carrot (*Daucus carota* L.) artificial seeds. *Sci. Hortic.* 51:1-11.
- Ma XM, Wu CF, Wang GR (2011). Application of artificial seeds in rapid multiplication of *Pseudostellaria heterophylla* Afr. *J. Biotechnol.* 10:15744-15748
- Machii H (1992). *In vitro* growth of adventitious buds in Mulberry, *Morus alba* L. *Jpn. J. Breed.* 42:553-559.
- Makowczynska J, Andrzejewska-golec E (2006). Somatic seeds of *Plantago asiatica* L. *Acta Soc. Bot. Poloniae*, 75:17-21.

- Malemnganba H, Ray BK, Bhattacharyya S, Deka PC (1996). Regeneration of encapsulated protocorms of *Phaius tankervilleae* stored at low temperature. Indian J. Exp. Biol. 34:802-805.
- Malek MA (2009). *In vitro* propagation of pointed gourd (*Trichosanthes dioica* Roxb.) through encapsulated shoot tips. Bangladesh J. Agric. Res. 34:555-563.
- Mamiya K, Sakamoto Y (2001). A method to produce encapsulatable units for synthetic seeds in *Asparagus officinalis*. Plant Cell Tiss. Org. Cult. 64:27-32.
- Mandal J, Pattnaik S, Chand PK (2000). Alginate encapsulation of axillary buds of *Ocimum americanum* L (Hoary Basil), *O. basilicum* L. (Sweet Basil), *O. gratissimum* L. (Shrubby Basil), and *O. sanctum* (Sacred Basil), *In Vitro Cell Dev. Biol. Plant* 36:267-292.
- Manjikhola S, Uppeandra D, Meena J (2005). Organogenesis, embryogenesis and synthetic seed production in *Arnebia euchroma* a critically endangered medicinal plant of the Himalaya. *In Vitro Cell Dev. Biol. Plant* 41: 244-248.
- Maqsood M, Mujib A, Siddiqui ZH (2012). Synthetic seed development and conversion to plantlet in *Catharanthus roseus* (L.) G. Don. *Biotechnology* 11:37-43.
- Mariani P (1992). Eggplant somatic embryogenesis combined with synthetic seed technology. *Capsicum Newslett.* pp. 289-294.
- Mariani TS, Latif S, Ginting G, Miyake H (2008) Somatic embryogenesis of oil palm (*Elaeis guineensis* Jacq.) for synthetic seed production. <http://www.sith.itb.ac.id/publikasi-totik/oil%20palm%20202.pdf>
- Marsolais AA, Wilson DPM, Tsujita MJ, Senaratna T (1991). Somatic embryogenesis and artificial seed production in Zonal (*Pelargonium x hortorum*) and Regal (*Pelargonium x domesticum*) geranium. *Can. J. Bot.* 69:1188-1193.
- Martin KP (2003). Clonal propagation, encapsulation and reintroduction of *Ipsea malabarica* (Reichb.f) Hook JD, An endangered orchid. *In vitro Cell Dev. Biol. Plant* 39:322-326.
- Maruyama E, Kinoshita I, Ishii K, Ohba K, Saito A (1997a). Germplasm conservation of the tropical forest trees, *Cedrela odorata* L., *Guazuma crinita* Mart. and *Jacaranda mimosaeifolia* D. Don by shoot tip encapsulation in calcium alginate and storage at 12-25° C. *Plant Cell Rep.* 16:393-396.
- Maruyama E, Kinoshita I, Ishii K, Shigenaga H, Ohba K (1997b). Alginate encapsulation technology for the propagation of the tropical forest trees: *Cedrela odorata* L., *Guazuma crinita* Mart., *Jacaranda mimosaeifolia* D. Don. *Silvae. Genet.* 46:17-23.
- Mathur J, Ahuja PS, Lal N, Mathur AK (1989). Propagation of *Valeriana wallichii* DC. using encapsulated apical and axial shoot buds. *Plant Sci.* 60:111-116.
- Matsumoto K, Hirao C, Teixeira J (1995). *In vitro* growth of encapsulated shoot tips in Banana (*Musa* Sp.). *Acta Hort.* 370:13-20
- McKersie BD, Brown DCW (1996). Somatic embryogenesis and artificial seeds in forage legumes. *Seed Sci. Res.* 6:109-126.
- McKersie BD, Senaratna T, Bowley SR, Brown DCW, Krochko JE, Bewley JD (1989). Application of artificial seed technology in the production of Hybrid alfalfa (*Medicago sativa* L.). *In vitro Cell Dev. Biol. Plant* 25:1183-1188.
- McKersie BD, Senaratna T, Bowley SR, Brown DCW, Kielly A, Krochko JE, Bewley JD (1990). Artificial seeds application in the production of hybrid alfalfa (*Medicago sativa* L.) in: *Abstr.Viith Int. Cong plant tissue culture.* Amsterdam p. 259.
- McKersie BD, Bowley SR (1993). Synthetic seeds of alfalfa. In K. Redenbaugh (ed), *Synthetic seeds; Application of synthetic seeds to crop improvement.* CRC press. Boca Raton. FL. pp. 231-255.
- Micheli M, Hafiz IA, Standardi A (2007). Encapsulation of *in vitro* derived explants of olive (*Olea europaea* L. cv. Moraiolo): effect of storage on capsule and derived shoot performance. *Sci. Hortic.* 113: 286-292.
- Micheli M, Pellegrino S, Piccioni E, Standardi A (2002). Effect of double encapsulation and coating on synthetic seed conversion in M 26 apple rootstock. *J. Microencapsulation* 19: 347-356.
- Mishra J, Singh M, Palni LMS, Nandi SK (2011). Assessment of genetic fidelity of encapsulated micro shoots of *Picrorhiza kurrooa*. *Plant Cell Tiss. Organ Cult.* 104: 181-186.
- Mohanraj R, Ananthan R, Bai VN (2009). Production and storage of synthetic seeds in *Coelogyne breviscapa* Lindl. *Asian J. Biotechnol.* 1:124-128.
- Molle K, Dupis JM, Ducos JP, Ansepm A, Croios-savidan I, Petiard V, Freyssinet G (1993). In K. Redenbaugh (ed), *Synthetic seeds. Application of synthetic seeds to crop improvement.* CRC Press pp. 257-270.
- Mondal PK, Bhattacharya A, Sood A, Ahuja PS (2002). Propagation of tea (*Camella sinensis* (L.) O. Kuntze ) by shoot proliferation of alginate-encapsulated axillary bud stored at 4°C. *Curr. Sci.* 83:941-944.
- Mukunthakumar S, Mathur J (1992). Artificial seed production in the male bamboo *Dendrocalamus strictus* L. *Plant Sci.* 87:109-113.
- Muralidharan EM, Mascarenhas AF (1995). Somatic embryogenesis *Eucalyptus*. In *Somatic embryogenesis In: Woody plants*, Mohan SJ, Pramod GK, Ronald JN (Ed), Kluwer Academic Publishers, The Netherlands 2:23-39.
- Murashige T (1977). Plant cell and organ culture as horticultural practice. *Acta Hort.* 78:17-30.
- Murthy KSR, Raghu Ramulu D, Chandrashekar J, Emmanuel, Pullaiah T (2006). *In vitro* flowering of *Spathoglottis plicata* Bl. (Orchidaceae). *Phytomorphology* 53: 117-120.
- Nagananda GS, Satishchandra N, Ragath S (2011). Regeneration of encapsulated protocorm like bodies of medicinally important vulnerable orchid *Flickingeria nodosa* (Dalz.) Seidenf. *Int. J. Bot.* 7:310-313.
- Naik GR, Chikkagouda NR (1997). Production and regeneration of synthetic seeds in sugarcane (*Saccharum officinarum* L.). *Indian Sugar* pp. 130-135.
- Naik SK, Chand PK (2006). Nutrient alginate encapsulation of *in vitro* nodal segments of Pomegranate (*Punica granatum* L.) for germplasm distribution and exchange. *Sci. Hortic.* 108:247-252.
- Nair RR, Gupta DS (2007). *In vitro* plant regeneration from encapsulated somatic embryos of black pepper (*Piper nigrum* L.). *J. Plant Sci.* 2: 283-292.
- Nakashimada Y, Uozumi N, Kobayashi T (1995). Production of plantlets for use as artificial seeds from horseradish hairy roots fragmented in a blender. *J. Ferment. Bioeng.* 79: 458-464.
- Narula A, Kumar S, Srivastava PS (2007). Genetic fidelity of *in vitro* regenerants, encapsulation of shoot tips and high diosgenin content in *Dioscorea bulbifera* L. a potential alternative source of diosgenin. *Biotechnol. Lett.* 29:623-629.
- Nassar AH (2003). Slow growth storage of encapsulated germplasm of *Coffea arabica* L. *Int. J. Agric. Biol.* 5:517-520.
- Nhut DT, Duy N, Ha-vy NN, Khue CD, Khiem DV, Hang NTT and Vinh DN (2004). Artificial seeds for propagation of *Anthurium 'TROPICAL'*. *J. Agric. Sci. Technol.* 4:73-78.
- Nhut DT, Tien TNT, Huong MTN, Hien NTT, Huyen PX, Luan VQ, Silva JATD (2005). Artificial seeds for propagation and preservation of *Cymbidium* spp. *Propagation Ornamental Plants* 5:67-73.
- Nieves N, Lorenzo JC, Blanco MA, Gonzalez J, Perralta H (1998). Artificial endosperm Cleopatra tangerine zygotic embryos, a model for somatic embryo encapsulation. *Plant Cell Tiss. Org. Cult.* 54:77-83.
- Nieves W, Zambrano Y, Tapia R, Cid M, Pina D, Castillo R (2003). Field performance of artificial seeds derived sugarcane plants. *Plant Cell Tiss. Org. Cult.* 75:279-282.
- Nirala NK, Das DK, Reddy MK, Srivastava PS, Sopory SK, Upadhyaya KC (2010). Encapsulated somatic embryos of grape (*Vitis vinifera* L.): An efficient way for storage, transport and multiplication of pathogen free plant material. *Asian. Pac J. Mol. Biol. Biotechnol.* 8:159-162.
- Nomura K, Komamine A (1985). Identification and isolation of single cells that produce somatic embryos at a high frequency in a carrot suspension culture. *Plant Physiol.* 79:988-991.
- Nunes EDC, Benson EE, Ultramari AC, Araujo PS, Moser JR, Viana AM (2003). *In vitro* conservation of *Cedrela fissilis* Vellozo (Meliaceae), A native tree of the Brazilian Atlantic forest. *Biodivers. Conserv.* 12:837-848.
- Nyende AB, Schittenhelm S, Mix-Wagner G, Greef J (2003). Production, storability, and regeneration of shoot tips of potato (*Solanum tuberosum* L.) encapsulated in calcium alginate hollow beads. *In vitro Cell. Dev. Biol. Plant* 39: 540-544.
- Nyende AB, Schittenhelm S, Mix-Wagner G, Greef JM (2005). Yield and canopy development of field grown potato plants derived from

- synthetic seeds. *Eur. J. Agron.* 22:175-184.
- Onay A, Jeffree CE, Yeoman MM (1996). Plant regeneration from encapsulated embryoids and embryogenic masses of pistachio, *Pistacia vera* L. *Plant Cell Rep.* 15:723-726.
- Onishi N, Mashiko T, Okamoto A (1992). Cultural system producing encapsulatable units of synthetic seeds in celery. *Acta. Hortic.* 319: 113-118.
- Onishi N, Sakamoto Y, Hirosawa T (1994). Synthetic seeds as an application of mass propagation of somatic embryos. *Plant Cell Tiss. Org. Cult.* 39:137-145.
- Ozden TY, Carlo AD, Gumusel F, Pigmattelli S, Lambardi M (2008). Development of encapsulation techniques for the production and conservation of synthetic seeds in ornamental plants. *Propagation Ornamental Plants* 8:17-22.
- Padmaja G, Reddy LR, Reddy GM (1995). Plant regeneration from synthetic seeds of groundnut, *Arachis hypogaea* L. *Indian J. Exp. Biol.* 33:967-971.
- Pandey A, Chand S (2005). Efficient plant regeneration from encapsulated somatic embryos of *Hyoscyamus muticus* L. *Indian J. Biotechnol.* 4:546-550.
- Patel AV, Pusch I, Mix-Wagner G, Vorlop KD (2000). A novel encapsulation technique for the production of artificial seeds. *Plant Cell Rep.* 19: 868-874
- Pattnaik SK, Chand PK (2000). Morphogenic response of the alginate-encapsulated axillary buds from *in vitro* shoot cultures of six mulberries. *Plant Cell Tiss. Org. Cult.* 60:177-185.
- Pattnaik SK, Sahoo Y, Chand PK (1995). Efficient plant retrieval from alginate encapsulated vegetative buds of mature Mulberry trees. *Sci Hortic.* 61: 227-239.
- Pereira JES, Guede RDS, Costa FHDS, Schmitz CCB (2008). Composicao da matriz de encapsulamento na formacao e conversao de sementes sinteticas de pimento-longa. *Horticultura Brasileira* 26: 63-96.
- Phonkajornyod P, Pawelzik E, Vearasilp S (2004). Dry synthetic seed production and desiccation tolerance induction in somatic embryo of sweet pepper. *Deutscher Tropentag*, October 5-7, Berlin.
- Piccioni E (1997). Plantlets from encapsulated micropropagated buds of M 26 apple rootstock. *Plant Cell Tiss. Org. Cult.* 47: 255-260.
- Piccioni E, Standardi A (1995). Encapsulation of micropropagated buds of six woody species. *Plant Cell Tiss. Org. Cult.* 42: 221-226.
- Pinker I, Abdel-rahman SSA (2005). Artificial seeds for propagation of *Dendranthema x Grandiflora* (RAMAT). *Propagation Ornamental Plants* 5: 186-191.
- Preece JE, West TP (2006). Greenhouse growth and acclimatization of encapsulated *Hibiscus moschatus* nodal segments. *Plant Cell Tiss. Organ Cult.* 87:127-138.
- Priya BT, Arumugam S (2003). Synthetic seed production in banana. *Adv. Plant Sci.* 16:219-222.
- Rady MR, Hanafy MS (2004). Synthetic seed technology for encapsulation and regrowth of *in vitro* derived *Gypsophila paniculata* L. shoot tips. *Arab J. Biotechnol.* 7:251-264.
- Rai MK, Asthana P, Singh SK, Jaiswal VS, Jaiswal U (2009). The encapsulation technology in fruit plants - A review. *Biotechnol. Adv.* 27: 671-679.
- Rai MK, Jaiswal VS (2008). Synthetic seeds of guava (*Psidium guajava* L.) from somatic embryos and plant regeneration. In: Arya ID, Arya Sarita, editors. *Utilization of biotechnology in plant sciences*. Dehradun, India: Microsoft Printech (I) Pvt. Ltd., pp. 217-225.
- Rai MK, Jaiswal VS, Jaiswal U (2008a). Encapsulation of shoot tips of guava (*Psidium guajava* L.) for short-term storage and germplasm exchange. *Sci. Hortic.* 118:33-38.
- Rai MK, Jaiswal VS, Jaiswal U (2008b). Alginate-encapsulation of nodal segments of guava (*Psidium guajava* L.) for germplasm exchange and distribution. *J. Hortic. Sci. Biotechnol.* 83:569-573.
- Rai MK, Jaiswal VS, Jaiswal U (2008c). Effect of ABA and sucrose on germination of encapsulated somatic embryos of guava (*Psidium guajava* L.). *Sci. Hortic.* 117:302-305.
- Ramakrishnappa (1998). Studies on somatic embryogenesis and production of synthetic seeds in *Gladiolus* cultivars. Ph. D. dissertation, University of Agriculture Science, Bangalore.
- Ramesh M, Marx R, Mathan G, Pandian SK (2009). Effect of bavistin on *in vitro* plant conversion from encapsulated uninodal microcuttings of micropropagated *Bacopa monnieri* (L) – An Ayurvedic Herb. *J. Environ. Biol.* 30:441-444.
- Rao PS, Ganapathi TR, Suprasanna P, Bapat VA (1993). Encapsulated shoot tips of banana a new propagation and delivery system. *Infomusa* 2:4-5.
- Rao PV, Singh B (1991). Plant regeneration from encapsulated somatic embryos of *Solanum melongena* L. *Plant Cell Rep.* 10:7-11.
- Ray A, Bhattacharya S (2008). Storage and plant regeneration from encapsulated shoot tips of *Rauvolfia serpentina*– an effective way of conservation and mass propagation. *South Afr. J. Bot.* 74:776-779.
- Reddy GS, Kumar DD, Babu RS, Madhav M (2005). Callus culture and synthetic seed production in *Rauvolfia serpentina*. *Indian J. Hortic.* 62: 102-103.
- Redenbaugh K (1993). Synthetic seeds. Application of synthetic seeds to crop improvement. CRC press. Boca Raton. FL. p. 481.
- Redenbaugh K, Fujii JA, Slade D, Viss PR, Kossler M (1991a). Artificial seeds- encapsulated somatic embryos. In: Bajaj YPS (ed), *Biotechnology in Agriculture and Forestry*. High-Tech and micropropagation. Springer. Berlin. 2:395-416.
- Redenbaugh K, Fujii JA, Slade D (1993). Hydrated coatings for synthetic seeds. In K. Redenbaugh (eds) *synseeds Application of synthetic seeds to crop improvement*. CRC press. Boca Raton. FL. pp. 35-46.
- Redenbaugh K, Fujii JA, Slade D (1991b). Synthetic seed technology. In I.K. Vasil (ed), *Cell culture and somatic cell genetics of plants, scale up and automation in plant propagation in plants*. Academic press. San diego, Ca. 8:35-74.
- Redenbaugh K, Nichol J, Kossler ME, Paasch B (1984). Encapsulation of somatic embryos for artificial seed production. *In vitro Cell Dev. Biol.* *Plant* 20:256-257.
- Redenbaugh K, Paasch BD, Nichol JW, Kossler ME, Viss PR, Walker KA (1986). Somatic seeds: encapsulation of asexual plant embryos. *Biotechnology* 4:797-801.
- Redenbaugh K, Slade D, Viss PR, Fujii J (1987). Encapsulation of somatic embryos in synthetic seed coats. *Hort. Sci.* 22:803-809.
- Redenbaugh K, Walker K (1990). Role of artificial seeds in alfalfa breeding. In S. S. Bhojwani (ed), *Plant tissue culture: Applications and Limitations*. Elsevier Amsterdam. pp. 102-135.
- Refouvelet E, Nours SL, Tallon C, Daguin F (1998). A new method for *in vitro* propagation of lilac (*Syringa vulgaris* L.): Regrowth and storage conditions for axillary buds encapsulated in alginate beads, development of a pre-acclimatization stage. *Sci. Hort.* 74:233-241.
- Repunte VP, Taya M, Tone S (1995). Conservation of root regeneration potential of cell aggregates from horseradish hairy roots used as artificial seeds. *J. Chem. Eng. Jpn.* 29:874-880.
- Romay AC, Gardi T, Standardi A (2002). Plantlets from encapsulated *in vitro*-derived microcuttings of kiwifruit (cv top Star®). *Agric Mediterr.* 132:246-252.
- Romay AC, Gardi T, Standardi A (2003). Preliminary experiments on the conversion of encapsulated *in vitro*-derived microcuttings of raspberry (*Rubus idaeus* L.). *Italus Hortus* 10:29-34.
- Roy B, Mandal AB (2008). Development of synthetic seeds involving androgenic and pro-embryos in elite indica rice. *Indian J. Biotechnol.* 7:515-519.
- Ruffoni B, Giovannini A, Semeria L, Massabò F, Costantino C, Allavena A (1993). Tissue culture in *Lasianthus*: a tool to create genetic variation. In: Schiva T and Mercuri A (eds.), *Proceedings of the XVIIth Eucarpia Symposium "Creating genetic variation in ornamentals"*, Ist. Sper. Floric., Sanremo pp. 289-295.
- Ruffoni B, Massabò F, Giovannini A (1994). Artificial seed technology in ornamental plants, *Lasianthus* and *Genista*. *Acta Hortic.* 362: 297-304.
- Saiprasad GVS (2001). Artificial seeds and their applications. *Resonance* 6:39-46.
- Saiprasad GVS, Polisetty R (2003). Propagation of three orchid genera using encapsulated protocorm like bodies. *In vitro Cell Dev. Biol. Plant* 39:42-48.
- Sajina A, Mino D, Geetha SP, Samsuddin K, Rena J, Nirmal babu K, Ravindran PN (1997). Production of synthetic seed in few spice crops. In Edition, Ramanna KV, Sasikumar B, Nirmal Babu K and Santosh J (eds), *Biotechnology of spices, medicinal and aromatic plants*. Eapen. Kerala. India. Indian society of spices. pp. 65-69.

- Sakamoto Y, Mashiko T, Suzuki A, Kawata H, Iwasaki A (1992). Development of encapsulation technology for synthetic seeds. *Acta Hort.* 319:71-76.
- Sandavol-Yugar EW, Vesco LLD, Steinmacher DA, Stolf EC, Guerra MP (2009). Microshoots encapsulation and plant conversion of *Musa* sp. cv. grand naine. *Ciencia Rural* 39:998-1004.
- Sandeep B (2008). Development and evaluation of synthetic seeds in sugarcane. Masters dissertation. University of Agricultural Sciences. Dharwad.
- Sarkar D, Naik PS (1998a). Nutrient encapsulation of potato nodal segments for germplasm exchange and distribution. *Biologia Plantarum* 40:285-290.
- Sarkar D, Naik PS (1998b). Synseeds in potato. An investigation using nutrient-encapsulated *in vitro* nodal segments. *Sci. Hortic.* 73:179-184.
- Sarmah DK, Borthakur M, Borua PK (2010). Artificial seed production from encapsulated PLBs regenerated from leaf base of *Vanda coerulea* Griff. ex Lindl. an endangered orchid. *Curr. Sci.* 98:686-690.
- Schafer-menuhr A, Mix-wagner G, Vorlop KD (2003). Regeneration of plants from cell suspension cultures and encapsulated cell suspension cultures of *Solanum tuberosum* L. cv *clarissa*. *Landbauforschung Volkenrode* 53:53-59.
- Schuller A., Reuther G, Geiger T (1989). Somatic embryogenesis for seeds explant of *Abies alba*. *Plant Cell Tiss. Org. Cult.* 17:53-58.
- Senaratna T, McKersie BD, Bowley SR (1990). Artificial seeds of alfalfa (*Medicago sativa* L.). Induction of desiccation tolerance in somatic embryos. *In vitro Cell Dev. Biol. Plant* 26:85-90.
- Senaratna T (1992). Artificial seeds. *Biotechnology advances*, 10:379-392.
- Senaratna T, McKersie BD, Bowley SR (1989). Desiccation tolerance of alfalfa (*Medicago sativa* L.) somatic embryogenesis: influence of abscisic acid, stress pretreatments and drying rates. *Plant Sci.* 65:253-259.
- Shaohu T, Min S, Kunpei L (1994). Studies on artificial seed of *Ipomoea aquatica* Forsk. Shoot performance. *Sci. Hortic.* 113:286-292.
- Sharma A, Tandon P, Kumar A (1992). Regeneration of *Dendrobium wardianum* Warner (Orchidaceae) from synthetic seeds. *Indian J. Exp. Biol.* 30:747-748.
- Sharma TR, Singh BM, Chauhan RS (1994). Production of disease free encapsulated buds of *Zingiber officinale* Rosc. *Plant Cell Rep.* 13:300-302.
- Shigeta J, Sato S (1994). Plant regeneration and encapsulation of somatic embryos of horseradish. *Plant Sci.* 102:109-115.
- Sicurani M, Piccioni E, Standardi A (2001). Micropropagation and preparation of synthetic seed in M 26 apple rootstock: Attempts towards saving labor in the production of adventitious shoot tips suitable for encapsulation. *Plant Cell Tiss. Org. Cult.* 66:207-216.
- Singh AK, Chand S (2010). Plant regeneration from alginate encapsulated somatic embryos of *Dalbergia sissoo* Roxb. *Indian J. Biotechnol.* 9:319-324.
- Singh AK, Sharma M, Varshney R, Agarwal SS, Bansal KC (2006a). Plant regeneration from alginate encapsulated shoot tips of *Phyllanthus amarus* Schum and Thom. A medicinally important plant species. *In vitro Cell Dev. Biol. Plant* 42:109-113.
- Singh AK, Varshney R, Sharma M, Agarwal SS, Bansal KC (2006b). Regeneration of plants from alginate encapsulated shoot tips of *Withania somnifera* (L.) Dunal. a medicinally important plant species. *J. Plant Physiol.* 163:220-223.
- Singh B, Sharma S, Rani G, Virk GS, Zaidi AA, Nagpal A (2007). *In vitro* response of encapsulated and non encapsulated embryos of Kinnow mandarin (*Citrus nobilis* Lour x *C. Deliciosa* Tenora). *Plant Biotechnol. Rep.* 1:101-107.
- Singh F (1991). Encapsulation of *Spathoglottis plicata* protocorms. *Lindleyana* 6:61-63.
- Soneji JR, Rao PS, Mhatre M (2002). Germination of synthetic seeds of pineapple (*Ananas comosus* (L.) Merr. *Plant Cell Rep.* 20:891-894.
- Srinivas L (2002). Morphogenic studies *in vitro* in banana (*Musa* spp). Masters dissertation, University of Mumbai. Mumbai. India.
- Srivastava V, Shamshand A, Khan, Banerjee S (2009). An evaluation of genetic fidelity of encapsulated Microshoots of the medicinal plant: *Cineraria maritima* following six months of storage. *Plant Cell Tiss. Org. Cult.* 99:193-198.
- Standardi A, Micheli M, Piccioni E (1995). Incapsulamento in alginato di espanti micropropagati. *Italus Hortus* 2:46-52.
- Standardi A, Piccioni E (1997). Rooting induction in encapsulated buds of M26 apple rootstock for synthetic seeds. Seed production; *Plant Physiol. Biochem.* 56:309-314.
- Stephen R, Jayabalan N (2000). Artificial seed production in coriander (*Coriandrum sativum* L.). *Plant Tissue Cult.* 10:45-49.
- Sudhakara K, Nagaraj BN, Santosh kumar AV, Sunil kumar KK, Vijaya kumar NK (2000). Studies on the production and storage potential of synthetic seeds in cocoa (*Theobroma cacao*). *Seed Res.* 28:119-125.
- Sundararaj SG, Agrawal A, Tyagi RK (2010). Encapsulation for *in vitro* short-term storage and exchange of ginger (*Zingiber officinale* Rosc.) germplasm. *Sci. Hortic.* 125:761-766.
- Sunilkumar KK, Sudhakara K, Vijaykumar NK (2000). An attempt to improve storage life of *Hopea parviflora* seeds through synthetic seed production. *Seed Res.* 28:126-130.
- Suprasanna P, Ganapathi TR, Rao PS (1996). Artificial seeds in rice (*Oryza sativa* L.): Encapsulation of somatic embryos from mature embryo callus culture. *Asia Pac. J. Mol. Biol. Biotechnol.* 4:90-93.
- Suprasanna P, Anupama S, Ganapathi TR, Bapat VA (2000). *In vitro* growth and development of encapsulated shoot tips of different banana and plantation cultivars. *J. New Seeds* 3:19-25.
- Suprasanna P, Bharati G, Ganapathi TR, Bapat VA (2002). *In vitro* development of encapsulated somatic embryos in rice. *Trop. Agric. Res. Extension* 5:76-78.
- Swamy MK, Balasubramanya S, Anuradha M (2009). Germplasm conservation of Patchouli (*Pogostemon cablin* Benth.) by encapsulation of *in vitro* derived nodal segments. *Int. J. Biodivers. Conserv.* 1:224-230.
- Tabassum B, Nasil IA, Farooq AM, Latif Z, Husmain T (2010). Viability assessment of *in vitro* produced synthetic seeds of cucumber. *Afr. J. Biotechnol.* 9:7026-7032.
- Taha RM, Daud N, Hasbullah NA, Awal A (2009b). Somatic embryogenesis and production of artificial seeds in *Saintpaulia ionantha* Wendl. *Acta Hort.* 829:331-336.
- Taha RM, Hasbullah NA, Awal A (2009a). Production of synthetic seeds from microshoots and somatic embryos of *Gerbera jamesonii* Bolus ex. Hook. f. *Acta Hort.* 829:91-98.
- Tan TK, Loon WS, Jhor E, Loh CS (1998). Infection of *Spathoglottis plicata* (Orchidaceae) seeds by mycorrhizal fungus. *Plant Cell Rep.* 18: 14-19.
- Tsai CJ, Saunders JW (1999). Encapsulation, germination, and conversion of somatic embryos in sugarbeet. *Encapsulated Somatic Embryos.* 36:11-36.
- Tejavathi DH, Gayathamma K, Sowmya R (2006). Production of plantlets from encapsulated *in vitro* shoot buds and somatic embryos of *Agave vera-cruz* Mill. *Plant Cell Biotechnol. Mol. Biol.* 7: 183-186.
- Thobunluepop P, Pawelzik E, Vearasilp S (2009). Possibility of sweet corn synthetic seed production. *Pak. J. Biol. Sci.* 12:1085-1089.
- Timbert R, Barbotin JN, Kersulec A, Bazinet C, Thomas D (1995). Physico-chemical properties of the encapsulation matrix and germination of carrot somatic embryos. *Biotechnol. Bioeng.* 46:573-578.
- Timbert R, Barbotin JN, Thomas D (1996). Enhancing carrot somatic embryos survival during slow dehydration, by encapsulation and control of dehydration. *Plant Sci.* 120:215-222.
- Tsvetkov I, Jouve L, Hausman JF (2006). Effect of alginate matrix on *in vitro* derived encapsulated apical microcuttings of hybrid aspen. *Biologia Plantarum*, 50:722-724.
- Ummi KH, Dayana AD, Norahimah AR, Natrisya Nur AR, Firdaus A (2011). Effects of different storage duration on encapsulated *Parkia speciosa* zygotic embryo germination. *Empowering Science, Technology and Innovation Towards a Better Tomorrow* 120: 758-763
- Uozumi N, Nakashimada Y, Kato Y, Kobayashi T (1992). Production of artificial seeds from horseradish hairy roots. *J. Ferment. Bioeng.* 73: 21-26.
- Van der Valk P, Zaal MACM, Creemers-Molannar J (1989). Somatic embryogenesis and plant regeneration in inflorescence and seed derived callus cultures of *Poa pratensis* L. (Kentucky blue-grass). *Plant Cell Rep.* 7: 644-647 .

- Vdovitchenko MY, Kuzovkina LN (2011). Artificial seed preparation as the efficient method for the storage and production of healthy cultured roots of medicinal plants. *Russian J. Plant Physiol.* 58:461-468.
- Verma SK, Raj MK, Asthana P, Jaiswal VS, Jaiswal U (2010). *In vitro* plantlets from alginate- encapsulated shoot tips of *Solanum nigrum* L. *Sci. Hort.* 124:517-521.
- Vij SP, Kaur P, Gupta A (2001). Synseeds and their utility in orchids: *Dendrobium densiflorum* Lindl. *Phytomorphology* 51:159-161.
- Wake H, Akasaka A, Umetsu H, Ozeki Y, Shimamura K, Matsunaga T (1992). Enhanced germination of artificial seeds by marine Cyanobacterial extract. *Appl. Microbiol. Biotechnol.* 36:684-688.
- Wang W, Wang S, Wu X, Jin X, Chen F (2007). High frequency plantlet regeneration from callus and artificial seed production of rock plant *Pogonatherum paniceum* (Lam.) Hack. (Poaceae). *Sci. Hort.* 113:196-201.
- Williams EG, Maheswaran G (1986). Somatic embryogenesis: Factors influencing coordinated behaviour of cells as an embryogenic group. *Ann. Bot.* 57:443-462.
- Winkelmann T, Meyer L, Serek M (2004). Germination of encapsulated somatic embryos of *Cyclamen persicum*. *Hort. Sci.* 39:1093-1097.
- Yu X, Liua H, Tongb L (2008) Feeding scenario of the silkworm *Bombyx mori* L. *Acta Astronautica* 63:1086-1092.
- Zhang M, Peng S, Yang X, Xu L (2009). Preparation technique of *Cremastra appendiculata* synthetic seed. (in Chinese with English abstract), China. *J. Chinese Materia. Med.* 34:1894-1897.
- Zych M, Furmanowa M, Krajewska-patan A, Lowicka A, Dreger M, Mendlewska S (2005). Micropropagation of *Rhodiola kirilowii* plants using encapsulated axillary buds and callus. *Acta Biologica Cracoviensia Series Botanica* 47:83-87.