

Full Length Research Paper

Ginger extract (*Zingiber officinale*) triggers apoptosis and G₀/G₁ cells arrest in HCT 116 and HT 29 colon cancer cell lines

Shailah Abdullah¹, Siti Amalina Zainal Abidin¹, Noor Azian Murad², Suzana Makpol¹, Wan Zurinah Wan Ngah¹ and Yasmin Anum Mohd Yusof^{1*}

¹Department of Biochemistry, Faculty of Medicine, Universiti Kebangsaan Malaysia, Jalan Raja Muda Abdul Aziz, 50300 Kuala Lumpur, Malaysia.

²Centre of Lipids and Engineering and Applied Research, Universiti Teknologi Malaysia, Jalan Semarak, 50300 Kuala Lumpur, Malaysia.

Accepted 16 April, 2010

Although many studies have shown the antitumor properties of ginger extract (*Zingiber officinale*), little is known regarding the mechanism of its effects. This study was conducted to determine the mechanism of antitumor effects of ginger extract by evaluating apoptosis rate and cell cycle progression status in colon cancer cell lines HCT 116 and p53 defective HT 29. HCT 116 and HT 29 cells were cultured in the presence of ginger extract at various concentrations for 24 h. The percentage of cell viability was determined by 3-(4, 5-dimethylthiazol-2-yl)-2, 5-di phenyl tetrazolium bromide (MTT) assay. Our results showed that ginger extract inhibited proliferation of HCT 116 and HT 29 cells with an IC₅₀ of 496 ± 34.2 µg/ml and 455 ± 18.6 µg/ml, respectively. We also found that ginger extract at increasing concentrations induced apoptosis dose dependently in both colon cancer cells. Apoptosis rates were 11.15, 35.05 and 57.49% for HCT 116 and 4.39, 19.81 and 28.09% for HT 29 at 200, 500 and 800 µg/ml of ginger extract, respectively. Ginger extract arrested HCT 116 and HT 29 cells at G₀/G₁ and G₂/M phases with corresponding decreased in S-phase. This study suggests that ginger extract may exert its antitumor effects on colon cancer cells by suppressing its growth, arresting the G₀/G₁-phase, reducing DNA synthesis and inducing apoptosis.

Key words: *Zingiber officinale*, HCT 116, HT 29, G₀/G₁ phase, S phase, apoptosis.

INTRODUCTION

Colorectal cancer accounts for 10 to 15% of all cancers and is the second leading cause of cancer-related death in industrialized countries (Virginie et al., 2007). According to National Cancer Registry of Malaysia (Gerard and Halimah, 2003), colon cancer is recognized as the commonest cancer among men and the third most common cancer among women. Colon cancer occurrence is commonly ascribed to the transformation of normal colon epithelium to adenomatous polyps and ultimately invasive cancer. Since cancer is a disorder of

deregulated cell proliferation and cell survival (Evan and Vousden, 2001), inhibiting cell proliferation and increasing apoptosis in tumors are effective strategies for preventing tumor growth (Kim et al., 2005). Recent studies showed that curcumin, an active component of turmeric inhibits the growth of human colon cancer cells independent of COX-2 expression (Hanif et al., 1997). Lee et al. (2009) showed that thiosulfinates from *Allium tuberosum* L. inhibited cell proliferation and activated both the caspase-dependent and caspase-independent apoptotic pathways in colon cancer cells (Lee et al., 2009). Other study demonstrated that luteolin induced G₁ and G₂/M cell-cycle arrest and promoted apoptosis through down regulation of several antiapoptotic proteins in colon cancer cells.

*Corresponding author. E-mail: anum@medic.ukm.my. Tel: 603-4040 5297. Fax: 603-2693 8037.

Emerging evidence has demonstrated that many natural products isolated from plant sources possess antitumor properties (Sakamoto et al., 1991). Since nature has provided many effective anticancer agents, plant derived drug research has made significant progress in anticancer therapies. Many components of medicinal or dietary plants have been identified as possessing potential chemopreventive properties capable of inhibiting, retarding or reversing the multi-stage carcinogenesis process (Surh et al., 1998; Surh et al., 2002). Plants of the ginger family (*Zingiber officinale*) are one of the most heavily consumed dietary substances in the world (Surh, 1999). It is one of the most widely used spices and has been used in traditional oriental medicines for centuries. Its extract and major pungent compounds have been shown to exhibit a variety of biological activities (Wei et al., 2005). The oleoresin from rhizome of ginger contains pungent ingredients including gingerol, shogaol and zingerone (Surh et al., 1998). Ginger's active components have been reported to exhibit cancer-preventive activity in several experimental carcinogenesis models including skin carcinogenesis (Murillo et al., 2008; Katiyar et al., 1996). Ginger supplementation suppresses colon carcinogenesis in the presence of procarcinogen (DMH) (Manju and Nalini, 2005). Recent study showed that [6]-gingerol a major phenolic compound derived from ginger, inhibited TRAIL-induced NF- κ B activation by impairing the nuclear translocation of NF- κ B, suppresses cIAP expression and increased TRAIL-induced caspase-3/7 activation in gastric cancer cells. On the other hand, [6]-shogaol alone reduced viability by damaging microtubules, arrested cell cycle in G₂/M phase and reduced viability in a caspase-3/7-independent manner in gastric cancer cells (Kazuhiro, 2007). Lee et al. (1998) showed that [6]-gingerol exerted inhibitory effects on the cell viability and DNA synthesis while inducing apoptosis of human promyelocytic leukemia HL-60 cells (Lee and Surh, 1998). Ginger was also shown to have anticancer effect by inducing apoptosis in rat liver cancer cells via up-regulation of the expression of pro-apoptotic protein, caspase-8 and downregulation of the expression of anti-apoptotic protein Bcl-2 (Yasmin et al., 2008). Exposure of Jurkat human T cell leukemia cells to various ginger constituents galanals A and B (isolated from the flower buds of Japanese ginger) resulted in apoptosis mediated through the mitochondrial pathway (Miyoshi et al., 2003). β -Elemene, an anticancer drug extracted from the ginger plant, triggered apoptosis in non-small-cell lung cancer cells through a mitochondrial release of the cytochrome c-mediated apoptotic pathway (Wang et al., 2005).

We have selected *Z. officinale* in this study to determine its mechanism as a potential chemopreventive agent against colon cancer cells HCT 116 and HT 29. Our results demonstrated that the inhibition of colon cancer cell growth by ginger involves both interference with cell cycle progression and apoptotic event.

MATERIALS AND METHODS

Ginger extract

Z. officinale (ginger) crude extract was obtained by ethanol extraction as provided by Noor Azian Murad from Center for Lipids Engineering Applied Research (CLEAR), Universiti Teknologi Malaysia. The sequence of pre-treatment for ginger extract include: peeling, slicing, washing of ginger rhizomes followed by bleaching, blanching, drying and grinding prior to solvent extraction. The sample calculation to obtain the moisture loss percent based on the initial and final weight of the sample is given by the equation:

$$\% \text{ moisture loss} = \frac{\text{Initial weight} - \text{current sample weight}}{\text{Initial weight}} \times 100\%$$

The dried ginger was fibrous and tough and therefore grinding of samples was done manually by pounding using 'mortar'. The pounded samples were then shredded manually. The ground ginger was dried prior to extraction using rotary evaporator in ethanol (1 L) for 6 h. The solvent was removed under vacuum at 500 mbar in the first hour, followed by 400 and 300 mbar in the next two hours to yield oleoresin, a brown viscous liquid (9.80 g, 4.9%). Refractive index reading of pure oleoresin is 1.5100.

Cell culture and treatment

HCT 116 and HT 29 cell lines were obtained from the American Type Culture Collection (Rockville, MD USA) and were cultured in RPMI 1640 medium (Flowlab, Australia) supplemented with 10% Fetal Calf Serum (FCS) (PAA Laboratories GmbH), 100 U/ml of penicillin and streptomycin (Thermo Scientific) at 37°C in 5% CO₂. Rates of cells proliferation, apoptosis and cell cycle were performed when cells reached 70% confluence density. Ginger extract was dissolved in 0.01% DMSO and added to cell lines after an overnight incubation.

Cell viability assay: 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide (MTT) assay

For cell viability assay, 2×10^4 cells (HCT 116 and HT 29)/well were plated in 100 μ l of RPMI 1640 media. HCT 116 and HT 29 cells were incubated overnight at 37°C in humidified atmosphere of 5% CO₂ for cells attachment. Ginger extract was added at various concentrations ranging from 0, 10, 50, 100, 200, 500, 800, 1000 and 1500 μ g/ml after 24 h incubation. After 24 h incubation, MTT solution (2 mg/ml) was added to the plate at the final concentration of 0.5 mg/ml. The resulting MTT-products were determined by measuring the absorbance at 550 nm with ELISA reader. Each point represents the mean of triplicate experiments.

$$\% \text{ viability} = \frac{\text{Optical density of sample}}{\text{Optical density of control}} \times 100$$

Cell cycle assay

The HCT 116 and HT 29 cells were seeded into 25 cm² flask (1×10^5 cells/ml) and were incubated with 200, 500 and 800 μ g/ml ginger extract for 24 h. Following treatment, cells were trypsinized and incubated for 30 min at room temperature in staining solution consisting of propidium iodide (PI; 50 μ g/ml), sodium citrate (0.1%), Triton X-100 (0.1%) and DNase-free RNase (20 μ g/ml). Stained cells were then analyzed for DNA content by flow cytometer within




Figure 1. Effect of ginger extract on the proliferation of HCT 116 and HT 29 cells. Cells were treated with different concentration of ginger extract, incubated at 37°C and harvested after 24 hour. Cell viability was determined by the MTT assay. Data represent the mean $\pm$ S.D. (n = 3).

3 h of staining. Percentages of cells in each phase were calculated using Cell Modfit software programs (Becton Dickson, Canada).

Apoptosis–annexin V-FITC Staining

Apoptosis was detected with an annexin V-FITC kit (Beckton-Dickinson, Canada). All adhering and floating cells were harvested after incubation for 24 h with 200, 500 and 800 µg/ml ginger extract. Cells were collected, washed with ice-cold PBS and centrifuged. The cell pellet was resuspended in ice-cold binding buffer (100 mM HEPES/ NaOH, pH 7.5 containing 1.4 M NaCl and 25 mM CaCl₂) at a density of 1×10^6 cells per ml. Five hundred µl of this cell suspension was transferred to a 5 ml culture tube, to which 5 µl of Annexin V-FITC conjugate and 10 µl of propidium iodide were added. The cells were gently vortexed and incubated for 10 min at room temperature in the dark. The fluorescence of the cells was immediately determined by flow cytometry (Becton Dickson, Canada). The results were analysed using Cell Modfit software programs (Becton Dickson, Canada).

Statistical analysis

The experiments were repeated at least 3 times and the results were expressed as mean $\pm$ S.D. Statistical evaluation was done using the ANOVA (SPSS 16.0) where $p < 0.05$ was considered significant.

RESULTS

Figure 1 showed that ginger extract significantly inhibited the proliferation of both colon cancer cells in a dose-dependent manner. Proliferation of HCT 116 cells decreased when treated with ginger extract resulting in a 50% reduction at 496 ± 34.2 µg/ml, while HT 29 cells

showed significant decrease in proliferation with 50% reduction at 455 ± 18.6 µg/ml of ginger extract.

To elucidate the mechanisms involved in the reduction in cell number by ginger, we examined whether such reduction was associated with cytostatic effect due to changes in cell cycle progression. Figures 2a and 2b summarized the relative percentages of HCT 116 and HT 29 cells in each phase of the cell cycle, following a 24-h treatment with varying ginger extract concentrations (200, 500 and 800 µg/ml). The cell cycle alterations were concentration dependent. As shown in Figures 2a and 2b, in the absence of ginger extract, 48.82% HCT 116 cells and 54.22% of HT 29 were in the G₀/G₁-phase. However, treatment with ginger extract arrested both cells at G₀/G₁-phase: 63.15% for HCT 116 and 70.98% for HT 29 at 200 µg/ml. Interestingly, we found a unique characteristic whereby higher doses of ginger extract (800 µg/ml) did not arrest cells at G₀/G₁-phase but instead they were arrested at G₂/M-phase. The G₀/G₁ and G₂/M arrests were accompanied by corresponding reduction in the percentages of cells in the S-phase as shown in Figure 2a and 2b. Reduction in DNA synthesis was dependent on the dose of ginger extract: 15.06, 9.52 and 8.66%, when treated with 200, 500 and 800 µg/ml of ginger extract for HCT 116 cells and 8.81, 14.63 and 3.21% for HT 29 cells when treated with the same concentrations of ginger extract, respectively.

Figure 3a showed apoptotic effect of ginger extract on both HCT 116 and HT 29 cells using double staining method FITC-conjugated annexin V and PI. Ginger treatment increased the number of early and late apoptotic cells, respectively. Ginger extract at increasing concentrations induced higher rate of poptosis




Figure 2a. Cell cycle profile of HCT 116 colon cancer cells in the presence of ginger extract. After a 24 h incubation with varying ginger extract concentrations. DNA content was evaluated with propidium iodide (PI) staining and fluorescence measured and analyzed. The values are the means $\pm$ SD of three independent experiments. a; significant compared to G0/G1 phase of control cell, b; significant compared to S phase of control cell, c; significant compared to G2/M phase of control cells..




Figure 2b. Distribution of cell cycle phases of HT 29 cells after treatment with ginger extract. After a 24 hr incubation with varying ginger extract concentrations (0, 200, 500 and 800 μ g/ml), DNA content were evaluated with propidium iodide staining and fluorescence measured and analyzed. The values are the means $\pm$ SD of three independent experiments. a; significant compared to G0/G1 phase of control cell, b; significant compared to S phase of control cell, c; significant compared to G2/M phase of control cells.




Figure 3a. Induction of apoptosis by ginger extract in human colon cancer cell lines, HT 29 and HCT 116. Percentages of apoptotic cells after a 24 h incubation with varying ginger extract concentrations. a: significant compared to control HT 29, b: significant compared to control HCT 116.

significantly ($p < 0.05$) compared to control for both HCT 116 and HT 29 cells as shown by cytograms of apoptosis for HCT 116 (Figure 3b) and HT 29 cells (Figure 3c).

DISCUSSION

There has been substantial interest in the search for potential chemopreventive agents in the past years for treatment against many cancers. Understanding how dietary components regulate proliferation and cell survival could play a critical role in the development of new agents that can prevent and treat cancer with low toxicity (Lee and Surh, 1998). Cancer chemoprevention was defined as the utilization of chemically active compounds to reverse, suppress and prevent progression of disease from pre-invasive cancer to frank malignancy (Sporn et al., 1976).

Numerous diet-derived agents are included among promising agents and agent combination that are being evaluated clinically as chemopreventive agents for major cancer targets including breast, prostate, colon and lung cancers (Kelloff et al., 2000). Ginger (*Z. officinalis* Roscoe, Zingiberaceae) is not only widely used as a dietary condiment but it has also been extensively utilized as a traditional oriental medicine (Taraphdar et al., 2001). Gingerols (that is, 6-gingerol, 8-gingerol and zingerone) and [6]-paradol have been identified as the main active ingredients of ginger and are responsible for the antioxidant activity and its characteristic pungent taste (Gruenwald et al., 2000; Keum et al., 2002). Antioxidative capacity of ginger has been associated with the ability of ginger to inhibit carcinogenesis by reducing oxidative stress and inducing apoptosis (Shukla and Singh, 2007; Katiyar, 1996). Bode and his co-workers, reported that

[6]-paradol exerted its primary inhibitory effect on cell transformation through the induction of apoptosis (Bode et al., 2001). [6]-Paradol and other structurally related derivatives induced apoptosis in oral squamous carcinoma cell line in a dose-dependent manner through a caspase-3-dependent mechanism (Keum et al., 2002). Ginger has been shown to inhibit cell growth in several tumor cells including liver (HepG2) (Yasmin et al., 2008), gastrointestinal (Yoshimi et al., 1992) and breast (Nagasawa et al., 2002). Ginger was also found to possess an anti-tumor promoting potential as determined by inhibition of phorbol ester-induced inhibition of Epstein-Barr virus (EBV) activation in Raji cells (Koshimizu et al., 1988).

The present study provides evidence that ginger may act as a potent growth inhibitory compound in human colon adenocarcinoma cells and supports the possibility of a chemopreventive potential in colon cancer cells. Our study showed that HT 29 cells were more sensitive to ginger treatment with an IC_{50} much lower [455 µg/ml] than HCT 116 cells [496 µg/ml]. The cytotoxic effect could be as a result of the active component, gingerol that has been reported to inhibit the growth of HCT 116 human colon colorectal and liver HepG₂ cancer cells (Bode, 2003; Harliansyah et al., 2007). Azoxymethane-induced intestinal carcinogenesis in rats was significantly suppressed by dietary administration of gingerol (Yoshimi et al., 1992).

We also reported in this study that the inhibition mechanism of colon cancer cells growth by ginger involved interference in both cell cycle progression and apoptosis. Ginger was able to induce cell cycle progression in G₀/G₁ phase and apoptotic death in a dose-dependent manner in both cells. Deregulated cell cycle progression driven by activation of growth-stimulating oncogenes, is one of the




Figure 3b. Flow cytometric analysis of apoptosis induction by ginger in HCT 116 cells. Representative cytograms of apoptosis in HCT 116 cells incubated for 24 hours in the absence (A) or presence of 200 (B), 500 (C) and 800 μ g (D) ginger. Within a cytogram, a quadrants 2 and 4 represent late and early apoptotic cells, respectively; quadrant 1, necrotic cells; quadrant 3, viable cells.

primary characteristics of cancer cells. Cell cycle progression is tightly controlled by the regulation of expression and activity of cyclin/cyclin-dependent kinase (CDK) complexes (Taraphdar et al., 2001). Dysregulation of the cell cycle check points and overexpression of growth-promoting cell cycle factors such as cyclin D1 and cyclin-dependent kinase (CDK) are associated with tumorigenesis (Deihl, 2002). Several dietary agents including curcumin (Mukhopadhyay et al., 2002), resveratrol (Estrov et al., 2003), genistein (Li et al., 2005), dietary isothiocyanates (Jakubikova et al., 2005), apigenin (Takagaki et al., 2005) and silibinin (Tyagi et al., 2002)

have been shown to block the deregulated cell cycle in cancers. In our study, ginger managed to arrest cell cycle at G_0/G_1 -phase for both colon cancer cells at concentration below IC_{50} , whereas at higher concentrations, both cells were arrested by ginger extract at G_2/M phase. The cell cycle arrest observed could be due to inhibitory effect of ginger on cyclin dependent kinases and activation of cell cycle check points. Palozza et al. (2002) showed that arrest of the cell cycle and accumulation of cells in G_2/M phase in colon cancer cells was followed by the decreased expression of cyclin A, a protein known to regulate cdc2 kinase activity in G_2/M phase after




Figure 3c. Flow cytometric analysis of apoptosis induction by ginger in HT 29 cells. Representative cytograms of apoptosis in HT 29 cells incubated for 24 hours in the absence (A) or presence of 200 (B), 500 (C) and 800 $\mu\text{g/ml}$ (D) ginger.

treatment with β -carotene. β -carotene also increased apoptosis through the downregulation of this protein.

In addition to regulation of the cell cycle, apoptosis plays an important role in the maintenance of tissue homeostasis whereby damaged cells are removed since impaired apoptosis contributes to development of cancer (Taraphdar et al., 2001). Compound that can elicit apoptosis is a good chemopreventive agent. In our study, we have shown that ginger at increasing concentration was not only able to inhibit DNA synthesis but also induced apoptosis especially at higher concentration and the percentage of cells that underwent apoptosis, increased dose dependently for both cells. Our

observation of apoptosis in mutant p53- expressing HT 29 is similar to the study by Park et al. (2006) who found that [6]-gingerol induced apoptotic death in pancreatic cells. The percentage of late apoptotic cells was low compared to early apoptotic cells as detected by annexin V staining, suggesting that apoptosis occurred rather slow in colon cancer cells after treatment (Huang and Pardee, 1999). This is consistent with the observation by other researchers whereby colon cancer cells (HCT 116 and HT 29 cells) did not undergo apoptosis rapidly (Goldwasser et al., 1996). Apoptosis that occurred could be due to the down-regulation of NF- κB . Expression of several NF- κB regulated genes including Bcl-2, Bcl-XL,

clAP, surviving, TRAF1 and TRAF2 have been reported to function primarily by blocking the apoptosis pathway (Aggarwal, 2004). Other than that, ginger also could induced apoptosis through mitochondrial pathway involving caspase-8, BID cleavage, cytochrome *c* release and caspase-3 activation like curcumin (Shishodia et al., 2005).

Conclusion

In this from our study, we have demonstrated that ginger extract inhibited the proliferation and induced apoptosis in HCT 116 and HT 29 colon cancer cells. The results suggest that ginger stimulates apoptosis through GoG₁ cell cycle arrest. Future our study will may thus be further investigate the possible use of ginger extract as a new alternative chemotherapeutic agent for human colon cancer.

Competing interests

To our knowledge, previous studies have shown the chemopreventive effect of *Z. officinale* in other cancer cells but less report evidence on colon cancer cells. Thus, our study represent among the first to show the mechanism of anticancer effect of *Z. officinale* in colon cancer cells HCT 116 and HT 29.

ACKNOWLEDGEMENTS

This research was funded by Fundamental Research Grant Science (FRGS), Universiti Kebangsaan Malaysia (UKM), Malaysia through UKM-FF-03-FRGS0007-2007. We also thank Dr. Noor Azian Murad from Center for Lipids Engineering Applied Research (CLEAR), Universiti Teknologi Malaysia, who has kindly provided the ginger extract.

REFERENCES

- Aggarwal BB (2004). Nuclear factor-kappaB: the enemy within. *Cancer Cell. growth factor-induced cell transformation and activator protein* 16(3): 203-208..
- Bode AM, Ma WY, Surh YJ, Dong Z (2001). Inhibition of epidermal activation by [6]-gingerol. *Cancer Res.* 61: 850-853.
- Bode A (2003). Ginger is an effective inhibitor of HCT 116 human colorectal carcinoma *in vivo*. Paper presented at the Frontier in Cancer Prevention Research Conference, Phoenix, AZ, October: pp. 26-30.
- Deihl JA (2002). Cycling to cancer within cyclin D1. *Cancer Biol. Ther.* 1(3): 226-231.
- Estrov Z, Shishodia S, Faderl S, Harris D, Van Q, Kantarjian HM (2003). Resveratrol blocks interleukin-1beta-induced activation of the nuclear transcription factor NF-kappaB, inhibits proliferation, causes S-phase arrest and induces apoptosis of acute myeloid leukemia cells. *Blood* 102(3): 987-995.
- Evan GI, Vousden KH (2001). Proliferation, cell cycle and apoptosis in cancer. *Nature* 411: 342-348.
- Gerard LCC, Halimah Y (2003). Second Report of the National Cancer Incidence in Malaysia National Cancer Registry, Ministry of Health Malaysia.
- Goldwasser F, Bae I, Fornace AJ, Pommier Y (1996). Differential GADD45, p21CIP1/WAF1, MCL-1 and topoisomerase II gene induction and secondary DNA fragmentation after camptothecin-induced DNA damage in two mutant p53 human colon cancer cell lines. *Oncol. Res.* 8: 317-323.
- Gruenewald J, Brendler T, Jaenicke C (2000). PDR for Herbal Medicines. Medical Economics Company, Montvale, New Jersey (USA).
- Hanif R, Qiao L, Shiff SJ (1997). Curcumin, a natural plant phenolic food additive, inhibits cell proliferation and induces cell cycle changes in colon adenocarcinoma cell lines by a prostaglandin-independent pathway. *J. Lab. Clin. Med.* 130: 576-584.
- Harliansyah NAM, Wan ZWN, Yasmin AMY (2007). Antiproliferative, antioxidant and apoptosis effects of *Zingiber officinale* and 6-Gingerol on HepG₂ cells. *Asian J. Biochem.* 2 (6): 421-426.
- Huang L, Pardee AB (1999). β -Lapachone induces cell cycle arrest and apoptosis in human colon cancer cells. *Mole. Med.* 5: 711-720.
- Jakubikova J, Bao Y, Sedlak J (2005). Isothiocyanates induce cell cycle arrest, apoptosis and mitochondrial potential depolarization in HL-60 and multidrug-resistant cell lines. *Anticancer Res.* 25(5): 3375-3386.
- Katiyar SK, Agarwal R, Mukhtar H (1996). Inhibition of tumor promotion in SENCAR mouse skin by ethanol extract of *Zingiber officinale* rhizome. *Cancer Res.* 56: 1023-1030.
- Kazuhiro I, Ando T, Maeda O, Ohmiya N, Niwa Y, Kadomatsu K, Goto H (2007). Ginger ingredients reduce viability of gastric cancer cells via mechanisms. *Biochem. Biophys. Res. Commun.* 362: 218-223.
- Kelloff GJ, Crowell JA, Steele VE, Lubet RA, Malone WA, Boone CW, Kopelovich L, Hawk ET, Lieberman R, Lawrence JA, Ali I, Viner JL, Sigman CC (2000). Progress in cancer chemoprevention: Development of diet-derived chemopreventive agents. *Nutri. Sci.* pp. 467-471.
- Keum YS, Kim J, Lee KH, Park KK, Surh YJ, Lee JM, Lee SS, Yoon JH, Joo SY, Cha IH, Yook JI (2002). Induction of apoptosis and caspase-3 activation by chemopreventive [6]-paradol and structurally related compounds in KB cells. *Cancer Lett.* 177: 41-47.
- Kim EJ, Lee YJ, Shin HK, Park JH (2005). Induction of apoptosis by the aqueous of *Rubus coreanus* in HT-29 human colon cancer cells. *Nutr.* 21: 1141-1148.
- Koshimizu K, Ohigashi H, Tokuda H, Kondo A, Yamaguchi K (1988). Screening of edible plants against possible anti-tumor promoting activity. *Cancer Lett.* 39: 247-257.
- Lee E, Surh YJ (1998). Induction of apoptosis in HL-60 cells by pungent vanilloids, [6]-gingerol and [6]-paradol. *Cancer Lett.* 134: 163-168.
- Lee JH, Yang HS, Park KW, Kim JY, Lee MK, Jeong IY, Shim KH, Kim YS, Yamada K, Seo KI (2009). Mechanisms of thiosulfinates from *Allium tuberosum* L.-induced apoptosis in HT-29 human colon cancer cells. *Toxicology Lett.* 188: 142-147.
- Li M, Zhang Z, Hill DL, Chen X, Wang H, Zhan R (2005). Genistein, a dietary isoflavone, down-regulates the MDM2 oncogene at both transcriptional levels. *Cancer Res.* 65(18): 8200-8208.
- Manju V, Nalini N (2005). Chemopreventive efficacy of ginger, a naturally occurring anticarcinogen during the initiation, post-initiation stages of 1,2 dimethylhydrazine-induced colon cancer. *Clinica Chimica Acta.* 358: 60-67.
- Miyoshi N, Nakamura Y, Ueda Y, Abe M, Ozawa Y, Uchida K, Osawa T (2003). Dietary ginger constituents, galanals A and B, are potent apoptosis inducers in Human T lymphoma Jurkat cells 199: 113-119.
- Mukhopadhyay A, Banarjee S, Stafford LJ, Xia C, Liu M, Aggarwal BB (2002). Curcumin-induced suppression of cell proliferation correlates with down-regulation of cyclin D1 expression and CDK4-mediated retinoblastoma protein phosphorylation. *Oncogene* 21(57): 8852-8861.
- Murillo G, Naithani R, Mehta RG (2008). Efficacy of Herbal Products in Colorectal Cancer Prevention. *Current Colorectal Cancer Reports* 4: 34-42.
- Nagasawa H, Watanabe K, Inatomi H (2002). Effects of bitter melon (*Momordica charantia* L) or ginger rhizome (*Zingiber officinale* rocs) on spontaneous mammary tumorigenesis in SHN mice. *Am. J. Chin. Med.* 30: 195-205.
- Palozza P, Serini S, Maggiano N, Angelini M, Boninsegna A, Fiorella

- DN, Ranelletti FO, Calviello G (2002). Induction of cell cycle arrest and apoptosis in human colon adenocarcinoma cell lines by β -carotene through regulation of cyclin A and Bcl-2 family proteins. *Carcinogenesis* 23: 11-18.
- Park YJ, Wen J, Bang S, Park SW, Song SY (2006). [6]-Gingerol induces cell cycle arrest and cell death of mutant p53-expressing pancreatic cancer cells. *Yonsei Med. J.* 47: 688-697.
- Sakamoto S, Kudo H, Kuwa K, Suzuki S, Kato T, Kawasaki T, Nakayama T, Kasahara N, Okamoto R (1991). Anticancer effects of a Chinese herbal medicine, juzu-taiho-to, in combination with or without 5-fluorouracil derivative on DNA-synthesizing enzymes in 1,2-dimethylhydrazine induced colonic cancer in rats. *Am. J. Chin. Med.* 19(3-4): 233-241.
- Shukla Y, Singh M (2007). Cancer preventive properties of ginger: A brief review. *Food Chem. Toxicol.* 45: 683-690.
- Sporn MB, Dunlop NM, Newton DL, Smith JM (1976). Prevention of chemical carcinogenesis by vitamin A and its synthetic analogs (retinoids). *Federation Proceedings* 35: 1332-1338.
- Surh Y (1999). Molecular mechanisms of chemopreventive effects of selected dietary and medicinal phenolic substances. *Mutation Res.* 428: 305-327.
- Surh YJ (2002). Anti-tumor promoting potential of selected spice ingredients with antioxidative and anti-inflammatory activities: a short review. *Food Chem. Toxicol.* 40: 1091-1097.
- Surh YJ, Lee E, Lee JM (1998). Chemopreventive properties of some pungent ingredients present in red pepper and ginger. *Mutation Res.* 402: 259-267.
- Takagaki N, Sowa Y, Oki T, Nakanishi R, Yogosawa S, Sakai T (2005). 26(1): 185-189.
- Taraphdar AK, Roy M, Bhattacharya RK (2001). Natural products as inducers of apoptosis: Implication for cancer therapy and prevention. *Current Sci.* 80: 10-11.
- Tyagi AK, Singh RP, Agarwal C, Chan DC, Agarwal R (2002). Silibinin strongly synergizes human prostate carcinoma DU145 cells to doxorubicin-induced growth inhibition, G2-M arrest and apoptosis. *Clin. Cancer Res.* 8(11): 3512-3519.
- Virginie DM, Pocard M, Richon S, Weiswald LB, Assayag F, Saulnier P, Judde JG, Janneau JL, Auger N, Validire P, Dutrillaux B, Praz FO, Bellet D, Poupon MF (2007). Establishment of Human Colon Cancer Cell Lines from Fresh Tumors versus Xenografts: Comparison of Success Rate and Cell Line Features. *Cancer Res.* 67(1): 398-407.
- Wang G, Li X, Huang F, Zhao J, Ding H, Cunningham C, Coad JE, Flynn DC, Reed E, Li QQ (2005). Antitumor effect of β -elemene in non-small-cell lung cancer cells is mediated via induction of cell cycle arrest and apoptotic cell death. *Cell. Mole. Life Sci.* 62: 881-893.
- Wei QY, Ma JP, Cai YJ, Yang L, Liu ZL (2005). Cytotoxic and apoptotic activities of diarylheptanoids and gingerol-related compounds from the rhizome of Chinese ginger. *J. Ethnopharm.* 102: 177-184.
- Yasmin AMY, Shahriza ZA, Looi ML, Shafina HMH, Harlianshah H, Noor AAH, Suzana M, Wan ZWN (2008). Ginger extract (*Zingiber officinale* Roscoe) triggers apoptosis in hepatocarcinogenesis induced rats. *Med. Health* 3(2): 263-274.
- Yoshimi N, Wang A, Morishita Y, Tanaka T, Sugie S, Kawai K, Yamahara J, Mori H (1992). Modifying effects of fungal and herb metabolites on azoxymethane-induced intestinal carcinogenesis in rats. *Japanese J. Cancer Res.* 83: 1273-1278.
- Yoshimi N, Wang A, Morishita Y, Tanaka T, Sugie S, Kawai K, Yamahara J, Mori H (1992). Modifying effects of fungal and herb metabolites on azoxymethane-induced intestinal carcinogenesis in rats. *J. Cancer Res.* 83: 1273-1278.