

ASCORBIC ACID AND TESTOSTERONE EFFICACY IN SPERMATOXIC AND HISTOMORPHOLOGICAL CHANGES IN *CORPORA CAVERNOSA* IN PRIAPISM INDUCED SPRAGUE-DAWLEY RATS

Igwe O¹, Aniah J. A², Godam E. T¹, Yama O, E³.

1. Department of Human Anatomy, Faculty of Basic Medical Sciences, Bingham University Karu, Nasarawa State, Nigeria
2. Department of Human Anatomy, Faculty of Medicine University of Abuja, Federal capital territory Abuja, Nigeria.
3. Department of Human Anatomy, Faculty of Basic Medical Sciences, Kaduna State University, Kaduna, Nigeria.

Correspondence: Godam Elvis Tams

E-mail: elvisgodam@gmail.com.

Abstract

Aim: The histomorphological changes in the corpora cavernosa following priapism remains an interesting observation due to its pathological and reproductive implications. We investigated the attenuating effect of testosterone and vitamin C on the corpora cavernosa induced fibrosis in priapism of rats.

Methods: Twenty-five Sprague – Dawley male rats were randomly allocated into 5 groups of rats each Group 1 served as the control. Group 2 had only priapism. Group 3 was induced with priapism and then treated with vitamin C. Group 4 was induced with priapism and then treated with testosterone. Group 5 was induced with priapism but were treated with both vitamin C and testosterone. Priapism was induced for one week followed by 6 hours' post priapism administration of testosterone and vitamin C.

Results: Results showed fibrotic corpora cavernosa in all groups except the control group. Sperm parameters indicated oligospermia and reduced sperm motility especially in group 2 and 3 when compared with the control group.

Conclusion: The results indicated that testosterone and vitamin C ameliorated priapism-induced ischemia – reperfusion injury at different time intervals in rats.

Key words: Priapism, Corpora cavernosa, Infertility, Testosterone, Vitamin C

INTRODUCTION

Priapism is a urological emergency characterised by an unrelenting and painful erection without sexual stimulation usually lasting for longer than 4h to 6h despite orgasm and basically involving only the corpora cavernosa, (Keoghane *et al.*, 2002, Burnett and Bivalacqua, 2007). It is most often drug induced and the antidepressant drug Traxodone has been frequently implicated (Igwe *et al.*, 2012). Other psychoactive substances involved in the aetiology of priapism include marijuana, cocaine, ethanol, the atypical antipsychotics; Risperidone, Olanzapine, Clozapine and Quetiapine. Erection is determined by both neuronal and vascular factors with the latter involving essential

components of arterial dilatation, relaxation of cavernosal smooth muscle and reduced venous outflow (Azadzo *et al.*, 1998). Divergence from the coordination of this mechanism alters the response of the erectile tissue and results in Erectile Dysfunction (Parivar F and Lue T.F, 1997). Ischemia, thrombosis, damaged blood vessels of the penis, impair erectile function or impotence has been associated with priapism. Ischemia-reperfusion injury is a complex phenomenon that causes destruction in both local and remote tissues (Sikka, 2016). Although reperfusion of ischemic tissue is necessary for reparative mechanisms, it has been shown to worsen the injury caused by ischemia via release of reactive oxygen species to the systemic

circulation (Munarriz *et al.*, 2003). Oxidative injury develops when there is excessive production of reactive oxygen species and/or free radicals, which exceeds the natural antioxidant defence mechanisms in the body (Munarriz *et al.*, 2003; Akunna *et al.*, 2013). Oxidative stress produces destructive changes in tissues by causing alterations and irreversible cellular damage, (Kaminski K.A, 2002). Ischemic reperfusion injury post treatment with various antioxidants has been evaluated in many organs, (Uluocak *et al.*, 2010). Concomitant administration of androgens and antioxidants has been a subject of debate as it relates to the management of priapism induced impotence. The controversy over the years has been on the efficacy of testosterone. This study was designed to investigate the role of testosterone and vitamin C administered 6 hours after the induction of priapism on the spermatozoa and histomorphology of the corpora cavernosa.

MATERIALS AND METHODS

ANIMALS

Twenty-five adult male Sprague–Dawley rats weighing between 250 and 300g were housed in solid plastic cages in animal house of Anatomy department of the College of Medicine, University of Lagos and allowed to acclimatize for 2 weeks. The animals were later grouped into five rats per cage and maintained at temperatures between 25 – 28°C. The rats were allowed to eat standard rodent chow and water *ad libitum* throughout the experimental period.

EXPERIMENTAL PROTOCOL

Induction of Priapism

It is a delicate procedure that demands intensive precautionary measures. The penis of rat is a very tender organ and could be easily injured. When traumatized, either by the use of too tight constriction band or rough handling could strangulate the penis and impair micturition process resulting in damming of urine within the urinary bladder, ureter or hydro nephrosis and uraemia. This destabilizes the kidneys and becomes a source of infection and renal failure which could lead to death of the rats, (Sanli *et al.*, 2004). Undoubtedly, the following procedure occurred earlier in this index study and 90% of all the first set of rats died. The subsequent use of 2mm slices of size 16F catheter and minimal trauma to the penis was successful. All operations were performed under sterile conditions. The animals were anesthetized with ketamine injection (50 mg/kg, ip). Priapism was induced with the method described by (Sanli *et al.*, 2004). The tip of a 60-cc syringe was applied

to the base of the flaccid penis, so a vacuum erection device was created. Before the application of vacuum to the penis, a constriction band, which was cut from 16 Fr Foley catheter in 2-mm slices, loaded around the tip of the vacuum erection device. Then the tip of the syringe was placed at the base of the penis and withdrawn gently to induce erection in the rat penis. After induction of erection in sufficient grade, the constriction band was then placed at the base of the penis by slipping off the syringe. Testosterone was administered intramuscularly at 2.5mg/kg body weight three times weekly (Huang H. F. S and Nieschlag E, 1984) and oral Vitamin C at 25mg/kg body weight daily (Emdex, 2006) throughout the one week of priapism induction.

Animal groupings

The animals were divided into five groups containing 5 rats each.

Group 1: Served as the control and was treated with 5ml/kg body weight of normal saline

Group 2: In this group, priapism was induced for one week but no drug was administered

Group 3: Priapism was induced and 6 hours later 25 mg/kg of Vitamin C was injected intraperitoneally.

Group 4: Priapism was induced and 6hrs later 2.5mg/kg body weight of testosterone was injected intramuscularly at three times weekly.

Group 5: Priapism was induced and 2.5mg/kg body weight of testosterone and 25mg/kg body weight of vitamin C injected via same route as above. The animals were sacrificed one week after priapism induction. The penis and the seminiferous tubules were harvested for histology and seminal fluid analysis.

Tissue preparation for histological analysis

All rats were penectomized 1 week after the induction of priapism and the penis were fixed in Bouin's solution for 24 hours and then dehydrated by passing through ascending grades of alcohol (70%, 80%, 90% and absolute alcohol). After dehydration, tissues were cleared in xylene, infiltrated, and then embedded in paraffin wax. Each penis was sectioned along horizontal axis in 5mm thickness. Two sections from each rat were blocked in paraffin. Two sections of each block (total 4 sections for each testis) were stained with Haematoxylin and Eosin (H & E) according to routine light microscopic procedures.

Protocol for Haematoxylin and Eosin stain (Ehrlich's Haematoxylin)

Tissue sections were taken to water and stained with Mayer's Haematoxylin for 10 minutes. The

sections were then washed in water, differentiated in 1% acid alcohol for 1 minute and blued in Scot's tap for 2 minutes. Section was washed in water and stained in aqueous eosin for 20 seconds. Tissues sections were washed in water, dehydrated in 2 changes of alcohol, cleared in 2 changes of xylene and mounted in Diselene Pthalate xylene (DPX) and air dried.

Sperm Count

Following penectomy, the caudal aspect of the epididymis of each rat was excised and dipped into a solution of normal saline to enable the sperm cells to swim out. Haemocytometer chambers (Biocotec 1280 Zhejiang, China mainland) were prepared for counting according to the WHO criteria for semen analysis. Spermatozoa were viewed and counted under a light microscope. The haemocytometer was divided into nine fields. Spermatozoa were counted and recorded for just five random fields and the values recorded in million (10^6) (Tomlinson *et al.*, 2001).

Sperm Motility

Semen was collected from the caudal epididymis and assessed for motility using rapid forward progression, medium forward progression and slow forward progression. Each sample was accessed twice. For consistency, all readings were carried out at 37°C, (WHO, 2000).

Sperm Morphology

After liquefaction and prior to staining, an aliquot of semen was washed with Quinn's Sperm Washing Medium, SAGE (USA) and centrifuged at 3000rpm for 10 minutes. The supernatant was removed and 0.5 mL of Quinn's medium was added to the remaining pellet. 10 μ L of washed semen was then spread onto a glass slide, fixed and air-dried. The smears were washed with distilled water and stained with Spermac stain, Ferti Pro (Beernem, Belgium). After staining, the smears were washed with distilled water. Two

different examiners counted 200 cells *per* smear using bright field illumination at final magnification of 100x and oil immersion. Strict criteria (Menkveld, *et al.*, 1990) were applied for the evaluation, according to which a spermatozoon is normal if it has an oval head, 4.0–5.0 μ m long and 2.5–3.5 μ m wide, measured with an ocular micrometer. The length-to-width ratio should be 1.50–1.75. A normal spermatozoon has a well-defined acrosome that covers 40%–70% of the head. The midpiece is thin, less than 1 μ m wide, about 1.5 times longer than the head. Cytoplasmic droplets, if present, should not be larger than half of the head width. The tail is thin, uniform, uncoiled and about 45 μ m long. According to this classification system, all borderline forms are considered as abnormal.

Statistics

Statistical analysis was done using SPSS statistical software (version 7).

Paired T test was used to compare mean values between the control group and other groups in the study. The comparison was done for sperm count, sperm motility, sperm morphology, liver enzymes and haematological indices.

RESULTS

Histomorphological study

Fig 1 and fig 2 shows result for control group 1 Demonstrating normal tissue and cellular microstructure of the penis. Fig 3 and 4 (Group 2) was the Priapism induced group without treatment. show cellular changes and tissue fragmentation in the penile tissues. Fig 5 and 6 (Group 3), demonstrates Spongiform corpora carvernosa with dilated vascular spaces (fig 5) and fibrotic corpora carvernosa tissues (fig 6). Fig 7 and 8 (Group 4) shows preserved tissue and cellular structures with mild fibrotic tissue while fig 9 and fig 10 (Group 5) also shows similar results.




Fig. 1: Penis in Control Group X100



Fig. 2: Penis in Control Group X400




Fig. 3: Penis in Group 2 (X100)



Figure 4: Penis in Group 2: X400
F: Copora carvernosa fibrosis, **C:** Connective tissue, **S:** Smooth muscle, **P:** Lamina propria, **B:** Dilated blood vessel




Figure 5: Penis in Group 3: X100
Arrows: Spongiform copora carvernosa with dilated vascular spaces




Figure 6: Penis in Group 3: X400
Black arrow: Urethra, **Red arrow:** Fibrotic copora cavernosa



Fig. 7: Penis in group 4 (X100)




Figure 8: Penis in group 4 (X400)
B: Blood vessels, **C:** Connective tissue, **U:** Urethra, **S:** Smooth muscle fibrosis, **V:** Venous spaces




Figure 9: Penis in Group 5 (X400)

Table 1: Sperm morphology (%) and Sperm count ($\times 10^6$ cell/ml) with mean and standard deviation for various groups

Groups	Sperm count ($\times 10^6$ cell/ml)	Normal (%)	Abnormal (%)
Group 1(Control)	154 \pm 2.71	80 \pm 3.23	20 \pm 1.66
Group 2 (P-alone)	1.9 \pm 0.25**	60 \pm 2.34 *	40 \pm 0.98 **
Group 3(P+Vitamin C)	4.5 \pm 0.60**	60 \pm 3.27 *	40 \pm 1.34 *
Group 4 (P+Testosterone)	146 \pm 6.88	70 \pm 2.35	30 \pm 3.44 *
Group 5 5 (P+VitaminC+Testosterone)	111 \pm 11.54*	60 \pm 2.43 *	40 \pm 2.55 *

Data was expressed as mean \pm S.D. Statistics involve the use of t-test *($p < 0.05$) and ** ($p < 0.005$)

Table 2: Sperm motility (%) with mean and standard deviation for various groups

GROUPS	Fully Active (%)	Slightly Active (%)	Dead (%)
Group 1(Control)	80 ± 1.33	10 ± 0.98	10 ± 0.67
Group 2(P-alone)	5 ± 0.56 **	45 ± 2.33	50 ± 1.89 **
Group 3(P+Vitamin C)	20 ± 0.78 *	40 ± 1.23	40 ± 0.88 **
Group 4 (P+Testosterone)	50 ± 1.22	40 ± 2.11 *	10 ± 0.46
Group 5 (P+Vitamin C + Testosterone)	60 ± 2.41	30 ± 0.55	10 ± 0.43

Data is expressed as mean ± S.D. Statistics involve the use of t-test *(p<0.05) and ** (p<0.005)

DISCUSSION

The pathophysiology of priapism can be simplistically viewed as dysfunctional hemodynamic process of the penis, whereby the genital organ excessively endures blood engorgement and the major point of this phenomenon is the anoxia of penile tissue (Burnett, 1995). In anoxic state, smooth muscle of corpus cavernosum has minimal basal tension and has no spontaneous contractile activity. From the therapeutic view, this study also focused on the responsiveness of corpora cavernosa to testosterone and vitamin C in priapism.

Histomorphological profiles

Histological sections of the penis of control group revealed two pale corpora cavernosa occupying most of the central portion surrounded closely by dense connective tissue, the tunica albuginea (Figure 1 and 2). Also noted is the corpus spongiosum situated ventromedially and containing the slit – like penile urethra surrounded by connective tissue trabeculae and muscle strands, as well as blood vessels and nerves. Histological assessment of group 2 rats (Figure 3 and 4) showed distortions evidenced by extensive fibrosis with thickening of elastic and collagen fibres. Some of these fibres were already transforming into fibroblast. However, testosterone and vitamin C administered 6 hours after priapism induction in P + vitamin C group, P + Testosterone group and P + vitamin C + Testosterone group showed similar fibrotic changes though at a very minimal extent (Figure 5-10). This revealed that testosterone and vitamin C has some level of protective function on the corpora cavernosa.

Seminal Fluid Analysis

As shown in table 1 semen fluid analysis revealed that testosterone administration had potential for boosting up sperm count and motility. The group that received testosterone (group 4 and 5) had optimal values of sperm count and motility when compared to the control group. The mean values of sperm count for group 1, 2, 3, 4 and 5 were 154 x 10⁶ cell/ml, 4.9 x 10⁶ cell/ml, 4.5 x 10⁶ cell/ml, 146

x 10⁶ cell/ml and 111 x 10⁶ cell/ml, respectively. The sperm motility also showed increase in the recorded values between group 1, 4 and 5. The mean values of fully active sperm motility for group 1, 2, 3, 4 and 5 were 80%, 5%, 20%, 50% and 60% respectively (Table 2). The mean values of normal and abnormal sperm morphology for group 1, 2, 3, 4, and 5 were 80% and 20%, 60% and 40%, 60% and 40%, 70% and 30%, 60% and 40% respectively (Table 3). This revealed that the Testosterone + priapism group had higher normal sperm morphology when compared with the control group. The above results indicate that testosterone has a positive effect on spermatogenesis hence counter the impotent predisposition of the said groups in priapism.

CONCLUSION

We provided evidences for marked cavernosal fibrosis of the penis, oligospermia and reactional anaemia as a result of experimentally induced priapism. This implies that besides erectile dysfunction resulting from cavernosal fibrosis, patient may suffer low sperm counts (oligospermia) and severe haemolytic anaemia. Concomitant administration of testosterone and vitamin C significantly reversed the fibrotic effect of priapism on corpora cavernosa and the changes in all of these parameters after testosterone and vitamin C administration were in agreement with previous studies (Allamaneni, *et al.*, 2005). These results demonstrated the beneficial effects of testosterone and vitamin C in spermatogenesis and ischaemic reperfusion injury, as evidenced by changes in seminal fluid analysis and biochemical parameters. This preliminary study will enlighten the researchers with erectile dysfunction to construct more comprehensive studies on the effects of fibrosis, ischemia and reperfusion in cavernosal tissue. Thus, the usage of antioxidant agents in such conditions and clinical implications of antioxidant administration can be improved in the future

REFERENCES

- Allamaneni SS, Agarwal A, Nallela KP, Sharma RK, Thomas AJ, Sikka SC (2005). Characterization of oxidative stress status by evaluation of reactive oxygen species levels in whole semen and isolated spermatozoa. *Fertil Steril.*, 83:800–803.
- Akunna GG, Saalu LC, Ogunlade B, Ogunmodede OS, Akingbade AM (2013). Anti-Fertility Role of Allethrin Based-Mosquito Coil on Animal Models. *International Journal of Biology, Pharmacy and Allied Science*, 2(2): 192-207.
- Azadzoï KM, Goldstein I, Siroky MB, Traish AM, Krane RJ, Saenz de Tejada (1998). *J Urol.*, 160(6 Pt 1):2216-22.
- Burnett AL, Bivalacqua T (2007). Priapism: current principles and practice. *Urol Clin North Am* 34:631–642.
- Burnett AL (1995). Role of nitric oxide in the physiology of erection. *Biol Reprod.*; 52: 485-9
- Emdex (2006). *The Complete Drug Formulary for Nigeria's Health Professionals*. Obi CC (ed), Lindoz, Lagos; p. 166.
- Huang HFS, Nieschlag E (1984). *Journal of Reproduction and Fertility* 70, 31 – 38.
- Igné S, Rune AK, Erik J (2012). Division of Psychiatry, Haukeland, Sandviksleitet 1, 5035 Bergen, Norway, *Case Reports in Psychiatry* Volume 2012, Article ID 496364, 4 pages doi:10.1155/2012/496364
- Kaminski KA (2002). Oxidative stress and neutrophil activation—the two keystones of ischemia/reperfusion injury. *Int J Cardiol.*, 86:41.
- Keoghane SR, Sullivan ME, Miller MA (2002). The etiology, pathogenesis and management of priapism. *BJU Int*; 90: 149–54. Burnett A.L. (1995). Role of nitric oxide in the physiology of erection. *Biol Reprod.*; 52: 485-9.
- Menkveld R, Stander FSH, Kotze TJVW, Kruger TF, van Zyl JA (1990). The evaluation of morphological characteristics of human spermatozoa according to stricter criteria. *Hum Reprod.*, 5:586-92.
- Munarriz R, Park K, Huang YH, Saenz de Tejada, Moreland RB, Goldstein I, Traish AM (2003). Reperfusion of ischemic corporal tissue: physiologic and biochemical changes in an animal model of ischemic priapism. *Urology* 62:760–764.
- Parivar F, Lue TF (1997). Priapism. In: Hellstrom WJG (ed.), *Male Infertility and Sexual Dysfunction*. Berlin, Springer-Verlag.; 401-8.
- Sanli O, Armagan A, Kandirali E, Ozerman B, Ahmedov I, Solakoglu S, Nurten A, Tunç M, Uysal V, Kadioglu A (2004). TGF- 1 neutralizing antibodies decrease the fibrotic effects of ischemic priapism. *International Journal of Impotence Research*. 16, 492–497
- Sikka SC (2001). Relative impact of oxidative stress on male reproductive function. *Curr Med Chem.*, 8:851–862.
- Tomlinson MJ, Moffatt O, Manicardi GC, Bizzaro D, Afnan M, Sakkas D (2001). Interrelationships between seminal parameters and sperm nuclear DNA damage before and after density gradient centrifugation: implications for assisted conception. *Hum Reprod.*, 16(10):2160–2165
- Uluocak N, Atilgan D, Erdemir F, Parlaktas BS, Yasar A, Erkorkmaz U, Akbas A. (2010). *Int Urol Nephrol.*, 42(4):889-95.
- WHO (2000). *Manual for the standardized investigation and diagnosis of the infertile couple*. Cambridge University Press.